

A POPULATION OF $z > 2$ FAR-INFRARED *HERSCHEL*-SPIRE SELECTED STARBURSTS

C.M. CASEY¹, S. BERTA², M. BÉTHERMIN^{3,4}, J. BOCK^{5,6}, C. BRIDGE⁵, D. BURGARELLA⁷, E. CHAPIN^{8,9}, S.C. CHAPMAN^{10,11},
D.L. CLEMENTS¹², A. CONLEY¹³, C.J. CONSELICE¹⁴, A. COORAY^{15,5}, D. FARRAH¹⁶, E. HATZIMINAOGLOU¹⁷, R.J. IVISON^{18,19},
E. LE FLOC'H³, D. LUTZ², G. MAGDIS^{3,20}, B. MAGNELLI², S.J. OLIVER²¹, M.J. PAGE²², F. POZZI²³, D. RIGOPOULOU^{20,24},
L. RIGUCCINI^{3,25}, I.G. ROSEBOOM^{21,19}, D.B. SANDERS¹, DOUGLAS SCOTT⁸, N. SEYMOUR^{22,26}, I. VALTCHANOV⁹,
J.D. VIEIRA⁵, M. VIERO⁵, J. WARDLOW¹⁵

Received 2012 July 20; Accepted 2012 October 9.

ABSTRACT

We present spectroscopic observations for a sample of 36 *Herschel*¹ -SPIRE 250–500 μ m selected galaxies (HSGs) at $2 < z < 5$ from the *Herschel* Multi-tiered Extragalactic Survey (HerMES). Redshifts are confirmed as part of a large redshift survey of *Herschel*-SPIRE-selected sources covering ~ 0.93 deg² in six extragalactic legacy fields. Observations were taken with the Keck I Low Resolution Imaging Spectrometer (LRIS) and the Keck II DEep Imaging Multi-Object Spectrograph (DEIMOS). Precise astrometry, needed for spectroscopic follow-up, is determined by identification of counterparts at 24 μ m or 1.4 GHz using a cross-identification likelihood matching method. Individual source luminosities range from $\log(L_{\text{IR}}/L_{\odot})=12.5\text{--}13.6$ (corresponding to star formation rates 500–9000 $M_{\odot} \text{ yr}^{-1}$, assuming a Salpeter IMF), constituting some of the most intrinsically luminous, distant infrared galaxies yet discovered. We present both individual and composite rest-frame ultraviolet spectra and infrared spectral energy distributions (SEDs). The selection of these HSGs is reproducible and well characterized across large areas of sky in contrast to most $z > 2$ HyLIRGs in the literature which are detected serendipitously or via tailored surveys searching only for high- z HyLIRGs; therefore, we can place *lower limits* on the contribution of HSGs to the cosmic star formation rate density at $(7 \pm 2) \times 10^{-3} M_{\odot} \text{ yr}^{-1} \text{ h}^3 \text{ Mpc}^{-3}$ at $z \sim 2.5$, which is $>10\%$ of the estimated total star formation rate density (SFRD) of the Universe from optical surveys. The contribution at $z \sim 4$ has a lower limit of $3 \times 10^{-3} M_{\odot} \text{ yr}^{-1} \text{ h}^3 \text{ Mpc}^{-3}$, $\gtrsim 20\%$ of the estimated total SFRD. This highlights the importance of extremely infrared-luminous galaxies with high star formation rates to the build-up of stellar mass, even at the earliest epochs.

Subject headings: galaxies: evolution – galaxies: high-redshift – galaxies: infrared – galaxies: starbursts – submillimeter: galaxies

¹ Institute for Astronomy, University of Hawaii, 2680 Woodlawn Drive, Honolulu, HI 96822

² Max-Planck-Institut für Extraterrestrische Physik, Giessenbachstrasse, 85748 Garching, Germany

³ Laboratoire AIM-Paris-Saclay, CEA/DSM/Irfu - CNRS - Université Paris Diderot, CE-Saclay, pt courrier 131, F-91191 Gif-sur-Yvette, France

⁴ Institut d'Astrophysique Spatiale (IAS), bâtiment 121, Université Paris-Sud 11 and CNRS (UMR 8617), 91405 Orsay, France

⁵ California Institute of Technology, 1200 E. California Blvd., Pasadena, CA 91125

⁶ Jet Propulsion Laboratory, 4800 Oak Grove Drive, Pasadena, CA 91109

⁷ Laboratoire d'Astrophysique de Marseille - LAM, Université d'Aix-Marseille & CNRS, UMR7326, 38 rue F. Joliot-Curie, 13388 Marseille Cedex 13, France

⁸ Department of Physics & Astronomy, University of British Columbia, 6224 Agricultural Road, Vancouver, BC V6T 1Z1, Canada

⁹ European Space Astronomy Centre, Villanueva de la Cañada, 28691 Madrid, Spain

¹⁰ Institute of Astronomy, University of Cambridge, Madingley Road, Cambridge CB3 0HA, UK

¹¹ Department of Physics and Atmospheric Science, Dalhousie University, 6310 Coburg Rd, Halifax, NS B3H 4R2, Canada

¹² Astrophysics Group, Imperial College London, Blackett Laboratory, Prince Consort Road, London SW7 2AZ, UK

¹³ Center for Astrophysics and Space Astronomy 389-UCB, University of Colorado, Boulder, CO 80309

¹⁴ School of Physics and Astronomy, University of Nottingham, NG7 2RD, UK

¹⁵ Dept. of Physics & Astronomy, University of California, Irvine, CA 92697

¹⁶ Department of Physics, Virginia Tech, Blacksburg, VA 24061

1. INTRODUCTION

Submillimeter galaxies (SMGs, often selected by 850 μ m–1mm flux densities $\gtrsim 2$ mJy; Smail, Ivison & Blain 1997; Hughes *et al.* 1998; Barger *et al.* 1998; Eales *et al.* 1999) are the most intrinsically luminous starburst galaxies that have been identified to date. SMGs are thought to evolve much like local ultraluminous in-

¹⁷ ESO, Karl-Schwarzschild-Str. 2, 85748 Garching bei München, Germany

¹⁸ UK Astronomy Technology Centre, Royal Observatory, Blackford Hill, Edinburgh EH9 3HJ, UK

¹⁹ Institute for Astronomy, University of Edinburgh, Royal Observatory, Blackford Hill, Edinburgh EH9 3HJ, UK

²⁰ Department of Astrophysics, Denys Wilkinson Building, University of Oxford, Keble Road, Oxford OX1 3RH, UK

²¹ Astronomy Centre, Dept. of Physics & Astronomy, University of Sussex, Brighton BN1 9QH, UK

²² Mullard Space Science Laboratory, University College London, Holmbury St. Mary, Dorking, Surrey RH5 6NT, UK

²³ Dipartimento di Fisica e Astronomia, Viale Bertini Pichat, 6/2, 40127 Bologna, Italy

²⁴ RAL Space, Rutherford Appleton Laboratory, Chilton, Didcot, Oxfordshire OX11 0QX, UK

²⁵ NASA Ames, Moffett Field, CA 94035

²⁶ CSIRO Astronomy & Space Science, PO Box 76, Epping, NSW 1710, Australia

¹ *Herschel* is an ESA space observatory with science instruments provided by European-led Principal Investigator consortia and with important participation from NASA.

frared galaxies (ULIRGs; Sanders *et al.* 1988; Sanders & Mirabel 1996) via major mergers. The “merger” evolutionary scenario starts with the collision of gas-rich disk galaxies igniting an intense, short-lived ($\tau \sim 100$ Myr) phase of gas consumption and dust production via a starburst, followed by the formation of a quasar and eventually (1–2 Gyr later) a massive, elliptical galaxy. In contrast to local ULIRGs, SMGs at $z \sim 2$ –3 are much more luminous and more massive (in M_* and M_{H_2}), and sometimes much larger (Chapman *et al.* 2004; Biggs & Ivison 2008), thus they have been dubbed “scaled-up” (Tacconi *et al.* 2008), providing evidence for cosmic downsizing (Cowie *et al.* 1996).

The observation that the most luminous infrared sources are at the highest redshifts (e.g. GN20; Daddi *et al.* 2009) poses a unique problem for galaxy evolution studies. How can these distant ULIRGs be formed so quickly after the Big Bang with such high star formation rates? Their extreme infrared luminosities might stem from different evolutionary histories than the local ULIRG mergers, i.e. secular gas accretion (Dekel *et al.* 2009; Davé *et al.* 2010), but solving the origin of infrared-luminous galaxies requires large, uniformly selected samples of ULIRGs across many epochs.

Unfortunately, most $z > 2$ infrared-luminous galaxy samples number ~ 30 galaxies selected in non-uniform, biased ways. It is well known that the selection of SMGs is severely biased, first against galaxies with warmer dust temperatures (Blain *et al.* 2004; Chapman *et al.* 2004; Casey *et al.* 2009, 2011; Chapman *et al.* 2010; Magdis *et al.* 2010) and second, against galaxies at higher redshifts since they are unlikely to have bright radio counterparts (Chapman *et al.* 2005) or $24\mu\text{m}$ counterparts (Ivison *et al.* 2007; Clements *et al.* 2008) due to the radio K -correction and surface brightness dimming. Third, SMGs at $z > 2$ are often detected in inhomogeneous, serendipitous studies with a range of detection thresholds at different wavelengths in the far-infrared. Fourth, the spectroscopic follow-up and redshift confirmation of these sources is non-uniform; their success rate could relate to their FIR properties, e.g. color or single-band flux density. The lack of SMGs with confirmed redshifts $z \gtrsim 3.5$ has been alleviated in recent years with the discovery of several systems at $4 < z < 5.3$ (Wang *et al.* 2007; Wang, Barger & Cowie 2009; Daddi *et al.* 2009; Coppin *et al.* 2009; Capak *et al.* 2011; Smolčić *et al.* 2011; Walter *et al.* 2012), however this work has been severely limited by the rarity of $z > 3.5$ sources and the small area, non-uniform coverage of existing ground-based submillimeter surveys with SCUBA, MAMBO, LABOCA, and AzTEC.

The *Herschel Space Observatory* (Pilbratt *et al.* 2010) has surveyed $\sim 200\text{ deg}^2$ down to the $\approx 5\text{ mJy}$ confusion limit of SPIRE (Griffin *et al.* 2010; Nguyen *et al.* 2010) as part of the *Herschel* Multi-tiered Extragalactic Survey (HerMES; Oliver *et al.* 2012) at 250, 350, and $500\mu\text{m}$. Although high- z infrared galaxies are spatially rare, SPIRE has mapped much larger sky areas than previous submillimeter surveys and thus can detect a statistically significant population of $z > 2$ starbursts with a well-characterized selection.

This paper presents redshifts and spectra for 36 $2 < z < 5$ *Herschel* SPIRE-selected galaxies identified within a large sample of ≈ 1600 SPIRE-selected galaxies spec-

troscopically surveyed over $\sim 1\text{ deg}^2$. In section 2, we describe the source selection, biases in $24\mu\text{m}$ and radio samples, and spectroscopic observations. In section 3, we present redshift identifications. In section 4, we present our results, from derived luminosities, dust temperatures, the FIR/radio correlation, to composite rest-frame ultraviolet and infrared spectra. In section 5, we discuss the context of our results by calculating the *Herschel* SPIRE-selected galaxy (HSG) contribution to the cosmic star formation rate density (SFRD), and the implications for infrared-luminous galaxy evolution in the early Universe. In section 6, we conclude. Throughout we use a flat ΛCDM cosmology (Hinshaw *et al.* 2009) with $H_0 = 71\text{ km s}^{-1}\text{ Mpc}^{-1}$ and $\Omega_M = 0.27$. When possible, we discuss distance and volume using the general unit, $h^{-1}\text{ Mpc}$.

2. SAMPLE AND OBSERVATIONS

2.1. The *Herschel*-SPIRE selected galaxy (HSG) sample

The sources described in this paper were detected by the SPIRE instrument (Griffin *et al.* 2010) onboard the *Herschel Space Observatory* as part of the *Herschel* Multi-tiered Extragalactic Survey (HerMES; Oliver *et al.* 2012). Sources were spectroscopically observed in a large redshift survey follow-up program described in detail in a parallel paper, Casey *et al.* (2012), hereafter C12. The results of the redshift survey have been split between two papers due to the significant differences in the comprehensive 731 source $z < 2$ sample, identified through [OII], [OIII], $H\beta$, and $H\alpha$ emission, than the 36 galaxies at $z > 2$ identified primarily through rest-frame ultraviolet features. We refer the reader to C12 for a detailed discussion of our source selection and completeness and only briefly summarize those results here.

Due to the large beamsize of SPIRE observations ($18''$, $25''$, and $36''$, respectively, at 250, 350, and $500\mu\text{m}$), counterpart identification and point source photometry is performed by extracting flux from SPIRE maps (Levenson *et al.* 2010) at known positions of *Spitzer*-MIPS $24\mu\text{m}$ and VLA 1.4 GHz sources (see C12 for more details on data). This cross-identification prior source extraction method (called “XID”) is described in detail in Roseboom *et al.* (2010) and Roseboom *et al.* (2012). The disadvantage of the XID method is that it relies on SPIRE-bright sources being detectable at $24\mu\text{m}$ and/or 1.4 GHz, an assumption which is known to sometimes fail at $z \gtrsim 3$, depending on the depth of $24\mu\text{m}$ or 1.4 GHz coverage. For this reason, our spectroscopic survey was conducted in the HerMES coverage areas with the deepest available ancillary data in six different legacy fields: Lockman Hole North (LHN; $\alpha \sim 10\text{h } 46'$, $\delta \sim 59^\circ$), Cosmic Evolution Survey field (COSMOS; $\delta \sim 10\text{h } 0'$, $\delta \sim 2^\circ$), Great Observatories Origins Deep Survey North field (GOODS-N; $\alpha \sim 12\text{h } 36'$, $\delta \sim 62^\circ$), Elais-N1 (EN1; $\alpha \sim 16\text{h } 0'$, $\delta \sim 54^\circ$), the UKIDSS Ultra-deep field (UDS; $\alpha \sim 2\text{h } 19'$, $\delta \sim -5^\circ$), and the Extended Chandra Deep Field South (CDFs; $\alpha \sim 3\text{h } 30'$, $\delta \sim -28^\circ$).

Sources were selected for spectroscopic follow-up by detection at $> 3\sigma$ significance in at least one of the three SPIRE bands. The absolute flux limit changes field to field depending on XID prior source density, but averages ~ 10 – 12 mJy across the three bands. Throughout the rest of this paper we refer to this population as *Herschel*-

SPIRE selected galaxies (HSGs) for convenience. Higher priority follow-up is given to sources detected in all three bands, however the source density of all SPIRE-detected sources is low enough such that >98% of all HSGs can be surveyed within one spectroscopic mask area, whether it be with the Keck I Low Resolution Imaging Spectrometer (LRIS; covering $5.5' \times 7.8'$) or with the Keck II DEep Imaging Multi-Object Spectrograph (DEIMOS; covering $5' \times 16.7'$).

2.2. Biases in the HSG Sample

Although the completeness of the XID source identification technique is >95% for sources at $z < 2$ (Magdis *et al.* 2010; Roseboom *et al.* 2010, 2012; Béthermin *et al.* 2011), the completeness at higher redshifts is difficult to estimate since an increasing, non-negligible fraction of SPIRE-bright sources drop out at $24\ \mu\text{m}$ and/or 1.4 GHz with increasing redshift. This is also a function of the $24\ \mu\text{m}$ and 1.4 GHz depth, which is different field to field. Since this paper focuses exclusively on the $z > 2$ SPIRE-bright population, it is important to emphasize that the sample here is incomplete, biased and is not representative of all SPIRE-bright galaxies at $z > 2$. Constraining the whole SPIRE population at $z > 2$ will require detailed high-resolution submillimeter follow-up, e.g. from ALMA, of a large population of SPIRE-bright systems, particularly those that are radio and $24\ \mu\text{m}$ -faint and for sources which fail to yield optical spectroscopic identifications.

Note that the purity of the XID technique in counterpart identification is not guaranteed. In other words, XID might be incorrect in its identification of the multiwavelength counterpart for a *Herschel* source (as mentioned in Roseboom *et al.* 2010). The purity of this sample is impossible to gauge without direct far-infrared interferometric observations (this does away with confusion noise, as in Wang *et al.* 2011). However, we do note that radio counterpart identifications are more robust than $24\ \mu\text{m}$ counterparts due to their source rarity and radio's direct scaling with FIR luminosity (Chapman *et al.* 2003, 2005). Half of our sample is radio identified, with the other half showing no overall bias or trends which would skew our results.

Another possible bias of the HSG sample is the method of spectroscopic targeting. Our Keck observations were centered around high-priority 'red' SPIRE sources (e.g. $S_{250} < S_{350} < S_{500}$) which are thought to be the highest-redshift SPIRE sources (e.g. Cox *et al.* 2011). While only 1-2 'red' targets were chosen per mask, this could skew the total redshift distribution higher than if masks were laid down arbitrarily on the sky. As C12 describes, and as we discuss later in section 5, we measure the impact of 'red' sources on the overall redshift distribution to be negligible since many of the 'red' targets failed to yield spectroscopic identifications.

2.3. Spectroscopic Observations

Spectroscopic observations were carried out at the W.M. Keck Observatory using the Low Resolution Imaging Spectrometer (LRIS) on Keck I and the DEep Imaging Multi-Object Spectrograph (DEIMOS) on Keck II in 2011 and 2012. LRIS observations were carried out on 2011-Feb-06, 2012-Jan-26, 2012-Jan-27, and 2012-Feb-27 with the 400/3400 grism, 560nm dichroic, and primarily

the 400/8500 grating in the red with central multi-slit wavelength 8000Å for the $5.5' \times 7.8'$ mask. This setup yields a 1.09Å dispersion in the blue ($R \sim 4000$) and 0.80Å dispersion in the red ($R \sim 9000$). Integration times varied from ~ 2700 to 5600s per mask depending on air-mass and weather. DEIMOS observations were carried out on 2011-May-28, 2011-May-29, 2011-Nov-28, 2012-Feb-16, and 2012-Feb-17. The 600 lines/mm grating and 7200Å blaze angle was used, resulting in a 0.65Å dispersion ($R \sim 11000$). The GG455 filter was used to block higher-order light, and typical integration times per mask were ~ 2700 to 4800s. The resolution in LRIS red and DEIMOS is sufficient to distinguish between a single emission line (e.g. Ly α) and the [OII] doublet (rest-frame separation of 3Å). Data reduction for LRIS was done using our own custom-built IDL routines, while we used the DEEP2 DEIMOS data reduction pipeline for DEIMOS data²⁷.

Twenty-five LRIS masks were observed (thirteen under photometric conditions), and twenty-nine DEIMOS masks were observed (sixteen under photometric conditions), surveying a total of 1594 SPIRE-selected sources in $0.93\ \text{deg}^2$ ($0.43\ \text{deg}^2$ observed in photometric conditions). Of 1594 sources surveyed, 767 have confirmed spectroscopic redshifts identified primarily by the [OII] doublet, [OIII], H β , H α , [NII], Ca H and Ca K absorption, the Balmer break, H γ , Ly α , CIV, CIII], HeII, and the Lyman break. The lower redshift sources identified through rest-frame optical signatures are discussed in C12. Of the 767 confirmed redshifts, 36 are above $z = 2$ and thus comprise the HSG sample discussed in this paper. Out of the 36, we categorize 22 as secure based on the signal-to-noise of Ly α and/or detection of multiple spectral features, discussed in the next section.

Note that the 36 source sample discussed in this paper was not selected in any special way in SPIRE color, photometric redshift, or optical characteristics. There was no special selection imposed which would yield more high-redshift identifications, and the selection differs in no way from the lower redshift confirmed sources. This paper is simply a description of all $z > 2$ sources confirmed in our large spectroscopic survey.

The DEIMOS wavelength coverage roughly spans 4500–9500Å, whereas LRIS coverage spans 3000–10000Å. The limited wavelength coverage of DEIMOS results in a gap in our redshift coverage between $1.6 < z < 3.2$ which does not occur for LRIS observations. Indeed, very few sources observed by DEIMOS are identified in that redshift range, with the exception of 1HERMES X1.4 J104642.89+585650.0 at $z = 2.841$ and 1HERMES X24 J160545.99+534544.4 at $z = 2.555$. These are both quasars with strong, broad CIV emission. We take the DEIMOS redshift desert into account when calculating the luminosity function and contribution to the cosmic star formation rate density later in Section 5. The redshift distribution for all *Herschel*-SPIRE galaxies is shown in Figure 1. The photometric redshift distribution shown comes from all HSGs in the COSMOS field, not limited to those targeted for spectroscopic observations.

²⁷ The analysis pipeline used to reduce the DEIMOS data was developed at UC Berkeley by Michael Cooper with support from NSF grant AST-0071048.




FIG. 1.— Redshift distribution of *Herschel*-SPIRE selected galaxies from our survey. This paper analyzes the $z > 2$ sample, which consists of 36 sources, while the $z < 2$ sample, consisting of 731 sources, is analyzed in a parallel paper (Casey et al. 2012; C12). The distribution of photometric redshifts comes from sources in the COSMOS field, where the quality of the photometric redshifts is high over a large area; this distribution includes sources not necessarily in the spectroscopic survey, but follow the same selection as spectroscopic targets. The hashed area from $1.6 < z < 3.2$ highlights the DEIMOS redshift range where there is a deficit of sources due to spectroscopic incompleteness.

3. REDSHIFT IDENTIFICATION AND SPECTRA

Table 1 summarizes the spectroscopic identifications and multi-wavelength properties of the 36 SPIRE-selected galaxies identified at $2 < z < 5$. In each case, the signal-to-noise of the identifying feature ($\text{Ly}\alpha$ emission in the majority of cases) is required to be >5 in the two-dimensional spectrum. The two-dimensional spectra for $\text{Ly}\alpha$ identified sources is shown in Figure 2. Note that the S/N of $\text{Ly}\alpha$ changes from the two-dimensional spectra to one-dimensional extractions of the spectra depending on the sources’ compactness, the compactness of continuum relative to $\text{Ly}\alpha$ emission, and the observed wavelength of the line. For example, one-dimensional extractions with $\text{Ly}\alpha$ longward of $\sim 6000\text{\AA}$ are prone to contamination by the OH forest, thus will be of significantly poorer quality in 1D than in 2D where OH features are more easily distinguished from real lines.

Initially, we split sources into two categories: sources with multiple spectroscopic features (e.g. $\text{Ly}\alpha$ emission, SiIV absorption, CIV emission) and single-line identifications (where $\text{Ly}\alpha$ is identified as the only emission line). The former group of identifications is naturally more secure than the latter. The latter source list is dominated by sources at the high redshift end, $3 \lesssim z \lesssim 5$, since multiple features are naturally more difficult to identify in higher redshift sources. However, there are clear cases where the identification of $\text{Ly}\alpha$ at $>5\sigma$ is more secure than the identification of multiple other features at low signal-to-noise ($S/N \lesssim 5\sigma$). An example of a secure single-line source is 1 HERMES X24 J160802.63+542638.1 at $z = 3.415$ where $\text{Ly}\alpha$ is detected at $>5\sigma$. An example of a less secure multi-feature source is 1 HERMES X1.4 J123622.58+620340.3 at $z = 3.579$

where both $\text{Ly}\alpha$ and CIV are detected at $S/N < 5\sigma$. In this paper, we choose to segregate sources with secure spectroscopic identifications from those with tentative identifications rather than sources with single- or multi-line identifications.

Sources are categorized as ‘tentative’ rather than ‘secure’ if (a) the signal-to-noise of the $\text{Ly}\alpha$ feature is $5 < S/N < 7$ as measured in the two-dimensional LRIS/DEIMOS spectrum using an appropriately sized, adjustable aperture, and if (b) out of five co-authors who did a thorough quality ranking of each source’s spectrum, at least three ranked the source as ‘tentative’ rather than ‘secure.’ Figure 3 show all rest-frame ultraviolet spectra for all 36 sources in the sample.

Tentative sources are marked with a dagger (†) in Table 1 and denoted as ‘tentative’ in Figures 2 and 3. Out of 36 sources, we classify 22 as secure and 14 as tentative. Note that one source, 1HERMES X1.4 J100024.00+021210.9 at $z = 3.553$ is classified as tentative *not* because the $\text{Ly}\alpha$ emission is of low- S/N , but because there is peculiarly strong emission from SiIV relative to $\text{Ly}\alpha$ and CIV; this could be an artifact of the noise, but since it is particularly unusual, we have categorized this source as tentative.

Photometric redshifts are used, when available, to verify spectroscopic redshifts to within $\Delta z/(1+z) \sim 1/2$ (Ilbert et al. 2010; Rowan-Robinson et al. 2008; Strazzullo et al. 2010; Cardamone et al. 2010). This threshold is not strict, but rather used to spot egregious disagreements between spectroscopy and photometry. These photometric redshift catalogs are of varying quality depending on the depth and number of optical and near-infrared bands available, from the limited and shallow coverage in EN1 to the extensive 30-band coverage in COSMOS. The loose constraint of agreement between photometric and spectroscopic redshift is based on the fact that photometric redshifts are notoriously unreliable for dusty starbursts. Even in the deepest multi-band fields like COSMOS, $\sim 30\%$ of SPIRE sources do not have photometric redshifts, (e.g. Mobasher et al. 2007; Ilbert et al. 2010).

Viewing geometry can have a very strong impact on the relative fractions and wavelengths of escaped ultraviolet/optical light (Siana et al. 2007, 2010; Scarlata et al. 2009). Heavy obscuration can have dramatic impact on photometric redshift reliability in two ways. First, by obscuring rest-frame ultraviolet and optical light so much that the source has an artificially high photometric redshift (i.e. the source is thought to drop out due to redshifting rather than reddening). Second, dusty sources often have differential absorption of $\text{Ly}\alpha$ photons and non-resonantly scattered continuum photons (e.g. the ‘UV chimney’ argument, see Neufeld 1991), resulting in artificially low photometric redshifts.

Even in the lower redshift samples (C12), sources with very confident photometric redshifts from the multi-band data of COSMOS and CDFS have photometric redshift accuracy of $\Delta z/(1+z) = 0.29$. This highlights the lack of reliability in photometric redshifts of dusty galaxies. Therefore, the primary purpose of the photometric redshift restriction in this paper is to verify that the detected emission line is in fact $\text{Ly}\alpha$ and not [OII] (which would correspond to redshifts between 0 and 1).




FIG. 2.— Two-dimensional spectra around the identified $\text{Ly}\alpha$ features for the sources only identified through a single emission line. This emission line is not thought to be $[\text{OII}]$ or MgII in these cases since there is no detection of commonly bright, accompanying emission lines, e.g. $[\text{OII}]$, $[\text{OIII}]$ and $\text{H}\beta$, within the wavelength coverage of LRIS/DEIMOS observations. Note the wide variety of $\text{Ly}\alpha$ morphologies, from very diffuse to very compact, with and without redward continuum. The sources at higher redshifts are observed at higher wavelengths, thus likely have one-dimensional extractions contaminated by the OH forest.




Figure 2 — continued.




FIG. 3.— One dimensional spectra of $z > 2$ SPIRE sources in observed wavelength shown in the left panels. The flux scaling is arbitrary. The wavelengths of rest-frame ultraviolet features are marked by vertical dotted gray lines with noted names. All lines are shown irrespective of whether or not they are seen in emission, absorption, and whether or not they are detected. Redshifts are measured off of the Ly α redshift in all cases where Ly α is detected (a minority of sources is identified by CIV emission). The Ly α redshifts are typically redshifted with respect to other spectral features, evidence of stellar winds (to be discussed at more length in a future paper). Right panels show best-fit far-infrared spectral energy distributions to infrared photometry. The total SED is a solid line while the underlying cold-dust modified blackbody is dashed. The derived IR luminosities and dust temperatures are shown.




Figure 3 — continued.




Figure 3 — continued.




Figure 3 — continued.




Figure 3 — continued.




Figure 3 — continued.




Figure 3 — continued.




Figure 3 — continued.




Figure 3 — continued.




FIG. 4.— Spectroscopic redshift against photometric redshift for our SPIRE-selected galaxy sample. Sources with identifications based on multiple spectroscopic features (emission lines, absorption lines, breaks) are shown as black circles, while identifications made from a single-feature (i.e. $\text{Ly}\alpha$ emission) are open circles. The photometric and spectroscopic redshifts for the full sample of 767 HSGs (C12) is shown as small gray points. Since the single-line source emission lines could have been mis-identified, we also mark the corresponding spectroscopic redshifts if [OII] is assumed and find generally poor agreement with the photometric redshift (black crosses). The distribution in spectroscopic redshifts for sources with *no* photometric redshifts is shown in gray; there is no obvious trend with redshift.

Fifteen of the 36 sources do not have photometric redshifts—even though some of them lie in areas of very deep ancillary data—a testament to their dusty and optically obscured nature. The sources with fewer identifiable spectral signatures in our spectra are less optically luminous than those with multiple features. Those which are optically fainter will have less reliable photometric redshifts and are more likely to be categorized here as tentative.

3.1. Alternate Redshift Identifications

Figure 4 plots photometric redshift against spectroscopic redshift where photometric redshifts are available. The distribution in spectroscopic redshift of sources without photometric redshift is the gray histogram. Alternate redshifts, assuming the emission lines are actually [OII] instead of $\text{Ly}\alpha$, are shown as crosses on Figure 4. The [OII]-implied spectroscopic redshifts are less consistent with photometric redshifts than the $\text{Ly}\alpha$ -implied spectroscopic redshifts. This makes sense in the context of the probable enhanced $\text{Ly}\alpha$ -to-continuum ratios in these galaxies.

While our identified $\text{Ly}\alpha$ emission lines could also be MgII (2798Å), MgII would be suggestive of a strong AGN. With AGN, other features (e.g. CIII] 1909Å and [OII]) would also be detectable. Given the expected relative strengths of MgII relative to [OII], CIII], CIV, and $\text{Ly}\alpha$, MgII is unlikely to be identified as the only discernible feature. In the case of [OII], both [OIII] and $\text{H}\beta$ emission lines and/or CaH & K absorption, or the 4000Å break features are expected. Also at wavelengths $>6000\text{Å}$, the [OII] doublet is resolvable with LRIS and DEIMOS. In comparison to our lower redshift sample

(C12), for which there are ~ 500 [OII] identified lines at the same wavelengths, the sources in this paper have none of the above signatures which would point to a mis-identification as [OII].

Also note that if the $\text{Ly}\alpha$ identified lines were actually [OIII] or $\text{H}\beta$ they would sit at even lower redshifts than if [OII]. Because there are no sufficiently bright emission lines between $\text{Ly}\alpha$ (1216Å) and [OII] (3727Å) for starbursts, it is straightforward to segregate between $z > 2$ and $0 < z < 1$ sources using photometric redshift.

We inspect the optical images of each source which does not have a photometric redshift to judge the plausibility of our identifications. All three GOODS-N sources are *u*-band drop-outs. Both 1HERMES X1.4 J123622.58 and 1HERMES X1.4 J123732.66 are also *b*-band drop-outs (the two higher redshift sources), and all three have optical photometry which is consistent with their spectroscopic identifications. Of the eight LHN sources without photometric redshifts, 1HERMES X1.4 J104557.12 is compact and has $i \approx 24.5$, 1HERMES X1.4 J104701.68 and 1HERMES X1.4 J104709.6 have $i \approx 24.3$, 1HERMES X1.4 J104722.6 and 1HERMES X1.4 J104707.7 have $i \approx 25$, and 1HERMES X1.4 J104620.4, 1HERMES X24 J104642.9, and 1HERMES X1.4 J104649.9 are *i*-band drop outs. These magnitudes do not rule out the possibility that these are low-redshift sources, but indicates consistency between our high- z spectroscopic identifications and photometry. In COSMOS, only one source, 1HERMES X24 J100133.36+023726.9, has no photometric redshift since it drops out in all images. Elais-N1 sources have much shallower photometric coverage than the other fields, hence more sources without photometric redshifts. While all sources are detectable in wide *i*-band imaging, multi-band imaging is not available across the whole field. None of the sources are sufficiently bright or extended at *i*-band to be a convincing $z < 1$ identification.

Note that 1HERMES X24 J161506.65+543846.9, the highest redshift source in our sample at $z = 4.95$ has an odd assortment of photometric measurements, dropping out in all wavebands (including *z*-band) except *i*-band, where it has a magnitude of 22.8 (AB). The low photometric redshift ($z_p = 1.94$) is likely caused by this peculiar optical SED, but is also perfectly consistent with $\text{Ly}\alpha$ in *i*-band at $z \sim 5$, and enhanced $\text{Ly}\alpha$ -to-continuum ratios (Neufeld 1991). We also note that this source is classified as tentative.

4. SAMPLE CHARACTERISTICS

The importance and context of the $2 < z < 5$ HSG population can only be judged with a basic understanding on the physical characteristics of the sample. Here we measure those physical characteristics, compare them to the properties of other galaxy populations, and assess the impact of infrared selection biases on our interpretation.

4.1. SED fits, luminosities and dust temperatures

The infrared photometry summarized in Table 1 is fit with a FIR spectral energy distribution (SED) consisting of a coupled single dust temperature modified blackbody and mid-infrared power law such that

$$S(\lambda) = N_{\text{bb}} \frac{(1 - e^{-(\frac{\lambda_0}{\lambda})^\beta})(\frac{c}{\lambda})^3}{e^{hc/\lambda kT} - 1} + N_{\text{pl}} \lambda^\alpha e^{-(\frac{\lambda}{\lambda_c})^2} \quad (1)$$

where $S(\lambda)$ is in units of Jy and T is the galaxy’s characteristic “cold” dust temperature (the dust temperature dominating most of the infrared luminosity and dust mass). The emissivity index is represented by β , and λ_0 is the wavelength at which optical depth is unity (here fixed at $\lambda_0 = 200\mu\text{m}$, as described in Conley *et al.* 2011). The slope of the mid-infrared powerlaw component is given by α , and λ_c is the wavelength where the gradient of the modified blackbody is equal to α . N_{bb} and N_{pl} are the coefficients of the modified blackbody and power laws respectively. N_{pl} is a fixed function of N_{bb} , α , and T such that the powerlaw and modified blackbody are continuous at λ_c . This SED fitting method is described fully in Casey (2012), and is also discussed in C12 as applied to the low-redshift population. It is given in related forms in Blain *et al.* (2002), Blain, Barnard & Chapman (2003), Younger *et al.* (2009), and Conley *et al.* (2011).

To reduce the number of free parameters, we fix $\alpha = 2.0$ for sources without PACS photometry and $\beta = 1.5$ for all (the measured values found for local IRAS and some distant ULIRGs Casey 2012). These leaves two free parameters: N_{bb} , which effectively scales with L_{IR} , and T , the temperature of the modified blackbody. We remeasure dust temperatures for each galaxy by determining the wavelength where the flux density peaks and convert that to a dust temperature via Wien’s Law. This provides a more consistent measure of dust temperature which can be used in comparisons between SEDs fit with alternate techniques, using model templates or direct fits (Chary & Elbaz 2001; Dale & Helou 2002; Blain, Barnard & Chapman 2003; Siebenmorgen & Krügel 2007; Draine & Li 2007; Kovács *et al.* 2010). We compute infrared luminosities by integrating the above best-fit spectral energy distribution between 8 and $1000\mu\text{m}$. The SED fits are shown alongside the sources’ optical spectra in Figure 3, and their infrared luminosities and dust temperatures are given in Table 1.

The luminosities of this sample range from 3.2×10^{12} – $6.3 \times 10^{13} L_{\odot}$, implying infrared star formation rates of 500 – $9000 M_{\odot} \text{yr}^{-1}$ (with one outlier at $1.6 \times 10^{14} L_{\odot}$, $26000 M_{\odot} \text{yr}^{-1}$, whose redshift is tentative). Most galaxies in the sample have SFRs an order of magnitude beyond the extreme activity seen in ULIRGs (which have $SFR \approx 200$ – $1000 M_{\odot} \text{yr}^{-1}$ by the scaling given in Kennicutt 1998). These starbursts are amongst the most extreme star forming galaxies seen in the Universe (amongst other HyLIRG populations, Rowan-Robinson 2000; Bridge *et al.* 2012).

The conspicuously high star formation rates (e.g. above $\sim 1000 M_{\odot} \text{yr}^{-1}$) might lead us to believe that AGN contaminate the FIR luminosity or rather, that there is potential variation in star formation laws at high redshift. There has been some recent discussion of whether or not the Kennicutt (1998) scaling between IR luminosity and SFR holds under ‘extreme’ conditions or at high redshifts (Swinbank *et al.* 2008). Assuming a modified IMF would produce more modest SFRs than the default Salpeter IMF. While this might change our interpretation and change the star formation rates we measure here, we use the Kennicutt scaling for SFRs in this paper to be consistent with literature work.

Figure 5 shows the infrared luminosity against dust temperature for the HSG sample. There is a noticeable




FIG. 5.— Infrared luminosity against dust temperature for the sample, color coded by redshift interval: $2 < z < 3$ (blue), $3 < z < 4$ (green), and $4 < z < 5$ (red). Overplotted are typical lower luminosity limit boundaries—a function of dust temperature—at a given wavelength, redshift, and flux density limit. A 12 mJy flux density limit is assumed for the SPIRE bands (dashed lines) and a 5 mJy limit is assumed at $850\mu\text{m}$ (solid lines). This illustrates how half of the SPIRE-selected sample have dust temperatures too warm to be $850\mu\text{m}$ -detectable. Sources with AGN optical signatures are marked with small white dots at their centers.

absence of very warm sources at lower luminosities. Similarly, there are very few cold sources at high luminosities. This is primarily a consequence of selection effects in the SPIRE bands. Warm-dust galaxies are selected against in the SPIRE bands, even at these high-redshifts, due to the sensitive variation of infrared flux density measurements with dust temperature; the dashed lines illustrate the lower luminosity detection limits as a function of dust temperature, for a galaxy at $z = 2$, $z = 3$, or $z = 4$. This dust-temperature selection bias is even more exaggerated at $850\mu\text{m}$. The luminosity detection limits for $850\mu\text{m}$ selection are shown as solid lines on Figure 5, nearly bisecting the SPIRE population so that about half would be $850\mu\text{m}$ -undetected). Before *Herschel*, the cold-dust temperature bias of submillimeter observations was the focus of many studies looking for the elusive “warm-dust” SMGs (Blain *et al.* 2004; Chapman *et al.* 2004; Casey *et al.* 2009, 2011; Chapman *et al.* 2010; Magdis *et al.* 2010; Chapin *et al.* 2011).

Figure 6 plots HSGs’ infrared luminosities against redshift relative to the detection limits of the selection wavelengths: the three 250, 350, and $500\mu\text{m}$ SPIRE bands, $24\mu\text{m}$ and 1.4 GHz. The detection limits in the SPIRE bands depend on the assumed SED shape and dust temperature which is not uniform for all SPIRE-selected galaxies. We use the distribution of measured dust temperatures in Figure 5 to set limits on the detection boundaries in $L_{\text{IR}}-z$ on Figure 6. In other words, we measure that the mean dust temperature for $250\mu\text{m}$ -selected galaxies to be 40 K, for $350\mu\text{m}$ -selected galaxies as 39 K, and for $500\mu\text{m}$ -selected galaxies as 37 K. Differences on the scale of a few degrees are negligible for these illustrative boundary lines in Figure 6, so we adopt a 40 K SED for all three selection boundaries. Most of the fields which have radio data have a detection limit $\gtrsim 40\mu\text{Jy}$, so




FIG. 6.— Integrated infrared luminosity (8–1000 μm) against spectroscopic redshift for our sample (black sources are secure and gray sources are tentative, as in Table 1). Luminosities span 2.4×10^{12} – $4.0 \times 10^{13} L_{\odot}$, with one outlier at $1.4 \times 10^{14} L_{\odot}$. The detection boundaries of each selection wavelength are also illustrated as a function of redshift. The SPIRE detection boundaries assume flux densities $> 12 \text{ mJy}$ and an SED dust temperature of 40 K. The radio boundary assumes that the FIR/radio correlation (Helou, Soifer & Rowan-Robinson 1985) holds and a $> 40 \mu\text{Jy}$ radio detection limit. The $24 \mu\text{m}$ limit assumes a $150 \mu\text{Jy}$ detection limit along with a typical $24 \mu\text{m}$ – L_{IR} scaling (e.g. Le Floch *et al.* 2005). Radio sources are marked by a magenta asterisk, while $24 \mu\text{m}$ sources are black.

we construct the radio detection boundary based on the FIR/radio correlation for starbursting galaxies (Helou, Soifer & Rowan-Robinson 1985; Condon 1992) with q_{IR} evolving as in Ivison *et al.* (2010). The $24 \mu\text{m}$ detection boundary is the least certain as it scales with L_{IR} ; many recent works note up to ~ 1 dex disagreement between extrapolated $24 \mu\text{m}$ infrared luminosities and direct measurements (e.g. Le Floch *et al.* 2005, 2009; Elbaz *et al.* 2011). This tells us that, although we can use the $24 \mu\text{m}$ detection boundary shown in Figure 6 as a rough guide, it should not be concerning that $24 \mu\text{m}$ -selected sources fall below the line by $\lesssim 0.3$ dex.

Although very few of these sources have existing $850 \mu\text{m}$ data, this work (especially Figure 5) suggests that half of the SPIRE-selected population would be undetectable in the original SCUBA $850 \mu\text{m}$ surveys *even* at these redshifts. Specifically, we estimate that only 31/36 sources would have $S_{850} > 2 \text{ mJy}$ (86%), while only 21/36 sources have $S_{850} > 5 \text{ mJy}$ (58%). The statistics of the temperature-bias selection effect are discussed in more detail in C12, as they relate to the lower redshift population where the statistics are more robust.

4.2. FIR/radio correlation

For the 23 radio-detected sources, we investigate the FIR/radio correlation for starburst galaxies (Helou, Soifer & Rowan-Robinson 1985; Condon 1992). Measuring the FIR/radio correlation in this sample is useful for checking that our sample is roughly consistent in FIR/radio given expectation from our measured redshifts (i.e. it is another reassurance on tentative identifications in particular). This correlation is measured via the ratio

of FIR luminosity to radio luminosity such that

$$q_{\text{IR}} = \log \left(\frac{1.02 \times 10^{18} L_{\text{FIR}}}{4\pi D_L^2} \left[\frac{\text{cm}^2}{L_{\odot}} \right] \right) - \log (1 \times 10^{-32} S_{1.4} (1+z)^{\alpha-1} [\mu\text{Jy}^{-1}]) \quad (2)$$

where L_{FIR} is the far-infrared luminosity measured in the range 40–120 μm given in L_{\odot} , D_L is the luminosity distance in cm, $S_{1.4}$ is the 1.4 GHz flux density in μJy , and α is the synchrotron slope, here set to 0.75 (Ibar *et al.* 2010; Ivison *et al.* 2010) and defined such that $S_{\nu} \propto \nu^{-\alpha}$.

Twenty-three of the 36 galaxies in our sample are radio detected (64%), and their measured q_{IR} ranges from 0.7 to 2.1 with mean value $\langle q_{\text{IR}} \rangle = 1.58 \pm 0.35$. Note that the infrared luminosity component in q_{IR} is $L_{\text{FIR}}(40 - 120)$, not $L_{\text{IR}}(8 - 1000)$. If $L_{\text{IR}}(8 - 1000)$ is used instead, luminosities and q increase by 0.40 ± 0.15 dex and the scatter in q grows; the increased scatter is caused by the contribution from the mid-infrared flux to L_{IR} . Only one galaxy in our sample is ‘radio-loud’ and indicative of AGN (the source at $z = 3.579$); this is consistent with the observed CIV emission in its rest-UV spectrum. Since we observe the rest of the radio-detected sample to agree within uncertainties with previous measures of $q_{\text{IR}}(z)$ in previous samples (Ivison *et al.* 2010; Kovács *et al.* 2010; Magnelli *et al.* 2012) lends additional credence to our redshift identifications. While each of the literature samples have different measures for q (ranging 1.3–2.2), all with uncertainties on the order of ~ 0.15 – 0.25 , the overall trend of an evolving q_{IR} is consistent between samples.

4.3. Composite Ultraviolet Spectra

Since the signal-to-noise on individual galaxy spectra shown in Figure 3 is quite low for most sources (except in the detection of $\text{Ly}\alpha$), we construct a composite rest-frame ultraviolet spectrum which can serve two purposes: it validates bulk redshift identification by way of detecting lower S/N spectral features around $\text{Ly}\alpha$, and it begins to shed light on the intrinsic rest-frame ultraviolet emission properties of extremely infrared luminous starbursts. Unfortunately, larger samples are necessary to perform the latter analysis; in this work, our primary goal is to help validate our redshift identifications through cross-correlation to the composite spectrum for sources with only single-line identifications (e.g. $\text{Ly}\alpha$).

It is clear from Figure 3 that these infrared-starbursts exhibit a wide range in spectral properties, from Lyman Break Galaxy (LBG) spectra, quasar spectra, starburst spectra, to those with very steep to very shallow UV slopes. We classify sources as LBGs if they exhibit a steep cutoff in continuum flux at rest-frame 1216\AA , as quasars if they exhibit broad, high-ionization emission lines, and starbursts as narrow-emission line galaxies whose emission line luminosity is more significant than continuum luminosity. Many sources are dust obscured and thus are noisy except for the detection of $\text{Ly}\alpha$ emission. The construction of a composite spectrum serves as a sanity check on the redshifts. The detection of lower S/N features in a composite spectrum does not *directly* confirm that every galaxy added in is correctly identified, but it does indicate that most of them are. Since co-adding spectra from higher S/N sources would wash out the low S/N features of the

other galaxies (in a sample of 36), we exclude the following sources from the composites (excluded on the basis of detection of non- $\text{Ly}\alpha$ spectral features at $>3\sigma$ significance, including continuum and CIV): 1HERMES X24 J033136.96 -275510.9 , 1HERMES X24 J095917.28 $+021300.4$, 1HERMES X24 J095948.00 $+024140.7$, 1HERMES X1.4 J100008.64 $+022043.1$, 1HERMES X24 J100020.16 $+021725.2$, 1HERMES X24 J100036.00 $+021127.6$, 1HERMES X1.4 J100024.00 $+021210.9$, 1HERMES X1.4 J100111.52 $+022841.3$, 1HERMES X24 J100146.56 $+024035.6$, 1HERMES X1.4 J104620.40 $+585933.4$, 1HERMES X1.4 J104636.00 $+585650.0$, 1HERMES X24 J160545.99 $+534544.4$, and 1HERMES X1.4 J033151.94 -275326.9 . In other words, the composite spectra are only made up of $\text{Ly}\alpha$ single-line detections, those with low $\sim 2\text{--}3\sigma$ CIV detections, and those without.

Two different composites are constructed based on the detection or non-detection of CIV emission at this low S/N level. Since most starbursts are expected to show absorption in CIV, the co-addition of sources with and without CIV might easily yield a null result and no absorption or emission. Six sources are co-added in the CIV emission composite (these are the remaining sources for which CIV emission is detected, as indicated in Table 1, attributed to an AGN). The remaining 17 sources are co-added to form the composite without CIV emission. Each composite is constructed by scaling the flux of each galaxy to an arbitrary fixed mean value in the wavelength range 1330–1400Å. This wavelength range is chosen for its proximity to $\text{Ly}\alpha$ and absence of spectral signatures. The two composite rest-frame ultraviolet spectra are shown in Figure 7.

Both composites show a very high-S/N $\text{Ly}\alpha$ line, $\text{Ly}\alpha$ break, SiIV absorption, and either CIV emission or absorption. The composite LBG spectrum from Shapley *et al.* (2003) is shown for comparison, although the relative sample sizes should be contrasted. The CIV emission composite, by design, consists of galaxies with non-negligible AGN emission; as a result, the width of the $\text{Ly}\alpha$ line is broader in the CIV emission composite than in the CIV absorption composite.

We test for consistency between the composite spectra and individual source spectra through cross-correlation in the off- $\text{Ly}\alpha$ wavelength regions (note that for each galaxy’s spectrum, we measure the cross-correlation with a composite *excluding* that source). This provides an indication of sources which might be contaminating the composite rather than boosting its signal-to-noise. All individual sources in the CIV composite have correlations > 0.4 at $\Delta\lambda = 0$ offsets. The individual source spectra making up the other composite are of lower signal-to-noise, and have cross-correlations ranging 0.2–0.7. However, four sources (1HERMES X24 J095830.24 $+015633.2$, 1HERMES X1.4 J104707.69 $+585149.1$, 1HERMES X1.4 J104649.92 $+590039.6$ and 1HERMES X1.4 J123536.28 $+623019.9$) have almost no correlation with the composite (0.05–0.15) which is caused by no continuum detection (since only the off- $\text{Ly}\alpha$ spectra are considered in the cross-correlation test). Three of these sources are considered ‘tentative’ in their spectroscopic identifications in Table 1. The fourth source, 1HERMES X24 J095830.24 $+015633.2$, is

secure as judged by the quality of the $\text{Ly}\alpha$ detection and the inconsistency of this line being incorrectly identified as [OII].

4.4. Spectral Signatures of AGN

Since many of the sources in our sample have clear AGN features in their optical spectra, one might think that the infrared luminosities are contaminated by significant AGN heating rather than starburst heating. Typically, the presence of an AGN warms dust to temperatures $\gtrsim 100\text{--}200$ K. In this sample the majority of galaxies have dust temperatures $\lesssim 70$ K. Furthermore, two QSOs and the 13 sources with CIV detections have dust temperatures in the 30–50 K range, perfectly consistent with star formation dominated infrared emission. While there is still potential for AGN contribution to L_{IR} , the lack of correlation with dust temperature indicates that the effect is small ($\lesssim 25\%$, the nominal contribution of mid-infrared powerlaw emission to L_{IR} , Casey 2012). This is consistent with prior measures of SMGs with AGN ranging $\sim 15\text{--}25\%$ (Swinbank *et al.* 2004; Pope *et al.* 2008; Menéndez-Delmestre *et al.* 2009; Laird *et al.* 2010; Coppin *et al.* 2010). Note that one study, Alexander *et al.* (2005), could be interpreted to disagree with this work (finding $\sim 75\%$ of SMGs have AGN), however a minority of the sources in that data have AGN which dominate the sources’ bolometric luminosity.

We can draw some basic conclusions from the CIV emission and spectral types in our sample to infer the overall AGN content of $z > 2$ HSGs. For sources of sufficient signal-to-noise ($>5\sigma$ in continuum), we can assess AGN spectral signatures source by source. Of the 20 galaxies which meet this S/N cut, there are three Lyman Break Galaxies (LBGs), three quasars, seven starbursts with AGN (e.g. CIV emission), and seven “pure” starbursts (see Table 1 for details). Of the remaining single-line identifications, none show AGN signatures. Of the 36 sources, 10 have AGN signatures, three of which are obvious quasars. Although very qualitative, this analysis implies an AGN fraction of $\sim 25\%$ for the $z > 2$ HSG sample. From the composite spectra (which was constructed from only the lower luminosity sources), our statistics agree by construction; in other words, 6 out of 23 sources had CIV emission, or $\sim 26\%$. In a series of detailed studies on the multiwavelength properties of 850 μm -selected SMGs, Alexander *et al.* (2005), Pope *et al.* (2008) and Menéndez-Delmestre *et al.* (2009) also measure AGN fractions $\approx 25\%$ for similarly luminous $z \sim 2$ starbursts.

Interestingly, the sources exhibiting AGN signatures in the optical do not show hotter dust temperature SEDs in the infrared. One might expect higher dust temperatures in the infrared with the presence of an AGN heating the surrounding material to temperatures $\sim 100\text{--}200$ K, exceeding normal heating from star formation, $\sim 30\text{--}50$ K. The sources with AGN signatures are marked with small white dots in Figure 5. The observation that the AGN does not seem to have significant impact on infrared luminosity or dust temperature is not surprising if you consider that the star formation activity is at least an order of magnitude more luminous.

Although the selections of the SMG population and the HSG population differ, finding 1/4 with AGN might suggest that HSGs are similar in most ways to SMGs




FIG. 7.— A comparison of the rest-frame ultraviolet composite spectra for *Herschel*-selected galaxies (HSGs) with the composite of Lyman Break Galaxies (LBGs; top spectrum) compiled in Shapley *et al.* (2003) from 811 individual galaxies. The composite spectra of HSGs are split into those with CIV emission (middle spectrum) and CIV absorption (bottom spectrum) and *exclude* all sources with high-S/N ($>3\sigma$) spectral features which are not Ly α emission. There are six galaxies in the CIV emission stack and 17 galaxies in the CIV absorption stack, both having a median redshift of $\langle z \rangle = 3.6$. Ly α and the Lyman break, SiIV absorption and CIV emission/absorption are detected in both composites. More observations of similar sources is needed to enhance the S/N of the composites, which exhibit a wide range of spectral properties (as seen here and in Figure 3). The spectra are arbitrarily offset in flux with marked zero-points.

without any enhanced AGN activity, despite slightly warmer overall dust temperatures and brighter mid-infrared fluxes in comparison (described in the next section). However, further detailed work on these samples is needed before any conclusion is drawn as to the evolutionary nature of these *Herschel*-selected galaxies relative to classic 850 μm -selected SMGs.

4.5. Composite Infrared Spectra

Figure 8 combines all SPIRE, *Spitzer* and PACS (where available) infrared photometry for all galaxies in our sample from rest-frame ≈ 40 to 150 μm and radio data. This includes observations at 24 μm , 70 μm , 100 μm , 160 μm , 250 μm , 350 μm , and 500 μm . At these redshifts, the SPIRE bands probe the Wien-side of the thermal dust emission peak. Mean SEDs are fit using the modified blackbody plus power law method described by Equation 1 for photometric data which are scaled to the mean infrared luminosity of the sample, $\langle L_{\text{IR}} \rangle = 1.8 \times 10^{13} L_{\odot}$, and then separately, scaled to the mean radio flux density of the sample, $\langle S_{1.4} \rangle = 95 \mu\text{Jy}$ (or rest-frame $S_{1.4} = 283 \mu\text{Jy}$ assuming $\alpha = 0.75$). While all 36 galaxies are used in the former SED fit (top panel of Figure 8), the radio-scaled SED fit only has contributions from radio-detected galaxies.

There are two notable aspects of these mean SED fits seen in Figure 8; the first is the difference between the observed 24 μm flux densities relative to predictions from a 850 μm -selected SMG template spectrum (Pope *et al.* 2008), and second is the difference in dust temperatures between luminosity- and radio-scaled SEDs. The issue of the discrepancy of mid-infrared emission relates to the ongoing discussion of suppression of PAH emission in infrared starburst galaxies (e.g. Elbaz *et al.* 2011), where it is suggested that most normal galaxies have a fixed ratio ($\equiv \text{IR8}$) between 7.7 μm emission (or $\approx 8 \mu\text{m}$

emission) and total integrated infrared luminosity, L_{IR} and that sources with enhanced infrared emission are called infrared starbursts. However, as Hainline *et al.* (2009) point out, the mid-infrared portion of the spectrum does not lend itself to simple interpretation in terms of what is or is not AGN dominated or starburst dominated. SMGs, known to be extreme starbursts, do in fact have enhanced infrared emission relative to PAH strength (e.g. see Pope *et al.* 2008; Menéndez-Delmestre *et al.* 2009), however in this sample—which is on average more distant than most 850 μm -selected SMGs—we see mid-infrared flux densities ~ 2 –5 times the SMG expectation, more consistent with the measured IR8 value for most “main sequence” galaxies. What does this suggest about the SPIRE sample’s evolutionary histories? Is PAH emission simply not suppressed in these distant starbursts, or could the 24 μm “excess” be due to AGN heating? Or could these high redshift infrared luminous galaxies be “main sequence” secularly evolving galaxies? While this might be a selection bias based on 24 μm or 1.4 GHz detectability, the radio-selected galaxies (bottom panel) still show a mid-infrared excess above expectation from the SMG template.

The differences in dust temperatures between the two mean SED fits ($42 \pm 3 \text{ K}$ vs. $54 \pm 4 \text{ K}$ for luminosity-scaled and radio-scaled, respectively) is traceable to a radio selection bias. For two galaxies with similar SPIRE flux densities, one with a warm temperature ($\sim 60 \text{ K}$) and one with a cold temperature ($\sim 30 \text{ K}$), the galaxy with the warm temperature is going to have a much higher integrated infrared luminosity and therefore much brighter 1.4 GHz detection at these redshifts. Therefore, when we consider just radio-detected galaxies, the average dust temperature increases due to the exclusion of cold, non-detectable galaxies.

As discussed earlier, we estimate that 16–43% of these




FIG. 8.— The mean infrared and radio SEDs for the sample re-normalized to the mean IR luminosity of the sample, $1.8 \times 10^{13} L_{\odot}$ (top) and to mean radio flux density, $95 \mu\text{Jy}$ (bottom). The mean normalized flux density in $\log(\lambda) = 0.1$ bins is shown as black squares, from rest-frame ≈ 4 – $150 \mu\text{m}$. Best-fit SEDs (solid black lines) are generated as described in the text according to Equation 1 with fixed $\beta = 1.5$. They comprise a cold-dust modified blackbody fit (dashed line) and a mid-infrared power law representative of warm dust emission. Radio synchrotron emission is added onto this best-fit infrared SED by assuming the FIR/radio correlation holds with a synchrotron slope of $\alpha = 0.75$. The composite SMG SED described in Pope *et al.* (2008) is shown as a dashed line. Both the luminosity-scaled SED and radio-scaled SED appear to have a $24 \mu\text{m}$ excess relative to the SMG expectation. The radio-scaled SED has a hotter characteristic dust temperature than the luminosity-scaled SED (both of which are uncertain by $\sim 3 \text{ K}$), likely driven by the bias against colder-dust galaxies of similar flux densities (less likely to be radio detected).

galaxies would be formally undetected at $850 \mu\text{m}$ at < 2 – 5 mJy . In other words, 16–43% of HSGs at $z > 2$ are consistent with the Submillimeter Faint Radio Galaxy (SFRG, formerly optically faint, “OFRG”) selection and not SMG selection (Chapman *et al.* 2004; Casey *et al.* 2009; Chapman *et al.* 2010; Magdis *et al.* 2010; Casey *et al.* 2011,?). The composite infrared SEDs from Figure 8 support this conclusion, since the range of observed $850 \mu\text{m}$ flux densities from the best-fit SEDs is in the $1 < S_{850} < 10 \text{ mJy}$ range, not as luminous at long wavelengths as the $850 \mu\text{m}$ -selected composite (Pope *et al.* 2008).

5. SFRD IMPLICATIONS AND DISCUSSION

To place these $z > 2$ SPIRE-selected galaxies in context with other high- z infrared galaxies and lower luminosity galaxies, we estimate their contribution to the cosmic star formation rate density (SFRD; Madau *et al.* 1996; Madau, Pozzetti & Dickinson 1998; Hopkins & Beacom 2006). The SFRD contribution allows a direct comparison of the importance of infrared-luminous galaxies to

the build-up of stellar mass in the Universe over a range of epochs. At lower redshifts $z \lesssim 1$, ultraluminous infrared galaxies are very rare and contribute little to the SFRD (Sanders *et al.* 2003) but towards $z \approx 1$, the importance of ULIRGs grows, and it is estimated that LIRGs and ULIRGs ($L_{\text{IR}} > 10^{11} L_{\odot}$) could contribute as much as $\sim 1/2$ of the total SFRD (see results from Spitzer; Le Floc’h *et al.* 2005, and work in C12). At $z > 1$, the contribution of infrared-luminous sources is much more difficult to measure, limited by small numbers of SMGs (Chapman *et al.* 2005; Wardlow *et al.* 2011) or complex selection biases or extrapolations from the mid-infrared (Caputi *et al.* 2007; Magnelli *et al.* 2011; Capak *et al.* 2011). The $2 < z < 5$ galaxies in this paper provide a unique sample to make this measurement, due to their well characterized selection over a relatively large sky area, $\sim 1 \text{ deg}^2$.

To arrive at SFRD estimates, we first compute the infrared luminosity function using a $1/V_{\text{max}}$ method, where each source is associated with the maximum volume in which it could be detected at its given luminosity, L_{IR} (Schmidt 1968). The number density of sources with luminosity between L and $L + \Delta L$ is given as $\Phi_z(L) \Delta L = \sum 1/(V_i(L) \times c_i)$ in units of $h^3 \text{ Mpc}^{-3} \log L^{-1}$. Here c_i is a completeness estimator which corrects for sample incompleteness at the selection wavelength, e.g. in this case, at 250 – $500 \mu\text{m}$. C12 presents a detailed discussion of this completeness factor as a function of selection wavelength flux density which varies field to field (based on the prior source catalog depths). Sources with flux densities $> 15 \text{ mJy}$ will be more than 90% complete (e.g. $c_i > 0.9$) in all fields.

The completeness estimator’s effect on the integrated SFRD is small compared to the uncertainty in the luminosity function itself from small number counts. Note, however, that this correction only pertains to one of the many sources of incompleteness of this sample; it is far more difficult to quantify and correct for incompleteness with respect to sources missing from the prior catalog list (e.g. 1.4 GHz or $24 \mu\text{m}$ faint), which is particularly a problem at $z > 2$, as well as spectroscopic incompleteness, i.e. the galaxies which have no detectable emission lines or are too optically obscured to be identified.

$V_i(L)$ is the maximum volume in which source i could reside and still be detectable by our survey. Since the detection limits of SPIRE alter between field catalogs, the detection limit determining the maximum volume is calculated source by source. This luminosity detection limit is determined much like the curves in Figure 6. For example, a source might have its highest S/N at $350 \mu\text{m}$, then its luminosity detection limit, thus maximal redshift limit z_{max} , is determined by setting a 3σ detection threshold at $350 \mu\text{m}$ where sigma is the local confusion plus instrumental noise in the $350 \mu\text{m}$ map. This z_{max} limit is then found across the entire survey area probed to determine accessible volume. The assumed dust temperature is that measured for the given source (since dust temperature does impact the steepness of the luminosity limit with redshift).

We split the luminosity function into two redshift bins: $2.0 < z < 3.2$ and $3.2 < z < 5.0$ with 15 sources in the former and 20 in the latter. Since LRIS and DEIMOS have different wavelength coverage, DEIMOS observations suffer from a redshift desert from $1.6 <$




FIG. 9.— The estimated luminosity function for $2 < z < 5$ *Herschel*-SPIRE selected galaxies compared to the luminosity function of $850\mu\text{m}$ -selected SMGs at $z \approx 2.5$ (Chapman *et al.* 2005; Wardlow *et al.* 2011). Up arrows denote the fact that this survey is *spectroscopically incomplete* and that the incompleteness is not well quantified at $z > 2$. Numbers next to each point indicate how many galaxies from our sample contribute to that luminosity bin; the numbers are comparable to those in the $z \approx 2$ SMG samples. The luminosity function for the whole sample at $3.2 < z < 5.0$ is shown in salmon, while the high confidence (h.c.) identifications’ luminosity function is shown in dark red.

$z < 3.2$ that LRIS observations do not, so we split the sample at $z = 3.2$ and only compute the density using LRIS observations between $z = 2.0$ and $z = 3.2$. This excludes two sources from the calculation (1HERMES X1.4 J104636.00+585650.0 at $z = 2.841$ and 1HERMES X24 J160545.99+534544.4 at $z = 2.555$) which were both surveyed with DEIMOS and detected in the redshift desert due to strong CIV emission caused by the presence of a quasar. Since high-redshift $z \gtrsim 2$ sources are only detectable on the masks observed in the best weather conditions, the effective area probed by LRIS for this calculation is 0.13 deg^2 over the range $2.0 < z < 5.0$ and for DEIMOS 0.30 deg^2 over the range $3.2 < z < 5.0$. The resulting *spectroscopically incomplete* luminosity functions are shown in Figure 9.

Since some of our identifications are less confident than others (e.g. those marked with a † in Table 1) we also compute the luminosity function excluding tentative identifications. The result is seen in Figure 9: while the $2.0 < z < 3.2$ luminosity function remains the same (differing only by two sources), the $3.2 < z < 5.0$ luminosity function drops by 11 sources (salmon vs dark red LFs).

The SFRD is then the luminosity weighted integral of the luminosity function, or the raw summation of the luminosity (converted to SFR) over accessible volume: $(7.0 \pm 2.0) \times 10^{-3} M_{\odot} \text{ yr}^{-1} h^3 \text{ Mpc}^{-3}$ at $2.0 < z < 3.2$ and $(5.5 \pm 0.6) \times 10^{-3} M_{\odot} \text{ yr}^{-1} h^3 \text{ Mpc}^{-3}$ at $3.2 < z < 5.0$ (full sample) and $(2^{+3}_{-1}) \times 10^{-3} M_{\odot} \text{ yr}^{-1} h^3 \text{ Mpc}^{-3}$ at $3.2 < z < 5.0$ (high confidence sample), shown in Figure 10 against other comparison populations. These points are lower limits since they do not include any sources which might be excluded from the prior catalogs at $24\mu\text{m}$ or 1.4 GHz , which is speculated to be a non-negligible frac-




FIG. 10.— Star formation rate density of *Herschel*-SPIRE selected galaxies (black points) relative to $850\mu\text{m}$ -selected SMGs (Chapman *et al.* 2005; Wardlow *et al.* 2011) and 1.2 mm -selected MMGs (Roseboom *et al.* 2012). The compilation of SFRD measurements from (Hopkins & Beacom 2006) is shown as a gray band, which is largely drawn from optical or rest-frame ultraviolet-selected galaxy populations corrected for dust extinction. The luminosity limits of integration are $10^{(12.4-13.4)} L_{\odot}$ at $z \sim 2.5$ and $10^{(12.8-13.6)} L_{\odot}$ at $z \sim 4$; the SFRD from sources sitting in the luminosity where the two redshift bins overlap ($10^{(12.8-13.2)} L_{\odot}$) is shown in green. The SFRD measurements for *Herschel*-selected galaxies at $z < 2$ are shown as gray points. The sharp drop in the SFRD of *Herschel*-selected samples at $z \sim 2$ is caused by the redshifting of the SED peak such that more infrared-luminous galaxies are SPIRE “drop-outs” and that only the warmest, most-luminous $> 10^{13} L_{\odot}$ systems are detectable with SPIRE at $z > 2$. Note that we observe an increase in the infrared-luminous contribution to the SFRD from $z \sim 2.6$ to $z \sim 4$.

tion ($\gtrsim 20\%$) at $z > 2$ (e.g. Magdis *et al.* 2010, Smail *et al.* in prep). Note also that the luminosity limits of the two redshift bins differ: the $z \sim 2.5$ bin covers $10^{12.4-13.2} L_{\odot}$ while the $z \sim 4$ bin covers $10^{12.8-13.6} L_{\odot}$. To assess luminosity evolution from $z \sim 2 - 5$, we compute the SFRD contributions in the overlapping luminosity range of both redshift bins, $10^{12.8-13.2} L_{\odot}$, shown as green points in Figure 10.

Although these measurements of the *Herschel* contribution to the SFRD at $2 < z < 5$ are lower limits due to our survey’s incompleteness, the effects of gravitational lensing and clustering could lead to an overestimation. Are these effects significant in this sample? For the former we use the conditional lensing probability as a function of SPIRE flux density (Francesco De Bernardis, private communication, Wardlow *et al.* 2012). For SPIRE flux densities $S_{500} < 80\text{ mJy}$ the distribution in number counts is dominated by a Schechter function rather than the flat-sloped source counts at $> 80\text{ mJy}$; galaxies with $S_{500} > 80\text{ mJy}$ have a high probability of being lensed by factors > 2 while this model predicts a mean lensing factor for this sample of $\langle \mu \rangle < 1.05$, which changes negligibly between $z = 2$ and $z = 5$.

The brightest source in our sample with $S_{250} = 73.1\text{ mJy}$, 1HERMES X24 J161506.65+543846.9, has the highest probability of being lensed (its expected lensing factor is $\langle \mu \rangle = 1.2$). It is the highest redshift source in our sample at $z = 4.952$. Due to its extreme luminosity compared to the rest of the sample and its tentative spectroscopic identification, we exclude this source from the star formation rate density (SFRD) measurement.

To assess the impact of clustering on the SFRD measurement, we need a good grasp of the spatial density

of $z > 2$ sources on our slit-masks and the possibility of biased placement of slit-masks around high- z clusters. The former can be gauged by the number of high- z confirmations per slit-mask; the 36 sources of this sample are distributed across 12 LRIS masks and 14 DEIMOS masks, with an additional 5 LRIS masks and 3 DEIMOS masks without any high- z sources; this averages to 0–2 galaxies per mask without any mask having more than 2 sources. Since the masks were distributed randomly with respect to one another in each field and none of the masks were close together, this demonstrates that these sources are indeed randomly distributed over the surveyed area, 0.93 deg^2 for the whole survey.

The possibility exists that there is an intrinsic bias of the placement of our slit-masks such that more high- z sources are observed than elsewhere. As explained in detail in C12, masks were placed around high-priority targets which were ‘red’ in their SPIRE colors (e.g. $S_{250} < S_{350} < S_{500}$ all with $S \gtrsim 15 \text{ mJy}$) and thought to be high redshift sources. Of the 36 confirmed $z > 2$ sources, seven ($\sim 19\%$) were originally high-priority targets. However, an additional 44 high-priority targets were identified at $z < 2$ and 78 were unidentified. Relative to the number of high-priority sources targeted, we measure $5 \pm 5\%$ as identified at $z > 2$, $34 \pm 5\%$ at $z < 2$ and $60 \pm 4\%$ unidentified. The same statistics for lower priority targets are $2 \pm 1\%$ at $z > 2$, $48 \pm 1\%$ at $z < 2$, and $50 \pm 1\%$ unidentified. Within uncertainties, the proportion of sources identified at $z > 2$ are the same between low-priority and high-priority targets, indicating no bias or advantage in targeting ‘red’ sources more than any other significant *Herschel*-SPIRE source. This implies that no clustering correction on the measured SFRD is necessary.

The lower limits to the SFRD set by SPIRE sources tells us that the early Universe potentially had a very substantial amount of star formation in short-lived, intense $> 1000 \text{ M}_\odot \text{ yr}^{-1}$ bursts as opposed to slow-progressing moderate levels of star formation. The contribution from SPIRE to the SFRD at these epochs is at least comparable to the contribution measured from longer wavelength-selected galaxies, like the $850 \mu\text{m}$ -selected SMGs (Chapman *et al.* 2005; Wardlow *et al.* 2011) or 1.2 mm -selected MMGs (Roseboom *et al.* 2012). This is made more interesting by the observation that the populations (SMG and HSG) only overlap by 21 out of 36 galaxies (58%). Further work aimed at confirming redshifts of $z > 2$ SPIRE sources, particularly those without radio or $24 \mu\text{m}$ counterparts, is needed to constrain these lower limits into real measurements so the importance of $\gtrsim 10^{13} \text{ L}_\odot$ activity in the first few Gyr of the Universe is understood.

6. CONCLUSIONS

The identification of submillimeter galaxies at early epochs in the Universe’s history is the key to understanding the limits of star formation and galaxy evolution on short timescales. This paper has presented new observations of 36 *Herschel*-SPIRE selected starburst galaxies between $2 < z < 5$, taken from a large Keck spectroscopic survey of 1594 SPIRE-selected galaxies covering 0.93 deg^2 .

We present the following conclusions:

- Our sample of 36 *Herschel*-selected galaxies (HSGs)

constitute some of the brightest, most extreme infrared starburst galaxies in the Universe. Spanning $2 < z < 5$, our sample has a mean luminosity $\langle L_{\text{IR}} \rangle = 1.8 \times 10^{13} \text{ L}_\odot$ ($\text{SFR} \approx 3100 \text{ M}_\odot \text{ yr}^{-1}$).

- These $2 < z < 5$ HSGs have a well characterized selection across 6 legacy fields and 0.93 deg^2 ; galaxies must be $> 3\sigma$ significant in one of the three *Herschel*-SPIRE bands and also be detected in deep $24 \mu\text{m}$ and/or 1.4 GHz survey coverage. Although it misses $24 \mu\text{m}$ or 1.4 GHz high- z dropouts, the selection is identical to low- z HSG selection. Sources at $z > 2$ comprise 5% of all galaxies selected via this method; although inefficient for finding high- z infrared galaxies, the selection is easily reproducible and well suited for volume density estimates.
- Our sample show a wide range of rest-frame ultraviolet spectral features: some galaxies classifiable as quasars, some as LBGs, and most as starbursts with a wide range of dust extinctions/reddening. The heterogeneous nature of their spectra provide additional evidence that the infrared-luminous stage might exist during a period when the host galaxy is rapidly evolving.
- The radio-detected subset of our sample (23/36) follow the FIR/radio correlation consistent with moderate evolution in q_{IR} from previous work (Ivison *et al.* 2010).
- We construct composite rest-frame ultraviolet spectra and rest-frame infrared spectral energy distributions to assess some aggregate properties of HSGs. In the rest-frame UV, we determine that 25% of HSGs exhibit CIV emission (a signature of AGN). In the infrared, HSGs exhibit a $24 \mu\text{m}$ excess relative to SMGs of similar L_{IR} ; without mid-IR spectra, it is impossible to know whether this is due to enhanced PAH emission (similar to “normal” galaxies) or AGN emission. The dust temperatures of radio-selected samples are warmer than those of the full sample.
- Our spectroscopic survey is incomplete due to selection bias at $24 \mu\text{m}$ and 1.4 GHz , as well as spectroscopic incompleteness caused by heavy dust obscuration in the rest-frame UV. Therefore, we are able to place *lower limits* on the contribution of $2 < z < 5$ HSGs to the cosmic star formation rate density, which is $> 7 \times 10^{-3} \text{ M}_\odot \text{ yr}^{-1} \text{ h}^3 \text{ Mpc}^{-3}$ at $z \approx 2.6$ and $> 3 \times 10^{-3} \text{ M}_\odot \text{ yr}^{-1} \text{ h}^3 \text{ Mpc}^{-3}$ at $z \approx 4$, corresponding to $> 10\%$ and $> 20\%$ of the best-estimates of the total SFRD at their respective epochs.

This work highlights the importance of extremely luminous FIR-bright galaxies to the build-up of stellar mass, particularly at early times in the Universe’s history. Further work on constraining completeness and the parent population of infrared-luminous galaxies at $z > 2$ is needed to understand the role that short-lived starbursts have in the context of galaxy evolution and formation.

ACKNOWLEDGEMENTS

We thank the anonymous referee for his/her constructive suggestions which improved the manuscript. CMC is generously supported by a Hubble Fellowship from Space Telescope Science Institute, grant HST-HF-51268.01-A. The data presented herein were obtained at the W.M. Keck Observatory, which is operated as a scientific partnership among the California Institute of Technology, the University of California and the National Aeronautics and Space Administration. The Observatory was made possible by the generous financial support of the W.M. Keck Foundation. The authors wish to recognize and acknowledge the very significant cultural role and reverence that the summit of Mauna Kea has always had within the indigenous Hawaiian community. We are most fortunate to have the opportunity to conduct observations from this mountain. This work would not be possible without the hard work and dedication of the Keck Observatory night and day staff; special thanks to Marc Kassis, Luca Rizzi and Greg Wirth for help and advice while observing. The analysis pipeline used to reduce the DEIMOS

data was developed at UC Berkeley with support from NSF grant AST-0071048.

SPiRE has been developed by a consortium of institutes led by Cardiff Univ. (UK) and including: Univ. Lethbridge (Canada); NAOC (China); CEA, LAM (France); IFSI, Univ. Padua (Italy); IAC (Spain); Stockholm Observatory (Sweden); Imperial College London, RAL, UCL-MSSL, UKATC, Univ. Sussex (UK); and Caltech, JPL, NHSC, Univ. Colorado (USA). This development has been supported by national funding agencies: CSA (Canada); NAOC (China); CEA, CNES, CNRS (France); ASI (Italy); MCINN (Spain); SNSB (Sweden); STFC, UKSA (UK); and NASA (USA).

This research has made use of data from the HerMES project (<http://hermes.sussex.ac.uk/>). HerMES is a *Herschel* Key Programme utilizing Guaranteed Time from the SPiRE instrument team, ESAC scientists and a mission scientist. HerMES is described in Oliver *et al.* (2012). The SPiRE data presented in this paper will be released through the HerMES Database in Marseille, HeDaM (<http://hedam.oamp.fr/HerMES>).

REFERENCES

- Alexander, D. M. *et al.* 2005, ApJ, 632, 736.
 Barger, A. J. *et al.* 1998, Nature, 394, 248.
 Béthermin, M. *et al.* 2011, A&A, 529, A4.
 Biggs, A. D. and Ivison, R. J. 2008, MNRAS, 385, 893.
 Blain, A. W., Barnard, V. E., and Chapman, S. C. 2003, MNRAS, 338, 733.
 Blain, A. W. *et al.* 2004, ApJ, 611, 725.
 Blain, A. W. *et al.* 2002, Phys. Rep., 369, 111.
 Bridge, C. R. *et al.* 2012, ArXiv e-prints.
 Capak, P. L. *et al.* 2011, Nature, 470, 233.
 Caputi, K. I. *et al.* 2007, ApJ, 660, 97.
 Cardamone, C. N. *et al.* 2010, ApJS, 189, 270.
 Casey, C. M. 2012, ArXiv e-prints.
 Casey, C. M. *et al.* 2009, MNRAS, 399, 121.
 Casey, C. M. *et al.* 2011a, MNRAS, 415, 2723.
 Casey, C. M. *et al.* 2011b, MNRAS, 411, 2739.
 Casey, C. M. *et al.* 2012, ApJ in press.
 Chapin, E. L. *et al.* 2011, MNRAS, 411, 505.
 Chapman, S. C. *et al.* 2005, ApJ, 622, 772.
 Chapman, S. C. *et al.* 2003, ApJ, 588, 186.
 Chapman, S. C. *et al.* 2010, MNRAS, 409, L13.
 Chapman, S. C. *et al.* 2004a, ApJ, 614, 671.
 Chapman, S. C. *et al.* 2004b, ApJ, 611, 732.
 Chary, R. and Elbaz, D. 2001, ApJ, 556, 562.
 Clements, D. L. *et al.* 2008, MNRAS, 387, 247.
 Condon, J. J. 1992, ARA&A, 30, 575.
 Conley, A. *et al.* 2011, ApJ, 732, L35.
 Coppin, K. *et al.* 2010, ApJ, 713, 503.
 Coppin, K. E. K. *et al.* 2009, MNRAS, 395, 1905.
 Cowie, L. L. *et al.* 1996, AJ, 112, 839.
 Cox, P. *et al.* 2011, ApJ, 740, 63.
 Daddi, E. *et al.* 2009, ApJ, 694, 1517.
 Dale, D. A. and Helou, G. 2002, ApJ, 576, 159.
 Davé, R. *et al.* 2010, MNRAS, 404, 1355.
 Dekel, A. *et al.* 2009, Nature, 457, 451.
 Draine, B. T. and Li, A. 2007, ApJ, 657, 810.
 Eales, S. *et al.* 1999, ApJ, 515, 518.
 Elbaz, D. *et al.* 2011, A&A, 533, A119+.
 Griffin, M. J. *et al.* 2010, A&A, 518, L3.
 Hainline, L. J. *et al.* 2009, ApJ, 699, 1610.
 Helou, G., Soifer, B. T., and Rowan-Robinson, M. 1985, ApJ, 298, L7.
 Hinshaw, G. *et al.* 2009, ApJS, 180, 225.
 Hopkins, A. M. and Beacom, J. F. 2006, ApJ, 651, 142.
 Hughes, D. H. *et al.* 1998, Nature, 394, 241.
 Ibar, E. *et al.* 2010, MNRAS, 401, L53.
 Ilbert, O. *et al.* 2010, ApJ, 709, 644.
 Ivison, R. J. *et al.* 2010a, MNRAS, 402, 245.
 Ivison, R. J. *et al.* 2007, ApJ, 660, L77.
 Ivison, R. J. *et al.* 2010b, A&A, 518, L31+.
 Kennicutt, Jr., R. C. 1998, ApJ, 498, 541.
 Kovács, A. *et al.* 2010, ApJ, 717, 29.
 Laird, E. S. *et al.* 2010, MNRAS, 401, 2763.
 Le Floc'h, E. *et al.* 2009, ApJ, 703, 222.
 Le Floc'h, E. *et al.* 2005, ApJ, 632, 169.
 Levenson, L. *et al.* 2010, MNRAS, 409, 83.
 Lutz, D. *et al.* 2011, A&A, 532(90).
 Madau, P. *et al.* 1996, MNRAS, 283, 1388.
 Madau, P., Pozzetti, L., and Dickinson, M. 1998, ApJ, 498, 106.
 Magdis, G. E. *et al.* 2010, MNRAS, 409, 22.
 Magnelli, B. *et al.* 2011, A&A, 528, A35.
 Magnelli, B. *et al.* 2012, A&A, 539(155).
 Menéndez-Delmestre, K. *et al.* 2009, ApJ, 699, 667.
 Mobasher, B. *et al.* 2007, ApJS, 172, 117.
 Neufeld, D. A. 1991, ApJ, 370, L85.
 Nguyen, H. T. *et al.* 2010, A&A, 518, L5.
 Oliver, S. J. *et al.* 2012, ArXiv e-prints.
 Pilbratt, G. L. *et al.* 2010, A&A, 518, L1.
 Pope, A. *et al.* 2008, ApJ, 689, 127.
 Roseboom, I. G. *et al.* 2012, MNRAS, 419, 2758.
 Roseboom, I. G. *et al.* 2010, MNRAS, 409, 48.
 Rowan-Robinson, M. 2000, MNRAS, 316, 885.
 Rowan-Robinson, M. *et al.* 2008, MNRAS, 386, 697.
 Sanders, D. B. *et al.* 2003, AJ, 126, 1607.
 Sanders, D. B. and Mirabel, I. F. 1996, ARA&A, 34, 749.
 Sanders, D. B. *et al.* 1988, ApJ, 325, 74.
 Scarlata, C. *et al.* 2009, ApJ, 704, L98.
 Schmidt, M. 1968, ApJ, 151, 393.
 Shapley, A. E. *et al.* 2003, ApJ, 588, 65.
 Siana, B. *et al.* 2007, ApJ, 668, 62.
 Siana, B. *et al.* 2010, ApJ, 723, 241.
 Siebenmorgen, R. and Krügel, E. 2007, A&A, 461, 445.
 Smail, I., Ivison, R. J., and Blain, A. W. 1997, ApJ, 490, L5+.
 Smolčić, V. *et al.* 2011, ApJ, 731, L27.
 Strazzullo, V. *et al.* 2010, ApJ, 714, 1305.
 Swinbank, A. M. *et al.* 2008, MNRAS, 391, 420.
 Swinbank, A. M. *et al.* 2004, ApJ, 617, 64.
 Tacconi, L. J. *et al.* 2008, ApJ, 680, 246.
 Walter, F. *et al.* 2012, Nature, 486, 233.
 Wang, W.-H., Barger, A. J., and Cowie, L. L. 2009, ApJ, 690, 319.
 Wang, W.-H. *et al.* 2011, ApJ, 726, L18.
 Wang, W.-H. *et al.* 2007, ApJ, 670, L89.
 Wardlow, J. L. *et al.* 2012, ArXiv e-prints.
 Wardlow, J. L. *et al.* 2011, MNRAS, 415, 1479.
 Younger, J. D. *et al.* 2009, MNRAS, 394, 1685.

TABLE 1
 $z > 2$ IDENTIFIED SPIRE-SELECTED GALAXIES

| NAME | z_{spec} | Comments | CLASS | z_{phot} | S_{24} (μJy) | S_{100} (mJy) | S_{160} (mJy) | S_{250} (mJy) | S_{350} (mJy) | S_{500} (mJy) | $S_{1.4\text{GHz}}$ (μJy) | L_{IR} (L_{\odot}) | T_{dust} (K) |
|-------------------------|-------------------|---|--------|-------------------|--------------------------------|--------------------|--------------------|--------------------|--------------------|--------------------|---|--|--------------------------|
| 1HERMES X24 J033136.96 | 275510.9... | 3.145 ^L Ly α -break | LBG | 3.44 ^d | 265 \pm 11 | — | — | 15.5 \pm 3.8 | 20.8 \pm 3.8 | 24.0 \pm 4.8 | ... | (1.0 $^{+0.3}_{-0.2}$) \times 10 ¹³ | 40.6 \pm 3.6 |
| 1HERMES X1.4 J033151.94 | 275326.9... | 2.938 ^L Ly α ; SiIV | SB | 2.96 ^d | 397 \pm 14 | — | — | 12.9 \pm 3.9 | 21.6 \pm 3.7 | 24.2 \pm 4.2 | 54.3 \pm 12.8 | (1.2 $^{+0.3}_{-0.2}$) \times 10 ¹³ | 43.8 \pm 3.4 |
| 1HERMES X24 J033319.58 | 274119.7... | 2.325 ^L Ly α | SB | 1.84 ^d | 329 \pm 13 | — | — | 16.0 \pm 3.8 | 19.0 \pm 3.8 | 15.2 \pm 4.5 | ... | (5.4 $^{+1.7}_{-1.3}$) \times 10 ¹² | 39.3 \pm 3.8 |
| 1HERMES X24 J095830.24 | +015633.2... | 2.327 ^L Diffuse Ly α | SB | 1.70 ^a | 418 \pm 16 | ... | ... | 19.2 \pm 2.2 | 23.4 \pm 3.0 | 23.0 \pm 3.3 | ... | (5.4 $^{+2.9}_{-1.9}$) \times 10 ¹² | 24.6 \pm 3.7 |
| 1HERMES X24 J095916.08 | +021215.3... | 4.454 ^L Ly α | SB | 2.97 ^a | 282 \pm 14 | ... | 10.1 \pm 3.4 | 25.8 \pm 2.2 | 24.1 \pm 2.9 | 14.7 \pm 3.2 | ... | (1.9 $^{+0.7}_{-0.7}$) \times 10 ¹³ | 45.9 \pm 10.7 |
| 1HERMES X24 J095917.28 | +021300.4... | 2.101 ^L Ly α ; SiIV; CIV; + | SB | 1.95 ^a | 472 \pm 17 | ... | 8.4 \pm 2.7 | 14.2 \pm 2.2 | 14.7 \pm 2.7 | 7.1 \pm 3.0 | ... | (4.4 $^{+1.4}_{-1.0}$) \times 10 ¹² | 51.5 \pm 5.2 |
| 1HERMES X1.4 J095934.08 | +021706.3... | 2.926 ^L Ly α | SB | 2.94 ^a | 1451 \pm 16 | ... | 11.5 \pm 2.9 | 11.4 \pm 2.2 | 12.9 \pm 3.4 | 0.2 \pm 4.7 | 170 \pm 15 | (2.0 $^{+0.4}_{-0.2}$) \times 10 ¹³ | 100 \pm 3.0 |
| 1HERMES X24 J095948.00 | +024140.7... | 2.600 ^L Ly α ; CIV | SB+AGN | 2.27 ^a | 909 \pm 16 | ... | 13.6 \pm 4.2 | 10.5 \pm 2.2 | 10.7 \pm 2.8 | 6.4 \pm 3.0 | ... | (5.9 $^{+5.3}_{-2.7}$) \times 10 ¹² | 39.4 \pm 12.2 |
| 1HERMES X1.4 J100008.64 | +022043.1... | 2.888 ^L Ly α -break; CIV | LBG | 2.18 ^a | 574 \pm 14 | ... | ... | 16.5 \pm 2.2 | 12.3 \pm 2.9 | 9.4 \pm 3.6 | 66 \pm 12 | (7.9 $^{+6.0}_{-3.4}$) \times 10 ¹² | 31.4 \pm 7.2 |
| 1HERMES X24 J100020.16 | +021725.2... | 2.105 ^L Ly α ; SiIV; CIV | QSO | 1.45 ^a | 401 \pm 16 | ... | ... | 20.0 \pm 2.2 | 12.6 \pm 3.5 | 22.3 \pm 3.7 | ... | (5.7 $^{+1.0}_{-1.0}$) \times 10 ¹² | 32.9 \pm 2.2 |
| 1HERMES X1.4 J100024.00 | +021210.9... | 3.553 ^L Ly α ; SiIV; CIV | SB+AGN | 2.97 ^a | 175 \pm 50 | 6.1 \pm 1.5 | 31.7 \pm 3.6 | 28.5 \pm 2.7 | 28.8 \pm 3.5 | 15.1 \pm 5.7 | 84 \pm 13 | (2.0 $^{+0.4}_{-0.3}$) \times 10 ¹³ | 54.0 \pm 5.3 |
| 1HERMES X24 J100036.00 | +021127.6... | 2.103 ^L Ly α ; SiIV; CIV | SB | 1.48 ^a | 158 \pm 17 | ... | 11.2 \pm 2.9 | 13.2 \pm 2.7 | 21.2 \pm 3.6 | 4.9 \pm 11.4 | ... | (2.7 $^{+1.3}_{-0.9}$) \times 10 ¹² | 29.0 \pm 3.9 |
| 1HERMES X1.4 J100111.52 | +022841.3... | 3.975 ^L Ly α -break; CIV | LBG | 2.60 ^a | 201 \pm 41 | ... | ... | 24.5 \pm 2.2 | 32.9 \pm 4.3 | 22.8 \pm 6.8 | 59 \pm 11 | (1.8 $^{+0.6}_{-0.5}$) \times 10 ¹³ | 46.7 \pm 7.2 |
| 1HERMES X24 J100133.36 | +023726.9... | 2.619 ^L Diffuse Ly α † | SB | —99 ^a | 245 \pm 15 | ... | ... | 10.1 \pm 2.2 | 9.1 \pm 2.9 | 0.0 \pm 9.9 | ... | (4.4 $^{+2.5}_{-1.9}$) \times 10 ¹² | 51.9 \pm 8.3 |
| 1HERMES X24 J100146.56 | +024035.6... | 2.050 ^L SiIV; CIV; HeII | SB+AGN | 1.88 ^a | 722 \pm 17 | 4.5 \pm 1.2 | ... | 11.7 \pm 2.2 | 6.9 \pm 2.8 | 2.2 \pm 3.1 | ... | (4.7 $^{+1.3}_{-1.0}$) \times 10 ¹² | 70.7 \pm 8.3 |
| 1HERMES X24 J100150.16 | +024017.2... | 2.883 ^L Ly α | SB | 2.57 ^a | 315 \pm 19 | ... | 17.5 \pm 4.6 | 13.8 \pm 2.2 | 12.2 \pm 2.9 | 8.9 \pm 4.0 | ... | (7.1 $^{+6.2}_{-3.3}$) \times 10 ¹² | 48.4 \pm 16.5 |
| 1HERMES X1.4 J100151.60 | +023909.5... | 4.538 ^L Ly α † | SB | 2.20 ^a | 433 \pm 102 | ... | ... | 17.8 \pm 2.2 | 0.5 \pm 5.4 | 22.1 \pm 3.2 | ... | (1.7 $^{+2.3}_{-0.9}$) \times 10 ¹³ | 35.0 \pm 11.3 |
| 1HERMES X1.4 J104557.12 | +590000.4... | 3.382 ^L Ly α ; CIV | SB | 3.38 ^c | 713 \pm 7 | 5.7 \pm 2.7 | 8.5 \pm 4.8 | 16.9 \pm 3.6 | 7.6 \pm 3.9 | 0.0 \pm 4.7 | 101.9 \pm 15.9 | (1.7 $^{+0.5}_{-0.4}$) \times 10 ¹³ | 87.6 \pm 8.9 |
| 1HERMES X1.4 J104620.40 | +585933.4... | 2.211 ^L Ly α ; CIV | SB+AGN | 2.12 ^c | 559 \pm 7 | — | ... | 13.1 \pm 3.5 | 3.4 \pm 3.9 | 0.0 \pm 4.7 | 64.5 \pm 11.8 | (5.1 $^{+2.7}_{-1.8}$) \times 10 ¹² | 63.1 \pm 9.7 |
| 1HERMES X1.4 J104636.00 | +585650.0... | 2.841 ^D CIV | QSO | 1.84 ^c | 166 \pm 7 | ... | ... | 11.3 \pm 3.5 | 5.4 \pm 4.0 | 0.0 \pm 4.9 | 19.4 \pm 3.6 | (4.1 $^{+2.0}_{-1.3}$) \times 10 ¹² | 48.0 \pm 10.8 |
| 1HERMES X24 J104642.89 | +585532.8... | 3.626 ^D Ly α ; CIV | SB+AGN | —99 ^c | 221 \pm 7 | — | ... | 9.5 \pm 3.5 | 14.1 \pm 4.0 | 0.0 \pm 8.4 | ... | (9.1 $^{+2.6}_{-2.6}$) \times 10 ¹² | 62.1 \pm 7.4 |
| 1HERMES X1.4 J104649.92 | +590039.6... | 4.710 ^L Ly α † | SB | 2.72 ^c | 580 \pm 7 | 3.6 \pm 2.5 | ... | 28.0 \pm 3.6 | 17.1 \pm 4.2 | 7.5 \pm 4.8 | 118.6 \pm 8.8 | (3.9 $^{+0.8}_{-0.7}$) \times 10 ¹³ | 85.6 \pm 6.1 |
| 1HERMES X1.4 J104701.68 | +590447.6... | 4.232 ^L Ly α † | SB | —99 ^c | 1000 \pm 8 | ... | ... | 26.2 \pm 3.5 | 12.0 \pm 4.2 | 4.7 \pm 4.8 | 81.8 \pm 12.2 | (4.1 $^{+0.9}_{-0.7}$) \times 10 ¹³ | 99.9 \pm 7.2 |
| 1HERMES X1.4 J104707.69 | +585149.1... | 4.677 ^D Ly α † | SB | 2.74 ^c | ... | 7.1 \pm 2.7 | ... | 14.5 \pm 3.6 | 31.5 \pm 8.5 | 0.0 \pm 10.7 | 36.0 \pm 6.3 | (2.6 $^{+1.2}_{-0.8}$) \times 10 ¹³ | 80.2 \pm 17.8 |
| 1HERMES X1.4 J104709.60 | +590951.1... | 2.942 ^L Ly α † | SB | —99 ^c | ... | 5.9 \pm 2.6 | 16.9 \pm 4.9 | 13.0 \pm 3.5 | 0.0 \pm 5.2 | 0.0 \pm 4.8 | 99.0 \pm 11.2 | (7.8 $^{+3.8}_{-2.8}$) \times 10 ¹² | 61.0 \pm 21.7 |
| 1HERMES X1.4 J104722.56 | +590111.7... | 4.521 ^D Ly α † | SB+AGN | 2.24 ^c | 444 \pm 7 | 2.8 \pm 2.5 | ... | 19.0 \pm 3.5 | 21.6 \pm 4.1 | 17.7 \pm 5.3 | 73.7 \pm 9.4 | (2.0 $^{+1.0}_{-1.0}$) \times 10 ¹³ | 45.2 \pm 15.6 |
| 1HERMES X1.4 J123536.28 | +623019.9... | 3.380 ^D Diffuse Ly α † | SB | — | — | — | — | 27.1 \pm 4.5 | 25.0 \pm 4.8 | 17.7 \pm 4.1 | 18.0 \pm 51.7 | (1.5 $^{+1.3}_{-0.7}$) \times 10 ¹³ | 47.1 \pm 18.3 |
| 1HERMES X1.4 J123622.58 | +620340.3... | 3.579 ^D Ly α ; CIV† | SB | — | — | — | — | 9.6 \pm 4.5 | 26.9 \pm 5.6 | 16.0 \pm 4.3 | 312.5 \pm 35.3 | (9.7 $^{+16.1}_{-6.1}$) \times 10 ¹² | 34.7 \pm 17.1 |
| 1HERMES X1.4 J123732.66 | +621013.4... | 4.019 ^D Ly α | SB | — | — | — | — | 16.4 \pm 4.5 | 9.7 \pm 4.9 | 1.1 \pm 5.3 | 36.5 \pm 8.4 | (2.9 $^{+0.8}_{-0.6}$) \times 10 ¹³ | 100 \pm 2.0 |
| 1HERMES X24 J160539.72 | +534450.3... | 3.546 ^D Ly α † | SB | —99 ^b | 251 \pm 16 | — | — | 16.2 \pm 4.3 | 16.9 \pm 4.4 | 20.4 \pm 4.6 | — | (1.3 $^{+0.3}_{-0.3}$) \times 10 ¹³ | 45.4 \pm 5.1 |
| 1HERMES X24 J160545.99 | +534544.4... | 2.555 ^D SiIV; CIV; CIII† | QSO | 2.31 ^b | 670 \pm 17 | — | — | 51.5 \pm 4.7 | 41.8 \pm 4.5 | 31.2 \pm 5.2 | — | (1.4 $^{+0.2}_{-0.2}$) \times 10 ¹³ | 39.6 \pm 2.0 |
| 1HERMES X24 J160603.63 | +541245.1... | 3.331 ^D Diffuse Ly α † | SB | —99 ^b | 323 \pm 16 | — | — | 15.5 \pm 4.3 | 16.0 \pm 4.4 | 8.3 \pm 4.9 | — | (1.0 $^{+0.3}_{-0.2}$) \times 10 ¹³ | 60.6 \pm 6.7 |
| 1HERMES X24 J160639.40 | +533558.4... | 3.801 ^D Ly α | SB | —99 ^b | 188 \pm 17 | — | — | 9.0 \pm 4.5 | 16.3 \pm 4.4 | 14.6 \pm 4.8 | — | (1.1 $^{+0.6}_{-0.4}$) \times 10 ¹³ | 47.1 \pm 7.3 |
| 1HERMES X24 J160802.63 | +542638.1... | 3.415 ^D Ly α | SB+AGN | —99 ^b | 442 \pm 18 | — | — | 32.7 \pm 4.3 | 35.9 \pm 4.6 | 37.9 \pm 4.5 | — | (2.2 $^{+0.3}_{-0.2}$) \times 10 ¹³ | 42.5 \pm 2.5 |
| 1HERMES X24 J160806.56 | +542301.6... | 3.721 ^D Ly α ; SiIV; CIV† | SB+AGN | —99 ^b | 246 \pm 16 | — | — | 9.8 \pm 4.3 | 7.4 \pm 4.6 | 16.8 \pm 4.6 | — | (1.3 $^{+0.4}_{-0.2}$) \times 10 ¹³ | 49.8 \pm 7.5 |
| 1HERMES X24 J161506.65 | +543846.9... | 4.952 ^D Ly α † | SB | 1.94 ^b | 720 \pm 174 | — | — | 73.1 \pm 4.3 | 49.6 \pm 4.4 | 22.8 \pm 4.8 | — | (1.2 $^{+0.2}_{-0.2}$) \times 10 ¹⁴ | 86.2 \pm 12.2 |

NOTE. — Superscripts in the z_{spec} column indicate the instrument with which the spectroscopic redshift was obtained, LRIS (L) or DEIMOS (D). *Herschel*-PACS photometry come from PEP (Lutz *et al.* 2011) in COSMOS and from HerMES (Oliver *et al.* 2012) in Lockman Hole North. Photometric redshifts come from: (a) Ilbert *et al.* (2010) in COSMOS; (b) Rowan-Robinson *et al.* (2008) in ELAIS-N1; (c) Strazzullo *et al.* (2010) in Lockman Hole North; and (d) Cardamone *et al.* (2010) in ECDF-S. The galaxy “CLASS” column describes the rest-frame ultraviolet spectrum as Lyman Break Galaxy (LBG), quasar (QSO), starburst with AGN (SB+AGN). Ellipsis in S24, S100, S160, S1.4, or z_{phot} columns denotes that the source is undetected at the corresponding wavelength or has no photometric redshift despite having the necessary optical imaging, whereas — denotes that no data exist. Sources marked with a † in the comments are classified as spectroscopically tentative, and excluded from the ‘high confidence’ SFRD estimate (see section 5).