

1 Perspective

2 **Nitrogen in the environment: inequalities and our changing relationship with nitrogen**

3 Carly J. Stevens

4 Lancaster Environment Centre, Lancaster University, Lancaster, LA1 4YQ. Email:

5 c.stevens@lancaster.ac.uk

6

7 Anthropogenic activities have greatly perturbed the global nitrogen (N) cycle. Planetary boundaries,
8 which describe a safe operating space for humanity, have already been exceeded for the N cycle (1).

9 In some parts of the world the environment has been effected by excess N, with negative impacts on
10 biological diversity, human health and climate. However, in other parts of the world shortages of N
11 mean that food needs cannot be met. Nitrogen is an abundant element on earth, it makes up 78.1%
12 of our atmosphere and is an essential nutrient for all forms of life. Much of this N is in the form of N₂
13 gas and is unreactive and not available for use by the majority of living organisms but a portion of it,
14 fixed by natural or man-made processes including the Haber-Bosch process, is in a reactive form (Nr
15 – including NO_y, NH_x, N₂O, HNO₃ and other organic and inorganic forms) available for use by living
16 organisms. Over the last century the amount of Nr from anthropogenic activities has increased to
17 such an extent that it now exceeds natural fixation and has more than doubled global cycling of N
18 (anthropogenic 210 Tg N yr⁻¹, natural 203 Tg N yr⁻¹) (2). As a consequence of this increase in the
19 fixation of N, N has become a major cause for concern in many parts of the world, polluting air,
20 water and soil (Figure 1).

21 A major cause of N pollution in the developed world is food production. Pollutant Nr, released to the
22 environment during food production and consumption, stems from a range of issues including the
23 over-use of relatively low-cost fertilisers, poor management of animal wastes, over consumption of
24 protein and food waste. Between 1961 and 2007 both N inputs and grain yields increased globally

25 but, the amount of added N recovered in the harvested crops remained relatively unchanged at
26 around 40%. This means that the amount of N lost to the environment has steadily increased.
27 However, there are considerable inequalities in N use globally. Countries outside the OECD and in
28 the major emerging economies the amount of N recovered in crops remains low. Not only are there
29 nutrient deficiencies but the nutrients that are available are often used inefficiently (3). Sub-Saharan
30 Africa provides a perfect example. Here, nutrient poor soils were yielding an average of 1 t ha⁻¹ for
31 grain crops in 2012, with fertiliser use averaging 9 kg ha⁻¹ of cultivated land. By contrast, in Asia, crop
32 yield reached 4.5 t ha⁻¹ but this was achieved with fertiliser application averaging 96 kg ha⁻¹ (4).
33 Shortages of N clearly lead to large problems in meeting population demands for food, but these
34 problems are just as intractable as the problems N pollution causes in other parts of the world.

35 One of the major consequences of increased reactive N availability has been an increase in
36 atmospheric deposition of Nr. Between 1900 and 1980 levels of oxidised N deposition in Europe
37 increased by 3-4 times whilst reduced N doubled (5). The concentrations of N in plant tissues, often
38 been considered an indicator of N status, declined between 1980 and 2017, despite globally
39 increasing availability of Nr. Examining 38,646 terrestrial plant samples collected from areas that had
40 not received any fertiliser input, Craine et al. (6) found global declines in foliar N concentrations. The
41 reduction in N content seems to indicate oligotrophication rather than eutrophication, i.e. there is
42 less N available for the plants than there was in the past. Whilst there have been declines in
43 deposition in some developed countries since 1980 this is not a global trend making this depletion of
44 plant N reserves hard to reconcile with increased Nr emissions. The authors suggest that this is
45 caused by increased levels of CO₂ and longer growing seasons, which allows greater levels of
46 biomass production. The authors also observe a decline in δ¹⁵N, although data are highly variable
47 and changes are small. Since stable isotopes are measured as a ratio of the heavy and light isotopes
48 in a sample, and atmospherically deposited sources of N would typically be light (the so-called
49 'Haber-Bosch effect' (7)), the observed decline in δ¹⁵N may therefore indicate increased atmospheric
50 deposition. Indeed, a recent modelling study has indicated increases in isotopically light N in global

51 oceans (7), confirmed by a study of coral at a remote reef (8). Combined with the global deposition
52 trends, this would suggest that Craine's findings (6) may be even more pressing because the declines
53 in plant N concentration are occurring despite a signal of increasing atmospheric deposition.

54 Craine et al. (6) use the results to question whether humanity has exceeded a true planetary
55 boundary for N availability because plant tissue N is falling. However, the extensive damage done to
56 ecosystems supports the argument that we have indeed exceeded a planetary boundary.

57 Atmospheric deposition of N has become a major driver of plant productivity globally (9) and is an
58 important driver of species richness and composition at a continental scale (10). Considerable
59 damage has already been done and many field experiments with simulated deposition have
60 demonstrated considerable inertia in the recovery of soil chemistry and species composition when
61 levels of N addition are reduced. For example, in an alpine grassland in the Rocky Mountains, USA,
62 12 years of simulated Nr deposition had resulted in significant changes in species composition,
63 including the decline of a previously dominant sedge and increases in other species. The study also
64 found changes in fungal to bacterial ratio, nitrification in the soil, soil pH, toxic metal concentrations
65 and cation concentrations. Nine years after applications of N were ceased many of these soil
66 variables had not returned to baseline levels and nor had biota (11). This type of finding is not
67 uncommon and it is possible, given the lack of recovery observed in some communities, that
68 alternative stable states may have been reached in some habitats.

69 Realisation of the extent of the damage caused by N deposition together with co-benefits from other
70 areas of environmental policy is beginning to result in reductions in emissions and deposition of N.

71 Deposition of oxidised N peaked in Europe in the 1980s and has since declined but there have been
72 much smaller declines in reduced N deposition (5). Similar trends have been observed in the USA
73 with recent reductions in deposition driven by reductions in emissions of oxidised N (12). However, if
74 we are to reduce the creation of Nr further we need wide ranging changes to agricultural practices
75 and our attitude to food. A recent paper highlighted the environmental pressures that the food

76 production system places on the environment and the need to make changes to our diet, combined
77 with technological improvements and reductions in food waste, if we are to stay within planetary
78 boundaries, including the boundary for N. The paper includes scenarios around dietary change
79 towards a healthier plant-based dietary pattern and not exceeding global dietary guidelines (13).
80 Meat consumption is particularly important in terms of driving our N footprint because of the large
81 amounts of Nr lost to the environment during meat production. Interestingly, in some countries, our
82 relationship with N in our diet is already beginning to change, whether we are aware of it or not.
83 Globally meat consumption continues to grow but there is some evidence that in some high-income
84 countries meat consumption per capita is beginning to decline (14).

85 Our N cycle has been hugely perturbed at a global scale and there is an urgent need to address the
86 problem of excess Nr in our environment. There are many potential approaches that can be taken
87 such as technical solutions to agricultural and industrial emissions and changes in practice in
88 polluting sectors, but these need to be widely adopted and supported with legislative limits. There is
89 also a need to address the lack of Nr in many regions of the world to ensure that food production is
90 sufficient to meet requirements. This is a complex problem with many societal considerations and
91 there is considerable debate around the role inorganic fertilisers should play (4). Balancing these two
92 contrasting issues presents a big challenge to the communication of Nr as an environmental problem
93 to the public and is one which can only be addressed through interdisciplinary collaboration
94 between a range of scientists, social scientists, governments and non-governmental organisations.
95 Nr excesses and shortages are set to continue to be major environmental issues into the future so
96 increasing awareness, changing behaviours and increasing regulation, particularly to reduce N
97 emissions, must all come together to address this global problem.

98

99 1. W. Steffen *et al.*, *Science* **347**, 736 (2015).

100 2. D. Fowler *et al.*, *Philosophical Transactions of The Royal Society B* **368**, 1-12 (2013).

- 101 3. R. Conant, A. B. Berdanier, P. R. Grace, *Global Biogeochemical Cycles* **27**, 558-566 (2013).
- 102 4. N. Gilbert, *Nature* **483**, 525-527 (2012).
- 103 5. M. Engardt, D. Simpson, M. Schwikowski, L. Granat, *Tellus B: Chemical and Physical*
104 *Meteorology* **69**, 1328945 (2017).
- 105 6. J. M. Craine *et al.*, *Nature Ecology and Evolution* **2**, 1735-1744 (2018).
- 106 7. S. Yang, N. Gruber, *Global Biogeochemical Cycles* **30**, 1418–1440 (2016).
- 107 8. S. Mii, D. M. Sigman, *Science* **356**, 749-752 (2017).
- 108 9. C. J. Stevens *et al.*, *Ecology* **96**, 1459-1465 (2015).
- 109 10. S. M. Simkin *et al.*, *Proceedings of the National Academy of Sciences of the United States of*
110 *America* **113**, 4086-4091 (2016).
- 111 11. W. D. Bowman *et al.*, *Ecological Applications* **28**, 1762-1772 (2018).
- 112 12. Y. Li *et al.*, *Proceedings of the National Academy of Sciences of the United States of America*
113 **113**, 5874-5879 (2016).
- 114 13. M. Springmann *et al.*, *Nature* **562**, 519-524 (2018).
- 115 14. H. C. J. Godfray *et al.*, *Science* **361**, eaam5324 (2018).

116

117 **Figure 1.** Problems associated with excess and insufficient nitrogen in terrestrial systems.

118 Information on the problems associated with insufficient nitrogen is based on (3).