

# Search for intermediate mass black hole binaries in the first and second observing runs of the Advanced LIGO and Virgo network

B. P. Abbott,<sup>1</sup> R. Abbott,<sup>1</sup> T. D. Abbott,<sup>2</sup> S. Abraham,<sup>3</sup> F. Acernese,<sup>4, 5</sup> K. Ackley,<sup>6</sup> A. Adams,<sup>7</sup> C. Adams,<sup>8</sup> R. X. Adhikari,<sup>1</sup> V. B. Adya,<sup>9</sup> C. Affeldt,<sup>10, 11</sup> M. Agathos,<sup>12, 13</sup> K. Agatsuma,<sup>14</sup> N. Aggarwal,<sup>15</sup> O. D. Aguiar,<sup>16</sup> L. Aiello,<sup>17, 18</sup> A. Ain,<sup>3</sup> P. Ajith,<sup>19</sup> G. Allen,<sup>20</sup> A. Allocca,<sup>21, 22</sup> M. A. Aloy,<sup>23</sup> P. A. Altin,<sup>9</sup> A. Amato,<sup>24</sup> S. Anand,<sup>1</sup> A. Ananyeva,<sup>1</sup> S. B. Anderson,<sup>1</sup> W. G. Anderson,<sup>25</sup> S. V. Angelova,<sup>26</sup> S. Antier,<sup>27</sup> S. Appert,<sup>1</sup> K. Arai,<sup>1</sup> M. C. Araya,<sup>1</sup> J. S. Areeda,<sup>28</sup> M. Arène,<sup>27</sup> N. Arnaud,<sup>29, 30</sup> S. M. Aronson,<sup>31</sup> K. G. Arun,<sup>32</sup> S. Ascenzi,<sup>17, 33</sup> G. Ashton,<sup>6</sup> S. M. Aston,<sup>8</sup> P. Astone,<sup>34</sup> F. Aubin,<sup>35</sup> P. Aufmuth,<sup>11</sup> K. AultONeal,<sup>36</sup> C. Austin,<sup>2</sup> V. Avendano,<sup>37</sup> A. Avila-Alvarez,<sup>28</sup> S. Babak,<sup>27</sup> P. Bacon,<sup>27</sup> F. Badaracco,<sup>17, 18</sup> M. K. M. Bader,<sup>38</sup> S. Bae,<sup>39</sup> A. M. Baer,<sup>7</sup> J. Baird,<sup>27</sup> P. T. Baker,<sup>40</sup> F. Baldaccini,<sup>41, 42</sup> G. Ballardin,<sup>30</sup> S. W. Ballmer,<sup>43</sup> A. Bals,<sup>36</sup> S. Banagiri,<sup>44</sup> J. C. Barayoga,<sup>1</sup> C. Barbieri,<sup>45, 46</sup> S. E. Barclay,<sup>47</sup> B. C. Barish,<sup>1</sup> D. Barker,<sup>48</sup> K. Barkett,<sup>49</sup> S. Barnum,<sup>15</sup> F. Barone,<sup>50, 5</sup> B. Barr,<sup>47</sup> L. Barsotti,<sup>15</sup> M. Barsuglia,<sup>27</sup> D. Barta,<sup>51</sup> J. Bartlett,<sup>48</sup> I. Bartos,<sup>31</sup> R. Bassiri,<sup>52</sup> A. Basti,<sup>21, 22</sup> M. Bawaj,<sup>53, 42</sup> J. C. Bayley,<sup>47</sup> M. Bazzan,<sup>54, 55</sup> B. Bécsy,<sup>56</sup> M. Bejger,<sup>27, 57</sup> I. Belahcene,<sup>29</sup> A. S. Bell,<sup>47</sup> D. Beniwal,<sup>58</sup> M. G. Benjamin,<sup>36</sup> B. K. Berger,<sup>52</sup> G. Bergmann,<sup>10, 11</sup> S. Bernuzzi,<sup>12</sup> C. P. L. Berry,<sup>59</sup> D. Bersanetti,<sup>60</sup> A. Bertolini,<sup>38</sup> J. Betzwieser,<sup>8</sup> R. Bhandare,<sup>61</sup> J. Bidler,<sup>28</sup> E. Biggs,<sup>25</sup> I. A. Bilenko,<sup>62</sup> S. A. Bilgili,<sup>40</sup> G. Billingsley,<sup>1</sup> R. Birney,<sup>26</sup> O. Birnholtz,<sup>63</sup> S. Biscans,<sup>1, 15</sup> M. Bischi,<sup>64, 65</sup> S. Biscoveanu,<sup>15</sup> A. Bisht,<sup>11</sup> M. Bitossi,<sup>30, 22</sup> M. A. Bizouard,<sup>66</sup> J. K. Blackburn,<sup>1</sup> J. Blackman,<sup>49</sup> C. D. Blair,<sup>8</sup> D. G. Blair,<sup>67</sup> R. M. Blair,<sup>48</sup> S. Bloemen,<sup>68</sup> F. Bobba,<sup>69, 70</sup> N. Bode,<sup>10, 11</sup> M. Boer,<sup>66</sup> Y. Boetzel,<sup>71</sup> G. Bogaert,<sup>66</sup> F. Bondu,<sup>72</sup> R. Bonnand,<sup>35</sup> P. Booker,<sup>10, 11</sup> B. A. Boom,<sup>38</sup> R. Bork,<sup>1</sup> V. Boschi,<sup>30</sup> S. Bose,<sup>3</sup> V. Bossilkov,<sup>67</sup> J. Bosveld,<sup>67</sup> Y. Bouffanais,<sup>54, 55</sup> A. Bozzi,<sup>30</sup> C. Bradaschia,<sup>22</sup> P. R. Brady,<sup>25</sup> A. Bramley,<sup>8</sup> M. Branchesi,<sup>17, 18</sup> J. E. Brau,<sup>73</sup> M. Breschi,<sup>12</sup> T. Briant,<sup>74</sup> J. H. Briggs,<sup>47</sup> F. Brighenti,<sup>64, 65</sup> A. Brillet,<sup>66</sup> M. Brinkmann,<sup>10, 11</sup> P. Brockill,<sup>25</sup> A. F. Brooks,<sup>1</sup> J. Brooks,<sup>30</sup> D. D. Brown,<sup>58</sup> S. Brunett,<sup>1</sup> A. Buikema,<sup>15</sup> T. Bulik,<sup>75</sup> H. J. Bulten,<sup>76, 38</sup> A. Buonanno,<sup>77, 78</sup> D. Buskulic,<sup>35</sup> C. Buy,<sup>27</sup> R. L. Byer,<sup>52</sup> M. Cabero,<sup>10, 11</sup> L. Cadonati,<sup>79</sup> G. Cagnoli,<sup>80</sup> C. Cahillane,<sup>1</sup> J. Calderón Bustillo,<sup>6</sup> T. A. Callister,<sup>1</sup> E. Calloni,<sup>81, 5</sup> J. B. Camp,<sup>82</sup> W. A. Campbell,<sup>6</sup> M. Canepa,<sup>83, 60</sup> K. C. Cannon,<sup>84</sup> H. Cao,<sup>58</sup> J. Cao,<sup>85</sup> G. Carapella,<sup>69, 70</sup> F. Carbognani,<sup>30</sup> S. Caride,<sup>86</sup> M. F. Carney,<sup>59</sup> G. Carullo,<sup>21, 22</sup> J. Casanueva Diaz,<sup>22</sup> C. Casentini,<sup>87, 33</sup> S. Caudill,<sup>38</sup> M. Cavaglià,<sup>88, 89</sup> F. Cavalier,<sup>29</sup> R. Cavalieri,<sup>30</sup> G. Celli,<sup>22</sup> P. Cerdá-Durán,<sup>23</sup> E. Cesarini,<sup>90, 33</sup> O. Chaibi,<sup>66</sup> K. Chakravarti,<sup>3</sup> S. J. Chamberlin,<sup>91</sup> M. Chan,<sup>47</sup> S. Chao,<sup>92</sup> P. Charlton,<sup>93</sup> E. A. Chase,<sup>59</sup> E. Chassande-Mottin,<sup>27</sup> D. Chatterjee,<sup>25</sup> M. Chaturvedi,<sup>61</sup> K. Chatzioannou,<sup>94</sup> B. D. Cheeseboro,<sup>40</sup> H. Y. Chen,<sup>95</sup> X. Chen,<sup>67</sup> Y. Chen,<sup>49</sup> H.-P. Cheng,<sup>31</sup> C. K. Cheong,<sup>96</sup> H. Y. Chia,<sup>31</sup> F. Chiadini,<sup>97, 70</sup> A. Chincarini,<sup>60</sup> A. Chiummo,<sup>30</sup> G. Cho,<sup>98</sup> H. S. Cho,<sup>99</sup> M. Cho,<sup>78</sup> N. Christensen,<sup>100, 66</sup> Q. Chu,<sup>67</sup> S. Chua,<sup>74</sup> K. W. Chung,<sup>96</sup> S. Chung,<sup>67</sup> G. Ciani,<sup>54, 55</sup> M. Cieślar,<sup>57</sup> A. A. Ciobanu,<sup>58</sup> R. Ciolfi,<sup>101, 55</sup> F. Cipriano,<sup>66</sup> A. Cirrone,<sup>83, 60</sup> F. Clara,<sup>48</sup> J. A. Clark,<sup>79</sup> P. Clearwater,<sup>102</sup> F. Cleva,<sup>66</sup> E. Coccia,<sup>17, 18</sup> P.-F. Cohadon,<sup>74</sup> D. Cohen,<sup>29</sup> M. Colleoni,<sup>103</sup> C. G. Collette,<sup>104</sup> C. Collins,<sup>14</sup> M. Colpi,<sup>45, 46</sup> L. R. Cominsky,<sup>105</sup> M. Constancio Jr.,<sup>16</sup> L. Conti,<sup>55</sup> S. J. Cooper,<sup>14</sup> P. Corban,<sup>8</sup> T. R. Corbitt,<sup>2</sup> I. Cordero-Carrión,<sup>106</sup> S. Corezzi,<sup>41, 42</sup> K. R. Corley,<sup>107</sup> N. Cornish,<sup>56</sup> D. Corre,<sup>29</sup> A. Corsi,<sup>86</sup> S. Cortese,<sup>30</sup> C. A. Costa,<sup>16</sup> R. Cotesta,<sup>77</sup> M. W. Coughlin,<sup>1</sup> S. B. Coughlin,<sup>108, 59</sup> J.-P. Coulon,<sup>66</sup> S. T. Countryman,<sup>107</sup> P. Couvares,<sup>1</sup> P. B. Covas,<sup>103</sup> E. E. Cowan,<sup>79</sup> D. M. Coward,<sup>67</sup> M. J. Cowart,<sup>8</sup> D. C. Coyne,<sup>1</sup> R. Coyne,<sup>109</sup> J. D. E. Creighton,<sup>25</sup> T. D. Creighton,<sup>110</sup> J. Cripe,<sup>2</sup> M. Croquette,<sup>74</sup> S. G. Crowder,<sup>111</sup> T. J. Cullen,<sup>2</sup> A. Cumming,<sup>47</sup> L. Cunningham,<sup>47</sup> E. Cuoco,<sup>30</sup> T. Dal Canton,<sup>82</sup> G. Dálya,<sup>112</sup> B. D'Angelo,<sup>83, 60</sup> S. L. Danilishin,<sup>10, 11</sup> S. D'Antonio,<sup>33</sup> K. Danzmann,<sup>11, 10</sup> A. Dasgupta,<sup>113</sup> C. F. Da Silva Costa,<sup>31</sup> L. E. H. Datrier,<sup>47</sup> V. Dattilo,<sup>30</sup> I. Dave,<sup>61</sup> M. Davier,<sup>29</sup> D. Davis,<sup>43</sup> E. J. Daw,<sup>114</sup> D. DeBra,<sup>52</sup> M. Deenadayalan,<sup>3</sup> J. Degallaix,<sup>24</sup> M. De Laurentis,<sup>81, 5</sup> S. Deléglise,<sup>74</sup> W. Del Pozzo,<sup>21, 22</sup> L. M. DeMarchi,<sup>59</sup> N. Demos,<sup>15</sup> T. Dent,<sup>115</sup> R. De Pietri,<sup>116, 117</sup> R. De Rosa,<sup>81, 5</sup> C. De Rossi,<sup>24, 30</sup> R. DeSalvo,<sup>118</sup> O. de Varona,<sup>10, 11</sup> S. Dhurandhar,<sup>3</sup> M. C. Díaz,<sup>110</sup> T. Dietrich,<sup>38</sup> L. Di Fiore,<sup>5</sup> C. DiFronzo,<sup>14</sup> C. Di Giorgio,<sup>69, 70</sup> F. Di Giovanni,<sup>23</sup> M. Di Giovanni,<sup>119, 120</sup> T. Di Girolamo,<sup>81, 5</sup> A. Di Lieto,<sup>21, 22</sup> B. Ding,<sup>104</sup> S. Di Pace,<sup>121, 34</sup> I. Di Palma,<sup>121, 34</sup> F. Di Renzo,<sup>21, 22</sup> A. K. Divakarla,<sup>31</sup> A. Dmitriev,<sup>14</sup> Z. Doctor,<sup>95</sup> F. Donovan,<sup>15</sup> K. L. Dooley,<sup>108, 88</sup> S. Doravari,<sup>3</sup> I. Dorrington,<sup>108</sup> T. P. Downes,<sup>25</sup> M. Drago,<sup>17, 18</sup> J. C. Driggers,<sup>48</sup> Z. Du,<sup>85</sup> J.-G. Ducoin,<sup>29</sup> P. Dupej,<sup>47</sup> O. Durante,<sup>69, 70</sup> S. E. Dwyer,<sup>48</sup> P. J. Easter,<sup>6</sup> G. Eddolls,<sup>47</sup> T. B. Edo,<sup>114</sup> A. Effler,<sup>8</sup> P. Ehrens,<sup>1</sup> J. Eichholz,<sup>9</sup> S. S. Eikenberry,<sup>31</sup> M. Eisenmann,<sup>35</sup> R. A. Eisenstein,<sup>15</sup> L. Errico,<sup>81, 5</sup> R. C. Essick,<sup>95</sup> H. Estelles,<sup>103</sup> D. Estevez,<sup>35</sup> Z. B. Etienne,<sup>40</sup> T. Etzel,<sup>1</sup> M. Evans,<sup>15</sup> T. M. Evans,<sup>8</sup> V. Fafone,<sup>87, 33, 17</sup> S. Fairhurst,<sup>108</sup> X. Fan,<sup>85</sup> S. Farinon,<sup>60</sup> B. Farr,<sup>73</sup> W. M. Farr,<sup>14</sup> E. J. Fauchon-Jones,<sup>108</sup> M. Favata,<sup>37</sup> M. Fays,<sup>114</sup> M. Fazio,<sup>122</sup> C. Fee,<sup>123</sup> J. Feicht,<sup>1</sup> M. M. Fejer,<sup>52</sup> F. Feng,<sup>27</sup> D. L. Ferguson,<sup>79</sup> A. Fernandez-Galiana,<sup>15</sup> I. Ferrante,<sup>21, 22</sup> E. C. Ferreira,<sup>16</sup> T. A. Ferreira,<sup>16</sup> F. Fidecaro,<sup>21, 22</sup> I. Fiori,<sup>30</sup> D. Fiorucci,<sup>17, 18</sup> M. Fishbach,<sup>95</sup> R. P. Fisher,<sup>7</sup> J. M. Fishner,<sup>15</sup>

- R. Fittipaldi,<sup>124, 70</sup> M. Fitz-Axen,<sup>44</sup> V. Fiumara,<sup>125, 70</sup> R. Flaminio,<sup>35, 126</sup> M. Fletcher,<sup>47</sup> E. Floden,<sup>44</sup> E. Flynn,<sup>28</sup> H. Fong,<sup>84</sup> J. A. Font,<sup>23, 127</sup> P. W. F. Forsyth,<sup>9</sup> J.-D. Fournier,<sup>66</sup> Francisco Hernandez Vivanco,<sup>6</sup> S. Frasca,<sup>121, 34</sup> F. Frasconi,<sup>22</sup> Z. Frei,<sup>112</sup> A. Freise,<sup>14</sup> R. Frey,<sup>73</sup> V. Frey,<sup>29</sup> P. Fritschel,<sup>15</sup> V. V. Frolov,<sup>8</sup> G. Fronzè,<sup>128</sup> P. Fulda,<sup>31</sup> M. Fyffe,<sup>8</sup> H. A. Gabbard,<sup>47</sup> B. U. Gadre,<sup>77</sup> S. M. Gaebel,<sup>14</sup> J. R. Gair,<sup>129</sup> L. Gammaitoni,<sup>41</sup> S. G. Gaonkar,<sup>3</sup> C. García-Quirós,<sup>103</sup> F. Garufi,<sup>81, 5</sup> B. Gateley,<sup>48</sup> S. Gaudio,<sup>36</sup> G. Gaur,<sup>130</sup> V. Gayathri,<sup>131</sup> G. Gemme,<sup>60</sup> E. Genin,<sup>30</sup> A. Gennai,<sup>22</sup> D. George,<sup>20</sup> J. George,<sup>61</sup> L. Gergely,<sup>132</sup> S. Ghonge,<sup>79</sup> Abhirup Ghosh,<sup>77</sup> Archisman Ghosh,<sup>38</sup> S. Ghosh,<sup>25</sup> B. Giacomazzo,<sup>119, 120</sup> J. A. Giaime,<sup>2, 8</sup> K. D. Giardina,<sup>8</sup> D. R. Gibson,<sup>133</sup> K. Gill,<sup>107</sup> L. Glover,<sup>134</sup> J. Griesmer,<sup>135</sup> P. Godwin,<sup>91</sup> E. Goetz,<sup>48</sup> R. Goetz,<sup>31</sup> B. Goncharov,<sup>6</sup> G. González,<sup>2</sup> J. M. Gonzalez Castro,<sup>21, 22</sup> A. Gopakumar,<sup>136</sup> S. E. Gossan,<sup>1</sup> M. Gosselin,<sup>30, 21, 22</sup> R. Gouaty,<sup>35</sup> B. Grace,<sup>9</sup> A. Grado,<sup>137, 5</sup> M. Granata,<sup>24</sup> A. Grant,<sup>47</sup> S. Gras,<sup>15</sup> P. Grassia,<sup>1</sup> C. Gray,<sup>48</sup> R. Gray,<sup>47</sup> G. Greco,<sup>64, 65</sup> A. C. Green,<sup>31</sup> R. Green,<sup>108</sup> E. M. Gretarsson,<sup>36</sup> A. Grimaldi,<sup>119, 120</sup> S. J. Grimm,<sup>17, 18</sup> P. Groot,<sup>68</sup> H. Grote,<sup>108</sup> S. Grunewald,<sup>77</sup> P. Gruning,<sup>29</sup> G. M. Guidi,<sup>64, 65</sup> H. K. Gulati,<sup>113</sup> Y. Guo,<sup>38</sup> A. Gupta,<sup>91</sup> Anchal Gupta,<sup>1</sup> P. Gupta,<sup>38</sup> E. K. Gustafson,<sup>1</sup> R. Gustafson,<sup>138</sup> L. Haegel,<sup>103</sup> O. Halim,<sup>18, 17</sup> B. R. Hall,<sup>139</sup> E. D. Hall,<sup>15</sup> E. Z. Hamilton,<sup>108</sup> G. Hammond,<sup>47</sup> M. Haney,<sup>71</sup> M. M. Hanke,<sup>10, 11</sup> J. Hanks,<sup>48</sup> C. Hanna,<sup>91</sup> M. D. Hannam,<sup>108</sup> O. A. Hannuksela,<sup>96</sup> T. J. Hansen,<sup>36</sup> J. Hanson,<sup>8</sup> T. Hardwick,<sup>2</sup> K. Haris,<sup>19</sup> J. Harms,<sup>17, 18</sup> G. M. Harry,<sup>140</sup> I. W. Harry,<sup>141</sup> R. K. Hasskew,<sup>8</sup> C. J. Haster,<sup>15</sup> K. Haughian,<sup>47</sup> F. J. Hayes,<sup>47</sup> J. Healy,<sup>63</sup> A. Heidmann,<sup>74</sup> M. C. Heintze,<sup>8</sup> H. Heitmann,<sup>66</sup> F. Hellman,<sup>142</sup> P. Hello,<sup>29</sup> G. Hemming,<sup>30</sup> M. Hendry,<sup>47</sup> I. S. Heng,<sup>47</sup> J. Hennig,<sup>10, 11</sup> M. Heurs,<sup>10, 11</sup> S. Hild,<sup>47</sup> T. Hinderer,<sup>143, 38, 144</sup> S. Hochheim,<sup>10, 11</sup> D. Hofman,<sup>24</sup> A. M. Holgado,<sup>20</sup> N. A. Holland,<sup>9</sup> K. Holt,<sup>8</sup> D. E. Holz,<sup>95</sup> P. Hopkins,<sup>108</sup> C. Horst,<sup>25</sup> J. Hough,<sup>47</sup> E. J. Howell,<sup>67</sup> C. G. Hoy,<sup>108</sup> Y. Huang,<sup>15</sup> M. T. Hübner,<sup>6</sup> E. A. Huerta,<sup>20</sup> D. Huet,<sup>29</sup> B. Hughey,<sup>36</sup> V. Hui,<sup>35</sup> S. Husa,<sup>103</sup> S. H. Huttner,<sup>47</sup> T. Huynh-Dinh,<sup>8</sup> B. Idzkowski,<sup>75</sup> A. Iess,<sup>87, 33</sup> H. Inchauspe,<sup>31</sup> C. Ingram,<sup>58</sup> R. Inta,<sup>86</sup> G. Intimi,<sup>121, 34</sup> B. Irwin,<sup>123</sup> H. N. Isa,<sup>47</sup> J.-M. Isac,<sup>74</sup> M. Isi,<sup>15</sup> B. R. Iyer,<sup>19</sup> T. Jacqmin,<sup>74</sup> S. J. Jadhav,<sup>145</sup> K. Jani,<sup>79</sup> N. N. Janthalur,<sup>145</sup> P. Jaradowski,<sup>146</sup> D. Jariwala,<sup>31</sup> A. C. Jenkins,<sup>147</sup> J. Jiang,<sup>31</sup> G. R. Johns,<sup>7</sup> D. S. Johnson,<sup>20</sup> A. W. Jones,<sup>14</sup> D. I. Jones,<sup>148</sup> J. D. Jones,<sup>48</sup> R. Jones,<sup>47</sup> R. J. G. Jonker,<sup>38</sup> L. Ju,<sup>67</sup> J. Junker,<sup>10, 11</sup> C. V. Kalaghatgi,<sup>108</sup> V. Kalogera,<sup>59</sup> B. Kamai,<sup>1</sup> S. Kandhasamy,<sup>3</sup> G. Kang,<sup>39</sup> J. B. Kanner,<sup>1</sup> S. J. Kapadia,<sup>25</sup> S. Karki,<sup>73</sup> R. Kashyap,<sup>19</sup> M. Kasprzack,<sup>1</sup> S. Katsanevas,<sup>30</sup> E. Katsavounidis,<sup>15</sup> W. Katzman,<sup>8</sup> S. Kaufer,<sup>11</sup> K. Kawabe,<sup>48</sup> N. V. Keerthana,<sup>3</sup> F. Kéfélian,<sup>66</sup> D. Keitel,<sup>141</sup> R. Kennedy,<sup>114</sup> J. S. Key,<sup>149</sup> F. Y. Khalili,<sup>62</sup> B. Khamesra,<sup>79</sup> I. Khan,<sup>17, 33</sup> S. Khan,<sup>10, 11</sup> E. A. Khazanov,<sup>150</sup> N. Khetan,<sup>17, 18</sup> M. Khursheed,<sup>61</sup> N. Kijbunchoo,<sup>9</sup> Chungle Kim,<sup>151</sup> G. J. Kim,<sup>79</sup> J. C. Kim,<sup>152</sup> K. Kim,<sup>96</sup> W. Kim,<sup>58</sup> W. S. Kim,<sup>153</sup> Y.-M. Kim,<sup>154</sup> C. Kimball,<sup>59</sup> P. J. King,<sup>48</sup> M. Kinley-Hanlon,<sup>47</sup> R. Kirchhoff,<sup>10, 11</sup> J. S. Kissel,<sup>48</sup> L. Kleybolte,<sup>135</sup> J. H. Klika,<sup>25</sup> S. Klimentko,<sup>31</sup> T. D. Knowles,<sup>40</sup> P. Koch,<sup>10, 11</sup> S. M. Koehlenbeck,<sup>10, 11</sup> G. Koekoek,<sup>38, 155</sup> S. Koley,<sup>38</sup> V. Kondrashov,<sup>1</sup> A. Kontos,<sup>156</sup> N. Koper,<sup>10, 11</sup> M. Korobko,<sup>135</sup> W. Z. Korth,<sup>1</sup> M. Kovalam,<sup>67</sup> D. B. Kozak,<sup>1</sup> C. Krämer,<sup>10, 11</sup> V. Krингель,<sup>10, 11</sup> N. Krishnendu,<sup>32</sup> A. Królak,<sup>157, 158</sup> N. Krupinski,<sup>25</sup> G. Kuehn,<sup>10, 11</sup> A. Kumar,<sup>145</sup> P. Kumar,<sup>159</sup> Rahul Kumar,<sup>48</sup> Rakesh Kumar,<sup>113</sup> L. Kuo,<sup>92</sup> A. Kutynia,<sup>157</sup> S. Kwang,<sup>25</sup> B. D. Lackey,<sup>77</sup> D. Laghi,<sup>21, 22</sup> P. Laguna,<sup>79</sup> K. H. Lai,<sup>96</sup> T. L. Lam,<sup>96</sup> M. Landry,<sup>48</sup> B. B. Lane,<sup>15</sup> R. N. Lang,<sup>160</sup> J. Lange,<sup>63</sup> B. Lantz,<sup>52</sup> R. K. Lanza,<sup>15</sup> A. Lartaux-Vollard,<sup>29</sup> P. D. Lasky,<sup>6</sup> M. Laxen,<sup>8</sup> A. Lazzarini,<sup>1</sup> C. Lazzaro,<sup>55</sup> P. Leaci,<sup>121, 34</sup> S. Leavey,<sup>10, 11</sup> Y. K. Lecoeuche,<sup>48</sup> C. H. Lee,<sup>99</sup> H. K. Lee,<sup>161</sup> H. M. Lee,<sup>162</sup> H. W. Lee,<sup>152</sup> J. Lee,<sup>98</sup> K. Lee,<sup>47</sup> J. Lehmann,<sup>10, 11</sup> A. K. Lenon,<sup>40</sup> N. Leroy,<sup>29</sup> N. Letendre,<sup>35</sup> Y. Levin,<sup>6</sup> A. Li,<sup>96</sup> J. Li,<sup>85</sup> K. J. L. Li,<sup>96</sup> T. G. F. Li,<sup>96</sup> X. Li,<sup>49</sup> F. Lin,<sup>6</sup> F. Linde,<sup>163, 38</sup> S. D. Linker,<sup>134</sup> T. B. Littenberg,<sup>164</sup> J. Liu,<sup>67</sup> X. Liu,<sup>25</sup> M. Llorens-Monteagudo,<sup>23</sup> R. K. L. Lo,<sup>96, 1</sup> L. T. London,<sup>15</sup> A. Longo,<sup>165, 166</sup> M. Lorenzini,<sup>17, 18</sup> V. Loriette,<sup>167</sup> M. Lormand,<sup>8</sup> G. Losurdo,<sup>22</sup> J. D. Lough,<sup>10, 11</sup> C. O. Lousto,<sup>63</sup> G. Lovelace,<sup>28</sup> M. E. Lower,<sup>168</sup> H. Lück,<sup>11, 10</sup> D. Lumaca,<sup>87, 33</sup> A. P. Lundgren,<sup>141</sup> R. Lynch,<sup>15</sup> Y. Ma,<sup>49</sup> R. Macas,<sup>108</sup> S. Macfoy,<sup>26</sup> M. MacInnis,<sup>15</sup> D. M. Macleod,<sup>108</sup> A. Macquet,<sup>66</sup> I. Magaña Hernandez,<sup>25</sup> F. Magaña-Sandoval,<sup>31</sup> R. M. Magee,<sup>91</sup> E. Majorana,<sup>34</sup> I. Maksimovic,<sup>167</sup> A. Malik,<sup>61</sup> N. Man,<sup>66</sup> V. Mandic,<sup>44</sup> V. Mangano,<sup>47, 121, 34</sup> G. L. Mansell,<sup>48, 15</sup> M. Manske,<sup>25</sup> M. Mantovani,<sup>30</sup> M. Mapelli,<sup>54, 55</sup> F. Marchesoni,<sup>53, 42</sup> F. Marion,<sup>35</sup> S. Márka,<sup>107</sup> Z. Márka,<sup>107</sup> C. Markakis,<sup>20</sup> A. S. Markosyan,<sup>52</sup> A. Markowitz,<sup>1</sup> E. Maros,<sup>1</sup> A. Marquina,<sup>106</sup> S. Marsat,<sup>27</sup> F. Martelli,<sup>64, 65</sup> I. W. Martin,<sup>47</sup> R. M. Martin,<sup>37</sup> V. Martinez,<sup>80</sup> D. V. Martynov,<sup>14</sup> H. Masalehdan,<sup>135</sup> K. Mason,<sup>15</sup> E. Massera,<sup>114</sup> A. Masserot,<sup>35</sup> T. J. Massinger,<sup>1</sup> M. Masso-Reid,<sup>47</sup> S. Mastrogiovanni,<sup>27</sup> A. Matas,<sup>77</sup> F. Matichard,<sup>1, 15</sup> L. Matone,<sup>107</sup> N. Mavalvala,<sup>15</sup> J. J. McCann,<sup>67</sup> R. McCarthy,<sup>48</sup> D. E. McClelland,<sup>9</sup> S. McCormick,<sup>8</sup> L. McCuller,<sup>15</sup> S. C. McGuire,<sup>169</sup> C. McIsaac,<sup>141</sup> J. McIver,<sup>1</sup> D. J. McManus,<sup>9</sup> T. McRae,<sup>9</sup> S. T. McWilliams,<sup>40</sup> D. Meacher,<sup>25</sup> G. D. Meadors,<sup>6</sup> M. Mehmet,<sup>10, 11</sup> A. K. Mehta,<sup>19</sup> J. Meidam,<sup>38</sup> E. Mejuto Villa,<sup>118, 70</sup> A. Melatos,<sup>102</sup> G. Mendell,<sup>48</sup> R. A. Mercer,<sup>25</sup> L. Mereni,<sup>24</sup> K. Merfeld,<sup>73</sup> E. L. Merilh,<sup>48</sup> M. Merzougui,<sup>66</sup> S. Meshkov,<sup>1</sup> C. Messenger,<sup>47</sup> C. Messick,<sup>91</sup> F. Messina,<sup>45, 46</sup> R. Metzdorff,<sup>74</sup> P. M. Meyers,<sup>102</sup> F. Meylahn,<sup>10, 11</sup> A. Miani,<sup>119, 120</sup> H. Miao,<sup>14</sup> C. Michel,<sup>24</sup> H. Middleton,<sup>102</sup> L. Milano,<sup>81, 5</sup> A. L. Miller,<sup>31, 121, 34</sup> M. Millhouse,<sup>102</sup>

- J. C. Mills,<sup>108</sup> M. C. Milovich-Goff,<sup>134</sup> O. Minazzoli,<sup>66, 170</sup> Y. Minenkov,<sup>33</sup> A. Mishkin,<sup>31</sup> C. Mishra,<sup>171</sup> T. Mistry,<sup>114</sup> S. Mitra,<sup>3</sup> V. P. Mitrofanov,<sup>62</sup> G. Mitselmakher,<sup>31</sup> R. Mittleman,<sup>15</sup> G. Mo,<sup>100</sup> D. Moffa,<sup>123</sup> K. Mogushi,<sup>88</sup> S. R. P. Mohapatra,<sup>15</sup> M. Molina-Ruiz,<sup>142</sup> M. Mondin,<sup>134</sup> M. Montani,<sup>64, 65</sup> C. J. Moore,<sup>14</sup> D. Moraru,<sup>48</sup> F. Morawski,<sup>57</sup> G. Moreno,<sup>48</sup> S. Morisaki,<sup>84</sup> B. Mouris,<sup>35</sup> C. M. Mow-Lowry,<sup>14</sup> F. Muciaccia,<sup>121, 34</sup> Arunava Mukherjee,<sup>10, 11</sup> D. Mukherjee,<sup>25</sup> S. Mukherjee,<sup>110</sup> Subroto Mukherjee,<sup>113</sup> N. Mukund,<sup>10, 11, 3</sup> A. Mullavey,<sup>8</sup> J. Munch,<sup>58</sup> E. A. Muñiz,<sup>43</sup> M. Muratore,<sup>36</sup> P. G. Murray,<sup>47</sup> A. Nagar,<sup>90, 128, 172</sup> I. Nardecchia,<sup>87, 33</sup> L. Naticchioni,<sup>121, 34</sup> R. K. Nayak,<sup>173</sup> B. F. Neil,<sup>67</sup> J. Neilson,<sup>118, 70</sup> G. Nelemans,<sup>68, 38</sup> T. J. N. Nelson,<sup>8</sup> M. Nery,<sup>10, 11</sup> A. Neunzert,<sup>138</sup> L. Nevin,<sup>1</sup> K. Y. Ng,<sup>15</sup> S. Ng,<sup>58</sup> C. Nguyen,<sup>27</sup> P. Nguyen,<sup>73</sup> D. Nichols,<sup>143, 38</sup> S. A. Nichols,<sup>2</sup> S. Nissanke,<sup>143, 38</sup> F. Nocera,<sup>30</sup> C. North,<sup>108</sup> L. K. Nuttall,<sup>141</sup> M. Obergaulinger,<sup>23, 174</sup> J. Oberling,<sup>48</sup> B. D. O'Brien,<sup>31</sup> G. Oganesyan,<sup>17, 18</sup> G. H. Ogin,<sup>175</sup> J. J. Oh,<sup>153</sup> S. H. Oh,<sup>153</sup> F. Ohme,<sup>10, 11</sup> H. Ohta,<sup>84</sup> M. A. Okada,<sup>16</sup> M. Oliver,<sup>103</sup> P. Oppermann,<sup>10, 11</sup> Richard J. Oram,<sup>8</sup> B. O'Reilly,<sup>8</sup> R. G. Ormiston,<sup>44</sup> L. F. Ortega,<sup>31</sup> R. O'Shaughnessy,<sup>63</sup> S. Ossokine,<sup>77</sup> D. J. Ottaway,<sup>58</sup> H. Overmier,<sup>8</sup> B. J. Owen,<sup>86</sup> A. E. Pace,<sup>91</sup> G. Pagano,<sup>21, 22</sup> M. A. Page,<sup>67</sup> G. Pagliaroli,<sup>17, 18</sup> A. Pai,<sup>131</sup> S. A. Pai,<sup>61</sup> J. R. Palamos,<sup>73</sup> O. Palashov,<sup>150</sup> C. Palomba,<sup>34</sup> H. Pan,<sup>92</sup> P. K. Panda,<sup>145</sup> P. T. H. Pang,<sup>96, 38</sup> C. Pankow,<sup>59</sup> F. Pannarale,<sup>121, 34</sup> B. C. Pant,<sup>61</sup> F. Paoletti,<sup>22</sup> A. Paoli,<sup>30</sup> A. Parida,<sup>3</sup> W. Parker,<sup>8, 169</sup> D. Pascucci,<sup>47, 38</sup> A. Pasqualetti,<sup>30</sup> R. Passaquieti,<sup>21, 22</sup> D. Passuello,<sup>22</sup> M. Patil,<sup>158</sup> B. Patricelli,<sup>21, 22</sup> E. Payne,<sup>6</sup> B. L. Pearlstone,<sup>47</sup> T. C. Pechsiri,<sup>31</sup> A. J. Pedersen,<sup>43</sup> M. Pedraza,<sup>1</sup> R. Pedurand,<sup>24, 176</sup> A. Pele,<sup>8</sup> S. Penn,<sup>177</sup> A. Perego,<sup>119, 120</sup> C. J. Perez,<sup>48</sup> C. Périgois,<sup>35</sup> A. Perreca,<sup>119, 120</sup> J. Petermann,<sup>135</sup> H. P. Pfeiffer,<sup>77</sup> M. Phelps,<sup>10, 11</sup> K. S. Phukon,<sup>3</sup> O. J. Piccinni,<sup>121, 34</sup> M. Pichot,<sup>66</sup> F. Piergiovanni,<sup>64, 65</sup> V. Pierro,<sup>118, 70</sup> G. Pillant,<sup>30</sup> L. Pinard,<sup>24</sup> I. M. Pinto,<sup>118, 70, 90</sup> M. Pirello,<sup>48</sup> M. Pitkin,<sup>47</sup> W. Plastino,<sup>165, 166</sup> R. Poggiani,<sup>21, 22</sup> D. Y. T. Pong,<sup>96</sup> S. Ponrathnam,<sup>3</sup> P. Popolizio,<sup>30</sup> E. K. Porter,<sup>27</sup> J. Powell,<sup>168</sup> A. K. Prajapati,<sup>113</sup> J. Prasad,<sup>3</sup> K. Prasai,<sup>52</sup> R. Prasanna,<sup>145</sup> G. Pratten,<sup>103</sup> T. Prestegard,<sup>25</sup> M. Principe,<sup>118, 90, 70</sup> G. A. Prodi,<sup>119, 120</sup> L. Prokhorov,<sup>14</sup> M. Punturo,<sup>42</sup> P. Puppo,<sup>34</sup> M. Pürer,<sup>77</sup> H. Qi,<sup>108</sup> V. Quetschke,<sup>110</sup> P. J. Quinonez,<sup>36</sup> F. J. Raab,<sup>48</sup> G. Raaijmakers,<sup>143, 38</sup> H. Radkins,<sup>48</sup> N. Radulesco,<sup>66</sup> P. Raffai,<sup>112</sup> S. Raja,<sup>61</sup> C. Rajan,<sup>61</sup> B. Rajbhandari,<sup>86</sup> M. Rakhamanov,<sup>110</sup> K. E. Ramirez,<sup>110</sup> A. Ramos-Buades,<sup>103</sup> Javed Rana,<sup>3</sup> K. Rao,<sup>59</sup> P. Rapagnani,<sup>121, 34</sup> V. Raymond,<sup>108</sup> M. Razzano,<sup>21, 22</sup> J. Read,<sup>28</sup> T. Regimbau,<sup>35</sup> L. Rei,<sup>60</sup> S. Reid,<sup>26</sup> D. H. Reitze,<sup>1, 31</sup> P. Rettegno,<sup>128, 178</sup> F. Ricci,<sup>121, 34</sup> C. J. Richardson,<sup>36</sup> J. W. Richardson,<sup>1</sup> P. M. Ricker,<sup>20</sup> G. Riemenschneider,<sup>178, 128</sup> K. Riles,<sup>138</sup> M. Rizzo,<sup>59</sup> N. A. Robertson,<sup>1, 47</sup> F. Robinet,<sup>29</sup> A. Rocchi,<sup>33</sup> L. Rolland,<sup>35</sup> J. G. Rollins,<sup>1</sup> V. J. Roma,<sup>73</sup> M. Romanelli,<sup>72</sup> R. Romano,<sup>4, 5</sup> C. L. Romel,<sup>48</sup> J. H. Romie,<sup>8</sup> C. A. Rose,<sup>25</sup> D. Rose,<sup>28</sup> K. Rose,<sup>123</sup> D. Rosińska,<sup>75</sup> S. G. Rosofsky,<sup>20</sup> M. P. Ross,<sup>179</sup> S. Rowan,<sup>47</sup> A. Rüdiger,<sup>10, 11, \*</sup> P. Ruggi,<sup>30</sup> G. Rutins,<sup>133</sup> K. Ryan,<sup>48</sup> S. Sachdev,<sup>91</sup> T. Sadecki,<sup>48</sup> M. Sakellariadou,<sup>147</sup> O. S. Salafia,<sup>180, 45, 46</sup> L. Salconi,<sup>30</sup> M. Saleem,<sup>32</sup> A. Samajdar,<sup>38</sup> L. Sammut,<sup>6</sup> E. J. Sanchez,<sup>1</sup> L. E. Sanchez,<sup>1</sup> N. Sanchis-Gual,<sup>181</sup> J. R. Sanders,<sup>182</sup> K. A. Santiago,<sup>37</sup> E. Santos,<sup>66</sup> N. Sarin,<sup>6</sup> B. Sassolas,<sup>24</sup> B. S. Sathyaprakash,<sup>91, 108</sup> O. Sauter,<sup>138, 35</sup> R. L. Savage,<sup>48</sup> P. Schale,<sup>73</sup> M. Scheel,<sup>49</sup> J. Scheuer,<sup>59</sup> P. Schmidt,<sup>14, 68</sup> R. Schnabel,<sup>135</sup> R. M. S. Schofield,<sup>73</sup> A. Schönbeck,<sup>135</sup> E. Schreiber,<sup>10, 11</sup> B. W. Schulte,<sup>10, 11</sup> B. F. Schutz,<sup>108</sup> J. Scott,<sup>47</sup> S. M. Scott,<sup>9</sup> E. Seidel,<sup>20</sup> D. Sellers,<sup>8</sup> A. S. Sengupta,<sup>183</sup> N. Sennett,<sup>77</sup> D. Sentenac,<sup>30</sup> V. Sequino,<sup>60</sup> A. Sergeev,<sup>150</sup> Y. Setyawati,<sup>10, 11</sup> D. A. Shaddock,<sup>9</sup> T. Shaffer,<sup>48</sup> M. S. Shahriar,<sup>59</sup> M. B. Shaner,<sup>134</sup> A. Sharma,<sup>17, 18</sup> P. Sharma,<sup>61</sup> P. Shawhan,<sup>78</sup> H. Shen,<sup>20</sup> R. Shink,<sup>184</sup> D. H. Shoemaker,<sup>15</sup> D. M. Shoemaker,<sup>79</sup> K. Shukla,<sup>142</sup> S. ShyamSundar,<sup>61</sup> K. Siellez,<sup>79</sup> M. Sieniawska,<sup>57</sup> D. Sigg,<sup>48</sup> L. P. Singer,<sup>82</sup> D. Singh,<sup>91</sup> N. Singh,<sup>75</sup> A. Singhal,<sup>17, 34</sup> A. M. Sintes,<sup>103</sup> S. Sitmukhambetov,<sup>110</sup> V. Skliris,<sup>108</sup> B. J. J. Slagmolen,<sup>9</sup> T. J. Slaven-Blair,<sup>67</sup> J. R. Smith,<sup>28</sup> R. J. E. Smith,<sup>6</sup> S. Somala,<sup>185</sup> E. J. Son,<sup>153</sup> S. Soni,<sup>2</sup> B. Sorazu,<sup>47</sup> F. Sorrentino,<sup>60</sup> T. Souradeep,<sup>3</sup> E. Sowell,<sup>86</sup> A. P. Spencer,<sup>47</sup> M. Spera,<sup>54, 55</sup> A. K. Srivastava,<sup>113</sup> V. Srivastava,<sup>43</sup> K. Staats,<sup>59</sup> C. Stachie,<sup>66</sup> M. Standke,<sup>10, 11</sup> D. A. Steer,<sup>27</sup> M. Steinke,<sup>10, 11</sup> J. Steinlechner,<sup>135, 47</sup> S. Steinlechner,<sup>135</sup> D. Steinmeyer,<sup>10, 11</sup> S. P. Stevenson,<sup>168</sup> D. Stocks,<sup>52</sup> R. Stone,<sup>110</sup> D. J. Stops,<sup>14</sup> K. A. Strain,<sup>47</sup> G. Stratta,<sup>186, 65</sup> S. E. Strigin,<sup>62</sup> A. Strunk,<sup>48</sup> R. Sturani,<sup>187</sup> A. L. Stuver,<sup>188</sup> V. Sudhir,<sup>15</sup> T. Z. Summerscales,<sup>189</sup> L. Sun,<sup>1</sup> S. Sunil,<sup>113</sup> A. Sur,<sup>57</sup> J. Suresh,<sup>84</sup> P. J. Sutton,<sup>108</sup> B. L. Swinkels,<sup>38</sup> M. J. Szczepańczyk,<sup>36</sup> M. Tacca,<sup>38</sup> S. C. Tait,<sup>47</sup> C. Talbot,<sup>6</sup> D. B. Tanner,<sup>31</sup> D. Tao,<sup>1</sup> M. Tápai,<sup>132</sup> A. Tapia,<sup>28</sup> J. D. Tasson,<sup>100</sup> R. Taylor,<sup>1</sup> R. Tenorio,<sup>103</sup> L. Terkowski,<sup>135</sup> M. Thomas,<sup>8</sup> P. Thomas,<sup>48</sup> S. R. Thondapu,<sup>61</sup> K. A. Thorne,<sup>8</sup> E. Thrane,<sup>6</sup> Shubhanshu Tiwari,<sup>119, 120</sup> Srishti Tiwari,<sup>136</sup> V. Tiwari,<sup>108</sup> K. Toland,<sup>47</sup> M. Tonelli,<sup>21, 22</sup> Z. Tornasi,<sup>47</sup> A. Torres-Forné,<sup>190</sup> C. I. Torrie,<sup>1</sup> D. Töyrä,<sup>14</sup> F. Travasso,<sup>30, 42</sup> G. Traylor,<sup>8</sup> M. C. Tringali,<sup>75</sup> A. Tripathee,<sup>138</sup> A. Trovato,<sup>27</sup> L. Trozzo,<sup>191, 22</sup> K. W. Tsang,<sup>38</sup> M. Tse,<sup>15</sup> R. Tso,<sup>49</sup> L. Tsukada,<sup>84</sup> D. Tsuna,<sup>84</sup> T. Tsutsui,<sup>84</sup> D. Tuyenbayev,<sup>110</sup> K. Ueno,<sup>84</sup> D. Ugolini,<sup>192</sup> C. S. Unnikrishnan,<sup>136</sup> A. L. Urban,<sup>2</sup> S. A. Usman,<sup>95</sup> H. Vahlbruch,<sup>11</sup> G. Vajente,<sup>1</sup> G. Valdes,<sup>2</sup> M. Valentini,<sup>119, 120</sup> N. van Bakel,<sup>38</sup> M. van Beuzekom,<sup>38</sup> J. F. J. van den Brand,<sup>76, 38</sup> C. Van Den Broeck,<sup>38, 193</sup> D. C. Vander-Hyde,<sup>43</sup> L. van der Schaaf,<sup>38</sup> J. V. VanHeijningen,<sup>67</sup> A. A. van Veggel,<sup>47</sup> M. Vardaro,<sup>54, 55</sup> V. Varma,<sup>49</sup> S. Vass,<sup>1</sup> M. Vasúth,<sup>51</sup> A. Vecchio,<sup>14</sup> G. Vedovato,<sup>55</sup> J. Veitch,<sup>47</sup> P. J. Veitch,<sup>58</sup> K. Venkateswara,<sup>179</sup>

G. Venugopalan,<sup>1</sup> D. Verkindt,<sup>35</sup> F. Vetrano,<sup>64, 65</sup> A. Viceré,<sup>64, 65</sup> A. D. Viets,<sup>25</sup> S. Vinciguerra,<sup>14</sup> D. J. Vine,<sup>133</sup> J.-Y. Vinet,<sup>66</sup> S. Vitale,<sup>15</sup> T. Vo,<sup>43</sup> H. Vocca,<sup>41, 42</sup> C. Vorwick,<sup>48</sup> S. P. Vyatchanin,<sup>62</sup> A. R. Wade,<sup>1</sup> L. E. Wade,<sup>123</sup> M. Wade,<sup>123</sup> R. Walet,<sup>38</sup> M. Walker,<sup>28</sup> L. Wallace,<sup>1</sup> S. Walsh,<sup>25</sup> H. Wang,<sup>14</sup> J. Z. Wang,<sup>138</sup> S. Wang,<sup>20</sup> W. H. Wang,<sup>110</sup> Y. F. Wang,<sup>96</sup> R. L. Ward,<sup>9</sup> Z. A. Warden,<sup>36</sup> J. Warner,<sup>48</sup> M. Was,<sup>35</sup> J. Watchi,<sup>104</sup> B. Weaver,<sup>48</sup> L.-W. Wei,<sup>10, 11</sup> M. Weinert,<sup>10, 11</sup> A. J. Weinstein,<sup>1</sup> R. Weiss,<sup>15</sup> F. Wellmann,<sup>10, 11</sup> L. Wen,<sup>67</sup> E. K. Wessel,<sup>20</sup> P. Webels,<sup>10, 11</sup> J. W. Westhouse,<sup>36</sup> K. Wette,<sup>9</sup> J. T. Whelan,<sup>63</sup> B. F. Whiting,<sup>31</sup> C. Whittle,<sup>15</sup> D. M. Wilken,<sup>10, 11</sup> D. Williams,<sup>47</sup> A. R. Williamson,<sup>143, 38</sup> J. L. Willis,<sup>1</sup> B. Willke,<sup>11, 10</sup> W. Winkler,<sup>10, 11</sup> C. C. Wipf,<sup>1</sup> H. Wittel,<sup>10, 11</sup> G. Woan,<sup>47</sup> J. Woehler,<sup>10, 11</sup> J. K. Wofford,<sup>63</sup> J. L. Wright,<sup>47</sup> D. S. Wu,<sup>10, 11</sup> D. M. Wysocki,<sup>63</sup> S. Xiao,<sup>1</sup> R. Xu,<sup>111</sup> H. Yamamoto,<sup>1</sup> C. C. Yancey,<sup>78</sup> L. Yang,<sup>122</sup> Y. Yang,<sup>31</sup> Z. Yang,<sup>44</sup> M. J. Yap,<sup>9</sup> M. Yazback,<sup>31</sup> D. W. Yeeles,<sup>108</sup> A. Yoon,<sup>7</sup> Hang Yu,<sup>15</sup> Haocun Yu,<sup>15</sup> S. H. R. Yuen,<sup>96</sup> A. K. Zadrożny,<sup>110</sup> A. Zadrożny,<sup>157</sup> M. Zanolini,<sup>36</sup> T. Zelenova,<sup>30</sup> J.-P. Zendri,<sup>55</sup> M. Zevin,<sup>59</sup> J. Zhang,<sup>67</sup> L. Zhang,<sup>1</sup> T. Zhang,<sup>47</sup> C. Zhao,<sup>67</sup> G. Zhao,<sup>104</sup> M. Zhou,<sup>59</sup> Z. Zhou,<sup>59</sup> X. J. Zhu,<sup>6</sup> A. B. Zimmerman,<sup>194</sup> M. E. Zucker,<sup>1, 15</sup> J. Zweizig,<sup>1</sup> F. Salemi,<sup>195</sup> and M. A. Papa<sup>196, 197, 195</sup>

(The LIGO Scientific Collaboration and the Virgo Collaboration)

<sup>1</sup>LIGO, California Institute of Technology, Pasadena, CA 91125, USA

<sup>2</sup>Louisiana State University, Baton Rouge, LA 70803, USA

<sup>3</sup>Inter-University Centre for Astronomy and Astrophysics, Pune 411007, India

<sup>4</sup>Dipartimento di Farmacia, Università di Salerno, I-84084 Fisciano, Salerno, Italy

<sup>5</sup>INFN, Sezione di Napoli, Complesso Universitario di Monte S.Angelo, I-80126 Napoli, Italy

<sup>6</sup>OzGrav, School of Physics & Astronomy, Monash University, Clayton 3800, Victoria, Australia

<sup>7</sup>Christopher Newport University, Newport News, VA 23606, USA

<sup>8</sup>LIGO Livingston Observatory, Livingston, LA 70754, USA

<sup>9</sup>OzGrav, Australian National University, Canberra, Australian Capital Territory 0200, Australia

<sup>10</sup>Max Planck Institute for Gravitational Physics (Albert Einstein Institute), D-30167 Hannover, Germany

<sup>11</sup>Leibniz Universität Hannover, D-30167 Hannover, Germany

<sup>12</sup>Theoretisch-Physikalisches Institut, Friedrich-Schiller-Universität Jena, D-07743 Jena, Germany

<sup>13</sup>University of Cambridge, Cambridge CB2 1TN, United Kingdom

<sup>14</sup>University of Birmingham, Birmingham B15 2TT, United Kingdom

<sup>15</sup>LIGO, Massachusetts Institute of Technology, Cambridge, MA 02139, USA

<sup>16</sup>Instituto Nacional de Pesquisas Espaciais, 12227-010 São José dos Campos, São Paulo, Brazil

<sup>17</sup>Gran Sasso Science Institute (GSSI), I-67100 L'Aquila, Italy

<sup>18</sup>INFN, Laboratori Nazionali del Gran Sasso, I-67100 Assergi, Italy

<sup>19</sup>International Centre for Theoretical Sciences, Tata Institute of Fundamental Research, Bengaluru 560089, India

<sup>20</sup>NCSA, University of Illinois at Urbana-Champaign, Urbana, IL 61801, USA

<sup>21</sup>Università di Pisa, I-56127 Pisa, Italy

<sup>22</sup>INFN, Sezione di Pisa, I-56127 Pisa, Italy

<sup>23</sup>Departamento de Astronomía y Astrofísica, Universitat de València, E-46100 Burjassot, València, Spain

<sup>24</sup>Laboratoire des Matériaux Avancés (LMA), CNRS/IN2P3, F-69622 Villeurbanne, France

<sup>25</sup>University of Wisconsin-Milwaukee, Milwaukee, WI 53201, USA

<sup>26</sup>SUPA, University of Strathclyde, Glasgow G1 1XQ, United Kingdom

<sup>27</sup>APC, Astroparticule et Cosmologie, Université Paris Diderot,

CNRS/IN2P3, CEA/Irfu, Observatoire de Paris,

Sorbonne Paris Cité, F-75205 Paris Cedex 13, France

<sup>28</sup>California State University Fullerton, Fullerton, CA 92831, USA

<sup>29</sup>LAL, Univ. Paris-Sud, CNRS/IN2P3, Université Paris-Saclay, F-91898 Orsay, France

<sup>30</sup>European Gravitational Observatory (EGO), I-56021 Cascina, Pisa, Italy

<sup>31</sup>University of Florida, Gainesville, FL 32611, USA

<sup>32</sup>Chennai Mathematical Institute, Chennai 603103, India

<sup>33</sup>INFN, Sezione di Roma Tor Vergata, I-00133 Roma, Italy

<sup>34</sup>INFN, Sezione di Roma, I-00185 Roma, Italy

<sup>35</sup>Laboratoire d'Annecy de Physique des Particules (LAPP), Univ. Grenoble Alpes,

Université Savoie Mont Blanc, CNRS/IN2P3, F-74941 Annecy, France

<sup>36</sup>Embry-Riddle Aeronautical University, Prescott, AZ 86301, USA

<sup>37</sup>Montclair State University, Montclair, NJ 07043, USA

<sup>38</sup>Nikhef, Science Park 105, 1098 XG Amsterdam, The Netherlands

<sup>39</sup>Korea Institute of Science and Technology Information, Daejeon 34141, South Korea

<sup>40</sup>West Virginia University, Morgantown, WV 26506, USA

<sup>41</sup>Università di Perugia, I-06123 Perugia, Italy

<sup>42</sup>INFN, Sezione di Perugia, I-06123 Perugia, Italy

<sup>43</sup>Syracuse University, Syracuse, NY 13244, USA

<sup>44</sup>University of Minnesota, Minneapolis, MN 55455, USA

<sup>45</sup>Università degli Studi di Milano-Bicocca, I-20126 Milano, Italy

- <sup>46</sup>INFN, Sezione di Milano-Bicocca, I-20126 Milano, Italy  
<sup>47</sup>SUPA, University of Glasgow, Glasgow G12 8QQ, United Kingdom  
<sup>48</sup>LIGO Hanford Observatory, Richland, WA 99352, USA  
<sup>49</sup>Caltech CaRT, Pasadena, CA 91125, USA  
<sup>50</sup>Dipartimento di Medicina, Chirurgia e Odontoiatria "Scuola Medica Salernitana," Università di Salerno, I-84081 Baronissi, Salerno, Italy  
<sup>51</sup>Wigner RCP, RMKI, H-1121 Budapest, Konkoly Thege Miklós út 29-33, Hungary  
<sup>52</sup>Stanford University, Stanford, CA 94305, USA  
<sup>53</sup>Università di Camerino, Dipartimento di Fisica, I-62032 Camerino, Italy  
<sup>54</sup>Università di Padova, Dipartimento di Fisica e Astronomia, I-35131 Padova, Italy  
<sup>55</sup>INFN, Sezione di Padova, I-35131 Padova, Italy  
<sup>56</sup>Montana State University, Bozeman, MT 59717, USA  
<sup>57</sup>Nicolaus Copernicus Astronomical Center, Polish Academy of Sciences, 00-716, Warsaw, Poland  
<sup>58</sup>OzGrav, University of Adelaide, Adelaide, South Australia 5005, Australia  
<sup>59</sup>Center for Interdisciplinary Exploration & Research in Astrophysics (CIERA), Northwestern University, Evanston, IL 60208, USA  
<sup>60</sup>INFN, Sezione di Genova, I-16146 Genova, Italy  
<sup>61</sup>RRCAT, Indore, Madhya Pradesh 452013, India  
<sup>62</sup>Faculty of Physics, Lomonosov Moscow State University, Moscow 119991, Russia  
<sup>63</sup>Rochester Institute of Technology, Rochester, NY 14623, USA  
<sup>64</sup>Università degli Studi di Urbino "Carlo Bo," I-61029 Urbino, Italy  
<sup>65</sup>INFN, Sezione di Firenze, I-50019 Sesto Fiorentino, Firenze, Italy  
<sup>66</sup>Artemis, Université Côte d'Azur, Observatoire Côte d'Azur, CNRS, CS 34229, F-06304 Nice Cedex 4, France  
<sup>67</sup>OzGrav, University of Western Australia, Crawley, Western Australia 6009, Australia  
<sup>68</sup>Department of Astrophysics/IMAPP, Radboud University Nijmegen, P.O. Box 9010, 6500 GL Nijmegen, The Netherlands  
<sup>69</sup>Dipartimento di Fisica "E.R. Caianiello," Università di Salerno, I-84084 Fisciano, Salerno, Italy  
<sup>70</sup>INFN, Sezione di Napoli, Gruppo Collegato di Salerno, Complesso Universitario di Monte S. Angelo, I-80126 Napoli, Italy  
<sup>71</sup>Physik-Institut, University of Zurich, Winterthurerstrasse 190, 8057 Zurich, Switzerland  
<sup>72</sup>Univ Rennes, CNRS, Institut FOTON - UMR6082, F-3500 Rennes, France  
<sup>73</sup>University of Oregon, Eugene, OR 97403, USA  
<sup>74</sup>Laboratoire Kastler Brossel, Sorbonne Université, CNRS, ENS-Université PSL, Collège de France, F-75005 Paris, France  
<sup>75</sup>Astronomical Observatory Warsaw University, 00-478 Warsaw, Poland  
<sup>76</sup>VU University Amsterdam, 1081 HV Amsterdam, The Netherlands  
<sup>77</sup>Max Planck Institute for Gravitational Physics (Albert Einstein Institute), D-14476 Potsdam-Golm, Germany  
<sup>78</sup>University of Maryland, College Park, MD 20742, USA  
<sup>79</sup>School of Physics, Georgia Institute of Technology, Atlanta, GA 30332, USA  
<sup>80</sup>Université de Lyon, Université Claude Bernard Lyon 1, CNRS, Institut Lumière Matière, F-69622 Villeurbanne, France  
<sup>81</sup>Università di Napoli "Federico II," Complesso Universitario di Monte S. Angelo, I-80126 Napoli, Italy  
<sup>82</sup>NASA Goddard Space Flight Center, Greenbelt, MD 20771, USA  
<sup>83</sup>Dipartimento di Fisica, Università degli Studi di Genova, I-16146 Genova, Italy  
<sup>84</sup>RESCEU, University of Tokyo, Tokyo, 113-0033, Japan  
<sup>85</sup>Tsinghua University, Beijing 100084, China  
<sup>86</sup>Texas Tech University, Lubbock, TX 79409, USA  
<sup>87</sup>Università di Roma Tor Vergata, I-00133 Roma, Italy  
<sup>88</sup>The University of Mississippi, University, MS 38677, USA  
<sup>89</sup>Missouri University of Science and Technology, Rolla, MO 65409, USA  
<sup>90</sup>Museo Storico della Fisica e Centro Studi e Ricerche "Enrico Fermi," I-00184 Roma, Italy  
<sup>91</sup>The Pennsylvania State University, University Park, PA 16802, USA  
<sup>92</sup>National Tsing Hua University, Hsinchu City, 30013 Taiwan, Republic of China  
<sup>93</sup>Charles Sturt University, Wagga Wagga, New South Wales 2678, Australia  
<sup>94</sup>Canadian Institute for Theoretical Astrophysics, University of Toronto, Toronto, Ontario M5S 3H8, Canada  
<sup>95</sup>University of Chicago, Chicago, IL 60637, USA  
<sup>96</sup>The Chinese University of Hong Kong, Shatin, NT, Hong Kong  
<sup>97</sup>Dipartimento di Ingegneria Industriale (DIIN), Università di Salerno, I-84084 Fisciano, Salerno, Italy  
<sup>98</sup>Seoul National University, Seoul 08826, South Korea  
<sup>99</sup>Pusan National University, Busan 46241, South Korea  
<sup>100</sup>Carleton College, Northfield, MN 55057, USA

- <sup>101</sup>INAF, Osservatorio Astronomico di Padova, I-35122 Padova, Italy  
<sup>102</sup>OzGrav, University of Melbourne, Parkville, Victoria 3010, Australia  
<sup>103</sup>Universitat de les Illes Balears, IAC3—IEEC, E-07122 Palma de Mallorca, Spain  
<sup>104</sup>Université Libre de Bruxelles, Brussels 1050, Belgium  
<sup>105</sup>Sonoma State University, Rohnert Park, CA 94928, USA  
<sup>106</sup>Departamento de Matemáticas, Universitat de València, E-46100 Burjassot, València, Spain  
<sup>107</sup>Columbia University, New York, NY 10027, USA  
<sup>108</sup>Cardiff University, Cardiff CF24 3AA, United Kingdom  
<sup>109</sup>University of Rhode Island, Kingston, RI 02881, USA  
<sup>110</sup>The University of Texas Rio Grande Valley, Brownsville, TX 78520, USA  
<sup>111</sup>Bellevue College, Bellevue, WA 98007, USA  
<sup>112</sup>MTA-ELTE Astrophysics Research Group, Institute of Physics, Eötvös University, Budapest 1117, Hungary  
<sup>113</sup>Institute for Plasma Research, Bhat, Gandhinagar 382428, India  
<sup>114</sup>The University of Sheffield, Sheffield S10 2TN, United Kingdom  
<sup>115</sup>IGFAE, Campus Sur, Universidade de Santiago de Compostela, 15782 Spain  
<sup>116</sup>Dipartimento di Scienze Matematiche, Fisiche e Informatiche, Università di Parma, I-43124 Parma, Italy  
<sup>117</sup>INFN, Sezione di Milano Bicocca, Gruppo Collegato di Parma, I-43124 Parma, Italy  
<sup>118</sup>Dipartimento di Ingegneria, Università del Sannio, I-82100 Benevento, Italy  
<sup>119</sup>Università di Trento, Dipartimento di Fisica, I-38123 Povo, Trento, Italy  
<sup>120</sup>INFN, Trento Institute for Fundamental Physics and Applications, I-38123 Povo, Trento, Italy  
<sup>121</sup>Università di Roma “La Sapienza,” I-00185 Roma, Italy  
<sup>122</sup>Colorado State University, Fort Collins, CO 80523, USA  
<sup>123</sup>Kenyon College, Gambier, OH 43022, USA  
<sup>124</sup>CNR-SPIN, c/o Università di Salerno, I-84084 Fisciano, Salerno, Italy  
<sup>125</sup>Scuola di Ingegneria, Università della Basilicata, I-85100 Potenza, Italy  
<sup>126</sup>National Astronomical Observatory of Japan, 2-21-1 Osawa, Mitaka, Tokyo 181-8588, Japan  
<sup>127</sup>Observatori Astronòmic, Universitat de València, E-46980 Paterna, València, Spain  
<sup>128</sup>INFN Sezione di Torino, I-10125 Torino, Italy  
<sup>129</sup>School of Mathematics, University of Edinburgh, Edinburgh EH9 3FD, United Kingdom  
<sup>130</sup>Institute Of Advanced Research, Gandhinagar 382426, India  
<sup>131</sup>Indian Institute of Technology Bombay, Powai, Mumbai 400 076, India  
<sup>132</sup>University of Szeged, Dóm tér 9, Szeged 6720, Hungary  
<sup>133</sup>SUPA, University of the West of Scotland, Paisley PA1 2BE, United Kingdom  
<sup>134</sup>California State University, Los Angeles, 5151 State University Dr, Los Angeles, CA 90032, USA  
<sup>135</sup>Universität Hamburg, D-22761 Hamburg, Germany  
<sup>136</sup>Tata Institute of Fundamental Research, Mumbai 400005, India  
<sup>137</sup>INAF, Osservatorio Astronomico di Capodimonte, I-80131 Napoli, Italy  
<sup>138</sup>University of Michigan, Ann Arbor, MI 48109, USA  
<sup>139</sup>Washington State University, Pullman, WA 99164, USA  
<sup>140</sup>American University, Washington, D.C. 20016, USA  
<sup>141</sup>University of Portsmouth, Portsmouth, PO1 3FX, United Kingdom  
<sup>142</sup>University of California, Berkeley, CA 94720, USA  
<sup>143</sup>GRAPPA, Anton Pannekoek Institute for Astronomy and Institute for High-Energy Physics, University of Amsterdam, Science Park 904, 1098 XH Amsterdam, The Netherlands  
<sup>144</sup>Delta Institute for Theoretical Physics, Science Park 904, 1090 GL Amsterdam, The Netherlands  
<sup>145</sup>Directorate of Construction, Services & Estate Management, Mumbai 400094 India  
<sup>146</sup>University of Białystok, 15-424 Białystok, Poland  
<sup>147</sup>King's College London, University of London, London WC2R 2LS, United Kingdom  
<sup>148</sup>University of Southampton, Southampton SO17 1BJ, United Kingdom  
<sup>149</sup>University of Washington Bothell, Bothell, WA 98011, USA  
<sup>150</sup>Institute of Applied Physics, Nizhny Novgorod, 603950, Russia  
<sup>151</sup>Ewha Womans University, Seoul 03760, South Korea  
<sup>152</sup>Inje University Gimhae, South Gyeongsang 50834, South Korea  
<sup>153</sup>National Institute for Mathematical Sciences, Daejeon 34047, South Korea  
<sup>154</sup>Ulsan National Institute of Science and Technology, Ulsan 44919, South Korea  
<sup>155</sup>Maastricht University, P.O. Box 616, 6200 MD Maastricht, The Netherlands  
<sup>156</sup>Bard College, 30 Campus Rd, Annandale-On-Hudson, NY 12504, USA  
<sup>157</sup>NCBJ, 05-400 Świerk-Otwock, Poland  
<sup>158</sup>Institute of Mathematics, Polish Academy of Sciences, 00656 Warsaw, Poland  
<sup>159</sup>Cornell University, Ithaca, NY 14850, USA  
<sup>160</sup>Hillsdale College, Hillsdale, MI 49242, USA  
<sup>161</sup>Hanyang University, Seoul 04763, South Korea  
<sup>162</sup>Korea Astronomy and Space Science Institute, Daejeon 34055, South Korea

- <sup>163</sup>*Institute for High-Energy Physics, University of Amsterdam,  
Science Park 904, 1098 XH Amsterdam, The Netherlands*
- <sup>164</sup>*NASA Marshall Space Flight Center, Huntsville, AL 35811, USA*
- <sup>165</sup>*Dipartimento di Matematica e Fisica, Università degli Studi Roma Tre, I-00146 Roma, Italy*
- <sup>166</sup>*INFN, Sezione di Roma Tre, I-00146 Roma, Italy*
- <sup>167</sup>*ESPCI, CNRS, F-75005 Paris, France*
- <sup>168</sup>*OzGrav, Swinburne University of Technology, Hawthorn VIC 3122, Australia*
- <sup>169</sup>*Southern University and A&M College, Baton Rouge, LA 70813, USA*
- <sup>170</sup>*Centre Scientifique de Monaco, 8 quai Antoine Ier, MC-98000, Monaco*
- <sup>171</sup>*Indian Institute of Technology Madras, Chennai 600036, India*
- <sup>172</sup>*Institut des Hautes Etudes Scientifiques, F-91440 Bures-sur-Yvette, France*
- <sup>173</sup>*IISER-Kolkata, Mohanpur, West Bengal 741252, India*
- <sup>174</sup>*Institut für Kernphysik, Theoriezentrum, 64289 Darmstadt, Germany*
- <sup>175</sup>*Whitman College, 345 Boyer Avenue, Walla Walla, WA 99362 USA*
- <sup>176</sup>*Université de Lyon, F-69361 Lyon, France*
- <sup>177</sup>*Hobart and William Smith Colleges, Geneva, NY 14456, USA*
- <sup>178</sup>*Dipartimento di Fisica, Università degli Studi di Torino, I-10125 Torino, Italy*
- <sup>179</sup>*University of Washington, Seattle, WA 98195, USA*
- <sup>180</sup>*INAF, Osservatorio Astronomico di Brera sede di Merate, I-23807 Merate, Lecco, Italy*
- <sup>181</sup>*Centro de Astrofísica e Gravitação(CENTRA),  
Departamento de Física, Instituto Superior Técnico, Universidade de Lisboa, 1049 – 001 Lisboa, Portugal*
- <sup>182</sup>*Marquette University, 11420 W. Clybourn St., Milwaukee, WI 53233, USA*
- <sup>183</sup>*Indian Institute of Technology, Gandhinagar Ahmedabad Gujarat 382424, India*
- <sup>184</sup>*Université de Montréal/Polytechnique, Montreal, Quebec H3T 1J4, Canada*
- <sup>185</sup>*Indian Institute of Technology Hyderabad, Sangareddy, Khamdi, Telangana 502285, India*
- <sup>186</sup>*INAF, Osservatorio di Astrofisica e Scienza dello Spazio, I-40129 Bologna, Italy*
- <sup>187</sup>*International Institute of Physics, Universidade Federal do Rio Grande do Norte, Natal RN 59078-970, Brazil*
- <sup>188</sup>*Villanova University, 800 Lancaster Ave, Villanova, PA 19085, USA*
- <sup>189</sup>*Andrews University, Berrien Springs, MI 49104, USA*
- <sup>190</sup>*Max Planck Institute for Gravitational Physik (Albert Einstein Institute), D-14476 Potsdam-Golm, Germany*
- <sup>191</sup>*Università di Siena, I-53100 Siena, Italy*
- <sup>192</sup>*Trinity University, San Antonio, TX 78212, USA*
- <sup>193</sup>*Van Swinderen Institute for Particle Physics and Gravity,  
University of Groningen, Nijenborgh 4, 9747 AG Groningen, The Netherlands*
- <sup>194</sup>*Department of Physics, University of Texas, Austin, TX 78712, USA*
- <sup>195</sup>*Max Planck Institute for Gravitational Physics (Albert Einstein Institute), D-30167 Hannover, Germany*
- <sup>196</sup>*Max Planck Institute for Gravitational Physics (Albert Einstein Institute), D-14476 Potsdam-Golm, Germany*
- <sup>197</sup>*University of Wisconsin-Milwaukee, Milwaukee, WI 53201, USA*

(Dated: June 18, 2019)

Gravitational wave astronomy has been firmly established with the detection of gravitational waves from the merger of ten stellar mass binary black holes and a neutron star binary. This paper reports on the all-sky search for gravitational waves from intermediate mass black hole binaries in the first and second observing runs of the Advanced LIGO and Virgo network. The search uses three independent algorithms: two based on matched filtering of the data with waveform templates of gravitational wave signals from compact binaries, and a third, model-independent algorithm that employs no signal model for the incoming signal. No intermediate mass black hole binary event was detected in this search. Consequently, we place upper limits on the merger rate density for a family of intermediate mass black hole binaries. In particular, we choose sources with total masses  $M = m_1 + m_2 \in [120, 800] M_{\odot}$  and mass ratios  $q = m_2/m_1 \in [0.1, 1.0]$ . For the first time, this calculation is done using numerical relativity waveforms (which include higher modes) as models of the real emitted signal. We place a most stringent upper limit of  $0.20 \text{ Gpc}^{-3}\text{yr}^{-1}$  (in co-moving units at the 90% confidence level) for equal-mass binaries with individual masses  $m_{1,2} = 100 M_{\odot}$  and dimensionless spins  $\chi_{1,2} = 0.8$  aligned with the orbital angular momentum of the binary. This improves by a factor of  $\sim 5$  that reported after Advanced LIGO’s first observing run.

PACS numbers: 04.80.Nn, 07.05.Kf, 95.55.Ym

## I. INTRODUCTION

The first two observing runs of Advanced LIGO and Virgo (O1 and O2 respectively) have significantly enhanced our understanding of black hole (BH) binaries in

\* Deceased, July 2018.

the universe. Gravitational waves (GWs) from 10 binary black hole mergers with total mass between  $18.6^{+3.1}_{-0.7} M_{\odot}$  and  $85.1^{+15.6}_{-10.9} M_{\odot}$  were detected during these two observing runs [1–8]. These observations have revealed a new population of heavy stellar mass BH components of up to  $50 M_{\odot}$ , for which we had no earlier electromagnetic observational evidence [8, 9]. This finding limit is consistent with the formation of heavier BHs from core collapse being prevented by a mechanism known as *pulsational pair-instability supernovae* (PISN) [10–13]. According to this idea, stars with helium core mass in the range  $\sim 32 - 64 M_{\odot}$  undergo pulsational pair instability leaving behind remnants of  $\lesssim 65 M_{\odot}$ . Stars with helium core mass in the range  $\sim 64 - 135 M_{\odot}$  undergo pair instability and leave no remnant, while stars with helium mass  $\gtrsim 135 M_{\odot}$  are thought to directly collapse to intermediate-mass black holes.

Intermediate mass black holes (IMBHs) are BHs heavier than stellar mass BHs but lighter than supermassive black holes (SMBHs), which places them roughly in the range of  $10^2 - 10^5 M_{\odot}$  [14, 15]. Currently there is only indirect observational evidence. Observations include probing the mass of the central BH in galaxies as well as massive star clusters with direct kinematical measurements which has led to recent claims for the presence of IMBHs [16–18]. Other observations come from the extrapolation of several scaling relations between the mass of the central SMBH and their host galaxies [19] to the mass range of globular clusters [20, 21]. In this way, several clusters have been found to be good candidates for having IMBHs in their centers [22–24]. If present, IMBHs would heat up the cores of these clusters, strongly influencing the distribution of the stars in the cluster and their dynamics, leaving a characteristic imprint in the surface brightness profile, as well as in the mass-to-light ratio [25]. Controversy exists regarding the interpretation of these observations, as some of them can also be explained by a high concentration of stellar-mass BHs or the presence of binaries [22–24, 26]. Empirical mass scaling relations of quasi-periodic oscillations [27] in luminous X-Ray sources have also provided evidence for IMBH [28]. Finally, IMBHs have been proposed as candidates to explain ultra-luminous X-Ray (ULX) sources in nearby galaxies, which are brighter than the accreting X-ray sources with stellar mass BHs [29, 30]. However, neutron stars or stellar-mass black holes emitting above their Eddington luminosity could also account for such observations. In summary, no definitive evidence of IMBHs has yet been obtained.

The possible astrophysical formation channels of IMBHs remain uncertain. Proposed channels include the direct collapse of massive first generation, low metallicity Population III stars [31–34] and mergers of stellar mass BHs in globular clusters [36] and multiple collisions of stars in dense young star clusters [18, 35, 37–39], among others [40]. Further, some astrophysical scenarios [14] indicate that SMBHs in galactic centers might be formed from hierarchical mergers of IMBHs [15, 41]. The di-

rect observation of IMBHs with gravitational waves could strengthen the possible evolutionary link between stellar mass BHs and SMBHs. Finally, observing an IMBH population would help to understand details of the pulsational pair-instability supernovae mechanism.

The GW observation of a coalescing binary consisting of at least one IMBH component or resulting in an IMBH remnant, which we will term an IMBHB, could provide the first definitive confirmation of the existence of IMBHs. In fact, IMBHBs are the sources that would emit the most gravitational-wave energy in the LIGO-Virgo frequency band, potentially making them detectable to distances (and redshifts) beyond that of any other LIGO-Virgo source [42]. Even in the absence of a detection, a search for IMBHBs provides stringent constraints on their merger rate density, which has implications for potential IMBHB and SMBH formation channels.

IMBHs are not only interesting from an astrophysical point of view, they are also excellent laboratories to test general relativity in the strong field regime [43–46]. Their large masses would lead to strong merger and ringdown signals in the Advanced LIGO-Virgo frequency band. Therefore, higher modes might be visible in IMBHB signals because those modes are especially strong in the merger and ringdown stages. The observation of multimodal merger and ringdown signals is paramount to understanding fundamental properties of general relativity, such as the no-hair theorem [47–50] and BH kick measurements [51–53].

The first search for GWs from IMBHBs was carried out in the data from initial LIGO and initial Virgo (2005–2010) [54, 55]. Owing to the large masses of IMBHBs, such systems are expected to merge at low frequencies where the initial detectors were less sensitive due to the presence of several noise sources, such as suspension noise, thermal noise and optical cavity control noise. As a result, the those detectors were sensitive to only the merger and ringdown phases of the IMBHB systems. Initial IMBHB analyses applied either the model-independent time-frequency searches [56] or ringdown searches. No IMBHB merger was detected in these searches.

Because of the improved low-frequency sensitivities of the Advanced LIGO and Advanced Virgo detectors [57, 58], IMBHB signals are visible in band for a longer period of time, which increases the effectiveness of modeled searches that use more than just the ringdown portion of an IMBHB’s waveform. Ref. [42] reports results from a combined search for IMBHBs that used two independent search algorithms: a matched-filter analysis, called GstLAL [59–61], which uses the inspiral, merger, and ringdown portions of the IMBHB waveform and the model-independent analysis coherent WaveBurst (cWB) [56]. No IMBHBs were found by these searches, and upper limits on the merger rate density for 12 targeted IMBHB sources with total mass between  $120 M_{\odot} - 600 M_{\odot}$  and mass ratios down to 1/10 were obtained. The most stringent upper limit on the

merger rate density from this combined analysis was<sup>189</sup>  
 0.93 Gpc<sup>-3</sup> yr<sup>-1</sup> for binaries consisting of two 100 M<sub>⊙</sub><sup>190</sup>  
 BHs with dimensionless spin magnitude 0.8 aligned with<sup>191</sup>  
 the system's orbital angular momentum.<sup>192</sup>

All upper limits on the IMBHB merger rate reported<sup>193</sup>  
 in past searches [42, 54, 55] were obtained using models<sup>194</sup>  
 for the GW signal that include only the dominant radiat<sup>195</sup>  
 ing mode, namely  $(\ell, m) = (2, \pm 2)$ , of the GW emission<sup>196</sup>  
 [62]. However, it has been shown that higher modes con<sup>197</sup>  
 tribute more substantially to signals emitted by heavy<sup>198</sup>  
 binaries. This impact increases as the system becomes<sup>199</sup>  
 more asymmetric in mass [63, 64], as the spin of the BHs  
 becomes more negative [65, 66], and as the precession in<sup>200</sup>  
 the binary becomes stronger [67]. As a consequence, the  
 omission of higher modes leads in general to more con<sup>201</sup>  
 servative upper limits on the IMBHB merger rate [68].<sup>201</sup>

In this work, we improve on past studies in two distinct<sup>202</sup>  
 ways. We use numerical relativity (NR) simulations with<sup>203</sup>  
 higher modes to model GW signals from IMBHBs for<sup>204</sup>  
 computing upper limit estimates. Additionally, our com<sup>205</sup>  
 bined analysis now includes the matched-filter search Py<sup>206</sup>  
 CBC [69, 70] in addition to GstLAL and cWB. Because<sup>207</sup>  
 of these novelties, we have, in addition to analyzing the<sup>208</sup>  
 O2 data set, reanalyzed the O1 data set and report here<sup>209</sup>  
 combined upper limits for the O1 and O2 observing runs.<sup>210</sup>  
 In this paper, we report upper limits on the merger rate<sup>211</sup>  
 density of 17 targeted (non-precessing) IMBHB sources.<sup>212</sup>  
 Our most stringent upper limit is 0.20 Gpc<sup>-3</sup> yr<sup>-1</sup> for<sup>213</sup>  
 equal-mass binaries with component spins aligned with<sup>214</sup>  
 the orbital angular momentum of the system and dimen<sup>215</sup>  
 sionless magnitudes  $\chi_{1,2} = 0.8$ .<sup>216</sup>

The rest of this paper is organized as follows. In Sec. II<sup>217</sup>  
 we describe the data set, outline the individual search al<sup>218</sup>  
 gorithms that make up the combined search, and report<sup>219</sup>  
 our search's null detection of IMBHBs. In Sec. III we<sup>220</sup>  
 describe the NR simulations that we use to compute up<sup>221</sup>  
 per limits on IMBHB merger rates report these for the<sup>222</sup>  
 case of 17 IMBHB sources. We draw final conclusions in<sup>223</sup>  
 Sec. IV.<sup>224</sup>

## II. IMBHB SEARCH IN O1 AND O2 DATA

### A. Data Summary

This analysis was carried out using O1 and O2 data<sup>228</sup>  
 sets from the two LIGO (Livingston and Hanford) de<sup>229</sup>  
 tectors and Virgo. We have used the final calibration,<sup>230</sup>  
 which was produced after the conclusion of the run, in<sup>231</sup>  
 cluding compensation for frequency-dependent fluctua<sup>232</sup>  
 tions in the calibration [71–73]. Well identified sources<sup>233</sup>  
 of noise have also been subtracted from the strain data<sup>234</sup>  
 as explained in Refs. [73, 74]. The maximum calibration<sup>235</sup>  
 uncertainty across the frequency band of [10–5,000] Hz for<sup>236</sup>  
 the two LIGO detectors is ∼10% in amplitude and ∼5<sup>237</sup>  
 degrees in phase for O1 and ∼4% in amplitude and ∼3<sup>238</sup>  
 degrees in phase for O2 [7, 71]. For Virgo we consider an<sup>239</sup>  
 uncertainty of 5.1% in amplitude and 2.3 degrees in phase<sup>240</sup>

[73]. After removing data with significant instrumental  
 disturbance, we use 48.6 days and 118.0 days of joint  
 Hanford-Livingston data from the O1 and O2 observing  
 runs respectively. The Virgo detector joined the LIGO  
 detectors during the last ∼15 days of O2, which provided  
 with an additional 4.0 days of coincident data with either  
 of Hanford-Virgo or Livingston-Virgo network. The data  
 from O1 and O2 was divided into 9 and 21 blocks respec-  
 tively with coincident time ranging from 4.7 – 7.0 days.  
 For more details, see Ref. [8].

### B. Search algorithms

We combine the two matched-filter searches, namely  
 GstLAL [59–61, 75] and PyCBC [69, 70], and one model-  
 independent analysis, cWB [76], into a single IMBHB  
 search. The two model-based matched filtering analy-  
 ses use a bank of templates made of pre-computed com-  
 pact binary merger GW waveforms. Matched filter based  
 analyses are optimal to extract known signals from sta-  
 tionary, Gaussian noise [77]. However, the templates we  
 use are limited to non-eccentric, aligned-spin systems.  
 They contain only the dominant waveform mode of the  
 GW emission and omit higher modes [64, 78]. Addi-  
 tionally, Advanced LIGO and Virgo data are known to  
 contain a large number of short noise transients [79],  
 which can mimic short GW signals like those emitted by  
 IMBHBs. While matched-filter searches use several tech-  
 niques to discriminate between noisy transients and real  
 GW events [61, 80, 81], they are known to lose significant  
 efficiency when looking for short signals like those from  
 IMBHBs. Therefore, the IMBHB search is carried out  
 jointly with an analysis that can identify short-duration  
 GW signals without a model for the morphology of the  
 GW waveform. In this search, all three analyses use O1  
 and O2 Advanced LIGO data. However, because of the  
 incomparable sensitivities between the Advanced LIGO  
 detector and Advanced Virgo detector, only the GstLAL  
 analysis uses Virgo data, as is done in Ref. [8].

#### 1. Modeled analyses

The matched-filter analyses GstLAL and PyCBC use  
 templates that span the parameter space of neutron stars,  
 stellar-mass BHs, and IMBHBs. In this study, we use the  
 same two searches reported in Ref. [8] to calculate upper  
 limits on the merger rate density of IMBHBs.

The matched-filter signal-to-noise ratio (SNR) time se-  
 ries is computed for every template. Triggers are pro-  
 duced when the SNR time series surpasses a predeter-  
 mined threshold, where clusters of triggers are trimmed  
 by maximizing the SNR within small time windows. In  
 addition, a signal consistency veto [61, 81, 82] is calcu-  
 lated for each trigger. A list of GW candidates is con-  
 structed from triggers generated by common templates  
 that are coincident in time across more than one detec-

241 tor, where the coincidence window takes into account the<sup>292</sup>  
 242 travel time between detectors. Next, a ranking statistic<sup>293</sup>  
 243 is calculated for each candidate that estimates a likeli-<sup>294</sup>  
 244 hood ratio that the candidate would be observed in the<sup>295</sup>  
 245 presence of a GW compared to a pure-noise expectation.<sup>296</sup>  
 246 Finally, a *p*-value<sup>1</sup>  $P$  is determined by comparing the<sup>297</sup>  
 247 value of its ranking statistic to that of triggers coming<sup>298</sup>  
 248 from background noise in the data. A detailed descrip-<sup>299</sup>  
 249 tion of the GstLAL and PyCBC pipelines can be found<sup>300</sup>  
 250 in Refs. [59–61, 75] and [69, 70], respectively; addition-<sup>301</sup>  
 251 ally, details outlining how candidates are ranked across<sup>302</sup>  
 252 observing runs can be found in Ref. [8].<sup>303</sup>

The GstLAL analysis uses the template bank described<sup>304</sup>  
 253 in Ref. [83]. The region of this bank that overlaps the<sup>305</sup>  
 254 IMBHB parameter space, which starts at a total mass<sup>306</sup>  
 255 of  $100 M_{\odot}$ , reaches up to a total mass of  $400 M_{\odot}$  in<sup>307</sup>  
 256 the detector frame and covers mass ratios in the range<sup>308</sup>  
 257 of  $1/98 < q < 1$ . The waveforms used are a reduced-<sup>309</sup>  
 258 order-model of the SEOBNRv4 approximant [84]. The<sup>310</sup>  
 259 spin of these templates are either aligned or anti-aligned<sup>311</sup>  
 260 with the orbital angular momentum of the system with<sup>312</sup>  
 261 dimensionless magnitudes less than 0.999.<sup>313</sup>

The PyCBC analysis uses the template bank described<sup>314</sup>  
 262 in Ref. [85]. The region of this bank that overlaps the<sup>315</sup>  
 263 IMBHB parameter space reaches up to a total mass<sup>316</sup>  
 264 of  $500 M_{\odot}$  in the detector frame, excluding templates<sup>317</sup>  
 265 with duration below 0.15 s, and covers the range of<sup>318</sup>  
 266  $1/98 < q < 1$ . The waveforms used are also a reduced-<sup>319</sup>  
 267 order-model of the SEOBNRv4 approximant, and the<sup>320</sup>  
 268 aligned or anti-aligned dimensionless spin magnitudes are<sup>321</sup>  
 269 less than 0.998.<sup>322</sup>

## 2. Un-modeled analysis

273 Coherent WaveBurst (cWB) is the GW transient de-<sup>323</sup>  
 274 tection algorithm designed to look for unmodeled short-<sup>324</sup>  
 275 duration GW transients in the multi-detector data from<sup>325</sup>  
 276 interferometric GW detector networks. Designed to oper-<sup>326</sup>  
 277 ate without a specific waveform model, cWB identifies co-<sup>327</sup>  
 278 incident excess power in the wavelet time-frequency rep-<sup>328</sup>  
 279 resentations of the detector strain data [86], for signal fre-<sup>329</sup>  
 280 quencies up to 1 kHz and durations up to a few seconds.<sup>330</sup>  
 The search identifies events that are coherent in multiple  
 281 detectors and reconstructs the source sky location and<sup>331</sup>  
 282 signal waveforms by using the constrained maximum like-<sup>332</sup>  
 283 lihood method [76]. The cWB detection statistic is based<sup>333</sup>  
 284 on the coherent energy  $E_c$  obtained by cross-correlating<sup>334</sup>  
 285 the signal waveforms reconstructed in multiple detectors.<sup>335</sup>  
 It is proportional to the network SNR and used to rank<sup>336</sup>  
 286 the events found by cWB.<sup>337</sup>

To improve the robustness of the algorithm against<sup>340</sup>  
 289 non-stationary detector noise, cWB uses signal-<sup>341</sup>  
 290 independent vetoes, which reduce the high rate of the<sup>342</sup>  
 291

initial excess power triggers. The primary veto cut is on the network correlation coefficient  $c_c = E_c/(E_c + E_n)$ , where  $E_n$  is the residual noise energy estimated after the reconstructed signal is subtracted from the data. Typically, for a GW signal  $c_c \approx 1$  and for instrumental glitches  $c_c \ll 1$ . Therefore, candidate events with  $c_c < 0.7$  are rejected as potential glitches.

To improve the detection efficiency of IMBHBs as well as to reduce the false alarm rates (FARs), the cWB analysis employs additional selection cuts based on the nature of IMBHB signals. IMBHB signals have two distinct features in the time-frequency representation. First, the signal frequencies lie below 250 Hz. We use this to exclude all the non-IMBHB events in the search, including noise events. Secondly, the inspiral signal duration in the detector band is relatively short, which leads to relatively low SNR in the inspiral phase as compared to the merger and ringdown phases. In the cWB framework, chirp mass ( $\mathcal{M} = (m_1 m_2)^{3/5} M^{-1/5}$ ) is estimated using the frequency evolution of a signal's inspiral. However, in the case of low SNRs, we cannot accurately estimate the chirp mass of the binary [87]; still, we use this framework to introduce additional cuts on the estimated chirp mass to reject non-IMBHB signals. The simulation studies show that IMBHB signals are recovered with  $|\mathcal{M}| > 10 M_{\odot}$  which we use in this search.<sup>2</sup> We apply this selection cut to reduce the noise background when producing the candidate events.

For estimation of the statistical significance of the candidate event, each event is ranked against a sample of background triggers obtained by repeating the analysis on time-shifted data [1]. To exclude astrophysical events from the background sample, the time shifts are selected to be much longer (1 second or more) than the expected signal time delay between the detectors. By using different time shifts, a sample of background events equivalent to approximately 500 years of background data is accumulated for each of the 30 blocks of data. The cWB candidate events that survived the cWB selection criteria, are assigned a FAR given by the rate of the corresponding background events with the coherent network SNR value larger than that of the candidate event.

## C. Combined search

Each of our three algorithms produces its own list of GW candidates, characterized by GPS time, FAR and associated *p*-value  $P$ . These three lists are then combined into a common single list of candidates. To avoid counting candidates more than once, candidates within a time window of 100 ms across different lists are assumed to be the same. To account for the use of three search

<sup>1</sup> The probability that noise would produce a trigger at least as significant as the observed candidate.

<sup>2</sup> Negative  $\mathcal{M}$  values correspond to frequencies decreasing with time, which could be due to the pixels corresponding to ringdown part.

algorithms, we apply a conservative trials factor of 3 and assign each candidate a new  $p$ -value given by

$$\bar{P} = 1 - (1 - P_{\min})^3, \quad (1)$$

where  $P_{\min}$  denotes the minimum  $p$ -value reported across the pipelines. This is equivalent to assuming that the three searches produce independent lists of candidates.<sup>3</sup> We note that while this choice of trials factor affects the significance of individual triggers, it will not change the numerical value of our upper limits. See Appendix B for a more detailed discussion.

No	Date	UTC time	Pipeline	FAR (yr <sup>-1</sup> )	SNR	$P_{\min}$
1	2017-05-02	04:08:44.9	cWB	0.34	11.6	0.14 <sup>75</sup>
2	2017-06-16	19:47:20.8	PyCBC	1.94	9.1	0.59 <sup>376</sup>
3	2015-11-26	04:11:02.7	cWB	2.56	7.5	0.68 <sup>377</sup>
4	2017-06-08	23:50:52.3	cWB	3.57	10.0	0.79 <sup>379</sup>
5	2017-04-05	11:04:52.7	GstLAL	4.55	9.3	0.88 <sup>380</sup>
6	2015-11-16	22:41:48.7	PyCBC	4.77	9.0	0.88 <sup>382</sup>
7	2016-12-02	03:53:44.9	GstLAL	6.00	10.5	0.94 <sup>383</sup>
8	2017-02-19	14:04:09.0	GstLAL	6.26	9.6	0.95 <sup>385</sup>
9	2017-04-23	12:10:45.0	GstLAL	6.47	8.9	0.95 <sup>386</sup>
10	2017-04-12	15:56:39.0	GstLAL	8.22	9.7	0.98 <sup>387</sup>

TABLE I: Details of the ten most significant events (excluding all published lower mass events). We report the date, UTC time, observing pipeline (individual analysis that observed the event with the highest significance), FAR, SNR, and  $P_{\min}$  for each event. The combined  $p$ -value  $\bar{P}$  of each event is calculated using Eq. 1. In the table, the events are tabulated in increasing value of  $P_{\min}$ .

have a FAR much larger than any of the GWs reported in Ref. [8]; no event in this list was found with enough significance to claim an IMBHB detection.

The top-ranked event<sup>4</sup> from Table I was observed by cWB in O2 data on May 2, 2017 at 04:08:44 UTC with a combined SNR of 11.6 in the two Advanced LIGO detectors and a significance of  $P_{\text{cWB}} = P_{\min} = 0.14$ . Applying eq. 1, this event has a combined  $p$ -value of  $\bar{P} = 0.36$ , too low to claim it as a gravitational wave detection.

Despite the low significance of this trigger, its characteristics were consistent with those of an IMBHB, and we decided to perform detailed data quality and parameter estimation follow-ups.<sup>5</sup> In order to check for the presence of environmental or instrumental noise, this event was vetted with the same procedure applied to triggers of marginal significance found in previous searches [8] in O1-O2 data. These checks identified a correlation between the trigger time and the glitching of optical lever lasers at the end of one arm of the Hanford detector. This is a known instrumental artifact previously observed to impact GW searches [88, 89]. The time of this trigger was not discarded by the pre-tuned data quality veto designed to mitigate the effects of these optical lever laser glitches. However, these vetoes are tuned for high efficiency and minimal impact on analyzable time rather than exhaustively removing all non-Gaussian features in the data. Further follow-up indicates that this instrumental artifact is likely contributing power to the gravitational wave strain channel at the time and frequency of the trigger. Given the SNR of the purported signal in the Hanford detector and the relatively low significance of the reported false alarm rate, we conclude that this trigger is likely explained by detector noise.

### III. UPPER LIMITS ON MERGER RATES

Given that no IMBHB signal was detected by our search, we proceed to place upper limits on the coalescence rate of these objects. This is done by estimating the sensitivity of our search to an astrophysically motivated population of simulated IMBHB signals that we inject in our detector data. However, given the absence of well motivated population estimates of IMBHBs, we opt for sampling the parameter space in a discrete manner (for details, please see Appendix C). As a consequence, in this section we estimate the sensitive distance reach as well as the upper limit on merger rate density for 17 selected fiducial IMBHB sources, tabulated in Table II of Appendix C using the loudest event method [90], following the procedure outlined in Ref. [42] and described again in Appendix A. For a given IMBHB

#### D. Search results

Here we report results from the combined cWB-GstLAL-PyCBC IMBHB search on full O1-O2 data. The top 21 most significant events from the combined search include the 11 GW events published in Ref. [8], namely GW150914 [1], GW151012, GW151226 [2], GW170104 [3], GW170608 [4], GW170729, GW170809, GW170814 [6], GW170817 [5], GW170818, and GW170823, and 10 events tabulated in Table I. All the events in Table I

<sup>3</sup> In general, correlations between searches would lead to a trials factor less than 3. However, at the time, we are not able to quantify this, and we choose to adopt the most conservative approach.

<sup>4</sup> We note that the most significant event in the O1 search reported in Ref. [42] is the third event in this Table.

<sup>5</sup> See Appendix D for further details regarding the parameter estimation investigations of this candidate.

409 source, gravitational waveforms from simulated systems<sup>459</sup>  
 410 scattered through space are injected into the data and<sup>460</sup>  
 411 recovered by each of the three analyses. In this section,<sup>461</sup>  
 412 we describe our simulation set and present our findings.<sup>462</sup>

463 to red shift  $z \sim 1$  (luminosity distance of 6.7 Gpc) us-  
 464 ing the TT+lowP+lensing+ext cosmological parameters  
 465 given in Table IV of Ref. [94], and individually redshifted  
 466 according to this cosmological model. The overall effect  
 467 of cosmological redshift is to shift each GW signal to  
 468 lower frequencies. At a given redshift, the mass of the  
 469 injection in the detector frame is  $(1+z)$  times larger than  
 470 the source frame mass, and the luminosity distance is  
 471  $(1+z)$  times the comoving distance. At redshifts of  $z=1$ ,  
 472 this results in a decrease in SNR, ranging from  $\sim 20\%$   
 473 for an equal-mass  $M = 100 M_{\odot}$  face-on system to  $\sim 50\%$   
 474 for an equal-mass,  $M = 200 M_{\odot}$  face-on system. The  
 475 injections are spaced roughly uniformly in time with an  
 476 interval of at least 80 seconds over the  $T_0 = 413.71$  days  
 477 of O1-O2 observing time, and each injection set covers a  
 478 total space-time volume  $\langle VT \rangle_{\text{tot}} = 110.68 \text{ Gpc}^3 \text{ yr}$ .

### 479 A. Injection set

480 Ref. [42] reports upper limits on the merger rate den-<sup>467</sup>  
 481 sity for 12 IMBHB systems in its Table I. The waveform<sup>468</sup>  
 482 simulations used to compute upper limits in that study<sup>469</sup>  
 483 contain only the dominant quadrupolar mode of the GW<sup>470</sup>  
 484 emission. In this work, we use highly accurate NR sim-<sup>471</sup>  
 485 ulations computed by the SXS [91], RIT [92], and Geor-<sup>472</sup>  
 486 giaTech [93] codes, which include higher modes. Since<sup>473</sup>  
 487 higher modes are particularly important for large asym-<sup>474</sup>  
 488 metries in mass and for high total mass binaries, in this  
 489 study we extend our parameter space to mass ratios as  
 490 low as  $q = 1/10$  and total masses as high as  $M = 800 M_{\odot}$ <sup>475</sup>  
 491 (see Appendix C Table II for a detailed list). In general,<sup>476</sup>  
 492 NR simulations can include modes of arbitrary  $(\ell, m)$  for  
 493 a given set of masses in the parameter space. However,<sup>477</sup>  
 494 weak modes are sometimes dominated by numerical noise<sup>478</sup>  
 495 and do not agree when compared across different numer-<sup>479</sup>  
 496 ical codes. In fact, we disregard numerical modes with<sup>480</sup>  
 497  $\ell \geq 5$ , because these have comparatively small ampli-<sup>481</sup>  
 498 tudes. The  $\ell \geq 5$  modes with  $m = \pm \ell$  have also particu-<sup>482</sup>  
 499 larly short wavelengths, which makes it more challenging<sup>483</sup>  
 500 for numerical relativity codes to resolve the propagation<sup>484</sup>  
 501 of these modes away from the binary. In order to assess<sup>485</sup>  
 502 the accuracy of the remaining modes, we only choose<sup>486</sup>  
 503 IMBHB simulations for which higher modes have been<sup>487</sup>  
 504 computed by at least two different NR codes. We select<sup>488</sup>  
 505 only those higher modes that agree to an overlap of at<sup>489</sup>  
 506 least 0.97 across all available NR codes for each of 17 the<sup>490</sup>  
 507 selected simulations.<sup>6</sup>

508 The higher modes that passed this criteria and were<sup>492</sup>  
 509 included in our analysis were the following:  $(\ell, m) =$ <sup>493</sup>  
 510  $\{(2, \pm 1), (2, \pm 2), (3, \pm 2), (3, \pm 3), (4, \pm 2), (4, \pm 3), (4, \pm 4)\}$ .<sup>494</sup>  
 511 Notably, the  $(2, 2)$  mode agrees across NR codes to an<sup>495</sup>  
 512 overlap  $> 0.995$  for every IMBHB source considered in<sup>496</sup>  
 513 this study; the two modes closest to the 0.97 overlap<sup>497</sup>  
 514 threshold were the  $(4, 4)$  and  $(4, 3)$  modes. We note<sup>498</sup>  
 515 that, similar to what was described in Ref. [68], omission<sup>499</sup>  
 516 of  $\ell > 4$  modes may lead to an underestimation of the<sup>500</sup>  
 517 power within the detector band radiated by the largest<sup>501</sup>  
 518 mass binary BHs.<sup>502</sup>


519 Of the 17 selected sources, we include three cases with<sup>503</sup>  
 520 spins aligned or anti-aligned with the total angular mo-<sup>504</sup>  
 521 mentum of the binary with dimensionless spin magni-<sup>505</sup>  
 522 tudes  $|\chi_{1,2}| = 0.8$ . The IMBHB injections are uni-<sup>506</sup>  
 523 formly distributed in the binary orientation parameters<sup>507</sup>  
 524  $(\varphi, \cos(\iota))$ , uniformly distributed in co-moving volume up<sup>508</sup>

### 525 B. Sensitive distance reach and merger rate 526 density estimate

527 We use the loudest event method [90] to calculate the  
 528 sensitive distance reach of our search and to place up-  
 529 per limits on the merger rate density of IMBHBs (see  
 530 Appendix C for a detailed description of our procedure).  
 531 The results of our combined search are reported using a  
 532 combined  $p$ -value of  $\bar{P} = 0.36$  given by that of our loudest  
 533 event in Table I.

534 The left panel of Figure 1 shows the sensitive distance  
 535 reach of our combined search toward our 17 targeted  
 536 IMBHB sources represented in the  $m_1 - m_2$  plane (see  
 537 Table II in Appendix C for a more detailed description).  
 538 We find an across-the-board improvement in the sensi-  
 539 tive distance of our combined search compared to the  
 540 12 targeted sources reported in Ref. [42]. In partic-  
 541 ular, we find that the combined search is most sensitive  
 542 to the  $(100 + 100) M_{\odot}$  aligned-spin source, which can  
 543 be observed up to 1.8 Gpc and is an increase of more  
 544 than 10% compared to the 1.6 Gpc obtained in Ref. [42].  
 545 These improvements are the result of better detector sen-  
 546 sitivity, the inclusion of higher modes in our injections,  
 547 and significant improvements to the cWB search algo-  
 548 rithm. As a general trend, our reach decreases for in-  
 549 creasing mass ratio and for increasing total mass once  
 550 this surpasses  $\sim 200 M_{\odot}$ . There are several reasons for  
 551 this behaviour. First, the intrinsic amplitude of IMBHB  
 552 signal decreases as the mass ratio decreases for a fixed to-  
 553 tal mass. Second, sources with small  $q$  have a significant  
 554 fraction of their power contained in their higher modes.  
 555 Consequently, they are not well matched by our search  
 556 templates, which only include the dominant quadrupole  
 557 mode. Last, although the intrinsic luminosity of IMBHBs  
 558 rises with total mass, the merger frequency decreases,  
 559 and so signals persist in the detector sensitive frequency  
 560 band for a very short duration. This makes it difficult  
 561 to distinguish them from noise transients. This effect  
 562 is evident from the roughly equivalent sensitive distances  
 563 obtained for the  $(60 + 60) M_{\odot}$  and  $(100 + 100) M_{\odot}$  sources

564 <sup>6</sup> We did allow for overlaps below 0.97 if the mode's contribution<sup>512</sup>  
 565 to the waveform was negligible.



**Figure 1.** The sensitive distance reach ( $D_{\langle VT \rangle_{\text{sen}}}$ ) in Gpc (left-panel) and 90% upper limit on merger rate density ( $R_{90\%}$ ) in  $\text{Gpc}^{-3} \text{yr}^{-1}$  (right-panel) for the 17 targeted IMBHB sources in the  $m_1 - m_2$  plane. Each circle corresponds to one class of IMBHBs in the source frame with a number in the circle indicating  $D_{\langle VT \rangle_{\text{sen}}}$  (left-panel) or  $R_{90\%}$  (right-panel). Spinning injection sets are labelled and shown as displaced circles. The blue and light blue shaded regions mark the template space encompassed by the GstLAL ( $M < 400 M_\odot$ ) and PyCBC ( $M < 500 M_\odot$ ) template banks respectively.

despite the significantly larger total mass of the latter.  
 The right panel of Figure 1 shows the upper limits on the merger rate density of our 17 targeted IMBHB sources, which improve on those reported after O1 in Ref. [42]. We set our most stringent upper limit at  $0.20 \text{ Gpc}^{-3} \text{yr}^{-1}$  for equal-mass spin-aligned IMBHBs with component masses of  $100 M_\odot$  and aligned dimensionless spins of 0.8. By assuming a redshift-independent globular cluster (GC) density of  $3 \text{ GC Mpc}^{-3}$  [95], we find that this upper limit is equivalent to  $0.07 \text{ GC}^{-1} \text{Gyr}^{-1}$ , an improvement of a factor of  $\sim 5$  over the  $0.31 \text{ GC}^{-1} \text{Gyr}^{-1}$  that was reported in Ref. [42]. We also observe that for all equal mass ratio IMBHB sources, the merger rate density upper limits are also improved. The sources with unequal masses show larger improvement in the merger rate density as compared to previous result.

the Advanced LIGO and Advanced Virgo detectors. No IMBHB detections were made in this search. The loudest candidate event was found with a marginal  $p$ -value  $\bar{P} = 0.36$  in our combined search. A detailed detector characterization study of this event suggested that it is likely explained by the detector noise.

Given the null detection, we place upper limits on the merger rate density for 17 IMBHB systems. For estimation of the rate upper limits, we use NR waveforms provided by the SXS, RIT, and Georgia Tech groups that include higher modes in the gravitational-wave emission. The reported rate limits are significantly more stringent than the previous result reported in Ref. [42]. In particular, the most stringent rate limit of  $0.20 \text{ Gpc}^{-3} \text{yr}^{-1}$  placed on  $(100+100) M_\odot$  aligned spin IMBHB systems is an improvement of a factor of  $\sim 5$ . This improvement is due to the combination of three factors: the increased sensitivity of our detector network, the improvements in the cWB search algorithm, and the incorporation of higher modes into our models for IMBHB signals.

We have conducted a search for IMBHBs in the data collected in the two observing runs of the Advanced LIGO and Virgo detectors. This search combined three analysis pipelines: two matched-filter algorithms GstLAL and PyCBC and the model-independent algorithm cWB. The PyCBC and cWB analyses use data from the Advanced LIGO detectors, and GstLAL uses data from

Anticipated increases of the network sensitivity in future runs, particularly at low frequency, and further improvement of the search algorithms will place more stringent upper limits on the merger rate density of IMBHBs and may even result in the first definitive detection of an IMBH.

565

## ACKNOWLEDGMENTS

619

620

The authors gratefully acknowledge the support of the United States National Science Foundation (NSF) for the construction and operation of the LIGO Laboratory and Advanced LIGO as well as the Science and Technology Facilities Council (STFC) of the United Kingdom, the Max-Planck-Society (MPS), and the State of Niedersachsen/Germany for support of the construction of Advanced LIGO and construction and operation of the GEO600 detector. Additional support for Advanced LIGO was provided by the Australian Research Council. The authors gratefully acknowledge the Italian Istituto Nazionale di Fisica Nucleare (INFN), the French Centre National de la Recherche Scientifique (CNRS) and the Foundation for Fundamental Research on Matter supported by the Netherlands Organisation for Scientific Research, for the construction and operation of the Virgo detector and the creation and support of the EGO consortium. The authors also gratefully acknowledge research support from these agencies as well as by the Council of Scientific and Industrial Research of India, the Department of Science and Technology, India, the Science & Engineering Research Board (SERB), India, the Ministry of Human Resource Development, India, the Spanish Agencia Estatal de Investigación, the Vicepresidència i Conselleria d'Innovació, Recerca i Turisme and the Conselleria d'Educació i Universitat del Govern de les Illes Balears, the Conselleria d'Educació, Investigació, Cultura i Esport de la Generalitat Valenciana, the National Science Centre of Poland, the Swiss National Science Foundation (SNSF), the Russian Foundation for Basic Research, the Russian Science Foundation, the European Commission, the European Regional Development Funds (ERDF), the Royal Society, the Scottish Funding Council, the Scottish Universities Physics Alliance, the Hungarian Scientific Research Fund (OTKA), the Lyon Institute of Origins (LIO), the Paris Île-de-France Region, the National Research, Development and Innovation Office Hungary (NKFIH), the National Research Foundation of Korea, Industry Canada and the Province of Ontario through the Ministry of Economic Development and Innovation, the Natural Science and Engineering Research Council Canada, the Canadian Institute for Advanced Research, the Brazilian Ministry of Science, Technology, Innovations, and Communications, the International Center for Theoretical Physics South American Institute for Fundamental Research (ICTP-SAIFR), the Research Grants Council of Hong Kong, the National Natural Science Foundation of China (NSFC), the Leverhulme Trust, the Research Corporation, the Ministry of Science and Technology (MOST), Taiwan and the Kavli Foundation. The authors gratefully acknowledge the support of the NSF, STFC, MPS, INFN, CNRS and the State of Niedersachsen/Germany for provision of computational resources.

## Appendix A: Sensitive Distance reach and merger rate

In this appendix we provide further details on our method to compute the averaged spacetime volume observed by a search and its corresponding sensitive distance at a given significance threshold. In general, the averaged spacetime volume to which our searches are sensitive is given by [96, 97]:

$$\langle VT \rangle_{\text{sen}} = T_0 \int dz \, d\boldsymbol{\theta} \, \frac{dV_c}{dz} \, \frac{1}{1+z} \, s(\boldsymbol{\theta}) \, f(z, \boldsymbol{\theta}). \quad (\text{A1})$$

Here,  $T_0$  is the length of the observation in the detector frame, and  $V_c(z)$  is the co-moving volume spanned by a sphere of redshift  $z$ . The function  $s(\boldsymbol{\theta})$  is the distribution of binary parameters  $\boldsymbol{\theta}$ , and  $0 \leq f(z, \boldsymbol{\theta}) \leq 1$ , where  $f(z, \boldsymbol{\theta})$  denotes the fraction of injections with parameters  $\boldsymbol{\theta}$  detected at a redshift  $z$ .

In this determination of sensitivity we have two main limitations. First, the true population of IMBHs in the Universe is unknown, so it prevents us from choosing a particular function  $s(\boldsymbol{\theta})$ . Second, numerical relativity waveforms cover a discrete parameter space in  $\boldsymbol{\theta}$ . For this reason, our study is focused on probing a discrete set of IMBH classes with parameters  $\{\theta_i\}$ , described in Table II. Then the averaged space-time volume sensitivity of Eq. A1 can be approximated using a Monte-Carlo technique via

$$\langle VT \rangle_{\text{sen}} \sim \frac{N_{\text{rec}}}{N_{\text{tot}}} \langle VT \rangle_{\text{tot}}. \quad (\text{A2})$$

Here,  $N_{\text{tot}}$  is the total number of injections in a given set, which are distributed in redshift and source orientations as indicated in Sec. IIIA.  $\langle VT \rangle_{\text{tot}}$  is the total spacetime volume into which injections were distributed.  $N_{\text{rec}}$  is the number of recovered injections by the search, i.e. the number of injections assigned a value  $\bar{P} \leq \bar{P}_0$ , where  $\bar{P}_0$  is in general some arbitrary threshold. In our case, we set  $\bar{P}_0 = 0.36$ , which is the  $\bar{P}$  of our most significant event in our combined search.

The corresponding sensitive distance reach is computed as

$$D_{\langle VT \rangle_{\text{sen}}} = \left( \frac{3 \langle VT \rangle_{\text{sen}}}{4\pi T_a} \right)^{1/3}, \quad (\text{A3})$$

where  $T_a$  is the amount of time analyzed by the search. We estimated the 90 % confidence upper limit in the merger rate density for selected simulated signal classes as given by

$$R_{90\%} = -\frac{\ln(0.1)}{\langle VT \rangle_{\text{sen}}}, \quad (\text{A4})$$

where  $\langle VT \rangle_{\text{sen}}$  is estimated using the loudest event method and Equation A2.

## Appendix B: Determining the p-value of the combined search

In general, the  $p$ -value of the triggers of our combined search is given by

$$\bar{P} = 1 - (1 - P_{\min})^m, \quad (B1)$$

where  $P_{\min}$  is the minimum  $p$ -value reported by any of our three searches, and  $m \in [1, 3]$  is the trials factor. The trials factor is  $m = 1$  if the three searches are fully correlated (for instance, if they are the same search) and three if they are fully independent. In this work we adopt a conservative approach and choose  $m = 3$ , omitting possible correlations between the three analysis pipelines. Indeed, excluding the eleven detected GWs mentioned in Sec. IID, none of the 123 events with  $\text{FAR} < 100/\text{yr}$  was common across the three pipelines.

We note that while the significance of individual triggers depends on our particular choice for the trials factor  $m$  applied in Eq. B1 (which we set to  $m = 3$ ),  $N_{\text{rec}}$  is independent of this choice. This is because every GW candidate output by the three analyses, including our loudest event, will have the same trials factor applied when combined into a single list, so that their relative ranking will remain unchanged (see Sec. IIC). Therefore, the numerical value of our upper limit is unaffected by our conservative choice of  $m = 3$ , since any choice would yield the same  $N_{\text{rec}}$  and  $\langle VT \rangle_{\text{sen}}$ .

As pointed out, since our choice of the trials factor affects the significance of individual triggers, our conservative approach may overly diminish the significance of prospective louder IMBHB triggers, and it may become important to make more accurate estimates of  $m$  in the future. Since the lowest  $p$ -value reported by any of our individual analyses was  $P = 0.14$ , we conclude that our choice of  $m$  does not impact our conclusion that no IMB-

HBs have been observed.

## Appendix C: Sensitive distance reach for individual search algorithms

In this appendix, we report and compare the sensitive distance reach of the three individual searches and the combined search at their respective loudest event thresholds (see Table II). For the case of the individual searches, this threshold is set to  $\bar{P}_0 = 0.14$ , equal to the loudest (most significant) event found by cWB; for the combined search, this is set to  $\bar{P}_0 = 0.36$ . We control for differences in the amount of analyzed time by only considering common observing times in Table II, which allows for a more direct comparison between the searches.

Table II shows that cWB reports the largest sensitive distance reach to every IMBHB source considered. This is expected for sources with  $M_{\text{tot}} > 500 M_{\odot}$ , since the GstLAL and PyCBC template banks are bounded by a total mass  $400 M_{\odot}$  and  $500 M_{\odot}$ , respectively. Additionally, since cWB is not limited by constraints on waveform morphology, it significantly outperforms matched-filter analyses in the large mass and small mass ratio regions of the parameter space that are covered by our analyses' template banks. This finding is consistent with Ref. [68], since in that region of parameter space, signals are shorter and higher modes are more important. Ref. [68] also found that matched-filter searches outperform cWB in the low mass end of our parameter space. Since then, however, cWB has undergone major improvements that have led to a sensitivity comparable to that of matched filter searches even for the lightest equal-mass systems considered in this analysis.

GstLAL reports sensitive distance reaches that are lower than those found in Ref. [42]. This is the result of using a large bank here that was not specifically tuned and targeted for IMBHBs. Future searches will benefit from investigations into optimal template placement and binning as well as a return to a dedicated IMBH bank.

## Appendix D: Loudest event parameter estimation

Despite the low significance of our loudest event, two characteristics motivated a detailed followup analysis. On the one hand, initial parameter estimation put this trigger in the IMBHB region of the parameter space. On the other, this trigger was observed by our matched filter analyses with an SNR of only  $\sim 6$ , much lower than that recovered by cWB. If this were a real GW, this difference might be indicative that the signal contained physics that our search templates omit (such as precession and higher modes), which would lead to a reduction of its SNR and significance.

To explore this possibility, we ran standard parameter

estimation on this event using the same approximants used in Ref. [8], namely SEOBNRv4 [84] and IMRPhenomPv2 [98]. Note that the latter approximant includes the effects of precession that our search templates omit. For the precessing IMRPhenomP run, we assumed a spin magnitude prior uniform between 0 and 0.99, and spin orientations were isotropically distributed on the sphere; for the spin-aligned SEOBNR waveforms, we used a spin prior such that the components of the spin aligned with the orbital angular momentum matched the prior used for the IMRPhenomP analysis. Remarkably, the two analyses not only report broadly consistent parameter posterior distributions but they also report consistent SNRs of  $\sim 6$ , in agreement with that reported by our

$m_1$ $M_\odot$	$m_2$ $M_\odot$	spin $\chi_{1,2}$	$M$ $M_\odot$	NR-simulation	$D_{\langle VT \rangle_{\text{sen}}} \text{ (Gpc)}$			
					cWB	GstLAL	PyCBC	combined
60	60	0	120	SXS:BBH:0180, RIT:BBH:0198:n140, GT:0905	1.2	1.2	1.2	1.3
60	60	0.8	120	SXS:BBH:0230, RIT:BBH:0063:n100, GT:0424	1.6	1.0	1.5	1.6
100	20	0	120	SXS:BBH:0056, RIT:BBH:0120:n140, GT:0906	0.72	0.69	0.70	0.76
100	50	0	150	SXS:BBH:0169, RIT:BBH:0117:n140, GT:0446	1.2	0.79	1.1	1.2
100	100	-0.8	200	SXS:BBH:0154, RIT:BBH:0068:n100	1.1	1.0	0.99	1.2
100	100	0	200	SXS:BBH:0180, RIT:BBH:0198:n140, GT:0905	1.4	0.90	1.3	1.4
100	100	0.8	200	SXS:BBH:0230, RIT:BBH:0063:n100, GT:0424	1.8	1.2	1.7	1.8
200	20	0	220	RIT:BBH:Q10:n173, GT:0568	0.48	0.30	0.36	0.49
200	50	0	250	SXS:BBH:0182, RIT:BBH:0119:n140, GT:0454	0.85	0.48	0.67	0.87
200	100	0	300	SXS:BBH:0169, RIT:BBH:0117:n140, GT:0446	1.1	0.59	0.86	1.1
300	50	0	350	SXS:BBH:0181, RIT:BBH:0121:n140, GT:0604	0.55	0.18	0.27	0.56
200	200	0	400	SXS:BBH:0180, RIT:BBH:0198:n140, GT:0905	1.0	0.47	0.72	1.0
300	100	0	400	SXS:BBH:0030, RIT:BBH:0102:n140, GT:0453	0.78	0.23	0.34	0.78
400	40	0	440	RIT:BBH:Q10:n173, GT:0568	0.35	0.10	0.16	0.35
300	200	0	500	RIT:BBH:0115:n140, GT:0477	0.79	0.16	0.14	0.79
300	300	0	600	SXS:BBH:0180, RIT:BBH:0198:n140, GT:0905	0.61	0.09	0.18	0.61
400	400	0	800	SXS:BBH:0180, RIT:BBH:0198:n140, GT:0905	0.31	0.10	0.23	0.31

TABLE II: The sensitive distance reach and the merger rate density calculated for the 17 targeted IMBHB sources considered in this study, whose intrinsic parameters are indicated in the first four columns. The fifth column indicates the numerical simulations used for each case, following the naming conventions of the corresponding NR groups. The next three columns report the sensitive distance reach for each of the individual analyses (cWB, GstLAL, and PyCBC), where we use the loudest event threshold of  $P = 0.14$  for each analysis for comparison purposes. The last column gives the sensitive distance reach from the combined search. To control for differences in the amount of analyzed time between individual analyses, we consider only common observed time across the three pipelines time, which yields  $T_a = 0.428$  years.

matched filter searches. The latter indicates that the low SNR obtained by our matched filter searches is not likely due to lack of precession in our templates. Assuming this event is a compact binary, we recover a source-frame chirp mass of  $70^{+24}_{-20} M_\odot$ , a source-frame total mass of  $171^{+68}_{-48} M_\odot$ , an effective inspiral spin of  $0.19^{+0.44}_{-0.46}$ , and a luminosity distance of  $7.0^{+8.0}_{-4.2}$  Gpc. We also note that, given the lack of information about the spins, spin results are sensitive to the choice of prior. Further parameter estimation was performed using the new SEOBNRv4HM [99] approximant, which includes the impact of higher order modes. The resulting consistent parameter posteriors and no increase of the SNR, suggesting that the low SNR obtained by our matched filter searches is not likely due to lack of higher modes in our templates either.

We further conducted parameter estimation of this

trigger by directly using numerical relativity waveforms of generic spin configurations and higher-modes with the RIFT algorithm [100, 101], which reported results consistent with those obtained by our waveform approximants. In addition, the event was also reconstructed using the model agnostic algorithm BayesWave [102, 103], which reported an SNR consistent with those obtained by our templates.

In summary, detailed followup of this event suggests that, in the most optimistic scenario, this trigger would be the combination of a weak IMBHB signal plus a noise transient with power detected by cWB (see Sec. II D), raising the significance of the underlying IMBHB signal. Since the resulting event has a marginal significance, the underlying IMBHB trigger would be even less significant. Hence, we conclude that this event is best explained by detector noise.

- [2] B. P. Abbott *et al.* (LIGO Scientific Collaboration, Virgo Collaboration), Phys. Rev. Lett. **116**, 241103 (2016), arXiv:1606.04855 [gr-qc].
- [3] B. P. Abbott *et al.* (LIGO Scientific Collaboration, Virgo Collaboration), Phys. Rev. Lett. **118**, 221101 (2017), arXiv:1706.01812 [gr-qc].
- [4] B. P. Abbott *et al.* (LIGO Scientific Collaboration, Virgo Collaboration), Astrophys. J. **851**, L35 (2017), arXiv:1711.05578 [astro-ph.HE].
- [5] B. Abbott *et al.* (LIGO Scientific Collaboration, Virgo Collaboration), Phys. Rev. Lett. **119**, 161101 (2017), arXiv:1710.05832 [gr-qc].
- [6] B. P. Abbott *et al.* (LIGO Scientific Collaboration, Virgo Collaboration), Phys. Rev. Lett. **119**, 141101 (2017), arXiv:1709.09660 [gr-qc].
- [7] B. P. Abbott *et al.* (LIGO Scientific Collaboration, Virgo Collaboration), Phys. Rev. **X6**, 041015 (2016), [erratum: Phys. Rev.X8,no.3,039903(2018)], arXiv:1606.04856 [gr-qc].
- [8] B. P. Abbott *et al.* (LIGO Scientific Collaboration and Virgo Collaboration), (2018), arXiv:1811.12907 [astro-ph.HE].
- [9] B. P. Abbott *et al.* (LIGO Scientific Collaboration and Virgo Collaboration), (2018), arXiv:1811.12940 [astro-ph.HE].
- [10] M. Spera and M. Mapelli, Mon. Not. Roy. Astron. Soc. **470**, 4739 (2017), arXiv:1706.06109 [astro-ph.SR].
- [11] S. E. Woosley, Astrophys. J. **836**, 244 (2017), arXiv:1608.08939 [astro-ph.HE].
- [12] N. Giacobbo, M. Mapelli, and M. Spera, Mon. Not. Roy. Astron. Soc. **474**, 2959 (2018), arXiv:1711.03556 [astro-ph.SR].
- [13] K. Belczynski *et al.*, Astron. Astrophys. **594**, A97 (2016), arXiv:1607.03116 [astro-ph.HE].
- [14] M. Mezcua, International Journal of Modern Physics D **26**, 1730021 (2017), arXiv:1705.09667.
- [15] F. Koliopanos, in *Proceedings of the XII Multifrequency Behaviour of High Energy Cosmic Sources Workshop*. 12-17 June (2017) p. 51, arXiv:1801.01095 [astro-ph.GA].
- [16] B. Kiziltan, H. Baumgardt, and A. Loeb, “An intermediate-mass black hole in the centre of the globular cluster 47 tucanae.”.
- [17] den Brok *et al.*, The Astrophysical Journal **809** (2015).
- [18] M. Atakan Gurkan, M. Freitag, and F. A. Rasio, Astrophys. J. **604**, 632 (2004), arXiv:astro-ph/0308449 [astro-ph].
- [19] J. Kormendy and L. C. Ho, Ann. Rev. Astron. Astrophys. **51**, 511 (2013), arXiv:1304.7762 [astro-ph.CO].
- [20] A. W. Graham, Astrophys. J. **746**, 113 (2012), arXiv:1202.1878 [astro-ph.CO].
- [21] A. W. Graham and N. Scott, Astrophys. J. **764**, 151 (2013), arXiv:1211.3199.
- [22] R. van den Bosch, T. de Zeeuw, K. Gebhardt, E. Noyola, and G. van de Ven, Astrophys. J. **641**, 852 (2006), arXiv:astro-ph/0512503 [astro-ph].
- [23] R. P. van der Marel and J. Anderson, Astrophys. J. **710**, 1063 (2010), arXiv:0905.0638 [astro-ph.GA].
- [24] J. Anderson and R. P. van der Marel, Astrophys. J. **710**, 1032 (2010), arXiv:0905.0627 [astro-ph.GA].
- [25] H. Baumgardt, “N-body modeling of globular clusters: Masses, mass-to-light ratios and intermediate-mass black holes.”.
- [26] H. Baumgardt, P. Hut, J. Makino, S. McMillan, and S. F. Portegies Zwart, Astrophys. J. **582**, L21 (2003), arXiv:astro-ph/0210133 [astro-ph].
- [27] R. A. Remillard and J. E. McClintock, Annual Review of Astronomy and Astrophysics **44**, 49 (2006), <https://doi.org/10.1146/annurev.astro.44.051905.092532>.
- [28] D. R. Pasham, T. E. Strohmayer, and R. F. Mushotzky, Nature **513**, 74 (2015), arXiv:1501.03180 [astro-ph.HE].
- [29] P. Kaaret, H. Feng, and T. P. Roberts, Ann. Rev. Astron. Astrophys. **55**, 303 (2017), arXiv:1703.10728 [astro-ph.HE].
- [30] S. Farrell, N. Webb, D. Barret, O. Godet, and J. Rodrigues, Nature **460**, 73 (2009), arXiv:1001.0567 [astro-ph.HE].
- [31] P. Madau and M. J. Rees, Astrophys. J. **551**, L27 (2001), arXiv:astro-ph/0101223 [astro-ph].
- [32] R. Schneider, A. Ferrara, P. Natarajan, and K. Omukai, Astrophys. J. **571**, 30 (2002), arXiv:astro-ph/0111341 [astro-ph].
- [33] T. Ryu, T. L. Tanaka, R. Perna, and Z. Haiman, Mon. Not. Roy. Astron. Soc. **460**, 4122 (2016), arXiv:1603.08513 [astro-ph.GA].
- [34] T. Kinugawa, K. Inayoshi, K. Hotokezaka, D. Nakauchi, and T. Nakamura, Mon. Not. Roy. Astron. Soc. **442**, 2963 (2014), arXiv:1402.6672 [astro-ph.HE].
- [35] M. C. Miller and D. P. Hamilton, Mon. Not. Roy. Astron. Soc. **330**, 232 (2002), arXiv:astro-ph/0106188 [astro-ph].
- [36] S. F. Portegies Zwart, H. Baumgardt, P. Hut, J. Makino, and S. L. W. McMillan, Nature **428**, 724 (2004), arXiv:astro-ph/0402622 [astro-ph].
- [37] S. F. Portegies Zwart and S. L. W. McMillan, Astrophys. J. **576**, 899 (2002), arXiv:astro-ph/0201055 [astro-ph].
- [38] M. Mapelli, Mon. Not. Roy. Astron. Soc. **459**, 3432 (2016), arXiv:1604.03559 [astro-ph.GA].
- [39] N. W. C. Leigh, N. Lützgendorf, A. M. Geller, T. J. Maccarone, C. Heinke, and A. Sesana, Mon. Not. Roy. Astron. Soc. **444**, 29 (2014), arXiv:1407.4459 [astro-ph.SR].
- [40] P. Marchant, N. Langer, P. Podsiadlowski, T. M. Tauris, and T. J. Moriya, Astron. Astrophys. **588**, A50 (2016), arXiv:1601.03718 [astro-ph.SR].
- [41] A. R. King and W. Dehnen, Mon. Not. Roy. Astron. Soc. **357**, 275 (2005), arXiv:astro-ph/0411489 [astro-ph].
- [42] B. P. Abbott *et al.* (LIGO Scientific Collaboration, Virgo Collaboration), Phys. Rev. **D96**, 022001 (2017), arXiv:1704.04628 [gr-qc].
- [43] B. P. Abbott *et al.* (LIGO Scientific Collaboration, Virgo Collaboration), Phys. Rev. Lett. **116**, 221101 (2016), [Erratum: Phys. Rev. Lett.121,no.12,129902(2018)], arXiv:1602.03841 [gr-qc].
- [44] N. Yunes, K. Yagi, and F. Pretorius, Phys. Rev. **D94**, 084002 (2016), arXiv:1603.08955 [gr-qc].
- [45] N. Yunes and X. Siemens, Living Rev. Rel. **16**, 9 (2013), arXiv:1304.3473 [gr-qc].
- [46] J. R. Gair, M. Vallisneri, S. L. Larson, and J. G. Baker, Living Rev. Rel. **16**, 7 (2013), arXiv:1212.5575 [gr-qc].
- [47] I. Kamaretsos, M. Hannam, S. Husa, and B. S. Sathyaprakash, Phys. Rev. **D85**, 024018 (2012), arXiv:1107.0854 [gr-qc].
- [48] J. Meidam, M. Agathos, C. Van Den Broeck, J. Veitch, and B. S. Sathyaprakash, Phys. Rev. **D90**, 064009 (2014), arXiv:1406.3201 [gr-qc].
- [49] E. Thrane, P. D. Lasky, and Y. Levin, Phys. Rev. **D96**,

- 102004 (2017), arXiv:1706.05152 [gr-qc].
- [50] G. Carullo *et al.*, Phys. Rev. **D98**, 104020 (2018), arXiv:1805.04760 [gr-qc].
- [51] J. A. Gonzalez, U. Sperhake, B. Bruegmann, M. Hannam, and S. Husa, Phys. Rev. Lett. **98**, 091101 (2007), arXiv:gr-qc/0610154 [gr-qc].
- [52] M. Campanelli, C. O. Lousto, Y. Zlochower, and D. Merritt, Phys. Rev. Lett. **98**, 231102 (2007), arXiv:gr-qc/0702133 [GR-QC].
- [53] J. Calderón Bustillo, J. A. Clark, P. Laguna, and D. Shoemaker, Phys. Rev. Lett. **121**, 191102 (2018), arXiv:1806.11160 [gr-qc].
- [54] J. Abadie *et al.* (LIGO Scientific Collaboration, Virgo Collaboration), Phys. Rev. **D85**, 102004 (2012), arXiv:1201.5999 [gr-qc].
- [55] J. Aasi *et al.* (LIGO Scientific Collaboration and Virgo Collaboration), Phys. Rev. **D89**, 122003 (2014), arXiv:1404.2199 [gr-qc].
- [56] S. Klimenko, S. Mohanty, M. Rakhmanov, and G. Mitselmakher, Phys. Rev. **D72**, 122002 (2005), arXiv:gr-qc/0508068 [gr-qc].
- [57] J. Aasi *et al.* (LIGO Scientific Collaboration), Class. Quant. Grav. **32**, 074001 (2015), arXiv:1411.4547 [gr-qc].
- [58] F. Acernese *et al.* (VIRGO), Class. Quant. Grav. **32**, 024001 (2015), arXiv:1408.3978 [gr-qc].
- [59] K. Cannon *et al.*, Astrophys. J. **748**, 136 (2012), arXiv:1107.2665 [astro-ph.IM].
- [60] S. Privitera, S. R. P. Mohapatra, P. Ajith, K. Cannon, N. Fotopoulos, M. A. Frei, C. Hanna, A. J. Weinstein, and J. T. Whelan, Phys. Rev. **D89**, 024003 (2014), arXiv:1310.5633 [gr-qc].
- [61] C. Messick *et al.*, Phys. Rev. **D95**, 042001 (2017), arXiv:1604.04324 [astro-ph.IM].
- [62] A. Taracchini, Y. Pan, A. Buonanno, E. Barausse, M. Boyle, T. Chu, G. Lovelace, H. P. Pfeiffer, and M. A. Scheel, Phys. Rev. **D86**, 024011 (2012), arXiv:1202.0790 [gr-qc].
- [63] L. Pekowsky, J. Healy, D. Shoemaker, and P. Laguna, Phys. Rev. **D87**, 084008 (2013), arXiv:1210.1891 [gr-qc].
- [64] V. Varma, P. Ajith, S. Husa, J. C. Bustillo, M. Hannam, and M. Pürrer, Phys. Rev. **D90**, 124004 (2014), arXiv:1409.2349 [gr-qc].
- [65] J. Calderón Bustillo, S. Husa, A. M. Sintes, and M. Pürrer, Phys. Rev. **D93**, 084019 (2016), arXiv:1511.02060 [gr-qc].
- [66] V. Varma and P. Ajith, (2016), arXiv:1612.05608 [gr-qc].
- [67] J. Calderón Bustillo, P. Laguna, and D. Shoemaker, Phys. Rev. **D95**, 104038 (2017), arXiv:1612.02340 [gr-qc].
- [68] J. Calderón Bustillo, F. Salemi, T. Dal Canton, and K. Jani, (2017), arXiv:1711.02009 [gr-qc].
- [69] T. Dal Canton *et al.*, Phys. Rev. **D90**, 082004 (2014), arXiv:1405.6731 [gr-qc].
- [70] S. A. Usman *et al.*, Class. Quant. Grav. **33**, 215004 (2016), arXiv:1508.02357 [gr-qc].
- [71] C. Cahillane *et al.* (LIGO Scientific), Phys. Rev. **D96**, 102001 (2017), arXiv:1708.03023 [astro-ph.IM].
- [72] A. Viets *et al.*, Class. Quant. Grav. **35**, 095015 (2018), arXiv:1710.09973 [astro-ph.IM].
- [73] F. Acernese *et al.* (Virgo), Class. Quant. Grav. **35**, 205004 (2018), arXiv:1807.03275 [gr-qc].
- [74] J. C. Driggers *et al.* (LIGO Scientific Collaboration), (2018), 1806.00532 [astro-ph.IM].
- [75] S. Sachdev, S. Caudill, H. Fong, R. K. Lo, C. Messick, D. Mukherjee, R. Magee, L. Tsukada, K. Blackburn, P. Brady, *et al.*, arXiv preprint arXiv:1901.08580 (2019).
- [76] S. Klimenko *et al.*, Phys. Rev. **D93**, 042004 (2016), arXiv:1511.05999 [gr-qc].
- [77] L. Wainstein and V. Zubakov, Prentice-Hall, Englewood Cliffs (1962).
- [78] C. Capano, Y. Pan, and A. Buonanno, Phys. Rev. **D89**, 102003 (2014), arXiv:1311.1286 [gr-qc].
- [79] M. Cabero *et al.*, (2019), arXiv:1901.05093 [physics.ins-det].
- [80] T. Dal Canton, S. Bhagwat, S. Dhurandhar, and A. Lundgren, Class. Quant. Grav. **31**, 015016 (2014), arXiv:1304.0008 [gr-qc].
- [81] A. H. Nitz, Class. Quant. Grav. **35**, 035016 (2018), arXiv:1709.08974 [gr-qc].
- [82] B. Allen, Phys. Rev. **D71**, 062001 (2005), arXiv:gr-qc/0405045 [gr-qc].
- [83] D. Mukherjee, S. Caudill, R. Magee, C. Messick, S. Privitera, S. Sachdev, K. Blackburn, P. Brady, P. Brockill, K. Cannon, S. J. Chamberlin, D. Chatterjee, J. D. E. Creighton, H. Fong, P. Godwin, C. Hanna, S. Kapadia, R. N. Lang, T. G. F. Li, R. K. L. Lo, D. Meacher, A. Pace, L. Sadeghian, L. Tsukada, L. Wade, M. Wade, A. Weinstein, and L. Xiao, arXiv e-prints , arXiv:1812.05121 (2018), arXiv:1812.05121 [astro-ph.IM].
- [84] A. Bohé *et al.*, Phys. Rev. **D95**, 044028 (2017), arXiv:1611.03703 [gr-qc].
- [85] T. Dal Canton and I. W. Harry, (2017), arXiv:1705.01845 [gr-qc].
- [86] V. Necula, S. Klimenko, and G. Mitselmakher, *Gravitational waves. Numerical relativity - data analysis. Proceedings, 9th Edoardo Amaldi Conference, Amaldi 9, and meeting, NRDA 2011, Cardiff, UK, July 10-15, 2011*, J. Phys. Conf. Ser. **363**, 012032 (2012).
- [87] V. Tiwari *et al.*, Phys. Rev. **D93**, 043007 (2016), arXiv:1511.09240 [gr-qc].
- [88] L. K. Nuttall, Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences **376**, 20170286 (2018).
- [89] B. P. Abbott *et al.* (LIGO Scientific Collaboration, Virgo Collaboration), Class. Quant. Grav. **35**, 065010 (2018), arXiv:1710.02185 [gr-qc].
- [90] R. Biswas, P. R. Brady, J. D. E. Creighton, and S. Fairhurst, Class. Quant. Grav. **26**, 175009 (2009), [Erratum: Class. Quant. Grav.30,079502(2013)], arXiv:0710.0465 [gr-qc].
- [91] A. H. Mroue *et al.*, Phys. Rev. Lett. **111**, 241104 (2013), arXiv:1304.6077 [gr-qc].
- [92] J. Healy, C. O. Lousto, Y. Zlochower, and M. Campanelli, Class. Quant. Grav. **34**, 224001 (2017), arXiv:1703.03423 [gr-qc].
- [93] K. Jani, J. Healy, J. A. Clark, L. London, P. Laguna, and D. Shoemaker, Class. Quant. Grav. **33**, 204001 (2016), arXiv:1605.03204 [gr-qc].
- [94] P. A. R. Ade *et al.* (Planck), Astron. Astrophys. **594**, A13 (2016), arXiv:1502.01589 [astro-ph.CO].
- [95] S. F. Portegies Zwart and S. McMillan, Astrophys. J. **528**, L17 (2000), arXiv:astro-ph/9910061 [astro-ph].
- [96] B. P. Abbott *et al.* (LIGO Scientific Collaboration, Virgo Collaboration), Astrophys. J. Suppl. **227**, 14

- (2016), arXiv:1606.03939 [astro-ph.HE].
- [97] B. P. Abbott *et al.* (LIGO Scientific Collaboration, Virgo Collaboration), *Astrophys. J.* **833**, L1 (2016), arXiv:1602.03842 [astro-ph.HE].
- [98] M. Hannam, P. Schmidt, A. Bohé, L. Haegel, S. Husa, *et al.*, *Phys. Rev. Lett.* **113**, 151101 (2014), arXiv:1308.3271 [gr-qc].
- [99] R. Cotesta, A. Buonanno, A. Bohé, A. Taracchini, I. Hinder, and S. Ossokine, *Phys. Rev.* **D98**, 084028 (2018), arXiv:1803.10701 [gr-qc].
- [100] J. Lange *et al.*, *Phys. Rev.* **D96**, 104041 (2017), arXiv:1705.09833 [gr-qc].
- [101] J. Lange, R. O’Shaughnessy, and M. Rizzo, arXiv e-prints (2018), arXiv:1805.10457 [gr-qc].
- [102] N. J. Cornish and T. B. Littenberg, *Class. Quant. Grav.* **32**, 135012 (2015), arXiv:1410.3835 [gr-qc].
- [103] LIGO Scientific Collaboration and Virgo Collaboration, “BayesWave,” <https://git.ligo.org/lscsoft/bayeswave/>, (2018).

DRAFT-V16