



41

42 **Abstract:** The diversity-productivity relationship suggests that increasing plant species could increase  
43 primary productivity, with this effect being explained in part by the suppression of plant antagonists.  
44 We conducted a global synthesis of 609 studies to investigate how plant diversity affects plants and  
45 their antagonists. Here we show that increasing plant species consistently promotes plant performance  
46 and suppresses antagonist performance in agroecosystems, grasslands and forests, for herbaceous and  
47 woody plants, across tropical and temperate zones, and for replacement series and additive experimental  
48 design studies. Crop diversification (e.g., intercropping and cover cropping) indirectly promotes crop  
49 production through the suppression of pests. This demonstrates that diversifying planting systems can  
50 increase productivity while reducing reliance on synthetic pesticides, offering a sustainable pathway for  
51 agriculture from subsistence to large scale agriculture. Overall, these results suggest crop diversification  
52 has considerable potential to support sustainable agroecosystems that benefit productivity while  
53 reducing reliance on synthetic pesticides.

54

55

56 Monocultures have been the heart of intensive agricultural and forestry systems since before  
57 the green revolution of the 1960's<sup>1</sup>. While monocultures can be high yielding<sup>2</sup>, they have had  
58 negative consequences, including loss of co-occurring native biodiversity<sup>3</sup> with knock-on  
59 consequences for biological pest control<sup>4</sup> and crop pollination<sup>5</sup>. Monocultures are linked to the  
60 heavy use of often synthetic pesticides and fertilizers, which contribute to impacts on  
61 agroecosystem biodiversity as well as having negative consequences in terms of wider diffuse  
62 pollution<sup>6,7</sup>. Increasing plant diversity can mitigate these negative effects by promoting  
63 multiple ecosystem functions and services<sup>8-11</sup>. This can be achieved through crop diversification  
64 (e.g., intercropping, cover cropping and sown field margins) in arable and horticultural systems,  
65 diversified grasslands, and mixed forest plantations. For example, intercropping has been  
66 shown to increase crop yield under low nitrogen availability<sup>12,13</sup>, to reduce pesticide and  
67 nitrogen fertilizer reliance while maintaining gross margins<sup>14</sup>, and to decrease pests<sup>15</sup>.

68 Mechanisms such as the resource-use complementarity hypothesis, based on negative  
69 interspecific interactions (competition theory), have been proposed to explain why increasing  
70 plant species can promote primary productivity<sup>8,16</sup>. However, an increase in productivity with  
71 increasing plant diversity may be explained by complementarity in top-down control. Here, the  
72 impact of specialist consumers (e.g. pest herbivores) on plants would be lessened as, thereby

73 promoting overall productivity<sup>8</sup>. Individual experiments have demonstrated that increasing  
74 plant diversity can suppress plant antagonists, as evidenced by decreases in the abundance or  
75 biomass of insect herbivores<sup>17,18</sup>, plant-feeding nematodes<sup>19</sup> and rodents<sup>20</sup>, decreases in  
76 competitive weeds (i.e. both agricultural as well as invasive non-native plants in natural  
77 ecosystems)<sup>21</sup>, and reductions in plant disease spread or damage<sup>22,23</sup>. Where the abundance of  
78 soil pathogens is not linearly related to the density of susceptible plant species there may also  
79 be a complementarity effect whereby plant-soil feedbacks negatively affect the growth of plants  
80 to a greater extent in monocultures than in diverse communities<sup>24-26</sup>. However, opposite results  
81 have also been reported in a few studies. For example, increasing plant species was shown to  
82 increase damage from herbivorous thrips<sup>27</sup>. If antagonists are suppressed, plant productivity  
83 may be enhanced indirectly by plant species richness resulting in increased yields in production  
84 systems such as crops, timber or fibre. The suppression of antagonists by plant diversity in  
85 agroecosystems (i.e., crop diversification) has significant potential to contribute to pesticide  
86 reduction strategies by maintaining antagonist populations below the threshold for economic  
87 injury<sup>6,28</sup>. However, there remains considerable debate over how generalized these plant  
88 diversity-antagonist relationships are as we currently lack a comprehensive understanding of  
89 their context dependence<sup>29</sup>.

90 Meta-analysis is an effective tool to support evidence-based practice by resolving  
91 seemingly contradictory research outcomes across the broader literature<sup>29</sup>. Previous meta-  
92 analyses have shown that plant species diversity has variable effects on invertebrate herbivore  
93 abundance and damage<sup>30-33</sup>. They have also shown that increasing plant species diversity can  
94 decrease arthropod pests<sup>34</sup>, and increase plant productivity<sup>33</sup> and crop yield<sup>34,35</sup>. In addition,  
95 past meta-analyses focusing on agroecosystem intercropping have also identified this can  
96 increase crop yields<sup>15,36-42</sup> and decrease pests<sup>43-46</sup>, including plant diseases<sup>15,47</sup>, weeds<sup>36,48,49</sup>,  
97 and pest nematodes<sup>15</sup>. Intercropping has also been shown to increase natural enemies of  
98 arthropod pests (i.e., predators and parasitoids)<sup>43-46</sup>. However, these previous meta-analyses  
99 have only considered specific plant antagonists or pests, and did not distinguish specific types  
100 of plant diversity or experimental designs. Further, they did not consider the indirect (mediation)  
101 effects of plant antagonists on the diversity-productivity relationship and how such mediation  
102 effects can vary globally across different ecosystems, plant diversity types, experimental design  
103 types (replacement series vs. additive), plant forms (herbaceous vs. woody), climatic zones  
104 (temperate vs. tropical), or types of experimental study (plot vs. pot).

105 Here, we conducted a global synthesis of 609 biodiversity experiments from which we  
106 derived 5,712 effect size observations on the impact of plant species diversity (i.e.,  
107 intercropping, cover cropping, sown field margins, diversified grasslands, and mixed forest  
108 plantations) on plant performance. We assessed the effects of plant species diversity on a wide  
109 range of antagonist groups, i.e., invertebrate herbivores, plant diseases (bacteria, fungi, and  
110 viruses), weeds, plant-feeding nematodes, and plant-consuming rodents (Fig. 1a; Extended  
111 Data Fig. 1; Supplementary Data 1). We focused on the consequences of increasing plant  
112 species on overall measures of: (i) plant growth, reproduction, and quality; (ii) invertebrate  
113 herbivore growth, reproduction, and damage that herbivores cause to plants; (iii) disease  
114 reproduction, spread and damage; (iv) weed growth, reproduction, and diversity; (v) nematode  
115 reproduction, and damage; (vi) rodent reproduction, and damage; and (vii) aggregate  
116 performance indicators<sup>29</sup> for each of these categories. From these response variables, we  
117 derived log ratio response (lnRR) effect sizes comparing diverse plant communities (i.e., high  
118 plant species richness) and simple plant communities or monocultures (i.e., low species  
119 richness). Piecewise structural equation modeling was then used to analyze the effects of plant  
120 species diversity on plant performance as mediated through plant antagonists across different  
121 levels of plant diversity. We hypothesized that plant species richness within agricultural (arable  
122 and horticulture), grassland, and forestry ecosystems (Fig.1b): (i) increases plant performance  
123 directly through either increased complementarity or decreased competition among different  
124 plant species (H1); (ii) decreases plant antagonist performance directly (H2), and (iii) increases  
125 plant performance indirectly by reducing the performance of plant antagonists (H3). We also  
126 propose that this diversity-antagonist-productivity relationship varies across different  
127 ecosystems, plant diversity types, experimental design types, plant life forms, and climatic  
128 zones. Together, we recommend a plant antagonist hypothesis and suggest that mediation  
129 effects of plant antagonists on plants can be another mechanism by which plant diversity  
130 influences plant performance both directly and indirectly.

131

## 132 **Results and Discussion**

### 133 **Direct effect of plant species richness on plant performance**

134 Our results were based on 5,712 observations from 609 articles and confirmed the biodiversity-  
135 productivity relationship (Supplementary Tables 1–3). However, agroecosystems (arable and  
136 horticulture; N=2,583) datasets were considerably larger than those for forests (N=118) and

137 grasslands (N=65). Overall, we found that increased plant species richness promoted plant  
138 performance across global terrestrial ecosystems, for metrics including plant growth, quality,  
139 and reproduction (Fig.1c) (CI=0.148 to 0.221, df=2390, ES (effect size)=0.184, P<0.001)  
140 (Supplementary Table 3). When we split the aggregate indicator of plant performance into plant  
141 growth, reproduction and quality, we also found each showed positive responses to increasing  
142 plant species (plant growth: CI=0.165 to 0.271, df=1160, ES=0.218, P<0.001; plant  
143 reproduction: CI=0.072 to 0.169, df=956, ES=0.120, P<0.001; plant quality: CI=0.068 to 0.170,  
144 df=272, ES=0.119, P<0.001; see Supplementary Table 3). All responses of plant performance  
145 to plant diversity were positive across different ecosystem types, with the largest effect size  
146 found in grasslands (Fig. 2a) (grasslands: CI=0.271 to 0.535, df=467, ES=0.403, P<0.001;  
147 agroecosystems: CI=0.107 to 0.179, df=1784, ES=0.143, P<0.001; forests: CI=0.041 to 0.208,  
148 df=137, ES=0.124, P=0.004) (Supplementary Table 4). We explain this pattern through five  
149 mechanisms by which increasing plant species could promote productivity: i) mixed planting  
150 (e.g., intercropping but not cover cropping) resulting in a complementarity effect (i.e., direct  
151 niche partitioning or facilitation between plants)<sup>8,16</sup> that impacts trophic interactions with other  
152 non-plant species<sup>24,25</sup>. This is relevant to our H1: complementarity or decreased competition  
153 hypothesis; ii) reduced pressure of plant antagonists (e.g., herbivores<sup>18,50</sup>, plant disease<sup>22</sup> and  
154 weeds<sup>21</sup>) indirectly facilitating plant productivity<sup>4,6</sup>. This has implication for our hypotheses  
155 H2: directly decreased plant antagonist performance, and H3: indirectly decreased plant  
156 antagonist performance; iii) strengthening of the top-down effects of natural enemies on crops  
157 by providing additional resources for natural enemies supporting increased population sizes<sup>35,51</sup>;  
158 iv) improved soil fertility<sup>13</sup>, field micro-climate<sup>52</sup>, and increasing nutrient use efficiency<sup>53</sup>; and  
159 v) direct positive effect on plant-plant interactions through the release by one or more plant  
160 species in a mixture of beneficial plant secondary compounds like allelochemicals<sup>54</sup>. We also  
161 found positive responses of plant performance in response to plant diversity for herbaceous and  
162 woody plants (Fig. 2b), temperate and tropical zones (Fig. 2c), plot and pot experiments (Fig.  
163 2d), and additive design and replacement series designs (Fig. 2f).

164 When we classified plant diversity in agroecosystems (i.e., crop diversification) into  
165 intercropping, cover cropping, and sown field margins, we found that intercropping increased  
166 plant performance through its action on plant growth, quality, and reproduction (growth:  
167 CI=0.034 to 0.159, df=517, ES=0.097, P=0.003; reproduction: CI=0.061 to 0.153, df=758,  
168 ES=0.107, P<0.001; quality: CI=0.054 to 0.198, df=201, ES=0.126, P=0.001; see Fig. 2e and

169 Supplementary Table 4). Increased crop production in intercropping might be due to a more  
170 efficient use of available resources through complementarity in niches, leading to a reduced  
171 reliance on external inputs that may be important in the context of sustainable crop  
172 production<sup>55,56</sup>. Likewise, we found that cover cropping increased plant performance (growth:  
173 CI=0.103 to 0.338, df=100, ES=0.220, P<0.001; reproduction: CI=0.080 to 0.402, df=148,  
174 ES=0.241, P=0.004; quality: CI=0.175 to 0.248, df=8, ES=0.212, P<0.001). Cover crops  
175 promote the production of main crops either by acting as green manures when incorporated  
176 into soils or directly through nitrogen release in the case of legumes<sup>57</sup>.

177 Possibly due to low sample sizes, the evidence for increased plant performance in  
178 response to sown field margins was equivocal with effect sizes more or less symmetrical  
179 around zero (growth: CI=-0.458 to 0.545, df=13, ES=0.044, P=0.853; reproduction:  
180 CI=-0.010 to 0.246, df=32, ES=0.118, P=0.069; quality: N=0; see Fig. 2d and  
181 Supplementary Table 4). However, field margins increase diversity at the scale of the whole  
182 field, and so direct interactions leading to and kind of complementarity with crop species are  
183 limited. Indeed, field margins likely play a principal role in increasing crop productivity is via a  
184 reduction in pest populations through spill over from invertebrate natural enemies or in the  
185 case of flowering crops through increased pollination<sup>4,6</sup>.

186

### 187 **Direct effects of plant species richness on plant antagonists**

188 Across all studies, plant species richness decreased the overall performance of plant antagonists  
189 (CI=-0.490 to -0.388, df=3318, ES=-0.439, P<0.001) as well as when considered as  
190 subgroups of herbivores (CI=-0.522 to -0.379, df=1809, ES=-0.451, P<0.001), plant diseases  
191 (CI=-0.465 to -0.323, df=601, ES=-0.394, P<0.001), weeds (CI=-0.632 to -0.396, df=600,  
192 ES=-0.514, P<0.001), plant-feeding nematodes (CI=-0.461 to -0.134, df=284, ES=-0.298,  
193 P<0.001) and plant-consuming rodents (CI=-0.677 to -0.270, df=20, ES=-0.474, P<0.001).  
194 This pattern of responses largely held when these aggregate performance indicators for each  
195 antagonist group were partitioned into individual components, i.e., when plant disease  
196 performance was considered in terms of its component metrics of plant disease reproduction,  
197 spread and damage (see Supplementary Table 3). When sample sizes were small, differences  
198 in the response to increased plant species richness tended to be not significant (i.e., herbivore  
199 growth: CI=-0.403 to 0.122, df=48, ES=-0.141, P=0.287, but for other cases see  
200 Supplementary Table 3).

201 We also tested whether responses differed among ecosystems. The responses of plant  
202 antagonist performance to increasing plant species was negative for agroecosystems  
203 (CI=-0.538 to -0.423, df=2777, ES=-0.481, P<0.001) and grasslands (CI=-0.466 to -0.147,  
204 df=315, ES=-0.306, P<0.001), but only marginally significant in forests (CI=-0.213 to 0.013,  
205 df=224, ES=-0.100, P=0.083). When plant antagonists were split into different subgroups (i.e.,  
206 invertebrate herbivores, plant diseases, nematodes, rodents, and weeds) and into different  
207 response categories (e.g., invertebrate herbivore growth, reproduction, and damage), the  
208 overall pattern described above was maintained in agroecosystems (Fig. 2a; Extended Data Fig.  
209 2a), but these responses varied in grasslands and forests (Extended Data Fig. 2b, c;  
210 Supplementary Table 4). We found that plant species richness reduced herbivore performance  
211 (i.e., herbivore growth, reproduction and damage) only in agroecosystems (CI=-0.602 to  
212 -0.443, df=1436, ES=-0.523, P<0.001) (Supplementary Table 4). This finding supports  
213 several key hypotheses, including: 1) the Enemies Hypothesis that suggests herbivore natural  
214 enemy performance is positively related to plant species richness<sup>6,33,58</sup>, 2) the Resource  
215 Concentration Hypothesis (RCH) which argues that the density of insect herbivores increases  
216 with monoculture host density and patch size<sup>58</sup>, and, finally, 3) the Insurance Hypothesis (IH)  
217 whereby more diverse crops provide insurance against pest damage<sup>59</sup>. However, there was little  
218 support for these hypotheses in grasslands (CI=-0.316 to 0.065, df=171, ES=-0.126, P=0.195)  
219 and forests (CI=-0.162 to 0.067, df=200, ES=-0.048, P=0.413) as the negative response of  
220 herbivores to plant species richness had confidence intervals that overlapped zero.

221 The observed differences may be due to the more common outbreaks (i.e., more severe  
222 impacts) of specialist herbivores in agroecosystems than grasslands, although for monospecific  
223 forest destructive outbreaks of specialist herbivore do occur. The prevalence of this effect may  
224 be exacerbated by lower crop diversity in arable or horticultural cropping systems (~2-3 plant  
225 species in general)<sup>18,23</sup> than would be seen in many grasslands (e.g. the Jena experiment in  
226 Germany<sup>10,11</sup> and Cedar Creek experiment in the USA<sup>9,11</sup>; both consider 60 plant species), and  
227 to a lesser extent for forests (e.g. a maximum of 24 tree species in a Chinese forestry study<sup>33</sup>).  
228 Increased plant species richness may also provide greater access to nutritionally superior or  
229 more variable food resources for insect herbivores in grasslands, particularly for oligophagous  
230 or polyphagous specie<sup>60</sup>.

231 Across all studies, we found that increased plant species richness decreased symptomatic  
232 disease expression in plants in agroecosystems (CI=-0.458 to -0.301, df=514, ES=-0.379,

233 P<0.001), grasslands (CI=-0.696 to -0.358, df=70, ES=-0.527, P<0.001) and forests  
234 (CI=-0.689 to -0.100, df=15, ES=-0.395, P=0.012), as well as weed performance in  
235 agroecosystems (CI=-0.610 to -0.369, df=547, ES=-0.490, P<0.001) and grasslands  
236 (CI=-1.411 to -0.072, df=52, ES=-0.742, P=0.031). As to different groups of plant  
237 antagonists in the three ecosystems, nematode performance only in agroecosystems  
238 (CI=-0.511 to -0.117, df=266, ES=-0.314, P=0.002), and rodent performance in both  
239 agroecosystems (CI=-0.712 to -0.428, df=10, ES=-0.570, P<0.001) and grasslands  
240 (CI=-0.924 to -0.160, df=6, ES=-0.542, P=0.013) also decreased with plant species richness.  
241 These findings strongly suggest that enhancing plant species richness is an effective method to  
242 promote antagonist control services in managed systems, such as agro-ecosystems, managed  
243 grasslands and production forest. Increased control of antagonists might be explained by the  
244 effects of plant species diversity on disease performance, resulting from altered wind and vector  
245 dispersal, modified microclimate, dilution of host density, decreased vectors of plant diseases  
246 (e.g., aphids and planthoppers), changes in host morphology and physiology and direct  
247 pathogen inhibition<sup>22</sup>. Suppressed weed growth in diverse mixtures may be linked to the spatio-  
248 temporal continuity of ground cover provided by the agricultural plants in diverse assemblages,  
249 with this providing little bare ground for weeds to establish<sup>61</sup>. It may also be related to the  
250 allelopathic effects of diversified plant species on weed growth and germination<sup>62</sup>, and to the  
251 physical and chemical properties of the mulches<sup>63</sup>. Decreased growth of nematodes in diverse  
252 plant communities may be a by-product of a greater diversity of allelopathic chemicals<sup>64</sup>.  
253 Alternatively, this may be due to many plant parasitic nematodes being relatively specialized  
254 such that a diverse assembly of crop plants is likely to result in a dilution of suitable plants  
255 across space and time<sup>65</sup>. Finally, increased rodent dispersal in monocrop plots may be driven  
256 by a response to limited habitat and resource diversity availability<sup>20</sup>. Overall, physical barriers,  
257 allelopathic chemicals, plant species resistance, natural enemies and dilution effects may be  
258 responsible for the negative effects of plant species richness on herbivores, plant diseases,  
259 nematodes and weeds<sup>22</sup>.

260 Next, we tested whether these responses of plant antagonists and their subgroups differed  
261 among plant life forms, climatic zones, experimental types, plant diversity types and  
262 experiment design types. We found qualitatively similar negative responses for both  
263 herbaceous and woody plants (Fig. 2b; Extended Data Fig. 3; Supplementary Table 5), for both  
264 temperate and tropical zones (Fig. 2c; Extended Data Fig. 4; Supplementary Table 6), and in

265 both additive (i.e. the densities of plant species increases as more species are added) and  
266 replacement series (i.e. plant density remains constant, even as more plant species are added)  
267 designs (Fig. 2f; Supplementary Table 7). Across types of experimental studies, we found  
268 stronger responses from the subgroup of plant antagonists to plant species richness in plot than  
269 in pot experiments, possibly due to smaller sample sizes in pot experiments (Fig. 2d; Extended  
270 Data Fig. 5; Supplementary Table 7). This would restrict below-ground components of these  
271 communities as well as dispersal parameters that would be encountered under field-based  
272 experiments. Consequently, they do not incorporate the intricate biotic interactions existing in  
273 natural communities.

274 Finally, we tested whether these responses of plant antagonists (i.e., pests in agroecosystems)  
275 and their subgroups differed among intercropping, cover cropping, and sown field margins. We  
276 found qualitatively similar negative responses for these crop diversification modes (Fig. 2e;  
277 Supplementary Table 4). The mechanisms to explain the decreased pests are proposed to be  
278 like those described above. Funnel plots for each trophic group was presented in Extended Data  
279 Fig. 6.

280

### 281 **Mediation analysis of plant species richness on plants and their antagonists**

282 We used mediation analysis (i.e., path analysis) to test our proposed diversity-antagonist-  
283 productivity relationship (Supplementary Tables 8–13). For our analysis, we collected 2,766  
284 estimates of interactions between pairs of plants and plant antagonists derived from 240 articles.  
285 First, we tested the effects of plant species richness (i.e., a binary variable) on plants and plant  
286 antagonists using multilevel piecewise structural equation models. In these models, we  
287 aggregated different plant antagonists together, i.e., invertebrate herbivores, plant diseases,  
288 weeds, plant-feeding nematodes, and plant-consuming rodents (Supplementary Methods). We  
289 found that plant species richness directly suppressed plant antagonists and increased plant  
290 performance across terrestrial ecosystems and in agroecosystems, grasslands, and forests (Fig.  
291 3). increasing plant species also increased plant performance indirectly by reducing the  
292 performance of plant antagonists across terrestrial ecosystems (supplementary Table 11) and  
293 specifically in agroecosystems (Fig. 3a). However, this indirect effect on plant performance via  
294 suppression of antagonists was less pronounced in grasslands (Fig. 3b) or forests (Fig. 3c).

295 When we classified plant diversity in agroecosystems into different types, we found that  
296 intercropping, cover cropping, and sown field margins directly suppressed plant antagonists

297 (i.e., pests) and increased plant performance (i.e., crop production) (Figs. 3d–f). Intercropping  
298 and cover cropping also increased crop production indirectly by reducing pests (Figs. 3d, e).  
299 The effect of sown field margins on crop production via the suppression of pests were not as  
300 clear as was seen for intercropping and cover crops (supplementary Table 11; Fig. 3f). Similarly,  
301 the direct effects of plant species richness on plant performance and plant antagonist  
302 performance were consistent in herbaceous-species dominated and woody-species dominated  
303 systems, temperate and tropical zones, and in plot and pot experiments. The mediation effects  
304 of plant antagonists on plant performance were significant for herbaceous plants ( $P=0.019$ ,  
305  $N=2,514$ ), plot experiments ( $P=0.027$ ,  $N=2,513$ ), and temperate zones ( $P=0.027$ ,  $N=1524$ ).  
306 This finding may in part be a product of the well-studied effects of intercropping research in  
307 field crops. However, mediation effects were not significant for woody plants ( $P=0.593$ ,  $N=252$ )  
308 or pot experiments ( $P=0.211$ ,  $N=253$ ), or in tropical zones ( $P=0.101$ ,  $N=936$ ) (Extended Data  
309 Figs.7; Supplementary Table 13). Indirect positive effects of plant species richness on plant  
310 performance were found to be mediated through a reduction of plant antagonist pressure when  
311 we separately tested for the effects of invertebrate herbivores, plant diseases and weed  
312 performance, although such an indirect effect was not evident for the nematodes (Fig. 4).

313 Our meta-analysis identifies that plant species richness enhances plant performance while  
314 suppressing the performance of various groups of plant antagonists, including invertebrate  
315 herbivores, plant diseases, weeds, plant-feeding nematodes and plant-consuming rodents.  
316 Further, across all studies our analysis indicated that in a comparison of low vs high plant  
317 diversity we see an increase in plant performance by 24.31% in agroecosystems, 69.59% in  
318 grasslands and 28.75% in forests. This can be explained by the decrease in plant antagonist  
319 performance by 30.18% in agroecosystems, 19.98% in grasslands and 8.85% in forests  
320 (Extended Data Fig. 8). Increased plant performance resulting from decreasing plant  
321 antagonist performance can be realized by adding only one plant species in agroecosystems,  
322 grasslands and forests (Tables 1, 2). These results highlight the negative effects of plant  
323 diversity on the plant antagonists, and in doing so contributes to the positive effects of the  
324 biodiversity-productivity relationship. The diversity-antagonist-productivity relationship for  
325 terrestrial ecosystems is highly dependent on the results from agroecosystems (studies from  
326 agroecosystems accounted for 93.38% of those from terrestrial ecosystems).

327 Piecewise structural equation model indicated that plant diversity can promote crop  
328 production through indirect effects (i.e., suppressing pests) in agroecosystems, as well as

329 indirect mediation effects driven by factors such as improved soil fertility<sup>13</sup>, field micro-  
330 climate<sup>52</sup> and increasing nutrient use efficiency<sup>53</sup>. While the data does not differentiate  
331 between specialist and generalist antagonists, theoretical predictions show that it is likely that  
332 specialists are mostly responsible for the observed effects.

333 From an applied perspective, our findings suggest that crop diversification, whether from  
334 intercropping, cover cropping, or sown field margins, can help to promote crop pest control  
335 and increase crop production<sup>33</sup>. Such crop diversification may also help to mitigate climate  
336 induced yield losses in the future by increasing system resilience<sup>66</sup>. Many other forms of crop  
337 diversification (e.g., crop rotations), landscape diversification, or cropping system  
338 diversification might also be beneficial for pest control and crop production<sup>67-69</sup>. It is  
339 important to note that this meta-analysis does not consider non-biological indicators (e.g.,  
340 profitability, or access to specific markets via accreditation) which may alter the practical  
341 viability of more diversified systems. Rather, the results of this study have identified the  
342 general benefits across diverse production systems for suppressing plant antagonists through  
343 increasing plant diversity. However, for each system the subtleties of which crops, varieties  
344 and unique traits to combine would need to be considered. This may include a need to  
345 consider other agronomic decisions such as the use of agrochemicals authorized for specific  
346 crops that may not be compatible with intercropping systems, as well as sowing density,  
347 timing and establishment patterns. This is beyond the scope of this analysis. Our results  
348 provide empirical support for more complex cropping systems that increase plant diversity in  
349 agricultural fields and management strategies that foster increased diversity species co-  
350 existence in grasslands and forests.

351

## 352 **Methods**

### 353 **Definition of increasing plant species and number of added plant species**

354 We considered “increasing plant species” as a binary variable (zero or one), indicating whether  
355 plant species richness was increased, but irrespective of the number of plant species added.  
356 Here “number of added plant species” is a continuous variable describing the increase in plant  
357 species richness between the control and the treatment. When comparing plant species richness  
358 of the control (i.e., pure, mono- or lowest plant species) with that of the treatment (i.e., higher  
359 plant species richness,  $\geq 2$  plant species richness), we ensured that comparisons were also  
360 between the same trophic groups. Sensitivity analysis for “increasing plant species” is provided

361 in Supplementary Table 14. Description for analyses of increasing plant species richness and  
362 number of added species richness is provided in Supplementary Table 15. Statistic values for  
363 the relationship between number of added species richness in the plant species richness  
364 treatment over the control is presented in Supplementary Table 16 and Supplementary Figs. 1–  
365 12. Detailed results for “increasing plant species” are shown in Fig. 1c, Figs. 2–4, Extended  
366 Data Figs. 2–8, Supplementary Tables 1–15, 17, and Supplementary Figs. 13–42. Detailed  
367 results for “number of added plant species” were shown in Tables 1 and 2, Supplementary  
368 Results, Supplementary Tables 9, 11, 13, 15 and 16.

369

### 370 **Study selection**

371 Studies were selected through a literature search of the Web of Science Core Collection,  
372 BIOSIS Previews, Derwent Innovations Index, KCI-Korean Journal Database, MEDLINE,  
373 Preprint Citation Index, ProQuest™ Dissertations & Theses Citation Index, and SciELO  
374 Citation Index. We used the Boolean search string based on the “TOPIC” searching: ["plant  
375 diversity" OR "crop diversity" OR "crop diversification" OR "plant species richness" OR  
376 "polyculture" OR "ground cover vegetation" OR "flower strip" OR "strip crop\*" OR "grassy  
377 field margin" OR "border crop" OR "intercrop\*" OR "interplant\*"] AND ["plant disease" OR  
378 "plant virus" OR "nematode" OR "weed" OR "herbivor\*" OR "pest" OR "biological control"  
379 OR "rodent" OR "yield" OR "productivity" OR "biomass"]. This literature search was initiated  
380 in June 2019, and finalized in August 2023. In total, the search yielded 386,895 articles (see  
381 Extended Data Fig. 1 for a PRISMA diagram). Articles were screened by Y.Q.W., L.F., J.L.,  
382 J.Z. and N.F.W.. Data were extracted from the articles by Y.Q.W., L.F., J.L. and N.F.W. during  
383 which regular cross-checking was performed to ensure consistency in extracted effect sizes.

384 We used data giving the mean values of multiple sampling dates or years. If these mean  
385 values were not presented, we used the data of the latest sampling period<sup>33</sup>. For articles that  
386 covered more than one experimental location, we considered these experimental results  
387 separately (see locations in Fig. 1a). When numeric values were not provided directly, we  
388 extracted them from figures using the “GetData Graph Digitizer” 2.26. However, where linear  
389 or non-linear relationships between plant species richness and one of the response variables  
390 was presented in a figure, we extracted the values by fitting regression equations<sup>29</sup>.

391 To avoid pseudoreplication of data, we excluded multiple comparisons conducted within a  
392 single experiment<sup>33</sup>. Observations with the lowest plant species richness were considered as

393 control groups, while those with higher plant species richness were considered as the treatment  
394 groups. When an article included different levels of plant species richness, measurements for  
395 the control groups (lowest plant species richness) were compared to all other treatments levels  
396 of plant species richness and treated as independent paired observations.

397

### 398 **Predictor variables**

399 We used eight categorical variables as predictor variables (see Supplementary Methods for  
400 details)—i) Trophic group: a categorical variable describing whether the target organisms were  
401 invertebrate herbivores, plant diseases (plant pathogenic viruses, fungi and bacteria that  
402 infested plants and cause damage to plants), weeds, plant-feeding nematodes, plant-consuming  
403 rodents (e.g., rats and mice that damage crops, pasture or trees), or plants (e.g., crops, fruits,  
404 grassland species and trees); and moreover, an aggregate categorical variable (i.e. plant  
405 antagonists) including invertebrate herbivores, plant diseases, weeds, plant-feeding nematodes  
406 and plant-consuming rodents. ii) Response category: growth, reproduction and damage of  
407 herbivores; reproduction, spread and damage of plant diseases; weed growth, reproduction and  
408 diversity (i.e., species richness and Shannon diversity of weeds); reproduction and damage of  
409 plant-feeding nematodes; reproduction and damage of plant-consuming rodents; and growth,  
410 reproduction and quality of plants. iii) Ecosystem type: agroecosystems, grasslands and forests.  
411 iv) Plant life form: herbaceous or woody plants. v) Climatic zone: temperate or tropical (data  
412 from greenhouse, indoor and laboratory experiments were removed from models including the  
413 climatic zone variable)<sup>29</sup>. vi) Experiment type: plots (i.e., field and common garden  
414 experiments) or pots (i.e., experiments with pots, containers, bottles, trays, boxes and tankers)  
415 (detailed description was presented in Supplementary Methods). vii) plant diversity types: crop  
416 diversification including intercropping, cover cropping and sown field margins in  
417 agroecosystems, mixed forest plantations, and diversified grasslands. viii) experimental design  
418 types: replacement series design and additive design. A replacement series design means that  
419 total plant density remains constant, even as more plant species are added.

420

### 421 **Definition of effect size and its measures**

422 To test the effect of plant species richness on the various groups (invertebrate herbivores,  
423 plant diseases, weeds, plant-feeding nematodes, plant-consuming rodents, or plants; and  
424 moreover, an aggregate plant antagonists categorical variable including invertebrate

425 herbivores, plant diseases, weeds, plant-feeding nematodes and plant-consuming rodents), we  
 426 calculated the effect size and lnRR of these groups. The first proposed formula as follows:

$$\ln RR_1 = \ln \left( \frac{m_1}{m_2} \right) \quad (\text{Eq.1})$$

$$v(\ln RR_1) = \frac{sd_1^2}{n_1 m_1^2} + \frac{sd_2^2}{n_2 m_2^2} = \frac{CV_1^2}{n_1} + \frac{CV_2^2}{n_2} \quad (\text{Eq.2})$$

427  
 428 where  $m_1$  and  $m_2$  are the observed mean value in the treatment and control groups,  $sd_1$  and  $sd_2$   
 429 are the standard deviations (SDs) in the treatment and control groups, and  $n_1$  and  $n_2$  are the  
 430 sample sizes in the treatment and control groups.  $m_1$ ,  $m_2$ ,  $sd_1$ ,  $sd_2$ ,  $n_1$  and  $n_2$  were extracted  
 431 from original articles. Namely, mean value, SD and sample sizes of both treatment and  
 432 control groups were included in our dataset to conduct meta-regression. This was done in  
 433 order to deal with missing standard deviations (SDs) in dataset, using the approach of  
 434 Nakagawa et al.<sup>70</sup> which is suited to accounting for missing SDs. Nakagawa et al.<sup>70</sup> proposed  
 435 this new method weighting average coefficients of variation estimated from studies that do  
 436 report SDs in the dataset. This is done by:

$$\ln RR_2 = \ln \left( \frac{m_1}{m_2} \right) + \frac{1}{2} \left( \frac{CV_1^2}{n_2} - \frac{CV_2^2}{n_1} \right) \quad (\text{Eq.3})$$

$$v(\ln RR_2) = \frac{CV_1^2}{n_1} + \frac{CV_2^2}{n_2} + \frac{CV_1^4}{2n_1^2} + \frac{CV_2^4}{2n_2^2} \quad (\text{Eq.4})$$

$$\ln RR_3 = \ln \left( \frac{m_1}{m_2} \right) + \frac{1}{2} \left( \frac{\left[ \frac{\sum_{i=1}^K n_{1i} CV_{1i}}{\sum_{i=1}^K n_{1i}} \right]^2}{n_1} - \frac{\left[ \frac{\sum_{i=1}^K n_{2i} CV_{2i}}{\sum_{i=1}^K n_{2i}} \right]^2}{n_2} \right) \quad (\text{Eq.5})$$

$$v(\ln RR_3) = \frac{\left[ \frac{\sum_{i=1}^K n_{1i} CV_{1i}}{\sum_{i=1}^K n_{1i}} \right]^2}{n_1} + \frac{\left[ \frac{\sum_{i=1}^K n_{2i} CV_{2i}}{\sum_{i=1}^K n_{2i}} \right]^2}{n_2} + \frac{\left[ \frac{\sum_{i=1}^K n_{1i} CV_{1i}}{\sum_{i=1}^K n_{1i}} \right]^4}{2n_1^2} + \frac{\left[ \frac{\sum_{i=1}^K n_{2i} CV_{2i}}{\sum_{i=1}^K n_{2i}} \right]^4}{2n_2^2} \quad (\text{Eq.6})$$

437 where  $CV = sd/m$  is the coefficient of variation;  $sd$  and  $n$  are the corresponding SDs and sample  
 438 size, respectively;  $CV_{1i}$  and  $CV_{2i}$  are the CVs from the  $i$ th study (study:  $i=1,2,\dots,K$ ). The  
 439 proposed Eq.3 and Eq.5 can improve the accuracy and precision of the overall mean estimate.

440 We can use Eq.5 and 6 to calculate the effect sizes and sample variances when SDs are  
441 missing and use Eq.3 and 4 to calculate the effect size and sample variances when SDs are  
442 not missing. This paper uses Eq.5 to calculate effect regardless of whether SDs are missing or  
443 not, which was refer as “All Cases” method<sup>70</sup> We used lnRR as the response variable for  
444 different models except for path analyses described below.

445

### 446 **Meta-regression**

447 Meta-regression<sup>71</sup> was applied to test whether the effect sizes of different trophic groups could  
448 be explained by increasing plant species and the various predictor variables. Specifically, we  
449 fitted three-level mixed-effects meta-regression models, using the R package metafor (version  
450 3.8-1). The effect size metric lnRR was calculated using the function “lnrr\_laj()” in R.file  
451 implemented using the package “func.R” developed by Nakagawa et al.<sup>70</sup>. This was used to  
452 calculate the effect size metric lnRR for each observation as well as the unbiased sample  
453 variance estimates as defined under Eq.6 (using function “v\_lnrr\_laj()” in R.file “func.R”).  
454 Trophic groups, trophic group response categories, ecosystem types, plant life forms, climatic  
455 zones, types of experimental study, types of plant diversity, and experimental design types,  
456 were included as moderators whose effects were assumed to be fixed. In all models, we treated  
457 ‘study’ as a random effect (see Supplementary Tables 1.1, 1.3, 2.1, 2.3). To handle non-  
458 independence in effect sizes of each study, we added a random effect as a unique identifier for  
459 each effect size in every study (EsID) which allows true effect sizes to vary within studies, and  
460 to account for the within-study effect and quantify within-study heterogeneity. To obtain robust  
461 results and account for differences in precision across studies and effect sizes, we weighted  
462 effect sizes using the inverse of the addition of the variance-covariance matrix, which explicitly  
463 captured the non-zero covariance arising from correlation between sampling variance within  
464 the same original articles and random effect variance<sup>72</sup>. Phylogenetic correction for plants was  
465 also undertaken to investigate the effects of plant species richness on trophic groups. To do  
466 this, plant species phylogenies were included as a random effect with phylogenetic relatedness  
467 as part of the correlation structure. In addition, phylogenetic trees of all crop species in  
468 agroecosystems were drawn to investigate the evolutionary relationships among different crop  
469 species (Supplementary Fig. 43). Specifically, we matched the crop species included in our  
470 analysis with the available synthetic tree in R package “rotl”, then the relationships between  
471 each matched crop species were returned so that they could be used to draw phylogenetic trees.

472 To adjust for repeated measurement of control values, we assigned the argument “V” in  
 473 “rma.mv()” function with the sampling variance-covariance matrix estimated as follows<sup>72</sup>:

$$v_1 = \frac{\left[ \frac{\sum_{i=1}^K n_{1i} CV_{1C,i}}{\sum_{i=1}^K n_{1C,i}} \right]^2}{n_{1C}} + \frac{\left[ \frac{\sum_{i=1}^K n_{2i} CV_{1T,i}}{\sum_{i=1}^K n_{1T,i}} \right]^2}{n_{1T}} + \frac{\left[ \frac{\sum_{i=1}^K n_{1i} CV_{1C,i}}{\sum_{i=1}^K n_{1C,i}} \right]^4}{2n_{1C}^2} + \frac{\left[ \frac{\sum_{i=1}^K n_{2i} CV_{1T,i}}{\sum_{i=1}^K n_{1T,i}} \right]^4}{2n_{1T}^2} \quad (\text{Eq.7})$$

$$v_2 = \frac{\left[ \frac{\sum_{i=1}^K n_{1i} CV_{1C,i}}{\sum_{i=1}^K n_{1C,i}} \right]^2}{n_{1C}} + \frac{\left[ \frac{\sum_{i=1}^K n_{2i} CV_{2T,i}}{\sum_{i=1}^K n_{2T,i}} \right]^2}{n_{2T}} + \frac{\left[ \frac{\sum_{i=1}^K n_{1i} CV_{1C,i}}{\sum_{i=1}^K n_{1C,i}} \right]^4}{2n_{1C}^2} + \frac{\left[ \frac{\sum_{i=1}^K n_{2i} CV_{2T,i}}{\sum_{i=1}^K n_{2T,i}} \right]^4}{2n_{2T}^2} \quad (\text{Eq.8})$$

$$v_{12} = \frac{\left[ \frac{\sum_{i=1}^K n_{1i} CV_{1C,i}}{\sum_{i=1}^K n_{1C,i}} \right]^2}{n_{1C}} + \frac{\left[ \frac{\sum_{i=1}^K n_{1i} CV_{1C,i}}{\sum_{i=1}^K n_{1C,i}} \right]^4}{2n_{1C}^2} \quad (\text{Eq.9})$$

474 where the subscripts 1T and 2T represent the different treatment group who share the control  
 475 value in a same group; the subscript 1C represents the 1st control group; *n* represents the  
 476 number of observations, *CV* represents the coefficients of variance of observed value. Taking  
 477 this approach 116 plant species defined the correlation structure based on the plant phylogeny  
 478 (see Supplementary Tables 1.2, 1.4, 2.2, 2.4; Supplementary Methods)<sup>73,74</sup>.

479 For each mixed-effects meta-regression model, we first fitted a base model by treating plant  
 480 species richness and trophic group as the fixed effect terms. Second, the interactions between  
 481 the trophic group and other predictor variables (types of ecosystems, plant life forms, climatic  
 482 zones, experiment study types, plant diversity types, and experimental design types) were also  
 483 included in the model to assess whether model fit was improved, using a likelihood-ratio test  
 484 (LRT). Third, the trophic group response category (nested within trophic group) and the  
 485 interactive effects between the response category and predictors were also included in the  
 486 model (using a LRT to allow model comparisons) (Supplementary Table 1). For example, the  
 487 model with “trophic group + ecosystem type” was compared to the base model with just trophic  
 488 group, and the model with “trophic group + trophic group × ecosystem type” would be  
 489 compared to a model with “trophic group + ecosystem type” (Supplementary Tables 1 and 2).

490 To examine whether the mean effect sizes of (added) plant species richness, response category  
491 and other predictors differed significantly from zero, we acquired estimations with their 95%  
492 confidence intervals, which were derived from the fitted meta-regression models. To assess the  
493 between-study heterogeneity in these models,  $I^2$  statistics were calculated<sup>72,75,76</sup>  
494 (Supplementary Tables 1–7) and orchard plots<sup>77</sup> are presented in Supplementary Figs. 13–42.

495

### 496 **Mediation analysis to test the effects of plant species richness**

497 To explore the trophic interactions between plants and various groups of plant antagonists, we  
498 established a new data subset: 1) comprising paired trophic observations (e.g., plant  
499 performance vs. herbivore performance) (see Supplementary Fig. 12); 2) encompassing the  
500 paired observations of plant performance vs. plant antagonist performance in all ecosystems,  
501 and separately for agroecosystems, grasslands and forests, and the paired observations of  
502 plant performance vs. plant antagonist performance in intercropping, cover cropping and  
503 sown field margin farming systems (Fig. 4; Supplementary Figs. 11a–d); 3) and comprising  
504 the paired observations of plant performance vs. plant antagonist performance within  
505 different plant life forms, climatic zones, and experimental type, respectively (Extended Data  
506 Fig. 7; Supplementary Figs. 11e–j). The lnRR effect sizes were derived from these pairwise  
507 data sets.

508 To analyze the direct effect of plant antagonists on plant performance in path analysis,  
509 residual regression was applied (Fig. 4 and Extended Data Fig. 7). In this analysis we used  
510 the lnRR as the response variable. Specifically, the direct effects of increasing plant species  
511 on plant antagonist were estimated using linear mixed-effects model (See details in  
512 Supplementary Methods):

$$513 \text{ plant antagonist performance } \ln\text{RR}_{ij} = \beta + r_i + \varepsilon_{ij}$$

514 Following Emmenegger & Bühlmann<sup>78</sup>. The direct effect of plant species richness on plant  
515 performance were estimated by regressing the Pearson residuals of the linear mixed-effects  
516 model:

$$517 \text{ plant performance } \ln\text{RR}_{ij} = \beta \times \text{plant antagonist performance } \ln\text{RR}_{ij} + r_i + \varepsilon_{ij}$$

518 on plant species richness. The direct effects of plant antagonists on plants were estimated by  
519 regressing the Pearson residuals extracted from the linear mixed-effects model:

$$520 \text{ plant performance } \ln\text{RR}_{ij} = \beta + r_i + \varepsilon_{ij}$$

521 on the Pearson residuals extracted from the linear mixed-effects model:

522 
$$\text{plant antagonist performance } \ln\text{RR}_{ij} = \beta + r_i + \varepsilon_{ij}$$

523 where  $\beta$  is the effect size,  $r_i$  is the random effect represents the heterogeneity between studies  
524 or the phylogenetic relatedness of the plant species within  $i$ th study,  $\varepsilon_{ij}$  is the random error  
525 term with variance equal to the  $\ln\text{RR}$ 's sample variance estimate.

526 In summary, for our path analyses, linear mixed-effects models were conducted with the R  
527 function "lme" of the package "nlme", with random intercepts for study IDs. Heteroscedasticity  
528 was accounted for by providing fixed variances based on  $\ln\text{RR}$ s and setting sigma to 1 in the  
529 lme call.

530 We extracted the z-values of corresponding coefficients to test the effects of plant species  
531 richness on each of the interactions between performance values of plants and herbivores,  
532 plants and diseases, plants and weeds, and plants and nematodes, respectively. Likewise, we  
533 extracted the z-values of corresponding coefficients to test the effects of plant species richness  
534 on the interactions between performance values of plants and the aggregate indicator (i.e., plant  
535 antagonists) in different ecosystems (Fig. 4), as well as the effects of plant species richness on  
536 the interactions between plant performance and plant antagonist performance for different plant  
537 life forms, climatic zones, experiment types, plant diversity types in agroecosystems  
538 (intercropping, cover cropping and sown field margins), and experimental design types,  
539 respectively (Extended Data Fig. 7). The estimations and test statistics were extracted using the  
540 R function "coef()" (Supplementary Tables 8, 10 and 12). The relative goodness-of-fit analyses  
541 for path analyses of predictor variables (plant species richness), were conducted by extracting  
542 AIC (Akaike information criterion), AICc (corrected Akaike information criterion), BIC  
543 (Bayesian information criterion) and log-likelihood from the fitted models, using R functions  
544 "AIC()", "AICc()",<sup>79</sup> "BIC()" and "logLik()" (Supplementary Table 15).

545

#### 546 **Publication bias test**

547 We assessed publication bias using regression tests<sup>80,81</sup> (Supplementary Table 2) which employ  
548 a partial slope test of association between effect size and the sample size. Here, a significant  
549 relationship ( $p < 0.05$ ) suggests publication bias. The trim-and-fill method was not employed as  
550 this is inappropriate for models with moderators<sup>82,83</sup>. Instead, we adopted the method suggested  
551 by Nakagawa et al.<sup>81</sup>, which uses a multilevel version of Egger regression to assess publication

552 bias in mixed-effects meta-regression analysis. Here, we considered lnRR as a response  
553 variable. Different categories of trophic groups, ecosystems, plant life forms, climatic zones,  
554 experimental studies, plant diversity types and experimental design types were considered as  
555 predictors, respectively, and sampling sizes were considered as an additional moderator in the  
556 mixed-effect model and the test statistics for coefficients of sampling sizes were used to test  
557 for publication bias (see Supplementary Methods).

558 We used R version 4.3.1<sup>84</sup> to conduct all statistical analyses, and used R package “metafor”  
559 3.8-1 to perform meta-regression and publication bias assessment<sup>85</sup>. In addition, we used R  
560 packages “nlme”<sup>86</sup> to residual regression in path analyses. A significance level of 0.05 was used  
561 for all tests. To test the lnRR is appropriate for our raw data, we conducted a Geary- Lajeunesse  
562 test<sup>87</sup>. The data and code used in this study are publicly available in Zenodo  
563 (<https://doi.org/10.5281/zenodo.17568135>) (ref. 88).

564

#### 565 **Data availability**

566 The raw and processed data used in this study is available and is deposited to Zenodo  
567 (<https://doi.org/10.5281/zenodo.17568135>) (ref. 88).

568

#### 569 **Code availability**

570 The code that supports the findings of this study has been deposited in Zenodo  
571 (<https://doi.org/10.5281/zenodo.17568135>) (ref. 88).

572

#### 573 **Acknowledgments**

574 We thank all of the researchers whose data and work have been included in this meta-analysis.  
575 N.F.W. was supported by Shanghai Agriculture Applied Technology Development Program,  
576 China (Grant No. 2023-02-08-00-12-F04586), Natural Science Foundation of Shanghai  
577 (22ZR1417200), National Natural Science Foundation of China (32172484), Fundamental  
578 Research Funds for the Central Universities (JKY01231718), Shanghai Science and  
579 Technology Innovation Action Plan from Shanghai Municipal Science and Technology  
580 Commission of China (22015821000) and National Ten Thousand Plan-Young Top Talents of  
581 China. L.F. was supported by the National Natural Science Foundation of China (82204063).  
582 Y.Q.H. was supported by National Key R&D Program of China (2023YFF1205101) and

583 National Natural Science Foundation of China (11971117). B.A.W is funded through the  
584 Natural Environment Research Council RestREco (NE/V006444/1) and AgZero+ (NE/  
585 W005050/1) projects. D.Z.D. was supported by the University of Zurich Research Priority  
586 program on Global Change and Biodiversity.

587

588 **Author contributions**

589 N.F.W. conceived the idea. N.F.W., Y.Q.W., L.F. and J.L. collected and analyzed data and  
590 drafted the article. N.F.W., Y.Q.W., L.F., J.L., B.A.W., Y.Q.H., A.E., A. H., M.L., Y.H., R.D.B.,  
591 P.K., D.Z.D., L.H.F., J.M.B., S.N., S.S., F.X., D.P.S., Z.L., J.Z. and C.S. wrote the manuscript.  
592 All authors prepared and edited the final drafts.

593

594

595 **Competing interests**

596 The authors declare no competing financial interests.

597

598

599 **Correspondence and requests for materials** should be addressed to N.F.W.

600

601

602

603

604

605

606

607

608

609

610

611  
612  
613  
614  
615  
616  
617  
618

**Table 1 | Quantified effects of number of added plant species on plant performance in different ecosystems.**  
Increased percentage of plant performance = (data of plant performance in treatment—data of plant performance in control) / data of plant performance in control×100%.

Number of added plant species over the control	Ecosystem types	Number of observations	Number of studies	Increased percentage of plant performance	Lower of 95% CI	Upper of 95% CI	t value	P value
1	Agroecosystem	1568	250	23.45%	23.42%	23.47%	12.5760	1.2657×10 <sup>-34</sup>
1	Grassland	96	40	35.02%	35.00%	35.04%	5.5417	2.6722×10 <sup>-7</sup>
1	Forest	39	13	5.71%	5.71%	5.72%	2.9297	0.0057
2	Agroecosystem	148	32	23.57%	23.56%	23.58%	9.2958	1.8594×10 <sup>-16</sup>
2	Grassland	14	9	14.87%	14.86%	14.88%	1.7463	0.1043
2	Forest	33	9	19.32%	19.31%	19.33%	3.2803	0.0025
3	Agroecosystem	51	15	44.60%	44.59%	44.62%	6.6179	2.3776×10 <sup>-8</sup>
3	Grassland	97	47	56.35%	56.30%	56.39%	4.6000	1.2903×10 <sup>-5</sup>
3	Forest	29	11	12.89%	12.89%	12.90%	3.6276	0.0011
4	Agroecosystem	6	5	20.21%	20.19%	20.23%	0.9271	0.3964
4	Grassland	10	3	61.86%	61.82%	61.90%	1.6249	0.1386
4	Forest	6	4	45.74%	45.71%	45.77%	1.2981	0.2509
5	Agroecosystem	5	3	77.77%	77.74%	77.79%	2.4287	0.0721
5	Grassland	9	7	227.52%	227.31%	227.72%	1.1663	0.2771
5	Forest	13	5	17.06%	17.05%	17.07%	2.5149	0.0272
6	Agroecosystem	2	2	9.87%	9.86%	9.88%	0.3540	0.7834
6	Grassland	2	2	95.62%	95.58%	95.67%	1.0300	0.4906
6	Forest	1	1	71.67%	NA	NA	NA	NA
7	Agroecosystem	1	1	226.93%	NA	NA	NA	NA
7	Grassland	72	34	61.98%	61.96%	62.00%	8.0216	1.5251×10 <sup>-11</sup>
7	Forest	6	3	61.79%	61.78%	61.80%	7.4426	0.0007
8	Grassland	7	5	27.22%	28.00%	27.23%	1.3044	0.2399
8	Forest	6	2	28.03%	28.01%	28.04%	1.7293	0.1443
9	Agroecosystem	1	1	89.47%	NA	NA	NA	NA
9	Grassland	5	2	195.45%	195.39%	195.51%	2.3752	0.0764
9	Forest	1	1	108.83%	NA	NA	NA	NA
10	Grassland	2	2	138.75%	138.72%	138.78%	2.2653	0.2646
10	Forest	1	1	120.35%	NA	NA	NA	NA
11	Grassland	4	3	69.81%	69.80%	69.81%	10.4638	0.0019
11	Forest	1	1	1217.65%	NA	NA	NA	NA
12	Grassland	5	1	-2.97%	-2.98%	-2.97%	-0.4319	0.6881
13	Agroecosystem	2	1	16.03%	16.01%	16.04%	0.6265	0.6437
13	Grassland	2	2	161.86%	161.82%	161.89%	2.1674	0.2752
14	Grassland	5	2	211.66%	211.58%	211.75%	1.9149	0.1280
15	Grassland	67	34	107.33%	107.29%	107.37%	7.9595	3.1282×10 <sup>-11</sup>
15	Forest	2	1	89.91%	89.90%	89.91%	7.6283	0.0830
19	Agroecosystem	1	1	-18.91%	NA	NA	NA	NA
20-56	Grassland	38	5	35.72%	35.71%	35.74%	4.4491	7.6204×10 <sup>-5</sup>
59	Grassland	33	12	138.53%	138.47%	138.60%	4.1016	2.6306×10 <sup>-4</sup>

619  
620  
621  
622  
623  
624

625  
626  
627  
628  
629

**Table 2 | Quantified effects of number of added plant species on plant antagonist performance in different ecosystems.** Decreased percentage of plant antagonist performance = (data of plant antagonist performance in control – data of plant antagonist performance in treatment) / data of plant antagonist performance in control×100%.

Number of added plant species over the control	Ecosystem types	Number of observations	Number of studies	Decreased percentage of plant antagonist performance	Lower of 95% CI	Upper of 95% CI	t value	P value
1	Agroecosystem	2468	381	29.54%	29.55%	29.52%	-31.0030	1.6622×10 <sup>-178</sup>
1	Grassland	94	36	6.76%	6.79%	6.73%	-0.7847	0.4346
1	Forest	50	21	12.28%	12.29%	12.26%	-2.4477	0.0180
2	Agroecosystem	183	49	38.35%	38.36%	38.34%	-16.3881	1.1383×10 <sup>-37</sup>
2	Grassland	8	3	38.77%	38.79%	38.75%	-1.7762	0.1190
2	Forest	29	14	10.48%	10.50%	10.46%	-1.0285	0.3125
3	Agroecosystem	57	25	25.95%	25.97%	25.94%	-4.3253	6.3300×10 <sup>-5</sup>
3	Grassland	64	33	20.67%	20.68%	20.65%	-3.7826	0.0003
3	Forest	41	19	12.74%	12.75%	12.73%	-2.7906	0.0080
4	Agroecosystem	27	11	27.75%	27.76%	27.74%	-3.6813	0.0011
4	Grassland	7	2	41.32%	41.33%	41.31%	-3.6973	0.0101
4	Forest	17	7	23.65%	23.66%	23.64%	-4.2638	0.0006
5	Agroecosystem	11	7	58.54%	58.55%	58.53%	-7.3694	2.3990×10 <sup>-5</sup>
5	Grassland	9	6	51.11%	51.12%	51.10%	-4.2763	0.0027
5	Forest	18	7	17.44%	17.45%	17.43%	-2.3623	0.0303
6	Agroecosystem	17	7	41.94%	41.96%	41.91%	-2.8634	0.0113
6	Grassland	4	1	0.75%	0.75%	0.74%	-0.1155	0.9153
7	Agroecosystem	4	3	-36.44%	-36.41%	-36.47%	0.7764	0.4941
7	Grassland	49	25	12.97%	12.99%	12.95%	-1.6799	0.0995
7	Forest	7	6	-23.22%	-23.21%	-23.23%	1.7877	0.1241
8	Agroecosystem	2	2	39.70%	39.71%	39.69%	-2.0239	0.2922
8	Grassland	6	3	72.18%	72.19%	72.18%	-9.3168	0.0002
8	Forest	2	2	39.89%	39.91%	39.87%	-1.0375	0.4883
9	Agroecosystem	1	1	38.76%	NA	NA	NA	NA
9	Grassland	4	1	35.55%	35.55%	35.54%	-3.4948	0.0396
9	Forest	2	2	37.52%	37.52%	37.52%	-42.1868	0.0151
10	Forest	6	3	-17.58%	-17.56%	-17.60%	0.7334	0.4962
11	Grassland	9	5	62.86%	62.87%	62.85%	-5.0456	0.0010
11	Forest	1	1	12.86%	NA	NA	NA	NA
12	Agroecosystem	3	1	19.44%	19.43%	19.43%	-135.4130	5.4531×10 <sup>-5</sup>
12	Forest	1	1	83.73%	NA	NA	NA	NA
13	Agroecosystem	2	1	86.00%	86.00%	86.00%	-39.8020	0.0160
13	Forest	4	2	27.24%	27.26%	27.23%	-1.3453	0.2712
14	Grassland	4	1	26.45%	26.46%	26.44%	-1.5537	0.2181
14	Forest	6	2	17.87%	17.88%	17.86%	-1.2576	0.2641
15	Agroecosystem	1	1	48.65%	NA	NA	NA	NA
15	Grassland	34	19	19.51%	19.54%	19.48%	-1.4517	0.1560
15	Forest	9	5	-49.00%	-48.98%	-49.01%	3.2985	0.0109
16	Forest	2	2	32.17%	32.18%	32.15%	-1.1270	0.4620
19	Agroecosystem	2	1	41.25%	41.26%	41.24%	-1.9508	0.3016
19	Forest	4	2	31.47%	31.49%	31.46%	-1.6193	0.2038
20-31	Grassland	9	6	45.85%	45.87%	45.83%	-2.3252	0.0485
21-44	Forest	26	6	-4.35%	-4.33%	-4.36%	0.4334	0.6684
59	Grassland	17	8	22.11%	22.13%	22.08%	-1.0508	0.3090

630

631

632

633

634

635

636

637

638

## 639 Figure legends

640

641 **Fig. 1 | Global distribution of 636 study locations of the responses of plants and their antagonists to plant species**  
642 **richness. a.** Study locations across global terrestrial ecosystems (world map in World Robinson projection; a literature search  
643 identified 536, 49 and 45 study locations for agroecosystems, grasslands and forests, respectively, from a total of 609  
644 published articles; seven articles included more than one study location (range 2–11)). **b.** Hypotheses (H1: direct effects of  
645 plant diversity on plant productivity; H2: direct effects of plant diversity on plant antagonists; H3: mediation effects of plant  
646 antagonists on plant diversity-productivity relationships). **c.** Responses of plants and their antagonists across all studies.  
647 Plant antagonists include invertebrate herbivores, plant diseases, weeds, plant-feeding nematodes and plant-consuming  
648 rodents. Weeds in forests were not found. Estimates for rodent damage ( $n < 3$ ) can be found in Supplementary Table 3. In Fig.  
649 1c, horizontal lines indicate the 95% confidence intervals around the means; numbers in brackets indicate the numbers of  
650 observations and articles; and the lines represent plant antagonist (black), invertebrate herbivore (red), plant disease (green),  
651 weed (blue), plant-feeding nematode (turquoise), plant-consuming rodent (purple), and plant (orange) performance  
652 responses, respectively. Map © ARCGIS Software.

653

654 **Fig. 2 | Mean effect sizes of response categories of plants and plant antagonists to plant species richness. a.** In  
655 agroecosystems, grasslands and forests. **b.** For herbaceous and woody plants. **c.** In temperate and tropical zones. **d.** For plot  
656 and pot experiments. **e.** In agroecosystems with intercropping, cover cropping and sown field margins. **f.** In additive design  
657 and replacement series design. Horizontal lines indicate the 95% confidence intervals around the means. Numbers in brackets  
658 indicate the numbers of observations and articles. All analyses used two-tailed t-tests, and no corrections for multiple  
659 comparisons were applied.

660

661 **Fig. 3 | Piecewise structural equation model for the effects of plant species richness on the performances of**  
662 **plants and plant antagonists. a.** In agroecosystems ( $N=2583$ ). **b.** In grasslands ( $N=65$ ). **c.** In forests ( $N=118$ ). **d.** In  
663 intercropping ( $N=2051$ ). **e.** In cover cropping ( $N=396$ ). **f.** In sown field margins ( $N=136$ ). Plant performance includes  
664 the growth, reproduction and quality of plants. Plant antagonists includes: i) invertebrate herbivores (herbivore growth  
665 and reproduction, and herbivory damage to plants); ii) plant diseases (reproduction, spread, and damage to plants); iii)  
666 weeds (growth, reproduction, and diversity); iv) nematodes (reproduction, and damage to plants); and v) rodents  
667 (reproduction, and damage to plants). \* $P < 0.05$ , \*\* $P < 0.01$ , \*\*\* $P < 0.001$ . The  $R^2$  was the proportion of variance of  
668 response explained by the corresponding predictor in each linear mixed model. Blue and red arrows denote positive and  
669 negative relationships, respectively. Numbers next to each arrow are the estimated coefficients from piecewise  
670 structural equation models, and line width is proportional to the magnitude of the coefficients (Supplementary Tables  
671 10, 11). The  $R^2$  measures the proportion of variance explained by the corresponding predictor in each linear mixed  
672 model. All analyses used two-tailed t-tests, and no corrections for multiple comparisons were applied.

673

674 **Fig. 4 | Piecewise structural equation model for the effects of plant species richness on the performances of**  
675 **plants and different groups of plant antagonists in global terrestrial ecosystems. a.** Effects on plant and antagonist  
676 performance ( $N=2766$ ). **b.** Effects on plant and invertebrate herbivore performance ( $N=1218$ ). **c.** Effects on plant and  
677 plant disease performance ( $N=638$ ). **d.** Effects on plant and weed performance ( $N=642$ ). **e.** Effects on plant and plant-  
678 feeding nematode performance ( $N=268$ ). Plant performance includes the growth, reproduction and quality of plants.  
679 Herbivore performance includes growth, reproduction and damage of invertebrate herbivores. Plant disease  
680 performance includes reproduction, spread and damage to plants. Weed performance includes growth, reproduction  
681 and diversity. Plant-feeding nematode performance includes reproduction and damage to plants. The data on the  
682 relationships between plant and plant-consuming rodent performances was not found. \* $P < 0.05$ , \*\* $P < 0.01$ , \*\*\* $P < 0.001$ .  
683 The  $R^2$  was the proportion of variance of response explained by the corresponding predictor in each linear mixed  
684 model. Blue and red arrows denote positive and negative relationships, respectively. Numbers next to each arrow are  
685 the estimated coefficients from piecewise structural equation models, and line width is proportional to the magnitude  
686 of the coefficients (Supplementary Tables 8, 9). The  $R^2$  measures the proportion of variance explained by the  
687 corresponding predictor in each linear mixed model. All analyses used two-tailed t-tests, and no corrections for  
688 multiple comparisons were applied.

689

690

691

692

## 693 **References**

- 694 1. Reckling, M., Watson, C. A., Whitbread, A. & Helming, K. Diversification for sustainable and resilient  
695 agricultural landscape systems. *Agron. Sustain. Dev.* **43**, 44 (2023).
- 696 2. Ray, D. K., Ramankutty, N., Mueller, N. D., West, P. C. & Foley, J. A. Recent patterns of crop yield growth  
697 and stagnation. *Nature Commu.* **3**, 1292 (2012).
- 698 3. Priyadarshana, T. S. et al. Crop and landscape heterogeneity increase biodiversity in agricultural landscapes:  
699 A global review and meta-analysis. *Ecol. Lett.* **27**, e14412 (2024).
- 700 4. Gurr, G. M. et al. Multi-country evidence that crop diversification promotes ecological intensification of  
701 agriculture. *Nat. Plants* **2**, 1–4 (2016).
- 702 5. Kovács-Hostyánszki, A., Espíndola, A., Vanbergen, A. J., Settele, J., Kremen, C. & Dicks, L. V. Ecological  
703 intensification to mitigate impacts of conventional intensive land use on pollinators and pollination. *Ecol. Lett.*  
704 **20**, 673–689 (2017).
- 705 6. Wan, N. F. et al. Increasing plant diversity with border crops reduces insecticide use and increases crop yield  
706 in urban agriculture. *eLife* **7**, e35103 (2018).
- 707 7. van der Werf, H. M. G., Knudsen, M. T. & Cederberg, C. Towards better representation of organic agriculture  
708 in life cycle assessment. *Nat. Sustain.* **3**, 419–425 (2020).
- 709 8. Loreau, M. & Hector, A. Partitioning selection and complementarity in biodiversity experiments. *Nature* **412**,  
710 72–76 (2001).
- 711 9. Tilman, D., Reich, P. B. & Knops, J. M. Biodiversity and ecosystem stability in a decade-long grassland  
712 experiment. *Nature* **441**, 629–632 (2006).
- 713 10. Scherber, C. et al. Bottom-up effects of plant diversity on multitrophic interactions in a biodiversity experiment.  
714 *Nature* **468**, 553–556 (2010).
- 715 11. Isbell, F. et al. High plant diversity is needed to maintain ecosystem services. *Nature* **477**, 199–202 (2011).
- 716 12. Tilman, D. Benefits of intensive agricultural intercropping. *Nat. Plants* **6**, 604–605 (2020).
- 717 13. Li, X. F. et al. Long-term increased grain yield and soil fertility from intercropping. *Nat. Sustain.* **4**, 943–950  
718 (2021).
- 719 14. Yan, E., Munier-Jolain, N., Martin, P. & Carozzi, M. Intercropping on French farms: Reducing pesticide and  
720 N fertiliser use while maintaining gross margins. *Eur. J. Agron.* **152**, 127036 (2024).
- 721 15. Chadfield, V. G. A., Hartley, S. E. & Redeker, K. R. Associational resistance through intercropping reduces  
722 yield losses to soil-borne pests and diseases. *New Phytol.* **235**, 2393–2405 (2022).
- 723 16. Prieto, I. et al. Complementary effects of species and genetic diversity on productivity and stability of sown  
724 grasslands. *Nat. Plants* **1**, 1–5 (2015).
- 725 17. Abdala-Robert, L. et al. Comparison of tree genotypic diversity and species diversity effects on different guilds  
726 of insect herbivores. *Oikos* **124**, 1527–1535 (2015).
- 727 18. Alarcon-Segura, V., Grass, I., Breustedt, G., Rohlf, M. & Tschamntke, T. Strip intercropping of wheat and  
728 oilseed rape enhances biodiversity and biological pest control in a conventionally managed farm scenario. *J.*  
729 *Appl. Ecol.* **59**, 1513–1523 (2022).

- 730 19. El-Hamawi, M. H., Youssef, M. M. A. & Zawam, H. S. Management of *Meloidogyne incognita*, the root-knot  
731 nematode, on soybean as affected by marigold and sea ambrosia (damsisa) plants. *J. Pest Sci.* **77**, 95–98 (2004).
- 732 20. Larsen, A. L. et al. Effects of habitat modification on cotton rat population dynamics and rodent community  
733 structure. *Forest Ecol. Manage.* **376**, 238–246 (2016).
- 734 21. Boetzel, F. A. et al. Undersowing oats with clovers supports pollinators and suppresses arable weeds without  
735 reducing yields. *J. Appl. Ecol.* **60**, 614–623 (2023).
- 736 22. Boudreau, M. A. Diseases in intercropping systems. *Annu. Rev. Phytopathol.* **51**, 499–519 (2013).
- 737 23. Brooker, R. W. et al. Improving intercropping: a synthesis of research in agronomy, plant physiology and  
738 ecology. *New Phytol.* **206**, 107–117 (2015).
- 739 24. Kulmatiski, A., Beard, K. H. & Heavilin, J. Plant-soil feedbacks provide an additional explanation for  
740 diversity-productivity relationships. *P. Roy. Soc. B-Biol. Sci.* **279**, 3020–3026 (2012).
- 741 25. van Ruijven, J., Ampt, E., Francioli, D. & Mommer, L. Do soil-borne fungal pathogens mediate plant diversity-  
742 productivity relationships? Evidence and future opportunities. *J. Ecol.* **108**, 1810–1821 (2020).
- 743 26. Forero, L. E., Kulmatiski, A., Grenzer, J. & Norton, J. Plant-soil feedbacks help explain plant community  
744 productivity. *Ecology* **103**, e3736 (2022).
- 745 27. Ziaie-Juybari, H., Pirdashti, H., Abo-Elyousr, K. A. M. & Mottaghian, A. Abiotic benefits of intercropping  
746 legumes and maize to reduce pests. *Arch. Phytopathol. Plant Prot.* **54**, 17–18 (2021).
- 747 28. Guinet, M. et al. Fostering temporal crop diversification to reduce pesticide use. *Nat. Commu.* **14**, 7416 (2023).
- 748 29. Wan, N. F. et al. Plant genetic diversity affects multiple trophic levels and trophic interactions. *Nat. Commu.*  
749 **13**, 7312 (2022).
- 750 30. Dassou, A. G. & Tixier, P. Response of pest control by generalist predators to local-scale plant diversity: a  
751 meta-analysis. *Ecol. Evol.* **6**, 1143–1153 (2016).
- 752 31. Vehviläinen, H., Koricheva, J. & Ruohomäki, K. Tree species diversity influences herbivore abundance and  
753 damage: meta-analysis of long-term forest experiments. *Oecologia* **152**, 287–298 (2007).
- 754 32. Letourneau, D. K. et al. Does plant diversity benefit agroecosystems? A synthetic review. *Ecol. Appl.* **21**, 9–21  
755 (2011).
- 756 33. Wan, N. F. et al. Global synthesis of effects of plant species diversity on trophic groups and interactions. *Nat.*  
757 *Plants* **6**, 503–510 (2020).
- 758 34. Iverson, A. L. et al. Do polycultures promote win-wins or trade-offs in agricultural ecosystem services? A  
759 meta-analysis. *J. Appl. Ecol.* **51**, 1593–1602 (2014).
- 760 35. Wan, N. F., Dainese, M., Wang, Y. Q. & Loreau, M. Cascading social-ecological benefits of biodiversity for  
761 agriculture. *Curr. Biol.* **34**, R587–R603 (2024).
- 762 36. Verret, V. et al. Can legume companion plants control weeds without decreasing crop yield? A meta-analysis.  
763 *Field Crop. Res.* **204**, 158–168 (2017).
- 764 37. Himmelstein, J., Ares, A., Gallagher, D. & Myers, J. A meta-analysis of intercropping in africa: impacts on  
765 crop yield, farmer income, and integrated pest management effects. *Int. J. Agric. Sustain.* **15**, 1–12 (2018).
- 766 38. Li, C. et al. Syndromes of production in intercropping impact yield gains. *Nat. Plants* **6**, 653–660 (2020a).
- 767 39. Li, C. et al. Yield gain, complementarity and competitive dominance in intercropping in China: A meta-

- 768 analysis of drivers of yield gain using additive partitioning. *Eur. J. Agron.* **113**, 125987 (2020b)
- 769 40. Li, C. et al. The productive performance of intercropping. *Proc. Natl Acad. Sci. USA* **120**, e2201886120 (2023).
- 770 41. Mudare, S. et al. Yield and fertilizer benefits of maize/grain legume intercropping in China and Africa: a meta-  
771 analysis. *Agron. Sustain. Dev.* **42**, 81 (2022).
- 772 42. Viaud, P. et al. Sugarcane yield response to legume intercropped: A meta-analysis. *Field Crop. Res.* **295**,  
773 108882 (2023).
- 774 43. Lopes, T. et al. Wheat (*Triticum aestivum* L.)-based intercropping systems for biological pest control. *Pest*  
775 *Manag. Sci.* **72**, 2193–2202 (2016).
- 776 44. Rakotomalala, A. A. N. A., Ficiciyan, A. M. & Tschardtke, T. Intercropping enhances beneficial arthropods  
777 and controls pests: A systematic review and meta-analysis. *Agric. Ecosyst. Environ.* **356**, 108617 (2023).
- 778 45. Yousefi, M. et al. The effectiveness of intercropping and agri-environmental schemes on ecosystem service of  
779 biological pest control: a meta-analysis. *Agron. Sustain. Dev.* **44**, 15 (2024).
- 780 46. Fernandez, A. R., Gleiser, G., Aizen, M. A. & Garibaldi, L. A. Intercropping functionally similar species  
781 reduces yield losses due to herbivory. A meta-analytical approach. *Agric. Ecosyst. Environ.* **361**, 108800 (2024).
- 782 47. Zhang, C. et al. Intercropping cereals with faba bean reduces plant disease incidence regardless of fertilizer  
783 input; a meta-analysis. *Eur. J. Plant Pathol.* **154**, 931–942 (2019).
- 784 48. Gu, C. F., Bastiaans, L., Anten, N. P. R., Makowski, D. & van der Werf, W. Annual intercropping suppresses  
785 weeds: A meta-analysis. *Agric. Ecosyst. Environ.* **322**, 107658 (2021).
- 786 49. Scott, D. & Freckleton, R. P. Crop diversification and parasitic weed abundance: a global meta-analysis. *Sci.*  
787 *Rep.* **12**, 19413 (2022).
- 788 50. Ramkat, R. C., Wangai, A. W., Ouma, J. P., Rapando, P. N. & Lelgut, D. K. Cropping system influences Tomato  
789 spotted wilt virus disease development, thrips population and yield of tomato (*Lycopersicon esculentum*). *Ann.*  
790 *Appl. Biol.* **153**, 373–380 (2008).
- 791 51. Wang, Y. Q. et al. Understanding biodiversity effects on trophic interactions with a robust approach to path  
792 analysis. *Cell Rep. Sustain.* **2**, 100362 (2025).
- 793 52. Wang, X. Y. et al. Microclimate, yield, and income of a jujube-cotton agroforestry system in Xinjiang, China.  
794 *Ind. Crop. Prod.* **182**, 114941 (2022).
- 795 53. Tang, X. Y. et al. Intercropping legumes and cereals increases phosphorus use efficiency; a meta-analysis.  
796 *Plant Soil* **460**, 89–104 (2021).
- 797 54. Inderjit, I. & Weiner, J. Plant allelopathic interference or soil chemical ecology? *Perspect. Plant Ecol. Evol.*  
798 *System.* **4**, 3–12 (2001).
- 799 55. Thorsted, M. D., Weiner, J. & Olesen, J. E. Above- and below-ground competition between intercropped winter  
800 wheat *Triticum aestivum* and white clover *Trifolium repens*. *J. Appl. Ecol.* **43**, 237–245 (2006).
- 801 56. Andersen, M. K., Hauggaard-Nielsen, H., Weiner, J. & Jensen, E. S. Evaluating competitive dynamics in two  
802 and three component intercrops. *J. Appl. Ecol.* **44**, 545–551 (2007).
- 803 57. dos Santos-Cordeiro, C. F., Rodrigues, D. R. & Echer, F. R. Cover crops and controlled-release urea decrease  
804 need for mineral nitrogen fertilizer for cotton in sandy soil. *Field Crop. Res.* **276**, 108387 (2022).
- 805 58. Root, R. B. Organization of a plant-arthropod association in simple and diverse habitats: the fauna of collards

- 806 (*Brassica oleracea*). *Ecol. Monogr.* **43**, 95–124 (1973).
- 807 59. Loreau, M. et al. Biodiversity as insurance: from concept to measurement and application. *Biol. Rev.* **96**, 2333–  
808 2354 (2021).
- 809 60. Schuldt, A. et al. Tree diversity promotes insect herbivory in subtropical forests of south-east China. *J. Ecol.*  
810 **98**, 917–926 (2010).
- 811 61. Osipitan, O. A., Dille, J. A., Assefa, Y. & Knezevic, S. Z. Cover crop for early season weed suppression in  
812 crops: systematic review and meta-analysis. *Agron. J.* **110**, 2211–2221 (2018).
- 813 62. Weiner, J. Applying plant ecological knowledge to increase agricultural sustainability. *J. Ecol.* **105**, 865–870  
814 (2017).
- 815 63. Kunz, C., Sturm, D. J., Varnholt, D., Walker, F. & Gerhards, R. Allelopathic effects and weed suppressive  
816 ability of cover crops. *Plant Soil Environ.* **62**, 60–66 (2016).
- 817 64. Hooks, C. R. R., Wang, K. H., Ploeg, A. & McSorley R. Using marigold (*Tagetes* spp.) as a cover crop to  
818 protect crops from plant-parasitic nematodes. *Appl. Soil Ecol.* **46**, 307–320 (2010).
- 819 65. Mishra, S. & DiGennaro, P. Root-knot nematodes demonstrate temporal variation in host penetration. *J.*  
820 *Nematol.* **52**, 1–8 (2020).
- 821 66. Costa, A. et al. Crop rotational diversity can mitigate climate-induced grain yield losses. *Global Change Biol.*  
822 **30**, e17298 (2024).
- 823 67. Tamburini, G. et al. Agricultural diversification promotes multiple ecosystem services without compromising  
824 yield. *Sci. Adv.* **6**, eaba1715 (2020).
- 825 68. Beillouin, D., Ben-Ari, T., Malézieux, E., Seufert, V. & Makowski D. Positive but variable effects of crop  
826 diversification on biodiversity and ecosystem services. *Global Change Biol.* **27**, 4697–4710 (2021).
- 827 69. MacLaren, C. et al. Long-term evidence for ecological intensification as a pathway to sustainable agriculture.  
828 *Nat. Sustain.* **5**, 770–779 (2022).
- 829 70. Nakagawa, S. et al. A robust and readily implementable method for the meta-analysis of response ratios with  
830 and without missing standard deviations. *Ecol. Lett.* **26**, 232–244 (2023).
- 831 71. van Houwelingen, H. C., Arends, L. R. & Stijnen, T. Advanced methods in meta-analysis: multivariate  
832 approach and meta-regression. *Statist. Med.* **21**, 589–624 (2002).
- 833 72. Nakagawa, S., Yang, Y., Macartney, E. L., Spake, R. & Lagisz, M. Quantitative evidence synthesis: a practical  
834 guide on meta-analysis, meta-regression, and publication bias tests for environmental sciences. *Environ. Evid.*  
835 **12**, 8 (2023).
- 836 73. Jin, Y. & Qian, H. V. PhyloMaker: an R package that can generate very large phylogenies for vascular plants.  
837 *Ecography* **42**, 1353–1359 (2019).
- 838 74. Jin, Y. & Qian, H. V. PhyloMaker2: An updated and enlarged R package that can generate very large  
839 phylogenies for vascular plants. *Plant Diversity* **44**, 335–339 (2022).
- 840 75. Ruppert, T. Meta-analysis: How to quantify and explain heterogeneity? *Eur. J. Cardiovas. Nur.* **19**, 646–652  
841 (2020).
- 842 76. Nakagawa, S. et al. orchaRd 2.0: An R package for visualising meta-analyses with orchard plots. *Methods Ecol.*  
843 *Evol.* **14**, 2003–2010 (2023).

- 844 77. Nakagawa, S. et al. The orchard plot: Cultivating a forest plot for use in ecology, evolution, and beyond. *Res.*  
845 *Syn. Meth.* **12**, 4–12 (2021).
- 846 78. Emmenegger, C. & Bühlmann, P. Plug-in machine learning for partially linear mixed-effects models with  
847 repeated measurements. *Scand. J. Statist.* **50**, 1553–1567 (2023).
- 848 79. Bartoń K. MuMIn: Multi-Model Inference. R package version 1.47.5, [https://CRAN.R-](https://CRAN.R-project.org/package=MuMIn)  
849 [project.org/package=MuMIn](https://CRAN.R-project.org/package=MuMIn) (2023).
- 850 80. Egger, M., Davey, S. G., Schneider, M. & Minder C. Bias in meta-analysis detected by a simple, graphical test.  
851 *BMJ.* **315**, 629–634 (1997).
- 852 81. Nakagawa, S. et al. Methods for testing publication bias in ecological and evolutionary meta-analyses.  
853 *Methods Ecol. Evol.* **13**, 4–21 (2022).
- 854 82. Duval, S. & Tweedie, R. Trim and fill: A simple funnel-plot-based method of testing and adjusting for  
855 publication bias in meta-analysis. *Biometrics* **56**, 455–463 (2000).
- 856 83. Nakagawa, S. & Santos, E. S. A. Methodological issues and advances in biological meta-analysis. *Evol. Ecol.*  
857 **26**, 1253–1274 (2012).
- 858 84. R Core Team. R: A Language and Environment for Statistical Computing. R Foundation for Statistical  
859 Computing, Vienna, Austria (2023).
- 860 85. Viechtbauer, W. Conducting Meta-Analyses in R with the metafor Package. *J. Stat. Soft.* **36**, 1–48 (2010).
- 861 86. Pinheiro, J., Bates, D. & R Core Team. nlme: Linear and Nonlinear Mixed Effects Models. R package version  
862 3.1-163. <https://CRAN.R-project.org/package=nlme> (2023).
- 863 87. Lajeunesse, M. J. Bias and correction for the log response ratio in ecological meta-analysis. *Ecology* **96**, 2056–  
864 2063 (2015).
- 865 88. Wan, N. F. Data and code for “Global evidence that plant diversity suppresses pests and promotes plant  
866 performance and crop production”. Zenodo <https://doi.org/10.5281/zenodo.17568135> (2025).