

BMJ Open

UNDERSTANDING REPEATED NON-ATTENDANCE IN HEALTH SERVICES - PILOT ANALYSIS OF ADMINISTRATIVE DATA AND FULL STUDY PROTOCOL FOR A NATIONAL RETROSPECTIVE COHORT

| | |
|---------------------------------|--|
| Journal: | <i>BMJ Open</i> |
| Manuscript ID | bmjopen-2016-014120.R1 |
| Article Type: | Protocol |
| Date Submitted by the Author: | 14-Dec-2016 |
| Complete List of Authors: | Williamson, Andrea; University of Glasgow, GPPC, School of Medicine, Dentistry and Nursing, MVLS Ellis, David; University of Lancaster Wilson, Philip; University of Aberdeen McQueenie, Ross; University of Glasgow, GPPC, School of Medicine, Dentistry and Nursing, MVLS McConnachie, Alex; University of Glasgow, Robertson Centre for Biostatistics |
| Primary Subject Heading: | Public health |
| Secondary Subject Heading: | General practice / Family practice, Epidemiology, Health services research |
| Keywords: | missed appointments, data linkage, administrative data, health utilisation, health inequalities, social vulnerability |
| | |

SCHOLARONE™
Manuscripts

only

1
2
3 **UNDERSTANDING REPEATED NON-ATTENDANCE IN HEALTH SERVICES**
4
5 **- PILOT ANALYSIS OF ADMINISTRATIVE DATA AND FULL STUDY**
6
7
8 **PROTOCOL FOR A NATIONAL RETROSPECTIVE COHORT**
9

10
11 Andrea E Williamson (corresponding author), General Practice and Primary Care,
12 School of Medicine, Dentistry and Nursing, MVLS, University of Glasgow.
13
14 andrea.williamson@glasgow.ac.uk Tel: 0141 330 8330
15
16

17
18
19 David A Ellis, Department of Psychology, University of Lancaster.
20
21 d.a.ellis@lancaster.ac.uk
22

23
24 Philip Wilson, Centre for Rural Health, Institute of Applied Health Sciences
25
26 University of Aberdeen. P.wilson@abdn.ac.uk
27

28
29
30 Ross McQueenie, General Practice and Primary Care, School of Medicine, Dentistry
31
32 and Nursing, MVLS, University of Glasgow. Ross.mcQueenie@glasgow.ac.uk
33

34
35 Alex McConnachie, Robertson Centre for Biostatistics, Institute of Health and
36
37 Wellbeing, MVLS University of Glasgow. Alex.McConnachie@glasgow.ac.uk
38
39

40
41
42
43 Word count 4869 excluding abstract, figures, qualitative data quotes and
44
45 references.
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

ABSTRACT

Introduction

Understanding the causes of low engagement in health care is a prerequisite for improving health services' contribution to tackling health inequalities. Low engagement includes missing health care appointments. Serially (having a pattern of) missing general practice appointments may provide a risk marker for vulnerability and poorer health outcomes.

Methods and analysis

A proof of concept pilot utilising general practice (GP) appointment data and a focus group with GPs informed the development of missed appointment categories: patients can be classified based on the number of appointments missed each year. The full study, using a retrospective cohort design, will link routine health service and education data to determine the relationship between general practice appointment attendance, health outcomes, health care utilization, preventive health activity, and social circumstances taking a life course approach and using data from the whole journey in NHS health care. 172 practices will be recruited (approximately 900,000 patients) across Scotland. The statistical analysis will focus on two key areas; factors that predict patients who serially miss appointments, and serial missed appointments as a predictor of future patient outcomes. Regression models will help understand how missed appointment patterns are associated with patient and practice characteristics.

2

1
2
3 We shall identify key factors associated with serial missed appointments and
4 potential interactions that might predict them.
5
6
7

8 **Ethics and dissemination**

9

10
11 The results of the project will inform debates concerning how best to reduce non-
12 attendance and increase patient engagement within health care systems.
13 Significant non-academic beneficiaries include governments, policy-makers and
14 medical practitioners. Results will be disseminated via a combination of academic
15 outputs (papers, conferences), social media, and through collaborative public
16 health/policy fora.
17
18
19
20
21
22
23

24 **STRENGTHS AND LIMITATIONS**

25

- 26 • This study will answer important question relating to the health service
27 component of tackling health inequalities
- 28 • A large dataset enables the researchers to follow patients' journey across the
29 whole health care system
- 30 • The study utilises data security and linkage capabilities in a sensitive and
31 robust manner
- 32 • The study has a clear yet flexible data analysis plan utilising the expertise of a
33 multi-disciplinary research team
- 34 • There are limitations of using administrative data from a range of data sources
35 of variable data quality.
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54

55 **KEYWORDS**

56
57
58
59
60

1
2
3 Missed appointments, data linkage, administrative data, primary care, health
4
5 utilisation, health promotion, health inequalities, social vulnerability
6
7
8
9

10 11 INTRODUCTION

12
13
14 Tackling health inequalities is a global health priority¹ and for health service
15
16 provision to have an effective role, we should understand better the reasons
17
18 behind, risks associated with, and needs of patients who do not engage effectively
19
20 with health care provision (even if it is free at the point of access); and tailor
21
22 services better to meet those needs. There remains a lack of published work
23
24 concerning repeated missed appointments, but previous research typically focuses
25
26 on the financial costs associated with non-attendance. One estimate has placed
27
28 the cost of missed United Kingdom (UK) general practice (community based family
29
30 medicine) appointments at £150 million per year². More recent Scottish
31
32 government data suggest that each missed hospital outpatient appointment costs
33
34 National Health Services (NHS) Scotland £120³. International data on costs to
35
36 health care systems are sparse. In a complex adaptive system such as health care,
37
38 the financial costs are contestable because clinicians will 'catch up' or get on with
39
40 other care or administrative tasks. What is important are the costs of opportunities
41
42 missed for improving patients' health and the potential for substantial long-term
43
44 savings to health systems.
45
46
47
48

49
50 To date studies investigating missed appointments have focused on single missed
51
52 appointments or single disease areas and have indicated they are associated with
53
54 poorer health outcomes³⁻⁶. Studies of single missed appointments have produced
55
56 conflicting results when it comes to designing effective interventions that can
57
58

59
60
4

1
2
3 increase attendance⁷⁻¹⁰. This may be due to a reliance on small samples in
4
5 disparate settings¹¹⁻¹⁵ and conflation of patients who occasionally miss
6
7 appointments with patients who have an established pattern of missing many.
8

9
10 The Health and Social Care Information Centre in England has recently published
11
12 data about repeated missed appointments. From their analysis of recorded missed
13
14 outpatient hospital appointments in England one in 50 patients (65,590 of 3.5
15
16 million) who missed an appointment failed to attend three or more further
17
18 appointments within three months¹⁶.
19

20
21 We hypothesise that repeated missed appointments reflect a pattern of behaviour.
22
23 We use the term 'serially' missing appointments to reflect this pattern, which may
24
25 be interrupted by attended appointments. Clinicians do report that patients who
26
27 serially miss appointments are of particular concern because they may have very
28
29 poor health, may be socially disadvantaged and are high users of unscheduled care
30
31 compared to patients who occasionally or never miss appointments¹⁷.
32
33

34
35 There is accumulating evidence that negative experiences in early life have
36
37 pervasive consequences for health over the life course including 'extensive
38
39 evidence of a strong link between early adversity and a wide range of health-
40
41 threatening behaviours'¹⁸. This body of work therefore provides a conceptual
42
43 framework for better understanding 'chaotic lives'¹⁹ as an explanatory factor in
44
45 health utilization behaviours such as missed appointments, and introduces the
46
47 possibility that serial missed appointments contribute to the inverse care law,
48
49 which states that health care provision is least likely to be provided to those that
50
51 need it most²⁰.
52
53

54
55 In the UK, publicly funded general practice (GP) provides almost universal
56
57 coverage for the population and generates around 90% of health contacts.
58

59
60
5

1
2
3 Appointment making is typically under the control of each patient directly.
4
5 General practice appointments therefore provide a sensible starting point for this
6
7 study of health and other outcomes across patients' life course. Subsequent results
8
9 will also have relevance for global health systems where patients have direct
10
11 access to a wider range of health care specialties.
12
13

14
15 Scotland has an established data linkage infrastructure which is under continuous
16
17 development. This pathfinder study will for the first time link large general
18
19 practice datasets (including appointment data) with data from across patients'
20
21 whole journey through health care.
22
23

24
25 The overarching study question is: is serially missing GP appointments a risk
26
27 marker for vulnerability and poorer health outcomes and thus a useful target for
28
29 developing interventions to improve engagement in health promoting care across
30
31 the health system?
32
33

34 35 36 **Aim and Research questions**

37
38 The overall aim of the study is to determine the relationship between general
39
40 practice appointment attendance, health care utilization, preventive health
41
42 activity, health outcomes, and social circumstances taking a life course approach
43
44 and using extracted health service and other relevant administrative data.
45
46

47
48 A pilot study sought to answer the first research question described below. The
49
50 subsequent questions underpin the full research protocol which compares cohorts
51
52 of Scottish patients (from birth to older age) who never, occasionally and serially
53
54 miss GP appointments.
55
56

1
2
3 Figure 1: Study research questions
4

5 An introduction to the full study protocol is described, followed by the methods
6 and results from a mixed methods pilot study that informed the protocol. A
7 description of protocol participants, data sources, variables and statistical analysis
8 then follows.
9
10
11
12

13 14 15 16 **METHODS AND ANALYSIS** 17

18
19 The full study protocol is for a retrospective cohort study of GP practice patient
20 records linked with secondary care and education administrative records in
21 Scotland.
22
23

24
25 The study commenced in July 2015 and will finish in December 2017. A pilot study
26 was conducted between July and September 2015 which is described next. The
27 cohort of 909,073 GP patient records for the full study was available in the
28 National Safehaven from September 2016 and analysis of these data is underway.
29 Permissions to access education data is secured, and the outcome of linkage
30 permissions for health data is not yet confirmed
31
32
33
34
35
36
37
38
39

40 41 **Pilot study** 42

43 The pilot study was separated into 2 sub-sections: a focus group to inform and
44 refine definition development (research question 1) and a 'proof of concept'
45 quantitative data analysis.
46
47
48

49 **Methods**

50 *Focus group* 51 52

53
54 A focus group was conducted in September 2015 with five GP participants. A focus
55 group was judged the most appropriate method to use because we aimed to
56
57
58
59
60

1
2
3 promote discussion of the topic such that participants would be able to compare
4 and contrast their own experiences with others from a range of practice and
5 professional experience settings²¹. Linked to this was the aim of asking participants
6 to make sense of, and provide feedback on the presented pilot data. The GPs
7 were a convenience, purposive sample based on two main principles. The first took
8 into account the evidence surrounding single missed appointments. This describes
9 missed appointments being more common in deprived, urban practices. The
10 sample therefore included GPs who worked in deprived and affluent urban areas
11 and a practice with a significant rural practice population from Scotland. Second,
12 the sample included the views of frontline GPs and GPs who had a range of
13 strategic roles in practice development and general practice management, locally
14 and nationally. AEW and PW utilised their professional knowledge of GP networks
15 and practice profiles to approach and recruit participants. Five out of twelve GPs
16 contacted were able to attend the focus group. Each GP contacted reported that
17 they felt this was an important topic worthy of attention. Barriers to attending
18 were location of the focus group (conducted in Glasgow) and managing time away
19 from other professional work. Additional file 1 describes each participant's
20 characteristics. Detailed information about participants' practice characteristics
21 was not collected. Three of the participants knew each other from their
22 professional roles outside of clinical practice. AEW conducted the focus group and
23 the analysis was conducted using Framework Analysis. Framework Analysis is a
24 useful thematic analysis approach especially when considering a focussed topic like
25 this one. Also in the context of being part of a larger mixed methods study,
26 epistemologically its use was a good fit²². DAE attended the focus group and
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 presented initial results from the 'proof of concept' pilot (described next) for
4 discussion. Additional file 2 describes the topics covered in the focus group.
5
6

7
8 *Proof of concept*
9

10 Research that uses GP appointment data has not previously been conducted using
11 the clinical recording systems in the Scottish NHS. A proof of concept pilot study
12 was undertaken utilising the NHS Trusted Third Party (TTP) Albasoft with 67,705
13 patient records to determine whether retrieving appointment data was feasible, to
14 refine other data parameters, and to inform the definition development as
15 described in research question 1. An additional confidentiality control ensured that
16 the research team did not know the identity of the recruited GP practices.
17
18
19
20
21
22
23
24

25 Additional file 3 describes the definition and role of TTPs.
26
27

28 Albasoft purposively recruited 10 Scottish practices on our behalf with the practice
29 characteristics illustrated in figure 2.
30
31
32

33
34 Figure 2: Pilot practice recruitment
35

36
37 Data were cleaned and appointment rules applied to categorise appointments as
38 attended or missed (DNA). Additional file 4 describes this process. This was
39 primarily based on the 'in' and 'out' time recorded for each appointment. If this
40 was recorded as '0' then the appointment was classified as Did Not Attend (DNA).
41
42
43
44
45
46 For each patient the total number of appointments made during the three-year
47 period was calculated as well as the number and percentage of appointments
48 missed. Appointments that were recorded incorrectly in the system were removed
49 at this stage. This included appointments where administrative records had
50 remained open for over 24 hours, making it difficult to confirm that these were
51
52
53
54
55
56
57
58
59
60

genuine appointments and not administrative errors. The pilot appointment rules are set out in table 1 below.

| Data description | Reason for removal |
|--------------------------------|------------------------------------|
| total appointment time < 0 min | Record open for more than 24 hours |
| total waiting time < 0 min | Record open for more than 24 hours |
| appointment <2 min | Not a medical appointment |
| administrator slot | Not a medical appointment |

Table 1 Rules to identify genuine appointments

Results

Focus group

Focus group participants reported making clear distinctions between patients who occasionally miss appointments and those who miss many. Patients who occasionally miss appointments do so because a crisis or another understandable circumstance has arisen; patients who serially miss appointments (SMA), described as missing more than two or three appointments can be easily identified by GPs. They were described as tending to have mental health, addiction, and/or social issues. They were described as high risk or vulnerable with concerns about their wider family. Patients who SMA were viewed as being different from the general GP population and being more likely to have ‘chaotic’ lifestyles associated with housing instability, money problems, a “*panicked lifestyle*”(P2). Patients who SMA were also described as being unable to manage GPs’ expectation of them and fit into GPs’ pre-determined slots. “*there’s the occasional DNA which are quite normal and often those are quite significant [in total numbers for the practice] but the serial people I think that’s a reflection of the chaos in their life whether that’s you know- mental health or issues with the social functioning- and inability to manage our expectation of them- to fit into our pre-defined slots.*” (P5)

1
2
3 All participants agreed with that view. However one participant also considered
4 that not all patients who SMA can be viewed as high risk; that instead some
5 patients do not value free health care. It was reported that some patients who SMA
6 go on to book another appointment the next day; *"I don't think it's the value of*
7 *the GP- I think it's the value of that appointment- I think the fact that it's, if you*
8 *don't miss it, if you miss it is no big deal you just make another one"* (P4).
9

10
11
12 Missed appointments were viewed as being more prevalent in practices in deprived
13 settings, but occurred in affluent areas too. In the affluent setting they were
14 important for a minority of marginalised, isolated patients with the same profile as
15 described above-who were viewed as living 'chaotic' lives.
16

17
18
19 Practices represented in the focus group do not have protocols for managing
20 patients who serially miss appointments (SMA) because response is dependent on
21 the patient's context. GPs understood that SMAs usually mean patients with
22 complex needs with workload implications for the practice. Strategies described
23 were varied, including allowing patients only to book on the day; *"my impression*
24 *is that deprived practices have a much higher percentage of on the day*
25 *appointments because they skew it towards people that don't attend"* (P3), seeing
26 the patient when they walk in, or the GP booking the follow up appointment for
27 the patient- a relationship building strategy. This could still lead to patients
28 missing an appointment, even just a couple of hours after it was made. It was
29 reported that some practices do remove patients from their list for SMA and this
30 created tension with other practices.
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51

52
53 The focus group were also asked to comment on the results from the proof of
54 concept initial data and they made recommendations about the full study design
55 described in Figure 3.
56
57
58
59
60

Figure 3 Focus group recommendations for the full study design

Proof of concept

A pilot analysis of 67,705 patient records showed that while just over 60% of our sample missed no appointments, over 30% missed one or more appointment during the three-year period with nearly 10% of patients missing three or more appointments.

Assuming that our final sample provides a similar distribution, we will classify patients based on the number of appointments missed as follows:

***Never* missed appointments: 0 per year average over 3 year period**

***Low* missed appointments: <1 per year average over 3 year period**

***Medium* missed appointments: 1-2 per year average over 3 year period**

***High* missed appointments: >2 per year average over 3 year period**

Our sampling both in the pilot data stage and the final full study sample was conducted such that we were likely to get a representative sample of Scottish patients and practices. Because our pilot sample was large, it is appropriate to assume that this will scale-up accordingly for the full study. The distribution of missed appointments also suggested useful categories based on integer numbers of missed appointments per year. This will be helpful for policy and clinical stakeholders.

FULL STUDY PROTOCOL

Participants and study size

12

1
2
3 Our target recruitment of GP practices seeks to ensure that a spread of urban and
4 rural practices, affluent and practices characterised by serving areas of blanket
5 high socio-economic (Deep End) deprivation. The information request made to
6 practices can be viewed in Figure 4.
7
8
9
10

11
12
13
14 Figure 4 Information request sent to target practices
15
16

17 18 19 **Data sources and variables**

20 21 **GP data**

22
23 The TTP has recruited the practices on the study team's behalf and will undertake
24 some specific data aggregation before transferring the data securely to the
25 National Safehaven for analysis. 'Safe Havens are specialised, secure environments
26 supported by trained, specialist staff where data in electronic patient records can
27 be processed and linked with other health data (and/or non-health-related data)
28 and made available for analysis to facilitate research while protecting patient
29 identity and privacy'²³. These are: calculating urban rural classification, SIMD
30 decile, categorising ethnicity into 'non BME (Black and Minority Ethnicity)', 'visibly
31 BME', and 'non visible BME' and rounding travel distance to practice/emergency
32 department for each patient record to the nearest kilometre. Once in a Safehaven,
33 additional steps will be taken to ensure that acceptable anonymization principles
34 are being applied, especially in relation to reporting of sensitive social
35 vulnerability data and reporting of rare conditions.
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50

51
52 A new data file containing the appointment history for each patient record will be
53 generated, which will be merged with individual patient information (Additional
54 file 4 describes this process based on our pilot data set).
55
56
57
58

59 13
60

Appointment validation and categorisation

Each appointment will be coded based on session type recorded by the practice (eg book on day appointments, or immunization clinic) and then further by professional type (eg GP partner, practice nurse). These descriptions are determined by individual practices so categorisation will be conducted by the GPs in the research team. The appointment rules set out in the pilot study will be applied. A sensitivity analysis based on the time the appointment takes will then also be conducted by comparing a random sample of patient appointments as described in figure 5.

Figure 5 Random sample of GP appointments for validation and sensitivity analysis

The appointment rules will be refined based on this. The time interval cut-off for apparently attended appointments will be determined by utilising the time interval that most accurately matches to actual attended appointments. Slots designated non face to face appointments will then be removed leaving only attended and non- attended face- to- face appointments. The appointment categories described from the pilot study regarding non- attendance for all patients will then be applied to the yearly average number of missed appointments over the three year period to generate the four categories of patients for further analysis. Using an average over three years takes account of what is recognised in the frequent attenders (rather than non- attenders) literature- that patients' appointment behaviour may vary over time in relation to illness episodes or social crises²⁴.

Health and education data linkage

1
2
3 Linkage will be conducted as access permissions and data sets become available.
4
5 Each administrative data source is available for different time periods (e.g.
6
7 hospital inpatients since 1981 and education outcomes since 2002) and this will be
8
9 made explicit when interpreting the results. The TTP will provide the Safe Haven
10
11 indexing team a file containing the GP dataset Community Health Index (CHI)
12
13 number and other patient identifiers. Every patient in the Scottish NHS has a CHI
14
15 number, a unique identifier that is used as such across all NHS services in Scotland.
16
17 This forms the cohort for the study. All data providers will supply identifiers to be
18
19 probability matched to the study cohort by the Safehaven linkage team (based on
20
21 CHI number and using other patient identifiers probabilistically for the small
22
23 number of records where it is anticipated CHI will be missing), who will return a
24
25 set of unique index numbers for those individuals successfully matched to the
26
27 study cohort; each data provider will receive a different set of unique index
28
29 numbers, and will use these index numbers as the basis of their data extract. Each
30
31 data extract will be submitted to the Safehaven linkage team, who will replace the
32
33 different index numbers with a common number across all files. This common
34
35 number is the unique patient identifier that the research team will work from
36
37 during analysis.
38
39
40
41
42

43
44 Figure 6 Proposed data sets for linkage with GP data
45

46 **Bias**

47 Accounting for patient turnover

48
49 This study seeks to ensure the inclusion of patients who are marginalised and who
50
51 are often missing from health service studies. There is evidence of overlap
52
53 between patients who miss appointments and those who are removed from
54
55 practice lists²⁵, a recognition of the impact that geographical boundary areas have
56
57
58
59
60

1
2
3 on patients who move around²⁶; notwithstanding the gap in the literature about
4
5 registration interruptions for patients who may go to prison or patients who remain
6
7 unregistered once they are removed from GP practice lists. We will therefore
8
9 summarise the numbers of patients joining and/or leaving their practice during the
10
11 study period; with reasons where this information is available. We will seek to
12
13 provide a full analysis of the data available for these patients and compare these
14
15 with the patients who are registered for the 3 year study period. Patients who are
16
17 not registered with participating practices, and are being seen as 'temporary
18
19 residents' by these practices, are excluded from the study. This is because these
20
21 patients full clinical record is held by their registered GP so very limited
22
23 information is available. Temporary residents tend to be people on holiday in the
24
25 practice area but will include some people who would be considered marginalised.
26
27
28

29 30 **Statistical methods**

31
32 Our statistical analysis is based on the study being a retrospective cohort study.
33
34 We will focus on two key areas; predictors of high rates of serial missed
35
36 appointments, and serial missed appointments as a predictor of future patient
37
38 outcomes.
39

40
41 Patient characteristics and practice characteristics may be associated with high
42
43 rates of serial missed appointments. Analyses will initially be descriptive,
44
45 summarising the rate of missed appointments in relation to the other factors
46
47 recorded at the point of entry to the study. Associations with patient
48
49 characteristics will be assessed as a whole, and in relation to different types of
50
51 practices (e.g. separately in rural and urban practices). Subsequently, we will build
52
53 regression models (Poisson or Negative Binomial),²⁷ to help understand how our
54
55
56
57
58
59
60

1
2
3 categories of missed appointments are associated with patient and practice
4 characteristics.
5
6

7
8 When considering other outcomes in relation to serial missed appointments, the
9 missed appointment rate category (none, <1, 1-2, or >2 per year) will be the
10 predictor variable. Appropriate regression models, according to the outcome, will
11 be used to assess whether any associations with serial missed appointment rates
12 are independent of other patient- or practice-level factors. Conflicting interactions
13 will be controlled for by using an 'offset term' in our negative binomial model
14 which accounts for number of appointments made or any other relevant factors.
15
16

17 We also plan to measure relevant quantitative variables (described next) recorded
18 *during* the study interval associated with having a lot of missed appointments. We
19 will explore whether these *differ* from the predictive factors already recorded at
20 entry to the study.
21
22

23 **Quantitative variables**

24
25 The following potential predictors of frequent non- attendance will be analysed:
26
27

28 **Demographics**

29 Patients' age, gender, minority ethnic group status (where available), deprivation
30 decile, rural/urban split, number of address moves, distance lived from GP
31 practice and distance from nearest A&E will be explored.
32

33 **Health conditions**

34 Health conditions will be reported using separate categories:
35

- 36 1. The incidence of multi-morbidity calculated from extracted Read codes based
37 on previous counts in Scotland²⁸
38
39

1
2
3 2. Descriptions of health conditions based on the priority 1 Read codes that GP
4 practices in Scotland use to populate patients' key information summaries (KIS) for
5 GP out of hours services. This is novel work as a coding structure has not previously
6 been applied to these Read codes. Read codes are the clinical coding system used
7 in UK general practice to record, clinical and administrative activity and diagnoses.
8

9
10
11
12
13 3. A count of psychotropic medicine prescriptions based on the British National
14 Formulary. This is in order to describe levels of psychological morbidity that are
15 not captured by diagnostic criteria.
16
17
18

19
20
21 4. These variables will then be compared to the ICD 10 coding data from patients'
22 secondary care linked data compiled from hospital admissions and outpatient
23 attendances. Diagnostic data from emergency department attendance was
24 deemed not of sufficient quality to utilise.
25
26
27
28

29 Social Vulnerability

30
31
32 One aspect of this study which is particularly ground-breaking is our investigation
33 of retrievable information about patients' social vulnerability. The Adverse
34 Childhood Experiences (ACE) questionnaire²⁹ will be utilised as a template to
35 match its nine descriptors of adversity to relevant Read codes in the patient's GP
36 record. In addition, coding that maps the consequences of ACE will be analysed. A
37 recent quantitative evaluation of Severe and Multiple Disadvantage will also be
38 matched to GP Read codes. This examines the overlap of patients being homeless,
39 in substance misuse services, or in prison over the preceding year³⁰. Further, an
40 exploration of additional Read codes that describe social vulnerability will be
41 mapped. An anonymised text search linked to Read codes from the dataset will
42 provide additional information about social vulnerability as it is recorded in the
43 free text portion of GP records. Taken together, these will provide the first
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 research evidence about the breadth and depth of social vulnerability recording by
4
5 GPs.
6

7 Health care utilisation

8
9
10 Read coding in relation to cervical, breast and bowel screening attendance will be
11
12 retrieved in addition to the proportion of patients who have had their blood
13
14 pressure checked and have participated in child health surveillance and
15
16 vaccination programmes across the life course. A sub-analysis of utilisation of
17
18 practice nurse and other health care professional's' appointments in the practice
19
20 will also be conducted and include an exploration of the relationship between
21
22 attending all primary care appointments and categories of non- attendance. This is
23
24 because data from the GP focus group suggested there is overlap between patients
25
26 who are serial non-attenders with patients who are very frequent attenders. We
27
28 will therefore consider the rate of attending appointments as a potential predictor
29
30 of the rate of non-attendance. Referrals that GPs make into other primary and
31
32 secondary care services will also be analysed. Outpatient attendances, hospital
33
34 admissions and utilisation of emergency departments, NHS 24 triage, GP out of
35
36 hours, and ambulance services will also be analysed when linked data become
37
38 available with a specific focus on how this relates to unmet need, for example how
39
40 might GP appointment category relate to patterns of other health care utilisation
41
42 between scheduled and unscheduled secondary care use.
43
44
45
46

47 Health care engagement

48
49 An analysis of GP Read codes and linked secondary care data will be carried out in
50
51 the following categories:
52

- 53 1. Patients not attending primary and secondary care appointments
- 54
- 55 2. Patients refusing screening
- 56
- 57
- 58

59 19
60

1
2
3 3. Patients being exception-reported (ie excluded from the denominator
4 population) from the Quality and Outcomes Framework (QOF) system for
5 performance measurement in general practice
6
7

8
9
10 4. Practices' measures of non-engagement with care for long term conditions
11

12 5. Patients taking 'irregular discharge' from hospital (when patients leave against
13 medical advice)
14

15
16 6. Patients not waiting to be seen in emergency departments
17

18
19 Family linkage

20
21 Diagnoses of children who are able to be linked through family linkage will be
22 analysed based on their mother's appointment category. This is contingent on the
23 child being included in the practice study population.
24
25

26
27 Education data

28
29 Attendance at school, exclusion from school, and educational attainment when
30 leaving school will be made with approximately a sixth of our patient cohort for
31 whom linked education data is available. This has the potential to inform future
32 planning around earlier interventions to reduce serial missed appointments.
33
34

35
36 Practice level data

37
38 Each patient record will be allocated a unique practice ID enabling the research
39 team to analyse each patient record output clustered by practice. This will be
40 proportion of patients aged over 75, by ethnicity (proportion BME), patient
41 rurality, patient level of deprivation decile, patient distance to practice, distance
42 to A&E appointments offered/engaged, days from when appointment is made,
43 multi-morbidity count, ACE score more than 4, Severe and Multiple Disadvantage
44 score, hospital referrals, and proportion of each appointment category by practice.
45
46

47
48 These analyses and output will be refined as the study proceeds taking patient
49
50

51
52
53
54
55
56
57
58
59 20
60

level findings and multilevel modelling that characterises the respective contributions of practice- and individual-level factors to missed appointment patterns.

Health outcomes

Mortality data regarding date and cause of death will be utilised from GP and linked data. This will sit alongside additional linked obstetric outcomes (from the Scottish Birth Record) for relevant women.

Table 2 summarise the quantitative variables for analysis

| Data categories | variables |
|-------------------------|--|
| Patient demographics | Age Sex ethnicity Count of address moves Distance to practice Distance to A&E SIMD decile Rural8 score |
| Health conditions | Multi-morbidity count Priority 1 read codes Psychotropic medication prescribing (BNF chapter) Secondary Health care diagnoses (inpatient and outpatient) |
| Social vulnerability | Adverse Childhood Experiences Severe and Multiple Disadvantage General social vulnerability coding frame |
| Health care utilisation | Breast screening Bowel screening Cervical screening BP checked Child health surveillance Vaccinations Practice nurse appointments Other health care professional appointments Primary care attendance distribution Hospital referrals Outpatient attendances Hospital admissions emergency departments attendance NHS 24 triage GP out of hours ambulance services callouts |

| | |
|-----------------------------|---|
| Health care engagement | DNA codes Refused screening QOF exempt Inappropriate use codes Self-discharge codes |
| Study exit | Patient death Patient moved practice |
| Family linkage | Secondary health care linkage with mother and child |
| Education data | School attendance School exclusion School attainment |
| Health outcomes | Cause of death |
| GP Practice characteristics | Practice list size Patient age distribution Ethnicity category distribution Patient rur8 score distribution Patient SIMD score distribution Patient distance to practice distribution Patient distance to A&E distribution Number of appointments offered/patients engaged past 3 years distribution Number of days since appointments made distribution Patient multi-morbidity score distribution Patient ACE score distribution Patient SMD score distribution Patient hospital referrals distribution Primary care attendance pattern distribution |

Table 2 Summary of quantitative categories and variables

ETHICS AND DISSEMINATION

This pathfinder linkage retrospective cohort study is necessarily complex in design and implementation because although cross-sectional it seeks to take a life course approach and follow the patients' journey through the health care system. Careful attention and significant resource has been devoted to the consideration of patient privacy and confidentiality. This has been integrated throughout the design of the study alongside the necessary data access and handling permissions. Additionally a study of this nature, which involves stakeholders across the NHS and other public services, requires a flexible time frame to allow access to raw data and to share

1
2
3 findings between members of the research team based in several institutions.
4
5 The proof of concept pilot did not require ethical approval because it was
6
7 considered service evaluation with the agreement we would not publish any results
8
9 about the practices who took part. Ethical permission to conduct the GP focus
10
11 group and publish the results was obtained by the MVLS ethics committee,
12
13 University of Glasgow (ref 200140181). A letter of comfort was obtained from the
14
15 West of Scotland NHS ethics committee and the MVLS ethics committee confirming
16
17 that the full study did not need health service ethics permissions. Multi- site NHS
18
19 R&D approval for the full study was obtained for all Scottish Health Boards
20
21 (NRS16/186358).
22
23

24
25 Due to the sensitive nature of administrative data from the NHS and public
26
27 education system in Scotland, the datasets generated and/or analysed during the
28
29 current study will not be publicly available. They have been made available to the
30
31 research team under controlled access and strictly for the purposes of this
32
33 research study only. Summary data, at the level of disclosure checked output from
34
35 the National Safehaven and statistical code, can be requested from the
36
37 corresponding author on reasonable request.
38
39

40 41 **Planned outputs**

42
43 Alongside peer reviewed academic papers reporting the findings described above,
44
45 the following additional outputs are planned.
46
47

48 49 **Data Visualisation**

50
51 Several web pages will be built to sit alongside key results. This will allow for the
52
53 rapid construction of interactive data visualisations which will be created using
54
55 “Shiny”³¹, a web application framework for R which is the statistical software used
56
57 for the study analysis. A simple platform will allow researchers and collaborators
58
59

1
2
3 to interact with the analyses in real-time and generate custom tables and graphs
4
5 as required. It can also provide non-experts with access to simple and complex
6
7 statistical analysis using a point-and-click interface. This will not rely on raw data
8
9 and will simply pull information from the summary descriptive analyses.
10

11 12 13 Case Studies

14
15 We also intend to use case studies to develop and illustrate our findings throughout
16
17 the course of all our analyses. For example, we will be able to identify typical
18
19 patient profiles of those who appear to miss many appointments in a very short
20
21 period of time and compare these events with short and long-term health
22
23 outcomes.
24

25 26 Conclusion

27
28
29 We shall identify key factors associated with serial missed appointments ranked in
30
31 order of importance as described above, but given the large sample size we shall
32
33 also be able to consider potential interactions that might predict serially missed
34
35 appointments.
36

37
38 Finally, this approach also explores the theory that low engagement with health
39
40 care should be viewed as a health harming behaviour, and will inform the debate
41
42 about tackling health inequalities at the health service delivery level. Moving from
43
44 theory into application, the results will allow us to better understand and develop
45
46 future interventions to reduce serial missed appointments.
47
48

49 REFERENCES

50
51
52 1. Commission on Social Determinants of Health. Closing the gap in a generation:
53
54 health equity through action on the social determinants of health. Final Report of
55

1
2
3 the Commission on Social Determinants of Health. Geneva: World Health
4 Organisation, 2008.

5
6
7 2. George A, Rubin G. Non-attendance in general practice: a systematic review and
8 its implications for access to primary health care. *Family Practice* 2003;20:178-84.

9
10
11 3. Campbell K, Millard A, McCartney G, et al. Who is least likely to attend? An
12 analysis of outpatient appointment 'Did Not Attend' (DNA) data in Scotland.

13
14
15 Edinburgh: NHS Health Scotland, 2015.

16
17
18 4. Neal RD, Hussain-Gambles M, Allgar VL, et al. Reasons for and consequences of
19 missed appointments in general practice in the UK: questionnaire survey and
20 prospective review of medical records. *BMC Family Practice* 2005;6(47)

21
22
23 5. Mitchell AJ, Selmes T. Why don't patients attend their appointments?

24
25
26 Maintaining engagement with psychiatric services. *Advances in Psychiatric*
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Treatment 2007;13:423-34.

6. Ciechanowski PS, Russo J, Katon W, et al. Where is the patient? The association
of psychosocial factors and missed primary care appointments in patients with
diabetes. *General Hospital Psychiatry* 2006;28:9-16.

7. Lehmann T, Lehmann A, Olivet M, et al. Missed appointments at a Swiss
university outpatient clinic. *Public Health* 2007;121:790-99.

8. Neilson KM, Faergeman O, Foldspang A, et al. Cardiac rehabilitation: health
characteristics and socio-economic status among those who do not attend.

European Journal of Public Health 2008;18:479-83.

9. Cashman S, Savageau J, Lemay C, et al. Patient health status and appointment
keeping in an urban community health center. *Journal of health care for the Poor*
and Underserved 2004;15:474-88.

- 1
2
3 10. Masuda Y, Kubo A, Kokase A, et al. Personal features and dropout from
4 diabetic care. *Environment, Health Prevention Medicine* 2006;11:115-9.
5
6
7 11. Gucciardia E. A systemic review of attrition from diabetes education services:
8 strategies to improve attrition and retention research. *Canadian Journal of*
9
10 *Diabetes Care* 2008;32(5365)
11
12
13 12. Murdock A, Rogers C, Lindsay H, et al. Why do patients not keep their
14 appointments? Prospective study in gastroenterology outpatient clinic. *Journal of*
15
16 *Royal Society of Medicine* 2002;95:284-6.
17
18
19 13. Farley R, Wade T, Birchmore L. Factors influencing attendance at cardiac
20 rehabilitation among coronary heart disease patients. *European Journal of*
21
22 *Cardiovascular Nursing* 2003;2:205-12.
23
24
25 14. Woodward V. Non attendance at a nurse-run asthma clinic. *Journal Clinical*
26
27 *Nurse* 1998;7:380-1.
28
29
30 15. Killaspy H, Banerjee S, King M, et al. Prospective controlled study of
31
32 psychiatric out-patient non- attendance, Characteristics and outcome. *British*
33
34 *Journal of Psychiatry* 2000;176:160-5.
35
36
37 16. Health and Social Care Information Centre England. Provisional monthly topic
38
39 of interest: outpatient did not attend (DNA) appointment. London, 2014.
40
41
42 17. Watt G. Deep End Report 19. Access to Specialists. The 19th meeting of
43
44 'General Practitioners at the Deep End'. University of Glasgow, 2013.
45
46
47 18. Shonkoff JP, Garner AS, The Committee On Psychosocial Aspects Of Child And
48
49 Family Health COEC, Adoption, And Dependent Care, And Section On
50
51 Developmental And Behavioral Pediatrics,. The Lifelong effects of Early Childhood
52
53 Adversity and Toxic Stress. *Pediatrics* 2012; 129.
54
55
56
57
58
59
60

- 1
2
3 19. Hussain-Gambles M, Neal RD, Dempsey O, et al. Missed appointments in
4 primary care: questionnaire and focus group study of health professionals. *British*
5 *Journal of General Practice* 2004;54:108-13.
6
7
8
9
10 20. Tudor Hart J. The Inverse Care Law. *The Lancet* 1971;297:405-12.
11
12 21. Kitzinger J. Qualitative Research: Introducing focus groups. *BMJ*
13 1995;311(7000):299-302.
14
15
16 22. Ritchie J, Lewis J. Qualitative Research Practice. A Guide for Social Science
17 Students and Researchers. first ed. London: Sage Publications 2003.
18
19
20 23. The Scottish Government. Charter for Safe Havens in Scotland: Handling
21 Unconsented Data from National Health Service Patient Records to Support
22 Research and Statistics. Edinburgh2015 [Available from:
23 <http://www.gov.scot/Publications/2015/11/4783/2>].
24
25
26
27
28
29
30 24. Smits FTM, Brouwer HJ, van Weert HCP, et al. Predictability of persistent
31 frequent attendance: a historic 3-year cohort study. *British Journal of General*
32 *Practice* 2009;59:114-19.
33
34
35
36 25. Williamson AE, Mullen K, Wilson P. Understanding “revolving door” patients in
37 general practice: a qualitative study. *BMC Family Practice* 2014;15(1):1-11.
38
39
40 26. O'Reilly D, Steele K. The removal of patients who live outside the practice
41 boundary. *British Journal of General Practice* 2005;55:384-86.
42
43
44
45 27. Zuur AF, Ieno EN. A protocol for conducting and presenting results of
46 regression-type analyses. *Methods in Ecology and Evolution* 2016;7(6):636-45.
47
48
49
50 28. Barnett K, Mercer SMN, Watt G, et al. Epidemiology of Multi-morbidity and
51 implications for health care, research and medical education: a cross sectional
52 study. *The Lancet* 2012;380((9836)):37-43.
53
54
55
56
57
58
59
60

1
2
3 29. Bellis MA, Ashton K, Hughes K, et al. Welsh Adverse Childhood Experiences
4 (ACE) Study. Adverse Childhood Experiences and their impact on health-harming
5 behaviours in the Welsh adult population. NHS Wales, 2015.
6
7

8
9
10 30. Bramley G, Fitzpatrick S, Edwards J, et al. Hard Edges Mapping Severe and
11 Multiple Disadvantage England. England: Lankelly Chase Foundation, 2015.
12

13
14 31. Ellis DA, Meridian HL. Thinking outside the box: developing dynamic data
15 visualizations for psychology with Shiny. *Frontiers in Psychology* 2015;6
16
17
18

19 20 21 **AUTHORS' CONTRIBUTIONS**

22
23 AEW is principal investigator for the study. DAE, PW and AMcC are co-investigators
24 on the study and RMcQ is the research assistant. AEW conceived and developed the
25 initial research proposal, reviewed the literature, conducted and analysed the
26 pilot focus group, contributed to analysis and interpretation of the quantitative
27 pilot data, developed the predictors, outcomes and associations of interest and led
28 on writing the paper. DAE supported the development of the initial research
29 proposal, reviewed the literature, conducted and analysed the quantitative pilot
30 data, developed the statistical and output plan, and contributed to writing the
31 paper. PW supported the development of the initial research proposal, reviewed
32 the qualitative and quantitative pilot results, reviewed the statistical and output
33 plan and contributed to writing the paper. RMcQ reviewed the statistical and
34 output plan and contributed to writing the paper. AMcC provided expert statistical
35 input to the study as it was developed, reviewed the statistical and output plan
36 and contributed to writing the paper.
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53

54 All authors read and approved the final manuscript.

55 56 57 **FUNDING STATEMENT**

58
59 28
60

1
2
3 This study is supported by a Scottish Government Chief Scientist Office research
4 grant (CZH/4/41118) with Safehaven and data linkage costs supported in lieu by
5 the DSLS at Scottish Government. These funding bodies had no role in the design of
6 the study, or collection, analysis, and interpretation of data or in writing the
7 manuscript.
8
9

10 11 12 13 **COMPETING INTERESTS**

14
15 The authors declare they have no competing interests.
16
17

18 19 **ACKNOWLEDGEMENTS**

20
21 We would like to acknowledge all GP practices and GPs who took part in the pilot
22 study. Also colleagues at Scottish Government who are supportive of the study in a
23 variety of ways especially Ellen Lynch in the Health Analytics Division. Dave Kelly's
24 technical and procedural expertise, wisdom and patience as director of our TTP
25 Albasoft Ltd, underpins all of what has been achieved to date.
26
27
28
29
30
31
32
33
34
35
36

37 38 **FIGURE LEGENDS**

39
40 Figure 1: Study research questions

41
42 Figure 2: Pilot practice recruitment

43
44 Figure 3 Focus group recommendations for the full study design

45
46 Figure 4 Information request sent to target practices

47
48 Figure 5 Random sample of GP appointments for validation and sensitivity analysis

49
50 Figure 6 Proposed data sets for linkage with GP data
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For peer review only

1. What is a useful definition of never, occasionally and serially missing GP appointments?
2. What are the differences in illness profile, including multi-morbidity across patients' life course between these categories of patients?
3. What are the differences in health service utilization across the primary, secondary, scheduled and unscheduled health services?
4. What are the differences in health outcomes across the whole health system?
5. What are the differences in social vulnerability?
6. Can missed appointments be used to develop a proxy for unmet health need?
7. Can conclusions be drawn to inform rational resource allocation?
8. Is there evidence that supports the future development of targeted interventions to reduce missed appointments?

Figure 1: study research questions
Figure 1: study research quest
131x134mm (300 x 300 DPI)



1. 6 practices in urban and 4 practices in rural settings based on 'rural 8' classification²² scores
2. 7 of those practices in areas of high deprivation - based on Scottish Index of Multiple Deprivation (SIMD)²³ average patient scores for the registered list
3. 2 practices have high proportions of minority ethnic group patients based on previous work by Albasoft.

Figure 2: pilot practice recruitment
Figure 2: pilot practice recru
164x41mm (300 x 300 DPI)

1. Participants thought that the most important aspect of the study was to work out whether missed appointments were predominantly a feature of practice behaviour (so the impact of adapted appointment systems that took account of patient behaviour) or a feature of the patients who missed appointments.
2. In terms of practices, participants thought that appointment systems and especially time from booking to appointment date was important.
3. They felt that it was important if SMA was a patient feature to identify the patients whose appointment behaviour could change and those whose could not - as GPs really do this already when they use strategies for managing patient's appointment behaviour.
4. Participants were astonished by the data presented that suggested some patients had missed 25-41 appointments over 3 years and viewed this as 'extreme'. They suggested these data need careful review and postulated it may be a data cleaning issue, an 'anomaly'. They also suggested it might be related to practice factors for example a very tolerant GP; or patient factors; if the data were accurate.
5. Participants also pointed out that the data presented also included patients who serially *attend* appointments. These patients are viewed as having similar characteristics to patients who serially miss appointments and would be a useful additional focus for the study.
6. Participants were surprised that patients who serially missed appointments were more likely to live close to the practice. Participants thought this may be because patients attributed lower value to their appointments because they had to make less effort to attend or it may be that this is a signal that SMA's are predominantly an urban problem as patients in urban areas tend to live closer to their GP practice.
7. The participants cautioned that care needs to be taken that real appointments are captured as they are not always accurately recorded in practice computer systems by appointment type.
8. Participants recommended that we take into account that patients will have clusters of missed appointments when specific events are happening in patient's lives such as a recent major bereavement.
9. The participants also described large variability in practice Read coding for vulnerability.
10. Participants thought it was important that the results of the study be illustrated by case studies of patients as this will be useful for practising clinicians' learning.

Figure 3: focus group recommendations
Figure 3: focus group recommen
157x197mm (300 x 300 DPI)

1
2
3
4
5
6
7 The data we are seeking are from the patient records of patients who have scheduled a
8 GP appointment in the 3 years preceding the data extract date:

9
10 1. Dates of GP appointments

11 a. with missed or attended codes, b. session type, c. type of practitioner (e.g. GP,
12 Practice Nurse), d. number of days since appointment made, e. unique practice ID

13
14 2. Patient demographic data

15 a. Age, b. Sex, c. Ethnicity, d. data zone, e. count of patient address moves, f.
16 distance to practice (rounded to km), g. distance to hospital with an emergency
17 department (rounded to km)

18
19 3. Clinical and prescribing data, Selected Read codes:

20 a. priority 1 coding (important health conditions that GPs code for export into the
21 electronic care summary), b. long term condition diagnoses² (ref) c. patient
22 vulnerability and adversity factors (ref), d. health screening (breast, bowel, cervical,
23 BP, child health surveillance), e. exception coding (hospital referrals, DNA codes,
24 refused screening, Quality and Outcomes (QOF, payment for targeted long term
25 conditions management) exemption reporting, inappropriate use codes, self- discharge
26 codes), f. specific prescribing information from BNF Central Nervous System (CNS)
27 chapter and additional prescriptions specific to 40 long term diagnoses in b.)

28
29 4. Exit codes:

30 a. patient death, b. patient moved practice.
31
32
33
34

35 Figure 4: information request to target practices
36 Figure 4: information request
37 146x120mm (300 x 300 DPI)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

After appointment category and rules have been applied:

Apparently attended appointments:

100 time interval more than 4 mins

200 less than 4 mins

200 less than 3 mins

200 less than 2 mins excluding 0 time appointments

200 less than 2 mins including 0 time appointments

100 apparently missed appointments

Figure 5: random sample of appointments for sensitivity analysis

Figure 5: random sample of app
130x55mm (300 x 300 DPI)

For peer review only

1. Deaths
2. SMR01 - hospital inpatients & day cases
3. SMR 25 -drug misuse database
4. SMR00 -hospital outpatients
5. SMR04- mental health admissions
6. A&E attendances
7. SMR02 -maternity services including a family index
8. Immunisation records
9. NHS24 - NHS advice help line contacts
10. SAS -Scottish Ambulance Service contacts
11. GP Out of Hours contacts
12. SQA education- attainment and attendance

Figure 6: proposed linkage datasets
Figure 6: proposed linkage dat
140x79mm (300 x 300 DPI)

Additional file 2: GP focus group participant characteristics

| Participant characteristics | Practice setting | Other work roles |
|--|--|---|
| <ul style="list-style-type: none"> • 4 male and 1 female GP • All aged 40-55 years old | <ul style="list-style-type: none"> • 3 high urban deprivation • 1 urban high affluence • 1 mixed semi-rural with pocket deprivation | <ul style="list-style-type: none"> • 1 clinical director of a Health and Social Care Partnership • 1 Local Medical Committee member • 1 clinical lead for a national innovation project • 2 with strategic Royal College of General Practitioner roles • 2 members of the 'GPs at the Deep End' steering group |

1
2
3 **1 Appendix 3: GP Focus Group Interview Schedule**
4

5
6 **2 Introductions:**
7

8
9 Name, how long in clinical practice, time in your job, brief description of practice
10
11 setting.
12

13
14 **5 A priori knowledge/experience of missed appointments**
15

16 Are missed appointments important? If so why? If not why not?
17

18
19 Can a distinction be made between patients who occasionally miss and those who
20
21 serially miss GP appointments? If so what are those distinctions? Are they
22
23 important?
24

25
26 How do you make that distinction in clinical practice? (probe distinctions between
27
28 individuals and practice settings)
29

30
31 What does it mean for you, your practice and patients? Specifically patients who
32
33 serially miss?
34

35 **14 Present proof of concept provisional data (data cut offs, patient profiles)**
36

37 What does this data tell us about the issue of serial missed appointments?
38

39
40 What are the obvious things it tells us? What are the surprises? Why?
41

42
43 Do you think it misses important aspects of what you think about the issue? Why
44
45 might that be?
46

47
48 If we present these options about what a definition of a patient who serially
49
50 missed appointments compared to one who occasionally does, which one do you
51
52 think is most accurate? Why?
53

54
55 Is there more information that we should look for before deciding we have a
56
57 definition? What should that be?
58

59 **24 Conclusion**
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

25 Finally, are there aspects of missed appointments and the definition development
26 we have worked on today that we have not yet covered and you would like to tell
27 us about?

For peer review only

Additional file 1: Definition and role of TTPs


1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

With the increasing demand for statistical, research and service planning information from primary care records a solution is required to reduce the exposure of patient and clinician information to the requesting organisations to a minimum. The recommended (Information Commissioners Office) method of achieving this is by using a trusted third party (TTP) as an intermediary between organisations, which significantly reduces the number of individuals with access to identifiable information. In this case the TTP's role is to provide the technical skills to extract the required information from the Data Controllers electronic records and process this into a form that is both fit for purpose and complies with principal 3 of the data protection act. This may require the removal/replacement of identifiers (anonymisation /pseudo-anonymisation) or the use of redaction techniques when only statistical information is required prior to release of information to the beneficiary.

A TTP is required to operate to strict guidelines as it may only processes data in accordance with instructions from the data controller and to a specification previously agreed by both data requestor and data provider. The TTP acts as a Data Processor on behalf of the Data Controller and abides by the principles defined in the data protection act. It is registered as a data processor with the ICO, provides a secure storage facility which operates procedural, physical and electronic access controls to protect the data it processes and has no specific interest in, not is affiliated to any organisation that has an interest in any data provided. Albasoft maintain a secure data processing and storage facility at the Centre for Health Science adjacent to Raigmore hospital in Inverness, this facility is solely hosted on the NHS network. No information is transferred out with the

1
2
3 NHS network. Its existing middleware platform Escro is an advanced practice based
4 reporting system and is used to securely process data locally at the practice before
5
6 transferring the results to their secure repository. Albasoft has an established track
7
8 record as a TTP for the Scottish Therapeutics Utility and increasingly in supporting
9
10 NHS research. In our study, Albasoft have established data sharing agreements with
11
12
13 Scottish GP practices for computerised access to the GP practice data.
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For peer review only



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

STROBE 2007 (v4) checklist of items to be included in reports of observational studies in epidemiology*
Checklist for cohort, case-control, and cross-sectional studies (combined)

| Section/Topic | Item # | Recommendation | Reported on page # |
|---------------------------|--------|--|-----------------------------|
| Title and abstract | 1 | (a) Indicate the study's design with a commonly used term in the title or the abstract | 1 |
| | | (b) Provide in the abstract an informative and balanced summary of what was done and what was found | 2-3 |
| Introduction | | | |
| Background/rationale | 2 | Explain the scientific background and rationale for the investigation being reported | 4-6 |
| Objectives | 3 | State specific objectives, including any pre-specified hypotheses | 6 |
| Methods | | | |
| Study design | 4 | Present key elements of study design early in the paper | 7 |
| Setting | 5 | Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection | 7 |
| Participants | 6 | (a) <i>Cohort study</i> —Give the eligibility criteria, and the sources and methods of selection of participants. Describe methods of follow-up <i>Case-control study</i> —Give the eligibility criteria, and the sources and methods of case ascertainment and control selection. Give the rationale for the choice of cases and controls <i>Cross-sectional study</i> —Give the eligibility criteria, and the sources and methods of selection of participants | Pilot 8-10 Full study 14 |
| | | (b) <i>Cohort study</i> —For matched studies, give matching criteria and number of exposed and unexposed <i>Case-control study</i> —For matched studies, give matching criteria and the number of controls per case | n/a |
| Variables | 7 | Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable | 15-17 |
| Data sources/ measurement | 8* | For each variable of interest, give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group | 15-17 |
| Bias | 9 | Describe any efforts to address potential sources of bias | 18 |
| Study size | 10 | Explain how the study size was arrived at | |
| Quantitative variables | 11 | Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen and why | 14 |
| Statistical methods | 12 | (a) Describe all statistical methods, including those used to control for confounding | 19-20 |
| | | (b) Describe any methods used to examine subgroups and interactions | 18, 19-20 |
| | | (c) Explain how missing data were addressed | 16-17 |
| | | (d) <i>Cohort study</i> —If applicable, explain how loss to follow-up was addressed <i>Case-control study</i> —If applicable, explain how matching of cases and controls was addressed | n/a |

| | | | |
|--------------------------|-----|--|-------|
| | | <i>Cross-sectional study</i> —If applicable, describe analytical methods taking account of sampling strategy 19-20 | |
| | | (e) Describe any sensitivity analyses | 16 |
| Results | | | N/A |
| Participants | 13* | (a) Report numbers of individuals at each stage of study—eg numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed | - |
| | | (b) Give reasons for non-participation at each stage | - |
| | | (c) Consider use of a flow diagram | - |
| Descriptive data | 14* | (a) Give characteristics of study participants (eg demographic, clinical, social) and information on exposures and potential confounders | - |
| | | (b) Indicate number of participants with missing data for each variable of interest | - |
| | | (c) <i>Cohort study</i> —Summarise follow-up time (eg, average and total amount) | - |
| Outcome data | 15* | <i>Cohort study</i> —Report numbers of outcome events or summary measures over time | - |
| | | <i>Case-control study</i> —Report numbers in each exposure category, or summary measures of exposure | - |
| | | <i>Cross-sectional study</i> —Report numbers of outcome events or summary measures | - |
| Main results | 16 | (a) Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (eg, 95% confidence interval). Make clear which confounders were adjusted for and why they were included | - |
| | | (b) Report category boundaries when continuous variables were categorized | - |
| | | (c) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period | - |
| Other analyses | 17 | Report other analyses done—eg analyses of subgroups and interactions, and sensitivity analyses | - |
| Discussion | | | - |
| Key results | 18 | Summarise key results with reference to study objectives | - |
| Limitations | 19 | Discuss limitations of the study, taking into account sources of potential bias or imprecision. Discuss both direction and magnitude of any potential bias | - |
| Interpretation | 20 | Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence | - |
| Generalisability | 21 | Discuss the generalisability (external validity) of the study results | - |
| Other information | | | - |
| Funding | 22 | Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based | 31-32 |

*Give information separately for cases and controls in case-control studies and, if applicable, for exposed and unexposed groups in cohort and cross-sectional studies.

Note: An Explanation and Elaboration article discusses each checklist item and gives methodological background and published examples of transparent reporting. The STROBE checklist is best used in conjunction with this article (freely available on the Web sites of PLoS Medicine at <http://www.plosmedicine.org/>, Annals of Internal Medicine at <http://www.annals.org/>, and Epidemiology at <http://www.epidem.com/>). Information on the STROBE Initiative is available at www.strobe-statement.org.

1
2
3 **UNDERSTANDING REPEATED NON-ATTENDANCE IN HEALTH SERVICES**
4
5 **- PILOT ANALYSIS OF ADMINISTRATIVE DATA AND FULL STUDY**
6
7
8 **PROTOCOL FOR A NATIONAL RETROSPECTIVE COHORT**
9

10
11 Andrea E Williamson (corresponding author), General Practice and Primary Care,
12 School of Medicine, Dentistry and Nursing, MVLS, University of Glasgow.
13
14 andrea.williamson@glasgow.ac.uk Tel: 0141 330 8330
15
16

17
18
19 David A Ellis, Department of Psychology, University of Lancaster.
20
21 d.a.ellis@lancaster.ac.uk
22

23
24 Philip Wilson, Centre for Rural Health, Institute of Applied Health Sciences
25
26 University of Aberdeen. P.wilson@abdn.ac.uk
27

28
29
30 Ross McQueenie, General Practice and Primary Care, School of Medicine, Dentistry
31
32 and Nursing, MVLS, University of Glasgow. Ross.mcQueenie@glasgow.ac.uk
33

34
35 Alex McConnachie, Robertson Centre for Biostatistics, Institute of Health and
36
37 Wellbeing, MVLS University of Glasgow. Alex.McConnachie@glasgow.ac.uk
38

39
40
41
42
43 Word count 4869 excluding abstract, figures, qualitative data quotes and
44
45 references.
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

ABSTRACT

Introduction

Understanding the causes of low engagement in health care is a prerequisite for improving health services' contribution to tackling health inequalities. Low engagement includes missing health care appointments. Serially (having a pattern of) missing general practice appointments may provide a risk marker for vulnerability and poorer health outcomes.

Methods and analysis

A proof of concept pilot utilising general practice (GP) appointment data and a focus group with GPs informed the development of missed appointment categories: patients can be classified based on the number of appointments missed each year. The full study, using a retrospective cohort design, will link routine health service and education data to determine the relationship between general practice appointment attendance, health outcomes, health care utilization, preventive health activity, and social circumstances taking a life course approach and using data from the whole journey in NHS health care. 172 practices will be recruited (approximately 900,000 patients) across Scotland. The statistical analysis will focus on two key areas; factors that predict patients who serially miss appointments, and serial missed appointments as a predictor of future patient outcomes. Regression models will help understand how missed appointment patterns are associated with patient and practice characteristics.

2

We shall identify key factors associated with serial missed appointments and potential interactions that might predict them.

~~A better understanding of these may also help inform future health promoting care across the health system.~~

Ethics and dissemination

The results of the project will inform debates concerning how best to reduce non-attendance and increase patient engagement within health care systems. Significant non-academic beneficiaries include governments, policy-makers and medical practitioners. Results will be disseminated via a combination of academic outputs (papers, conferences), social media, and through collaborative public health/policy fora.

STRENGTHS AND LIMITATIONS

- ~~This study will answer~~ important question relating to the health service component of tackling health inequalities
- ~~A large dataset enables the researchers to~~ ~~Power of a large dataset~~ following patients' journey across the whole health care system
- ~~The study~~ Utilising data security and linkage capabilities in a sensitive and robust manner
- ~~The study has~~ Having a clear yet flexible data analysis plan utilising the expertise of a multi-disciplinary research team
- ~~There are~~ Limitations of using administrative data from a range of data sources of variable data quality.

KEYWORDS

Missed appointments, data linkage, administrative data, primary care, health utilisation, health promotion, health inequalities, social vulnerability

INTRODUCTION

Tackling health inequalities is a global health priority¹ and for health service provision to have an effective role, we should understand better the reasons behind, risks associated with, and needs of patients who do not engage effectively with health care provision (even if it is free at the point of access); and tailor services better to meet those needs. There remains a lack of published work concerning repeated missed appointments, but previous research typically focuses on the financial costs associated with non-attendance. One estimate has placed the cost of missed [United Kingdom \(UK\)](#) general practice (community based family medicine) appointments at £150 million per year². More recent Scottish government data suggest that each missed hospital outpatient appointment costs National Health Services (NHS) Scotland £120³. International data on costs to health care systems ~~is~~are sparse. In a complex adaptive system such as health care, the financial costs are contestable because clinicians will ‘catch up’ or get on with other care or administrative tasks. What is important are the costs of opportunities missed for improving patients’ health and the potential for substantial long-term savings to health systems⁴.

To date studies investigating missed appointments have focused on single missed appointments or single disease ~~areas outcomes~~—and have indicated they are associated with poorer health outcomes³⁻⁶. Studies of single missed appointments

4

1
2
3 have produced conflicting results when it comes to designing effective
4 interventions that can increase attendance⁷⁻¹⁰. This may be due to a reliance on
5 small samples in disparate settings¹¹⁻¹⁵ and conflation of patients who occasionally
6 miss appointments with patients who have an established pattern of missing many.
7
8 The Health and Social Care Information Centre in England has recently published
9 data about repeated missed appointments. From their analysis of recorded missed
10 outpatient hospital appointments in England one in 50 patients (65,590 of 3.5
11 million) who missed an appointment failed to attend three or more further
12 appointments within three months¹⁶.

13
14 We hypothesise that repeated missed appointments reflect a pattern of behaviour.
15 We use the term 'serially' missing appointments to reflect this pattern, which may
16 be interrupted by attended appointments. Clinicians do report that patients who
17 serially miss appointments are of particular concern because they may have very
18 poor health, may be socially disadvantaged and are high users of unscheduled care
19 compared to patients who occasionally or never miss appointments¹⁷.

20
21 There is accumulating evidence that negative experiences in early life have
22 pervasive consequences for health over the life course including 'extensive
23 evidence of a strong link between early adversity and a wide range of health-
24 threatening behaviours'¹⁸. This body of work therefore provides a conceptual
25 framework for better understanding 'chaotic lives'¹⁹ as an explanatory factor in
26 health utilization behaviours such as missed appointments, and introduces the
27 possibility that serial missed appointments contribute to the inverse care law,
28 which states that health care provision is least likely to be provided to those that
29 need it most²⁰.

1
2
3 In the UK, publicly funded general practice (GP) provides almost universal
4 coverage for the population and generates around 90% of health contacts.
5
6 Appointment making is typically under the control of each patient directly.
7

8
9
10 General practice appointments ~~are~~ therefore provide a sensible starting point for
11 this study of health and other outcomes across patients' life course. Subsequent
12 results will also, ~~and~~ have relevance for global health systems where patients have
13
14 direct access to a wider range of health care specialties.
15
16
17

18
19
20 Scotland has an established data linkage infrastructure which is under continuous
21 development. This pathfinder study will for the first time link large general
22 practice datasets (including appointment data) with data from across patients'
23 whole journey through health care.
24
25
26
27

28
29 The overarching study question is: is serially missing GP appointments a risk
30 marker for vulnerability and poorer health outcomes and thus a useful target for
31 developing interventions to improve engagement in health promoting care across
32 the health system?
33
34
35
36
37

38
39 ~~We describe the aims and research questions for this study, the pilot work that was~~
40 ~~undertaken to inform it, and the resultant research protocol for the full study~~
41 ~~based in GP practices in Scotland.~~
42
43
44
45
46
47

48 **Aim and Research questions**

49
50 The overall aim of the study is to determine the relationship between general
51 practice appointment attendance, health care utilization, preventive health
52 activity, health outcomes, and social circumstances taking a life course approach
53 and using extracted health service and other relevant administrative data.
54
55
56
57
58

1
2
3 A pilot study sought to answer the first research question described below. The
4
5 subsequent questions underpin the full research protocol which compares cohorts
6
7 of Scottish patients (from birth to older peopleage) who never, occasionally and
8
9 serially miss GP appointments.
10

11
12
13
14 Figure 1: Study research questions

15
16 An introduction to the full study protocol is described, followed by the methods
17 and results from a mixed methods pilot study that informed the protocol. A
18 description of— protocol participants, data sources, variables and statistical
19 analysis then follows.
20
21
22
23
24
25

26 27 28 METHODS AND ANALYSIS

29
30
31
32
33 The full study protocol is for a retrospective cohort study of GP practice patient
34 records linked with secondary care and education administrative records in
35 Scotland.
36
37

38
39 The study commenced in July 2015 and will finish in December 2017. A pilot study
40 was conducted between July and September 2015 which is described next. The
41 cohort of 909,073 GP patient records for the full study was available in the
42 National Safehaven from September 2016 and analysis -of these data is underway.
43
44 Permissions to access education data is secured, and the outcome of linkage
45 permissions for health data is not yet confirmed
46
47
48
49
50
51
52
53
54

55 Pilot study

56
57
58
59 7
60

1
2
3 The pilot study was separated into 2 sub-sections: ~~a 'proof of concept'~~
4 ~~quantitative data analysis and~~ a focus group to inform and refine definition
5 development (research question 1) and a 'proof of concept' quantitative data
6 analysis-
7
8
9
10

11 Methods

12 ~~Proof of concept~~

13
14 ~~Research that uses GP appointment data has not previously been conducted using~~
15 ~~the clinical recording systems in the Scottish NHS. A proof of concept pilot study~~
16 ~~was undertaken utilising the NHS Trusted Third Party (TTP) Albasoft with 67,705~~
17 ~~patient records to determine whether retrieving appointment data was feasible, to~~
18 ~~refine other data parameters, and to inform the definition development as~~
19 ~~described in research question 1. An additional confidentiality control means that~~
20 ~~the research team do not know the identity of the recruited GP practices.~~
21
22

23 ~~Additional file 1 describes the definition and role of TTPs.~~

24 ~~Albasoft purposively recruited 10 Scottish practices on our behalf with the~~
25 ~~following characteristics:~~
26
27

28 ~~Figure 2: Pilot practice recruitment~~

29 Focus group

30
31 A focus group was conducted in September 2015 with five GP participants. A focus
32 group was judged the most appropriate method to use because we aimed to
33 promote discussion of the topic such that participants would be able to compare
34 and contrast their own experiences with others from a range of practice and
35 professional experience settings²¹. Linked to this was the aim of asking participants
36
37
38
39
40
41
42
43
44

1
2
3 to make sense of, and provide feedback on the presented pilot data. The GPs
4
5 were a convenience, purposive sample based on two main principles. The first took
6
7 into account the evidence surrounding single missed appointments. This describes
8
9 missed appointments being more common in deprived, urban practices. The
10
11 sample therefore included GPs who worked in deprived and affluent urban areas
12
13 and a practice with a significant rural practice population from Scotland. Second,
14
15 the sample included the views of frontline GPs and GPs who had a range of
16
17 strategic roles in practice development and general practice management, locally
18
19 and nationally. AEW and PW utilised their professional knowledge of GP networks
20
21 and practice profiles to approach and recruit participants. Five out of twelve GPs
22
23 contacted were able to attend the focus group. Each GP contacted reported that
24
25 they felt this was an important topic worthy of attention. Barriers to attending
26
27 were location of the focus group (conducted in Glasgow) and managing time away
28
29 from other professional work. Additional file 12 describes each participant's
30
31 characteristics. Detailed information about participants' practice characteristics
32
33 was not collected. Three of the participants knew each other from their
34
35 professional roles outside of clinical practice. AEW conducted the focus group and
36
37 the analysis was conducted using Framework Analysis. Framework Analysis is a
38
39 useful thematic analysis approach especially when considering a focussed topic like
40
41 this one. Also in the context of being part of a larger mixed methods study,
42
43 epistemologically its use was a good fit²². DAE attended the focus group and
44
45 presented initial results from the ~~pilot~~ 'proof of concept' pilot (described
46
47 next) quantitative data for discussion. Additional file 23 describes the topics
48
49 covered in the focus group.

50
51
52 Proof of concept

53
54
55
56
57
58
59
60
9

1
2
3 Research that uses GP appointment data has not previously been conducted using
4 the clinical recording systems in the Scottish NHS. A proof of concept pilot study
5 was undertaken utilising the NHS Trusted Third Party (TTP) Albasoft with 67,705
6 patient records to determine whether retrieving appointment data was feasible, to
7 refine other data parameters, and to inform the definition development as
8 described in research question 1. An additional confidentiality control
9 meansured that the research team did not know the identity of the recruited
10 GP practices.

11 Additional file 34 describes the definition and role of TTPs.

12 Albasoft purposively recruited 10 Scottish practices on our behalf with the practice
13 characteristics illustrated in figure 2. following characteristics:

14 Figure 2: Pilot practice recruitment

15 Data were cleaned and appointment rules applied to categorise appointments as
16 attended or missed (DNA). Additional file 4 describes this process. This was
17 primarily based on the 'in' and 'out' time recorded for each appointment. If this
18 was recorded as '0' then the appointment was classified as Did Not Attend (DNA).
19 For each patient the total number of appointments made during the three-year
20 period was calculated as well as the number and percentage of appointments
21 missed. Appointments that were recorded incorrectly in the system were removed
22 at this stage. This included appointments where administrative records had
23 remained open for over 24 hours, making it difficult to confirm that these were
24 genuine appointments and not administrative errors. The pilot appointment rules
25 are set out in table 1 below.

| <u>Data description</u> | <u>Reason for removal</u> |
|--|---|
| <u>total appointment time < 0 min</u> | <u>Record open for more than 24 hours</u> |
| <u>total waiting time < 0 min</u> | <u>Record open for more than 24 hours</u> |
| <u>appointment <2 min</u> | <u>Not a medical appointment</u> |
| <u>administrator slot</u> | <u>Not a medical appointment</u> |

Table 1 Rules to identify genuine appointments

Results

Focus group

Focus group participants reported making clear distinctions between patients who occasionally miss appointments and those who miss many. Patients who occasionally miss appointments do so because a crisis or another understandable circumstance has arisen; patients who serially miss appointments (SMA), described as missing more than two or three appointments can be easily identified by GPs.

They were described as tending to have mental health, addiction, and/or social issues. They were described as high risk or vulnerable with concerns about their wider family. Patients who SMA were viewed as being different from the general GP population and being more likely to have ‘chaotic’ lifestyles associated with housing instability, money problems, a “panicked lifestyle”(P2). Patients who SMA were also described as being unable to manage GPs’ expectation of them and fit into GPs’ pre-determined slots. “there’s the occasional DNA which are quite normal and often those are quite significant [in total numbers for the practice] but the serial people I think that’s a reflection of the chaos in their life whether that’s you know- mental health or issues with the social functioning- and inability to manage our expectation of them- to fit into our pre-defined slots.” P5

—All participants agreed with that view. However one participant also considered that not all patients who SMA can be viewed as high risk; that instead some patients do not value free health care. It was reported—that some patients who

11

1
2
3 SMA go on to book another appointment the next day; *“I don’t think it’s the value*
4 *of the GP- I think it’s the value of that appointment- I think the fact that it’s, if*
5 *you don’t miss it, if you miss it is no big deal you just make another one”* P4.
6
7

8
9 Missed appointments were viewed as being more prevalent in practices in deprived
10 settings, but occurred in affluent areas too. In the affluent setting they were
11 important for a minority of marginalised, isolated patients with the same profile as
12 described above-who were viewed as living ‘chaotic’ lives.
13

14 Practices represented in the focus group do not have protocols for managing
15 patients who serially miss appointments (SMA) because response is dependent on
16 the patient’s context. GPs understood that SMAs usually mean patients with
17 complex needs with workload implications for the practice. Strategies described
18 were varied, including allowing patients only to book on the day; *“my impression*
19 *is that deprived practices have a much higher percentage of on the day*
20 *appointments because they skew it towards people that don’t attend”* P3, seeing
21 the patient when they walk in, or the GP booking the follow up appointment for
22 the patient- a relationship building strategy. This could still lead to patients
23 missing an appointment, even just a couple of hours after it was made. It was
24 reported that some ~~Some~~ practices do remove patients from their list for SMA and
25 this created tension with other practices. ~~Some practices have a negative view of~~
26 ~~patients who SMA.~~
27

28 The focus group were also asked to comment on the results from the proof of
29 concept initial data and they made recommendations about the full study design
30 described in Figure 3.
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48

49
50
51
52
53
54
55
56
57
58 Figure 3 Focus group recommendations for the full study design

59 12

*Results—definition of serial missed appointments**Proof of concept*

Following the pilot analysis, data were cleaned and appointment rules applied to categorise appointments as attended or missed (DNA). Appendix 4 describes this process. This was primarily based on the ‘in’ and ‘out’ time recorded for that appointment. If this was recorded as ‘0’ then the appointment was classified as Did Not Attend (DNA). For each patient the total number of appointments made during the three-year period was calculated as well as the number and percentage of appointments missed. Appointments that were recorded incorrectly in the system were removed at this stage. This included appointments where administrative records had remained open for over 24 hours, making it difficult to confirm that these were genuine appointments and not administrative errors. The pilot appointment rules are set out in table 1 below.

| Data description | Reason for removal |
|--------------------------------|------------------------------------|
| total appointment time < 0 min | Record open for more than 24 hours |
| total waiting time < 0 min | Record open for more than 24 hours |
| appointment < 2 min | Not a medical appointment |
| administrator slot | Not a medical appointment |

Table 1 Rules to identify genuine appointments

A pilot analysis of 67,705 patient records showed that while just over 60% of our sample missed no appointments, over 30% missed one or more appointment during the three-year period with nearly 10% of patients missing three or more appointments.

1
2
3 Assuming that our final sample provides a similar distribution, we will classify
4 patients based on the number of appointments missed as follows:

5
6
7
8 ~~over the last three years as follows:~~

9
10
11 ~~Never missed appointments per year, 0~~

12
13 ~~Low missed appointments per year, <1~~

14
15 ~~Medium missed appointments per year, 1-2~~

16
17 ~~High missed appointments per year, 2 or more~~

18
19
20 ~~Never missed appointments: 0 per year average over 3 year period~~

21
22 ~~Low missed appointments: <1 per year average over 3 year period~~

23
24 ~~Medium missed appointments: 1-2 per year average over 3 year period~~

25
26 ~~High missed appointments: >2 per year average over 3 year period~~

27
28
29 Our sampling both in the pilot data stage and the final full study sample was
30 conducted such that we were likely to get a representative sample of Scottish
31 patients and practices. Because our pilot sample was large, it is appropriate to
32 assume that this will scale-up accordingly for the full study. The distribution of
33 missed appointments also suggested useful categories based on integer numbers of
34 missed appointments per year. ~~Tand this which~~ will be helpful for policy and
35 clinical stakeholders.

36 37 38 39 40 41 42 43 44 45 FULL STUDY PROTOCOL

46 47 ~~Recruitment~~ Participants and study size

48
49 Our target recruitment of GP practices is 172 GP seeks to practices from across
50 Scotland ensuring that we had a spread of urban, and rural practices, affluent
51 and practices characterised by serving areas of blanket high socio-economic (Deep
52
53
54
55
56
57 End) deprivation practices. This will provide approximately 900,000 patient

58
59
60
14

1
2
3 ~~records for inclusion in the study.~~ The following is the information request made to
4
5 practices can be viewed in Figure 4.
6
7

8
9
10 Figure 4 Information request sent to target practices
11

12 Data Handling

13 Data sources and variables

14 GP data

15
16
17
18
19
20
21 The TTP ~~hasis~~ recruited the practices on the study team's behalf and will
22
23 undertake some specific data aggregation before transferring the data securely to
24
25 the National Safehaven for analysis. 'Safe Havens are specialised, secure
26
27 environments supported by trained, specialist staff where data in electronic
28
29 patient records can be processed and linked with other health data (and/or non-
30
31 health-related data) and made available for analysis to facilitate research while
32
33 protecting patient identity and privacy'²³. These are: calculating urban rural
34
35 classification, SIMD decile, categorising ethnicity into 'non BME (Black and Minority
36
37 Ethnicity)', 'visibly BME', and 'non visible BME' and rounding travel distance to
38
39 practice/emergency department ~~to the nearest kilometre~~ for each patient record
40
41 to the nearest kilometre. Once in a Safehaven, additional steps will be taken to
42
43 ensure that acceptable anonymization principles are being applied, especially in
44
45 relation to reporting of sensitive social vulnerability data and reporting of rare
46
47 conditions.
48
49

50
51
52 A new data file containing the appointment history for each patient record will be
53
54 generated, which will be merged with individual patient information (Additional
55
56 file 4 describes this process ~~sets this out~~ based on our pilot data set).
57
58

59
60
15

Appointment validation and categorisation

Each appointment will be coded by the based on session type recorded by the practice (eg book on day appointments, or immunization clinic) and then further by professional type (eg GP partner, practice nurse). These descriptions are determined by individual practices so categorisation will be conducted by the GPs in the research team. The appointment rules set out in the pilot study will be applied. A sensitivity analysis based on the time the appointment takes will then also be conducted by comparing a random sample of patient appointments as described in ccording to figure 5.

Figure 56 Random sample of GP appointments for validation and sensitivity analysis

The appointment rules will be refined based on this. The time interval cut-off for apparently attended appointments will be determined by utilising the time interval that most accurately matches to actual attended appointments. Slots designated non face to face appointments will then be removed leaving only attended and non- attended face- to- face appointments. The appointment categories described from the pilot study regarding non- attendance for all patients will then be applied to the yearly average number of missed appointments over the three year extract period to generate the four categories of patients for further analysis. Using an average over three years takes account of what is recognised in the frequent attenders (rather than non- attenders) literature- that patients' appointment behaviour may vary over time in relation to illness episodes or social crises²⁴.

Health and education data linkage

Linkage This will be conducted as access permissions and data sets become available ~~for linkage so will be incremental~~. Each administrative data source is available for different time periods (e.g. hospital inpatients since 1981 and education outcomes since 2002) and this will be made explicit when interpreting the results. The TTP will provide the Safe Haven indexing team a file containing the GP dataset Community Health Index (CHI) number and other patient identifiers. Every patient in the Scottish NHS has a CHI number, a unique identifier that is used as such across all NHS services in Scotland. This forms the cohort for the study. All data providers will ~~provide~~ supply identifiers to be probability matched to the study cohort by the Safehaven linkage team (based on CHI number and using other patient identifiers probabilistically for the small number of records where it is anticipated CHI will be missing), who will return a set of unique index numbers for those individuals successfully matched to the study cohort; each data provider will receive a different set of unique index numbers, and will use these index numbers as the basis of their data extract. Each data extract will be submitted to the Safehaven linkage team, who will replace the different index numbers with a common number across all files. This common number is the unique patient identifier that the research team will work from ~~to analyse the linked data during analysis.~~

Figure 65 Proposed data sets for linkage with GP data

~~Appointment validation and categorisation~~

~~Each appointment will be coded by the session type recorded by the practice (eg book on day appointments, or immunization clinic) and then further by professional type (eg GP partner, practice nurse). These descriptions are determined by individual practices so categorisation will be conducted by the GPs in the research team. The appointment rules set out in the pilot study will be applied. A sensitivity analysis based on the time the appointment takes will then also be conducted by comparing a random sample of patient appointments according to~~

~~Figure 6 Random sample of GP appointments for validation and sensitivity analysis~~

~~The appointment rules will be refined based on this. The time interval cut off for apparently attended appointments will be determined by utilising the time interval that most accurately matches to actual attended appointments. Slots designated non face to face appointments will then be removed leaving only attended and non attended face to face appointments. The appointment categories described from the pilot study regarding non attendance for all patients will then be applied to the yearly average number of missed appointments over the three year extract period to generate the four categories of patients for further analysis. Using an average over three years takes account of what is recognised in the frequent attenders (rather than non attenders) literature that patients' appointment behaviour may vary over time in relation to illness episodes or social crises²⁴.~~

Bias

18

Accounting for patient turnover

This study seeks to ensure the inclusion of patients who are marginalised and who are often missing from health service studies. There is evidence of overlap between patients who miss appointments and those who are removed from practice lists²⁵, a recognition of the impact that geographical boundary areas has^{ve} on patients who move around²⁶; notwithstanding the gap in the literature about registration interruptions for patients who may go to prison or patients who remain unregistered once they are removed from GP practice lists. We will therefore summarise the numbers of patients joining and/or leaving their practice during the study period; with reasons where this information is available. We will seek to provide a full analysis of the data available for these patients and compare these with the patients who are registered for the 3 year study period. Patients who are not registered with participating practices, and are being seen as 'temporary residents' by these practices, are excluded from the study. This is because these patients full clinical record is held by their registered GP so very limited information is available. Temporary residents tend to be people on holiday in the practice area but will include some people who would be considered marginalised.

Statistical ~~methods~~ analysis

Our statistical analysis is based on the study being a retrospective cohort study. We will focus on two key areas; predictors of high rates of serial missed appointments, and serial missed appointments as a predictor of future patient outcomes.

Patient characteristics and practice characteristics may be associated with high rates of serial missed appointments. Analyses will initially be descriptive,

1
2
3 summarising the rate of missed appointments in relation to the other factors
4 recorded at the point of entry to the study. —Associations with patient
5 characteristics will be assessed as a whole, and in relation to different types of
6 practices (e.g. separately in rural and urban practices). Subsequently, we will ~~use~~
7 ~~build~~ regression models (Poisson or Negative Binomial),²⁷ to help ~~us~~ understand
8 how our categories of missed appointments are associated with patient and
9 practice characteristics. ~~Specifically, Poisson or Negative Binomial regression will~~
10 ~~be used.~~

11
12
13
14
15
16
17
18
19
20
21
22 When considering other outcomes in relation to serial missed appointments, the
23 missed appointment rate category (none, <1, 1-2, or >2 per year) will be
24 ~~considered as~~ the predictor variable. Appropriate regression models, according to
25 the outcome ~~variable~~, will be used to assess whether any associations with serial
26 missed appointment rates are independent of other patient- or practice-level
27 factors. Conflicting interactions will be controlled for by using an 'offset term' in
28 our negative binomial model which accounts for number of appointments made or
29 any other relevant factors.

30
31
32
33
34
35
36
37
38
39
40
41 We also plan to measure relevant the quantitative variables (described next)
42 factors recorded *during* the study interval associated with having a lot of missed
43 appointments. We will explore whether these *differ* from the predictive factors
44 already recorded at entry to the study.

Quantitative variables

45
46
47
48
49
50
51
52
53
54 The following pPotential pPredictors of frequent non- attendance will be analysed:

55
56
57 Demographics

58
59 20
60

1
2
3 Patients' age, gender, minority ethnic group status (where available), deprivation
4
5 decile, rural/urban split, number of address moves, distance lived from GP
6
7 practice and distance from nearest A&E will be explored.
8

9 Health conditions

10 Health conditions will be reported using separate categories/variables:

11
12
13
14 1. The ~~Firstly by the~~ incidence of multi-morbidity calculated from extracted Read
15
16 codes based on previous counts in Scotland²⁸

17
18 2. , secondly D descriptions of health conditions based on the priority 1 Read codes
19
20 that GP practices in Scotland use to populate patients' key information summaries
21
22 (KIS) for GP out of hours services. This is novel work as a coding structure has not
23
24 previously been applied to these Read codes. Read codes are the clinical coding
25
26 system used in UK general practice to record, clinical and administrative activity
27
28 and diagnoses.

29
30
31
32 3. Thirdly, a count of psychotropic medicine prescriptions based on the British
33
34 National Formulary. will be generated. This is in order to describe levels of
35
36 psychological morbidity that are not captured by diagnostic criteria.

37
38
39 4. These three variables will then be compared to the ICD 10 coding data from
40
41 patients' secondary care linked data compiled from hospital admissions and
42
43 outpatient attendances. Diagnostic data from emergency department attendance
44
45 was deemed not of sufficient quality to utilise.
46

47 Social Vulnerability

48
49
50 One aspect of this study which is particularly ground-breaking is our investigation
51
52 of retrievable information about patients' social vulnerability. The Adverse
53
54 Childhood Experiences (ACE) questionnaire²⁹ will be utilised as a template to
55
56 match its nine descriptors of adversity to relevant Read codes in the patient's GP
57
58

59 21
60

1
2
3 record. In addition, coding that maps the consequences of ACE will be analysed. A
4
5 recent quantitative evaluation of Severe and Multiple Disadvantage will also be
6
7 matched to GP Read codes. This examines the overlap of patients being homeless,
8
9 in substance misuse services, or in prison over the preceding year³⁰. Further, an
10
11 exploration of additional Read codes that describe social vulnerability will be
12
13 mapped. An anonymised text search linked to Read codes from the dataset will
14
15 provide additional information about social vulnerability as it is recorded in the
16
17 free text portion of GP records. ~~Both of these taken~~ Taken together, these will
18
19 provide the first research evidence about the breadth and depth of social
20
21 vulnerability recording by GPs.
22
23

24
25 Health ~~carescreening and~~ utilisation

26
27 Read coding in relation to cervical, breast and bowel screening attendance will be
28
29 retrieved in addition to the proportion of patients who have had their blood
30
31 pressure checked and, have participated in child health surveillance and
32
33 vaccination programmes across the life course. A sub-analysis of utilisation of
34
35 practice nurse and other health care professional's' appointments in the practice
36
37 will also be conducted and include an exploration of the relationship between
38
39 attending all primary care appointments and categories of non- attendance. This is
40
41 because data from the GP focus group suggested there is overlap between patients
42
43 who are serial non-attenders with patients who are very frequent attenders. We
44
45 will therefore consider the rate of attending appointments as a potential predictor
46
47 of the rate of non-attendance. Referrals that GPs make into other primary and
48
49 secondary care services will also be analysed. Outpatient attendances, hospital
50
51 admissions and utilisation of emergency departments, NHS 24 triage, GP out of
52
53 hours, and ambulance services will also be ~~conducted analysed~~ when linked data
54
55
56
57
58
59
60

22

1
2
3 become available with a specific focus on how this relates to unmet need, for
4
5 example how might GP appointment category relates to patterns of other health
6
7 care utilisation between scheduled and unscheduled secondary care use
8

9
10
11
12 ~~Engagement with h~~Health_care engagement

13
14 An analysis of GP Read codes and linked secondary care data will be carried out in
15
16 the following categories: that relate to patients

17
18 1. patients not attending primary and secondary care appointments

19
20 2. patients refusing screening

21
22 3. patients—being exception-reported (ie excluded from the denominator
23
24 population) from the Quality and Outcomes Framework (QOF) system for
25
26 performance measurement in general practice

27
28 4. practices' measures of non-engagement with care for long term conditions

29
30 5. patients taking 'irregular discharge' from hospital (when patients leave against
31
32 medical advice)

33
34 6. patients and not waiting to be seen in emergency departments ~~will be conducted.~~

35
36
37
38 Family linkage

39
40 Diagnoses of children who are able to be linked through family linkage will be
41
42 analysed by based on their mother's appointment category. This is contingent on
43
44 the child being included in the practice study population.

45
46
47
48 Education data

49
50 Attendance at school, exclusion from school, and educational attainment when
51
52 leaving school will be made with approximately a sixth of our patient cohort for
53
54 whom linked education data is available. This has the potential to inform future
55
56 planning about around earlier interventions to reduce serial missed appointments.

Practice level data

Each patient record will be allocated a unique practice ID enabling the research team to analyse each patient record output clustered by practice. This will be proportion of patients aged over 75, by ethnicity_(proportion BME), patient rurality, patient level of deprivation decile, patient distance to practice, distance to A&E appointments offered/engaged, days from when appointment is made, multi-morbidity count, ACE score more than 4, Severe and Multiple Disadvantage score, hospital referrals, and proportion of each appointment category by practice.

These analyses and output will be refined as the study proceeds taking patient level findings and ~~multilevel modelling that takes the interaction between the patient and the practice into account.~~ multilevel modelling that characterises the respective contributions of practice- and individual-level factors to missed appointment patterns.

Health outcomes

Mortality data regarding date and cause of death will be utilised ~~both~~ from ~~the~~ GP and linked data. This will sit alongside ~~additional~~ ~~and~~ ~~from~~ linked obstetric outcomes (from the Scottish Birth Record) for relevant women.

Table 2 summarise the quantitative variables for analysis

| <u>Data categories</u> | <u>variables</u> |
|-----------------------------|---|
| <u>Patient demographics</u> | <u>Age</u> <u>Sex</u> <u>ethnicity</u> <u>Count of address moves</u> <u>Distance to practice</u> <u>Distance to A&E</u> <u>SIMD decile</u> <u>Rural8 score</u> |
| <u>Health conditions</u> | <u>Multi-morbidity count</u> <u>Priority 1 read codes</u> <u>Psychotropic medication prescribing (BNF chapter)</u> <u>Secondary Health care diagnoses (inpatient and</u> |

| | |
|---|---|
| | outpatient) |
| Social vulnerability | Adverse Childhood Experiences Severe and Multiple Disadvantage General social vulnerability coding frame |
| Health care utilisation | Breast screening Bowel screening Cervical screening BP checked Child health surveillance Vaccinations Practice nurse appointments Other health care professional appointments Primary care attendance distribution Hospital referrals Outpatient attendances Hospital admissions emergency departments attendance NHS 24 triage GP out of hours ambulance services callouts |
| Health care engagement | DNA codes Refused screening QOF exempt Inappropriate use codes Self-discharge codes |
| Study exit | Patient death Patient moved practice |
| Family linkage | Secondary health care linkage with mother and child |
| Education data | School attendance School exclusion School attainment |
| Health outcomes | Cause of death |
| GP Practice characteristics | Practice list size Patient age distribution Ethnicity category distribution Patient rur8 score distribution Patient SIMD score distribution Patient distance to practice distribution Patient distance to A&E distribution Number of appointments offered/patients engaged past 3 years distribution Number of days since appointments made distribution Patient multi-morbidity score distribution Patient ACE score distribution Patient SMD score distribution Patient hospital referrals distribution Primary care attendance pattern distribution |

[Table 2 Summary of quantitative categories and variables](#)

25

ETHICS AND DISSEMINATION

This pathfinder linkage retrospective cohort study is necessarily complex in design and implementation because although cross-sectional it seeks to take a life course approach and follow the patients' journey through [the](#) health care [system](#). Careful attention and significant resource has been devoted to the consideration of patient privacy and confidentiality. This has been integrated throughout the design of the study alongside the necessary data access and handling permissions. Additionally a study of this nature, which involves stakeholders across the NHS and other public services, requires a flexible time frame to allow access to raw data and to share findings between members of the research team based in several institutions.

The proof of concept pilot did not require ethical [permission approval](#) because it was considered service evaluation with the agreement we would ~~not~~ publish [any](#) results about the practices ~~which~~ [who](#) took part. ~~E~~thical permission to conduct the GP focus group and publish the results was obtained by the MVLS ethics committee, University of Glasgow (ref 200140181). A letter of comfort was obtained from the West of Scotland NHS ethics committee and the MVLS ethics committee ~~that~~ [confirming that](#) the full study did not need health service ethics permissions. Multi- site NHS R&D approval for the full study was obtained for all Scottish Health Boards (NRS16/186358).

Due to the sensitive nature of administrative data from the NHS and public education system in Scotland, the datasets generated and/or analysed during the current study will not be publicly available. They have been made available to the research team under controlled access ~~conditions~~ and strictly for the purposes of this research study only. Summary data, ~~at~~ the level of disclosure checked output

1
2
3 from the National Safehaven and statistical code, can be requested from the
4
5 corresponding author on reasonable request.
6

7 **Planned outputs**

8
9 Alongside peer reviewed academic papers reporting the findings described above,
10
11 the following ~~are~~ [additional outputs](#) ~~are~~ planned.
12
13

14 Data Visualisation

15
16 Several web pages will be built to sit alongside key results. This will allow for the
17
18 rapid construction of interactive data visualisations which will be created using
19
20 “Shiny”³¹, a web application framework for R [which is the statistical software used](#)
21
22 [for the study analysis](#). A simple platform will allow researchers and collaborators
23
24 to interact with the analyses in real-time and generate custom tables and graphs
25
26 as required. It can also provide non-experts with access to simple and complex
27
28 statistical analysis using a point-and-click interface. This will not rely on raw data
29
30 and will simply pull information from the summary descriptive analyses.
31
32
33

34 Case Studies

35
36 We also intend to use case studies to develop and illustrate our findings throughout
37
38 the course of all our analyses. For example, we will be able to identify typical
39
40 patient profiles of those who appear to miss many appointments in a very short
41
42 period of time and compare these events with short and long-term health
43
44 outcomes.
45
46
47

48 Conclusion

49
50 We shall identify key factors associated with serial missed appointments ranked in
51
52 order of importance as described [above](#), but given the large sample size we shall
53
54
55
56
57
58
59
60

1
2
3 also be able to consider potential interactions that might predict serially missed
4
5 appointments.
6

7
8 Finally, this approach also explores the theory that low engagement with health
9
10 care should be viewed as a health harming behaviour, and will inform the debate
11
12 about tackling health inequalities at the health service delivery level. [Moving from
13
14 theory into application, the results](#)This will allow us to better understand and
15
16 develop future interventions to reduce serial missed appointments.
17

18 REFERENCES

- 20
21 1. Commission on Social Determinants of Health. Closing the gap in a generation:
22
23 health equity through action on the social determinants of health. Final Report of
24
25 the Commission on Social Determinants of Health. Geneva: World Health
26
27 Organisation, 2008.
- 28
29 2. George A, Rubin G. Non-attendance in general practice: a systematic review and
30
31 its implications for access to primary health care. *Family Practice* 2003;20:178-84.
- 32
33 3. Campbell K, Millard A, McCartney G, et al. Who is least likely to attend? An
34
35 analysis of outpatient appointment 'Did Not Attend' (DNA) data in Scotland.
36
37 Edinburgh: NHS Health Scotland, 2015.
- 38
39 4. Neal RD, Hussain-Gambles M, Allgar VL, et al. Reasons for and consequences of
40
41 missed appointments in general practice in the UK: questionnaire survey and
42
43 prospective review of medical records. *BMC Family Practice* 2005;6(47)
- 44
45 5. Mitchell AJ, Selmes T. Why don't patients attend their appointments?
46
47 Maintaining engagement with psychiatric services. *Advances in Psychiatric
48
49 Treatment* 2007;13:423-34.
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 6. Ciechanowski PS, Russo J, Katon W, et al. Where is the patient? The association
4 of psychosocial factors and missed primary care appointments in patients with
5 diabetes. *General Hospital Psychiatry* 2006;28:9-16.
6
7
- 8
9 7. Lehmann T, Lehmann A, Olivet M, et al. Missed appointments at a Swiss
10 university outpatient clinic. *Public Health* 2007;121:790-99.
11
12
- 13 8. Neilson KM, Faergeman O, Foldspang A, et al. Cardiac rehabilitation: health
14 characteristics and socio-economic status among those who do not attend.
15
16 *European Journal of Public Health* 2008;18:479-83.
17
18
- 19 9. Cashman S, Savageau J, Lemay C, et al. Patient health status and appointment
20 keeping in an urban community health center. *Journal of health care for the Poor*
21 *and Underserved* 2004;15:474-88.
22
23
- 24 10. Masuda Y, Kubo A, Kokase A, et al. Personal features and dropout from
25 diabetic care. *Environment, Health Prevention Medicine* 2006;11:115-9.
26
27
- 28 11. Gucciardia E. A systemic review of attrition from diabetes education services:
29 strategies to improve attrition and retention research. *Canadian Journal of*
30 *Diabetes Care* 2008;32(5365)
31
32
- 33 12. Murdock A, Rogers C, Lindsay H, et al. Why do patients not keep their
34 appointments? Prospective study in gastroenterology outpatient clinic. *Journal of*
35 *Royal Society of Medicine* 2002;95:284-6.
36
37
- 38 13. Farley R, Wade T, Birchmore L. Factors influencing attendance at cardiac
39 rehabilitation among coronary heart disease patients. *European Journal of*
40 *Cardiovascular Nursing* 2003;2:205-12.
41
42
- 43 14. Woodward V. Non attendance at a nurse-run asthma clinic. *Journal Clinical*
44 *Nurse* 1998;7:380-1.
45
46
47
48
49
50
51
52
53
54
55

- 1
2
3 15. Killaspy H, Banerjee S, King M, et al. Prospective controlled study of
4 psychiatric out-patient non- attendance, Characteristics and outcome. *British*
5 *Journal of Psychiatry* 2000;176:160-5.
6
7
8
9
10 16. Health and Social Care Information Centre England. Provisional monthly topic
11 of interest: outpatient did not attend (DNA) appointment. London, 2014.
12
13
14 17. Watt G. Deep End Report 19. Access to Specialists. The 19th meeting of
15 'General Practitioners at the Deep End'. University of Glasgow, 2013.
16
17
18 18. Shonkoff JP, Garner AS, The Committee On Psychosocial Aspects Of Child And
19 Family Health COEC, Adoption, And Dependent Care, And Section On
20 Developmental And Behavioral Pediatrics,. The Lifelong effects of Early Childhood
21 Adversity and Toxic Stress. *Pediatrics* 2012; 129.
22
23
24
25
26
27 19. Hussain-Gambles M, Neal RD, Dempsey O, et al. Missed appointments in
28 primary care: questionnaire and focus group study of health professionals. *British*
29 *Journal of General Practice* 2004;54:108-13.
30
31
32
33
34 20. Tudor Hart J. The Inverse Care Law. *The Lancet* 1971;297:405-12.
35
36
37 21. Kitzinger J. Qualitative Research: Introducing focus groups. *BMJ*
38 1995;311(7000):299-302.
39
40
41 22. Ritchie J, Lewis J. Qualitative Research Practice. A Guide for Social Science
42 Students and Researchers. first ed. London: Sage Publications 2003.
43
44
45 23. The Scottish Government. Charter for Safe Havens in Scotland: Handling
46 Unconsented Data from National Health Service Patient Records to Support
47 Research and Statistics. Edinburgh2015 [Available from:
48 <http://www.gov.scot/Publications/2015/11/4783/2>].
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 24. Smits FTM, Brouwer HJ, van Weert HCP, et al. Predictability of persistent
4 frequent attendance: a historic 3-year cohort study. *British Journal of General*
5 *Practice* 2009;59:114-19.
6
7
8
9
10 25. Williamson AE, Mullen K, Wilson P. Understanding “revolving door” patients in
11 general practice: a qualitative study. *BMC Family Practice* 2014;15(1):1-11.
12
13
14 26. O'Reilly D, Steele K. The removal of patients who live outside the practice
15 boundary. *British Journal of General Practice* 2005;55:384-86.
16
17
18
19 27. Zuur AF, Ieno EN. A protocol for conducting and presenting results of
20 regression-type analyses. *Methods in Ecology and Evolution* 2016;7(6):636-45.
21
22
23 28. Barnett K, Mercer SMN, Watt G, et al. Epidemiology of Multi-morbidity and
24 implications for health care, research and medical education: a cross sectional
25 study. *The Lancet* 2012;380((9836)):37-43.
26
27
28
29 29. Bellis MA, Ashton K, Hughes K, et al. Welsh Adverse Childhood Experiences
30 (ACE) Study. Adverse Childhood Experiences and their impact on health-harming
31 behaviours in the Welsh adult population. NHS Wales, 2015.
32
33
34
35 30. Bramley G, Fitzpatrick S, Edwards J, et al. Hard Edges Mapping Severe and
36 Multiple Disadvantage England. England: Lankelly Chase Foundation, 2015.
37
38
39
40 31. Ellis DA, Meridian HL. Thinking outside the box: developing dynamic data
41 visualizations for psychology with Shiny. *Frontiers in Psychology* 2015;6
42
43
44
45
46
47

AUTHORS' CONTRIBUTIONS

48
49
50 AEW is principal investigator for the study. DAE, PW and AMcC are co-investigators
51 on the study and RMcQ is the research assistant. AEW conceived and developed the
52 initial research proposal, reviewed the literature, conducted and analysed the
53 pilot focus group, contributed to analysis and interpretation of the quantitative
54
55
56
57
58
59

60
31

1
2
3 pilot data, developed the predictors, outcomes and associations of interest and led
4 on writing the paper. DAE supported the development of the initial research
5 proposal, reviewed the literature, conducted and analysed the quantitative pilot
6 data, developed the statistical and output plan, and contributed to writing the
7 paper. PW supported the development of the initial research proposal, reviewed
8 the qualitative and quantitative pilot results, reviewed the statistical and output
9 plan and contributed to writing the paper. RMcQ reviewed the statistical and
10 output plan and contributed to writing the paper. AMcC provided expert statistical
11 input to the study as it was developed, reviewed the statistical and output plan
12 and contributed to writing the paper.

13
14
15
16
17
18
19
20
21
22
23
24
25 All authors read and approved the final manuscript.

26 27 **FUNDING STATEMENT**

28
29 This study is supported by a Scottish Government Chief Scientist Office research
30 grant (CZH/4/41118) with Safehaven and data linkage costs supported in lieu by
31 the DSLS at Scottish Government. These funding bodies had no role in the design of
32 the study, or collection, analysis, and interpretation of data or in writing the
33 manuscript.

34 35 36 37 38 39 40 **COMPETING INTERESTS**

41
42 The authors declare they have no competing interests.

43 44 45 46 **ACKNOWLEDGEMENTS**

47
48 We would like to acknowledge all GP practices and GPs who took part in the pilot
49 study. Also colleagues at Scottish Government who are supportive of the study in a
50 variety of ways especially Ellen Lynch in the Health Analytics Division. Dave Kelly's
51 technical and procedural expertise, wisdom and patience as director of our TTP

52
53
54
55
56
57
58 | Albasoft Ltd, underpins all of what has been achieved to date.

59
60
32

1
2
3
4
5 **FIGURE LEGENDS**
6

7
8 Figure 1: Study research questions
9

10 Figure 2: Pilot practice recruitment
11

12
13 Figure 3 Focus group recommendations for the full study design
14

15
16 Figure 4 Information request sent to target practices
17

18
19 Figure 5 Random sample of GP appointments for validation and sensitivity analysis
20

21 Figure 6 Proposed data sets for linkage with GP data
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60