

Weak c-ideals of a Lie algebra

Zekiye CILOGLU SAHIN^{1*}; David Anthony TOWERS²

¹Department of Mathematics, Faculty of Arts and Sciences, Suleyman Demirel University, Isparta, Turkey.

²Department of Mathematics and Statistics, Fylde College, Lancaster University, Lancaster, United Kingdom.

Received: .201

• Accepted/Published Online: .201

• Final Version: .201

Abstract: A subalgebra B of a Lie algebra L is called a weak c-ideal of L if there is a subideal C of L such that $L = B + C$ and $B \cap C \leq B_L$ where B_L is the largest ideal of L contained in B . This is analogous to the concept of weakly c-normal subgroups, which has been studied by a number of authors. We obtain some properties of weak c-ideals and use them to give some characterisations of solvable and supersolvable Lie algebras. We also note that one-dimensional weak c-ideals are c-ideals.

Key words: Weak c-ideal, Frattini ideal, Lie algebras, Nilpotent, Solvable, Supersolvable.

1. Introduction

Throughout L will denote a finite-dimensional Lie algebra over a field F . If B is a subalgebra of L we define B_L , the *core* (with respect to L) of B to be the largest ideal of L contained in B . We say that a subalgebra B of L is a *weak c-ideal* of L if there is a subideal C of L such that $L = B + C$ and $B \cap C \leq B_L$. This is a generalisation of the concept of a c-ideal which was studied in [9]. It is analogous to the concept of weakly c-normal subgroup as introduced by Zhu, Guo and Shum in [15]; this concept has since been further studied by a number of authors, including Zhong and Yang ([14]), Zhong, Yang, Ma and Lin ([13]), Tashtoush ([7]) and Jehad ([4]) who called them c-subnormal subgroups.

The maximal subalgebras of a Lie algebra L and their relationship to the structure of L have been studied extensively. It is well known that L is nilpotent if and only if every maximal subalgebra of L is an ideal of L (see [1]). A further result is that if L is solvable then every maximal subalgebra of L has codimension one in L if and only if L is supersolvable (see [2]). In [9] similar characterisations of solvable and supersolvable Lie algebras were obtained in terms of c-ideals. The purpose here is to generalise these results to ones relating to weak c-ideals.

In section two we give some basic properties of weak c-ideals; in particular, it is shown that weak c-ideals inside the Frattini subalgebra of a Lie algebra L are necessarily ideals of L . In section three we first show that all maximal subalgebras of L are weak c-ideals of L if and only if L is solvable and that L has a solvable maximal subalgebra that is a weak c-ideal if and only if L is solvable. Unlike the corresponding results for c-ideals, it is necessary to restrict the underlying field to characteristic zero, as is shown by an example. Finally we have that if all maximal nilpotent subalgebras of L are weak c-ideals, or if all Cartan subalgebras of L are

*Correspondence: zekiyeciloglu@sdu.edu.tr

2010 *AMS Mathematics Subject Classification*: 17B05, 17B20, 17B30, 17B50

1 weak c -ideals and F has characteristic zero, then L is solvable.

2 In section four we show that if L is a solvable Lie algebra over a general field and every maximal
 3 subalgebra of each maximal nilpotent subalgebra of L is a weak c -ideal of L then L is supersolvable. If each
 4 of the maximal nilpotent subalgebras of L has dimension at least two then the assumption of solvability can be
 5 removed. Similarly if the field has characteristic zero and L is not three-dimensional simple then this restriction
 6 can be removed. In the final section we see that every one-dimensional subalgebra is a weak c -ideal if and only
 7 if it is a c -ideal.

8 If A and B are subalgebras of L for which $L = A + B$ and $A \cap B = 0$ we will write $L = A \oplus B$. The
 9 ideals $L^{(k)}$ and L^k are defined inductively by $L^{(1)} = L^1 = L$, $L^{(k+1)} = [L^{(k)}, L^{(k)}]$, $L^{k+1} = [L, L^k]$ for $k \geq 1$.
 10 If A is a subalgebra of L , the *centralizer* of A in L is $C_L(A) = \{x \in L : [x, A] = 0\}$.

11 **2. Preliminary Results**

Definition 2.1 Let I be a subalgebra of L . We call I a subideal of L if there is a chain of subalgebras

$$I = I_0 < I_1 < \dots < I_n = L,$$

12 where I_j is an ideal of I_{j+1} for each $0 \leq j \leq n - 1$.

Definition 2.2 A subalgebra B of a Lie algebra L is a weak c -ideal of L if there exists a subideal C of L
 such that

$$L = B + C \text{ and } B \cap C \leq B_L,$$

13 where B_L , the core of B , is the largest ideal of L contained in B .

14 **Definition 2.3** A Lie algebra L is called weak c -simple if L does not contain any weak c -ideals except the
 15 trivial subalgebra and L itself.

16 **Lemma 2.4** Let L be a Lie algebra. Then the following statements hold:

- 17 (1) Let B be a subalgebra of L . If B is a c -ideal of L then B is a weak c -ideal of L .
- 18 (2) L is weak c -simple if and only if L is simple.
- 19 (3) If B is a weak c -ideal of L and K is a subalgebra with $B \leq K \leq L$, then B is a weak c -ideal of
 20 K .
- 21 (4) If I is an ideal of L and $I \leq B$, then B is a weak c -ideal of L if and only if B/I is a weak c -ideal
 22 of L/I .

23 **Proof** (1) By the definition every ideal is a c -ideal and every c -ideal is a weak c -ideal so the proof is obvious.

(2) Suppose first that L is simple and let B be a weak c -ideal with $B \neq L$. Then

$$L = B + C \text{ and } B \cap C \leq B_L$$

24 where C is a subideal of L . But, since L is simple, B_L must be 0. Moreover, $C \neq 0$ so $C = L$. Hence $B = 0$
 25 and L is weak c -simple.

26 Conversely, suppose L is weak c -simple. Then, since every ideal of L is a weak c -ideal, L must be simple.

(3) If B is a weak c -ideal of L then there exists a subideal C of L such that

$$L = B + C \text{ and } B \cap C \leq B_L$$

Then $K = K \cap L = K \cap (B + C) = B + (K \cap C)$. Since C is a subideal of L there exists a chain of subalgebras

$$C = C_0 < C_1 < \dots < C_n = L$$

where C_j is an ideal of C_{j+1} for each $0 \leq j \leq n-1$. If we intersect this chain with K we get

$$C \cap K = C_0 \cap K < C_1 \cap K < \dots < C_n \cap K = L \cap K = K$$

and obviously $C_j \cap K$ is an ideal of $C_{j+1} \cap K$ for each $0 \leq j \leq n-1$. Hence $C \cap K$ is a subideal of K . Also,

$$B \cap (C \cap K) \leq B_K$$

1 so that B is a weak c -ideal of L .

(4) Suppose first that B/I is a weak c -ideal of L/I . Then there exists a subideal C/I of L/I such that

$$L/I = B/I + C/I \text{ and } B/I \cap C/I \leq (B/I)_{L/I} = B_L/I$$

2 It follows that $L = B + C$ and $B \cap C \leq B_L$ where C is a subideal of L .

Suppose conversely that I is an ideal of L with $I \leq B$ and B is a weak c -ideal of L . Then there exists a C subideal of L such that

$$L = B + C \text{ and } B \cap C \leq B_L.$$

Since I is an ideal and $I \leq B$ the factor algebra

$$L/I = (B + C)/I = B/I + (C + I)/I$$

where $(C + I)/I$ is a subideal of L/I and

$$(B/I) \cap (C + I)/I = (B \cap (C + I))/I = (I + B \cap C)/I \leq B_L/I = (B/I)_{L/I}$$

3 so B/I is a weak c -ideal of L/I . □

4 The *Frattini subalgebra* of L , $F(L)$, is the intersection of all of the maximal subalgebras of L . The
5 *Frattini ideal*, $\varphi(L)$, of L is $F(L)_L$. The next result is a generalisation of [9, Proposition 2.2]. The same
6 proof works but we will include it for completeness.

7 **Proposition 2.5** Let B, C be subalgebras of L with $B \leq F(C)$. If B is a weak c -ideal of L then B is an
8 ideal of L and $B \leq \varphi(L)$.

9 **Proof** Suppose that $L = B + K$ where K is a subideal of L and $B \cap K \leq B_L$. Then $C = C \cap L =$
10 $C \cap (B + K) = B + C \cap K = C \cap K$ since $B \leq F(C)$. Hence $B \leq C \leq K$, giving $B = B \cap K \leq B_L$ and B is
11 an ideal of L . It then follows from [8, Lemma 4.1] that $B \leq \varphi(L)$. □

12 An ideal A is *complemented* in L if there is a subalgebra U of L such that $L = A + U$ and $A \cap U = 0$.
13 We adapt this to define a complemented weak c -ideal as follows.

14 **Definition 2.6** Let L be a Lie algebra and B is a weak c -ideal of L . A weak c -ideal B is complemented
15 in L if there is a subideal C of L such that $L = B + C$ and $B \cap C = 0$.

16 Then we can give the following lemma:

1 **Lemma 2.7** If B is a *weak c-ideal* of a Lie algebra L , then B/B_L has a subideal complement in L/B_L , *i.e.*, there
 2 exists a subideal subalgebra C/B_L of L/B_L such that L/B_L is semidirect sum of C/B_L and B/B_L . Con-
 3 versely, if B is a subalgebra of L such that B/B_L has a subideal complement in L/B_L then B is a *weak c-ideal*
 4 of L .

5 **Proof** Let B be a weak c-ideal of L . Then there exists a subideal C of L such that $B + C = L$ and
 6 $B \cap C \leq B_L$. If $B_L = 0$ then $B \cap C = 0$ and so that C is a subideal complement of B in L . Assume that
 7 $B_L \neq 0$, then we can construct the factor algebras B/B_L and $(C + B_L)/B_L$. If we intersect these two factor
 8 algebras we have

$$\begin{aligned} \frac{B}{B_L} \cap \frac{C + B_L}{B_L} &= \frac{B \cap (C + B_L)}{B_L} \\ &= \frac{B_L + (B \cap C)}{B_L} \\ &= \frac{B_L}{B_L} = 0 \end{aligned}$$

Hence, $(C + B_L)/B_L$ is a subideal complement of B/B_L in L/B_L . Conversely, if K is a subideal of L such
 that K/B_L is a subideal complement of B/B_L in L/B_L then we have that

$$L/B_L = (B/B_L) + (K/B_L) \text{ and } (B/B_L) \cap (K/B_L) = 0$$

9 Then $L = B + K$ and $B \cap K \leq B_L$. Therefore B is a weak c-ideal of L . □

10 3. Some characterisations of solvable algebras

11 We will use the following Lemma which is due to Stewart [6, Lemma 4.2.5]

12 **Lemma 3.1** Let L be a Lie algebra over any field having two subideals H and K such that K is simple and
 13 not abelian. Suppose that $H \cap K = 0$. Then $[H, K] = 0$

14 **Theorem 3.2** Let L be a Lie-algebra over a field F of characteristic zero and let B be an ideal of L . Then
 15 B is solvable if and only if every maximal subalgebra of L not containing B is a weak c-ideal of L .

16 **Proof** Suppose every maximal subalgebra of L not containing B is a weak c-ideal of L . Then we need to
 17 show B is solvable. Assume that this is false and let L be a minimal counter-example. Let A be a minimal
 18 ideal of L and assume that M/A is a maximal subalgebra of L/A such that $(B + A)/A \not\subseteq M/A$. Then M is a
 19 maximal subalgebra of L with $B \not\subseteq M$, so M is a weak c-ideal of L . It follows that M/A is a weak c-ideal of
 20 L/A , and hence that $(B + A)/A$ is solvable. If $B \cap A = 0$, then $B \cong B/B \cap A \cong (B + A)/A$ is solvable. So we
 21 can assume that every minimal ideal of L is contained in B . Moreover, B/A is solvable for each such minimal
 22 ideal. If L has two distinct minimal ideals A_1 and A_2 then $B \cong B/A_1 \cap A_2$ is solvable, so L is monolithic
 23 with monolith A , say.

24 If A is abelian then B is solvable, so we must have that A is simple. Clearly, $B \not\subseteq \varphi(L)$, since $\varphi(L)$ is
 25 nilpotent, so there is a maximal subalgebra M of L such that $B \not\subseteq M$. Then M must be a weak c-ideal of L ,
 26 so there is a subideal C of L such that $L = M + C$ and $M \cap C \subseteq M_L$. Since $B \not\subseteq M_L$ we have that $M_L = 0$.

1 It follows that L is primitive of type 2 and hence that $C_L(A) = 0$, by [10, Theorem 1.1]. But $[C, A] = 0$ by
 2 Lemma 3.1, so $C = 0$, a contradiction. Hence B is solvable. So suppose now that B is solvable and let M
 3 be a maximal ideal of L not containing B . Then there exists $k \in \mathbb{N}$ such that $B^{(k+1)} \subseteq M$, but $B^{(k)} \not\subseteq M$.
 4 Clearly $L = M + B^{(k)}$ and $B^{(k)} \cap M$ is an ideal of L , so $B^{(k)} \cap M \subseteq M_L$. It follows that M is a c -ideal and
 5 hence a weak c -ideal of L . \square

6 **Corollary 3.3** Let L be a Lie algebra over a field F of characteristic zero. Then L is solvable if and only if
 7 every maximal subalgebra of L is a weak c -ideal of L .

8 Unlike the corresponding results for c -ideals, the above two results do not hold in characteristic $p > 0$,
 9 as the following example shows.

10 **Example 3.4** Let $L = sl(2) \otimes \mathcal{O}_1 + 1 \otimes F(\frac{\partial}{\partial x} + x\frac{\partial}{\partial x})$, where $\mathcal{O}_1 = F[x]$ with $x^p = 0$ is the truncated
 11 polynomial algebra in 1 indeterminate and the ground field, F , is algebraically closed of characteristic $p > 2$.
 12 Then $A = sl(2) \otimes \mathcal{O}_1$ is the unique minimal ideal of L . Put $S = sl(2) = Fu_{-1} + Fu_0 + Fu_1$ with $[u_{-1}, u_0] = u_{-1}$,
 13 $[u_{-1}, u_1] = u_0$, $[u_0, u_1] = u_1$ and let $M = (Fu_0 + Fu_1) \otimes \mathcal{O}_1 + 1 \otimes F(\frac{\partial}{\partial x} + x\frac{\partial}{\partial x})$. This is a maximal subalgebra
 14 of L which doesn't contain A . Suppose that it is a weak c -ideal of L . Then there is a subideal C of L such
 15 that $L = C + M$ and $C \cap M \subseteq M_L = 0$.

Let

$$C = C_0 < C_1 < \dots < C_n = L$$

16 where C_j is an ideal of C_{j+1} for each $0 \leq j \leq n-1$. Then $A \subseteq C_{n-1}$, so $A = C_{n-1}$ or $C_{n-1} = A + 1 \otimes F\frac{\partial}{\partial x}$.
 17 In the latter case it is straightforward to check that $C_{n-2} \subseteq A$. In either case, C must be inside a proper ideal
 18 of A , and hence inside $S \oplus O_1^+$, where O_1^+ is spanned by x, x^2, \dots, x^{p-1} . But now $u_{-1} \otimes 1 \notin C + M$. Hence
 19 M is not a weak c -ideal of L .

20 **Lemma 3.5** Let $L = U + C$ be a Lie algebra, where U is a solvable subalgebra of L and C is a subideal of
 21 L . Then there exists $n_0 \in \mathbb{N}$ such that $L^{(n_0)} \subseteq C$.

22 **Proof** Let $C = C_0 < C_1 < \dots < C_k = L$ where C_i is an ideal of C_{i+1} for $0 \leq i \leq k-1$. Then L/C_{k-1} is
 23 solvable and so there exists n_{k-1} such that $L^{(n_{k-1})} \subseteq C_{k-1}$. Suppose that $L^{(n_i)} \subseteq C_i$ for some $0 \leq i \leq k-1$.
 24 Now C_i/C_{i-1} is solvable, and so there is r_i such that $C_i^{(r_i)} \subseteq C_{i-1}$. Hence $L^{(n_i+r_i)} = (L^{(n_i)})^{(r_i)} \subseteq C_{i-1}$. Put
 25 $n_{i-1} = n_i + r_i$. The result now follows by induction. \square

26 **Theorem 3.6** Let L be a Lie algebra over a field F of characteristic zero. Then L has a solvable maximal
 27 subalgebra that is a weak c -ideal of L if and only if L is solvable.

28 **Proof** Suppose first that L has a solvable maximal subalgebra M that is a weak c -ideal of L . We show that
 29 L is solvable. Let L be a minimal counter-example. Then there is a subideal K of L such that $L = M + K$
 30 and $M \cap K \subseteq M_L$. If $M_L \neq 0$ then L/M_L is solvable, by the minimality assumption, and M_L is solvable,
 31 whence L is solvable, a contradiction. It follows that $M_L = 0$ and $L = M + K$. If R is the solvable radical
 32 of L then $R \subseteq M_L = 0$, so L is semisimple. But now, for all $n \geq 1$, $L = L^{(n)} \subseteq K \neq L$, by Lemma 3.5, a
 33 contradiction. The result follows. The converse follows from Corollary 3.3. \square

1 **Theorem 3.7** Let L be a Lie algebra over a field of characteristic zero such that all maximal nilpotent
 2 subalgebras are weak c -ideals of L . Then L is solvable.

3 **Proof** Suppose that L is not solvable but that all maximal nilpotent subalgebras of L are weak c -ideals of
 4 L . Let $L = R \oplus S$ be the Levi decomposition of L , where $S \neq 0$. Let B be a maximal nilpotent subalgebra
 5 of S and U be a maximal nilpotent subalgebra of L containing it. Then there is a subideal C of L such that
 6 $L = U + C$ and $U \cap C \subseteq U_L$. It follows from Lemma 3.5 that $S = S^{(n_0)} \subseteq L^{(n_0)} \subseteq C$, and so $B \subseteq U \cap C \subseteq U_L$,
 7 whence $S \cap U_L \neq 0$. But $S \cap U_L$ is an ideal of S and so is semisimple. Since U is nilpotent this is a contradiction.
 8 \square

9 **Theorem 3.8** Let L be a Lie algebra, over a field F of characteristic zero, in which every Cartan subalgebra
 10 of L is a weak c -ideal of L . Then L is solvable.

11 **Proof** Suppose that every Cartan subalgebra of L is a weak c -ideal of L , and that L has a non-zero Levi
 12 factor S . Let H be a Cartan subalgebra of S and let B be a Cartan subalgebra of its centralizer in the solvable
 13 radical of L . Then $C = H + B$ is a Cartan subalgebra of L (see [3]) and there is a subideal K of L such that
 14 $L = C + K$ and $C \cap K \subseteq C_L$. Now there is an $r \geq 2$ such that $L^{(r)} \subseteq K$, by Lemma 3.5. But $S \leq L^{(r)} \subseteq K$,
 15 so $C \cap S \subseteq C \cap K \subseteq C_L$ giving $C \cap S \subseteq C_L \cap S = 0$, a contradiction. It follows that $S = 0$ and hence that L
 16 is solvable. \square

17 4. Some characterisations of supersolvable algebras

18 The following is proved in [9, Lemma 4.1]

19 **Lemma 4.1** Let L be a Lie algebra over any field F , let A be an ideal of L and let U/A be a maximal
 20 nilpotent subalgebra of L/A . Then $U = C + A$, where C is a maximal nilpotent subalgebra of L .

21 We will also need the following result.

22 **Lemma 4.2** Let L be a Lie algebra over any field F and suppose that $L = B + K$, where B is a nilpotent
 23 subalgebra and K is a subideal of L . Then there exists $s \in \mathbb{N}$ such that $L^s \subseteq K$. Moreover, if A is a minimal
 24 ideal of L then either $A \subseteq K$ or $[L, A] = 0$.

25 **Proof** Since K is a subideal of L , there exists $r \in \mathbb{N}$ such that $L(\text{ad } K)^r \subseteq K$. As B is nilpotent, there
 26 exists $s \in \mathbb{N}$ such that $L^s = (B + K)^s \subseteq K$. Now $[L, A] = A$ or $[L, A] = 0$ and the former implies that
 27 $A \subseteq L^s \subseteq K$. \square

28 **Lemma 4.3** Let L be a Lie algebra, over any field F , in which every maximal subalgebra of each maximal
 29 nilpotent subalgebra of L is a weak c -ideal of L , and let A be a minimal abelian ideal of L . Then every
 30 maximal subalgebra of each maximal nilpotent subalgebra of L/A is a weak c -ideal of L/A .

31 **Proof** Suppose that U/A is a maximal nilpotent subalgebra of L/A . Then $U = C + A$ where C is
 32 a maximal nilpotent subalgebra of L by Lemma 4.1. Let B/A be a maximal subalgebra of U/A . Then
 33 $B = B \cap (C + A) = B \cap C + A = D + A$ where D is a maximal subalgebra of C with $B \cap C \subseteq D$. Now D is
 34 a weak c -ideal of L so there is a subideal K of L with $L = D + K$ and $D \cap K \subseteq D_L$.

If $A \leq K$ we have

$$\frac{L}{A} = \frac{D+K}{A} = \frac{D+A}{A} + \frac{K}{A} = \frac{B}{A} + \frac{K}{A},$$

and

$$\frac{B}{A} \cap \frac{K}{A} = \frac{B \cap K}{A} = \frac{(D+A) \cap K}{A} = \frac{D \cap K + A}{A} \leq \frac{D_L + A}{A} \leq \left(\frac{B}{A} \right)_{L/A}.$$

So suppose that $A \not\leq K$. Then Lemma 4.2 shows that $[L, A] = 0$. It follows that $A \leq C$ and $B = D$. We have $L = B + K$ and $B \cap K \leq B_L$, so

$$\frac{L}{A} = \frac{B}{A} + \frac{K+A}{A}$$

and

$$\frac{B}{A} \cap \frac{K+A}{A} = \frac{B \cap (K+A)}{A} = \frac{B \cap K + A}{A} \leq \frac{B_L + A}{A} \leq \left(\frac{B}{A} \right)_{L/A}.$$

□

2 **Lemma 4.4** Let L be a Lie algebra over any field F , in which every maximal nilpotent subalgebra of L is a
 3 weak c -ideal of L , and suppose that A is a minimal abelian ideal of L and M is a core-free maximal subalgebra
 4 of L . Then A is one dimensional.

5 **Proof** We have that $L = A \dot{+} M$ and A is the unique minimal ideal of L , by [10, Theorem 1.1]. Let C be a
 6 maximal nilpotent subalgebra of L with $A \leq C$. If $A = C$, choose B to be a maximal subalgebra of A , so that
 7 $A = B + Fa$ and $B_L = 0$. Then B is a weak c -ideal of L . So there is a subideal of K of L with $L = B + K$
 8 and $B \cap K \leq B_L = 0$. Now $L = B + K = B + K^L = K^L$, since $B \leq A \leq K^L$. It follows that $K = L$, whence
 9 $B = 0$ and $A = Fa$ is one dimensional.

10 So suppose that $C \neq A$. Then $C = A + M \cap C$. Let B be a maximal subalgebra of C containing $M \cap C$.
 11 Then B is a weak c -ideal of L , so there is a subideal K of L with $L = B + K$ and $B \cap K \leq B_L$. If $A \leq B_L \leq B$,
 12 we have $C = A + M \cap C \leq B$, a contradiction. Hence $B_L = 0$ and $L = B \dot{+} K$. Now $C = B + C \cap K$ and
 13 $B \cap C \cap K = B \cap K = 0$. As C is nilpotent this means that $\dim(C \cap K) = 1$. If $A \subseteq K$ we have that $A \leq C \cap K$,
 14 so $\dim A = 1$, as required. Otherwise, $[L, A] = 0$, by Lemma 4.2 and again $\dim A = 1$. □

15 We can now prove our main result.

16 **Theorem 4.5** Let L be a solvable Lie algebra over any field F in which every maximal subalgebra of each
 17 maximal nilpotent subalgebra of L is a weak c -ideal of L . Then L is supersolvable.

18 **Proof** Let L be a minimal counter-example and let A be a minimal abelian ideal of L . Then L/A satisfies the
 19 same hypothesis by Lemma 4.3 We thus have that L/A is supersolvable and it remains to show that $\dim A = 1$.

If there is another minimal ideal I of L , then

$$A \cong (A + I)/I \leq L/I$$

20 which is supersolvable and so $\dim A = 1$. So we can assume that A is the unique minimal ideal of L . Also, if
 21 $A \leq \varphi(L)$, we have that $L/\varphi(L)$ is supersolvable, whence L is supersolvable by [2, Theorem 7]. We therefore,
 22 further assume that $A \not\leq \varphi(L)$. It follows that $L = A \dot{+} M$, where M is a core-free maximal subalgebra of L .
 23 The result now follows from Lemma 4.4. □

1 If L has no one-dimensional maximal nilpotent subalgebras, we can remove the solvability assumption
 2 from the above result provided that F has characteristic zero.

3 **Corollary 4.6** Let L be a Lie algebra over a field F of characteristic zero in which every maximal nilpotent
 4 subalgebra has dimension at least two. If every maximal subalgebra of each maximal nilpotent subalgebra of L
 5 is a weak c -ideal of L , then L is supersolvable.

6 **Proof** Let N be the nilradical of L , and let $x \notin N$. Then $x \in C$ for some maximal nilpotent subalgebra C
 7 of L . Since $\dim C > 1$, there is a maximal subalgebra B of C with $x \in B$. Then there is a subideal K of L
 8 such that $L = B + K$ and $B \cap K \subseteq B_L \subseteq C_L \subseteq N$. Clearly, $x \notin K$, since otherwise $x \in B \cap K \subseteq N$. Moreover,
 9 $L^r \subseteq K$ for some $r \in \mathbb{N}$, by Lemma 4.2. We have shown that if $x \notin N$ there is a subideal K of L with $x \notin K$
 10 and $L^r \subseteq K$.

11 Suppose that L is not solvable. Then there is a semisimple Levi factor S of L . Choose $x \in S$. Then
 12 $x \in S = S^r \subseteq K$, a contradiction. Thus L is solvable and the result follows from Theorem 4.5. \square

13 If L has a one-dimensional maximal nilpotent subalgebra, then we can also remove the solvability
 14 assumption from Theorem 4.4, provided that underlying field F has again characteristic zero and L is not
 15 three-dimensional simple.

16 **Corollary 4.7** Let L be a Lie algebra over a field F of characteristic zero. If every maximal subalgebra of
 17 each maximal nilpotent subalgebra of L is a weak c -ideal of L , then L is supersolvable or three dimensional
 18 simple.

19 **Proof** If every maximal nilpotent subalgebra of L has dimension at least two, then L is supersolvable by
 20 Corollary 4.6. So we need only consider the case where L has a one-dimensional maximal nilpotent subalgebra
 21 say Fx . Suppose first that L is semisimple, so $L = S_1 \oplus \dots \oplus S_n$, where S_i is a simple ideal of L for $1 \leq i \leq n$.
 22 Let $n > 1$. If $x \in S_i$, then choosing $s \in S_j$ with $j \neq i$, we have that $Fx + Fs$ is a two dimensional abelian
 23 subalgebra, which contradicts the maximality of Fx . If $x \notin S_i$, for every $1 \leq i \leq n$, then x has nonzero
 24 projections in at least two of the S_k 's, say $s_i \in S_i$ and $s_j \in S_j$. But then $Fx + Fs_i$ is a two-dimensional
 25 abelian subalgebra, a contradiction again. It follows that L is simple. But then Fx is a Cartan subalgebra of
 26 L , which yields that L has rank one and thus is three dimensional.

27 So now let L be a minimal-counter example. We have seen that L is not semisimple, so it has a minimal
 28 abelian ideal A . By Lemma 4.3, L/A is supersolvable or three-dimensional simple. In the former case, L is
 29 solvable and so is supersolvable, by Theorem 4.5.

30 In the latter case, $L = A \oplus S$ where S is three-dimensional simple, and so a core-free maximal
 31 subalgebra of L . It follows from Lemma 4.4 that $\dim A = 1$. But now $C_L(A) = A$ or L . In the former
 32 case $S \cong L/A = L/C_L(A) \cong Inn(A)$, a subalgebra of $Der(A)$, which is impossible. Hence $L = A \oplus S$, where
 33 A and S are both ideals of L and again L has no one-dimensional maximal nilpotent subalgebras. \square

34 5. One dimensional weak c -ideals

35 **Lemma 5.1** Let L be a Lie algebra over any field F . Then the one-dimensional subalgebra Fx of L is a weak
 36 c -ideal of L if and only if it is a c -ideal of L .

37 **Proof** Let Fx be a weak c -ideal of L . Then there is a subideal K of L such that $L = Fx + K$ and
 38 $Fx \cap K \leq (Fx)_L$. Since either $K = L$ or K has codimension one in L , it is an ideal of L and Fx is a c -ideal

1 of L . □

2 We say that L is *almost abelian* if $L = L^2 \oplus Fx$, where L^2 is abelian and $[x, y] = y$ for all $y \in L^2$. Then
 3 the following result follows from Lemma 5.1 and [9, Theorem 5.2].

4 **Theorem 5.2** Let L be a Lie algebra over any field F . Then all one-dimensional subalgebras of L are weak
 5 c -ideals of L if and only if:

6 (i) $L^3 = 0$; or

7 (ii) $L = A \oplus B$, where A is an abelian ideal of L and B is an almost abelian ideal of L .

8 Acknowledgment

9 The authors would like to thank the referees for their valuable comments.

10 References

- 11 [1] Barnes, DW. Nilpotency of Lie algebras. *Mathematische Zeitschrift* 1962;79:237-238
- 12 [2] Barnes, DW. On the cohomology of soluble Lie algebras. *Mathematische Zeitschrift* 1967;101:343-349
- 13 [3] Dixmier, J. Sous-algèbres de Cartan et décompositions de Levi dans les algèbres de Lie (French). *Trans. Roy. Soc.*
 14 *Canada Sect. III* (3)1956; 50:17-21
- 15 [4] Jehad, J. Some conditions for solubility. *Mathematics Journal of Okayama University* 2000;42:1-5
- 16 [5] Skiba, AN. A note on c -normal subgroups of finite groups. *Algebra and Discrete Mathematics* 2005;3:85-95
- 17 [6] Stewart, IN. Subideals of Lie Algebras. University of Warwick institutional repository Ph.D. thesis 1969
- 18 [7] Tashtoush, M. Weakly c -normal and cs -normal subgroups of finite groups. *Jordan Journal of Mathematics and*
 19 *Statistics* 2008;1:123-132
- 20 [8] Towers, DA. A Frattini Theory for Algebras. *Proceedings of the London Mathematical Society* 1973; (3), 27:440-462
- 21 [9] Towers, DA. C -ideals of Lie algebras. *Communications in Algebra*, 2009;37:4366-4373.
 22 <https://doi.org/10.1080/00927870902829023>
- 23 [10] Towers, DA. Maximal subalgebras and chief factors of Lie algebras. *Journal of Pure and Applied Algebra* 2016;
 24 220: 482-493. <https://doi.org/10.1016/j.jpaa.2015.07.005>
- 25 [11] Towers, DA. The generalised nilradical of a Lie algebra. *Journal of Algebra* 2017; 470: 197-
 26 218. <https://doi.org/10.1016/j.jalgebra.2016.08.037>
- 27 [12] Wang, Y. C -normality of groups and its properties. *Journal of Algebra* 1996; 180:954-965
- 28 [13] Zhong G, Yang L, Ma X, and Lin S. On s -Permutably embedded and weakly c -normal subgroups of finite groups.
 29 *International Electronic Journal of Algebra* 2013; 14, 44-52
- 30 [14] Zhong G, Yang L. Finite groups with weakly c -normal subgroups. *Southeast Asian Bulletin of Mathematics* 2013;
 31 37:297-305
- 32 [15] Zhu L, Guo W, Shum KP. Weakly c -normal subgroups of finite groups and their properties. *Communications in*
 33 *Algebra* 2002; 30: 5505-5512. <https://doi.org/10.1081/AGB-120015666>
- 34 [16] Zusmanovich, P. Deformations of $W_1(n) \otimes A$ and modular semisimple Lie algebras with a solvable maximal
 35 subalgebra. *Journal of Algebra* 2003; 268: 603-635. [https://doi.org/10.1016/S0021-8693\(03\)00295-3](https://doi.org/10.1016/S0021-8693(03)00295-3) th a
 36 solvable maximal subalgebra”, *Journal of Algebra*, 268: 603-635, 2003.