

1 **When did the Indus River take on its “modern” drainage configuration?**

2
3 Yani Najman,¹ Guangsheng Zhuang,^{1,2} Andy Carter³, Lorenzo Gemignani^{4,5}, Ian Millar⁶, Jan Wijbrans⁴.

4
5 ¹LEC, Lancaster University, LA1 4YQ, UK

6 ²Department of Geology and Geophysics, Louisiana State University, Baton Rouge, LA 70803, USA

7 ³Dept of Earth and Planetary Sciences, Birkbeck College, University of London, UK.

8 ⁴Institut für Geologische Wissenschaften, Freie Universität Berlin, D-12249 Berlin, Germany.

9 ⁵ Department of Earth Sciences, Vrije Universiteit Amsterdam, 1081 HV, Amsterdam, Netherlands

10 ⁶NERC Isotope Geosciences Lab, BGS Keyworth, Nottingham, UK.

11
12 **Abstract:**

13 In order for sedimentary archives to be used as a record of hinterland evolution, the factors affecting the
14 archive must be known. In addition to tectonics, a number of factors such as changes in climate and
15 palaeo-drainage, as well as the degree of diagenesis, influence basin sediments. The Indus River-delta-fan
16 system records a history of Himalayan evolution, and both the onshore and offshore sedimentary
17 repositories have been extensively studied to research orogenesis. However, a number of unknowns
18 remain with regards to this system. This paper seeks to elucidate the palaeodrainage of the Indus River,
19 in particular when it took on its modern drainage configuration with respect to conjoinment of the main
20 Himalayan (Punjabi) tributary system with the trunk Indus River. We leverage the fact that the Punjabi
21 tributary system has a significantly different provenance signature to the main trunk Indus, draining
22 mainly the Indian plate. Therefore, after the time when the Punjabi tributary system joined the main trunk
23 Indus, the proportion of Indian plate material in the repositories downstream of the confluence should
24 have a higher proportion of Indian plate material compared to the upstream repository. We compared

25 bulk Sr-Nd data and detrital zircon U-Pb data from the Cenozoic upstream peripheral foreland basin and
26 downstream Indus delta and Indus Fan repositories. We determined that repositories below the
27 confluence had a higher proportion of Indian plate material compared to repositories above the
28 confluence, throughout Neogene times. We therefore conclude that the Indus River took on its current
29 configuration with the Punjabi tributary system draining into the Indus trunk river in the Paleogene, early
30 in the history of the orogen: Pinpointing the exact time when the tributary system joined the Indus should
31 be determinable from a shift to more Indian plate input in the downstream repositories only. Whilst the
32 upstream repository records no change in Indian plate input from Eocene to Neogene times, a shift to
33 increased Indian plate material occurs at the Eocene-Oligocene boundary in the delta but sometime
34 between 50-40 Ma in the fan. Further work is therefore required to understand the discrepancy between
35 the two downstream repositories but nevertheless we can conclude that the tributary system joined the
36 trunk Indus at or before the start of the Oligocene.

37

38 **Keywords:** Himalaya; Indus River; provenance; Sr-Nd analyses, zircon U-Pb analyses, mica Ar-Ar analyses.

39

40 **1. Introduction**

41 The Himalaya, as the largest orogen on Earth, garners significant interest from researchers in a variety of
42 disciplines. Whilst considerable information on the mountain belt's evolution can be determined from its
43 hard rock geology, its early history is often destroyed in these rocks by later tectonism, metamorphism
44 and/or erosion. In these circumstances, researchers turn to information recorded in the sediment archive
45 of material eroded from the mountain belt and preserved in surrounding sedimentary basins, both
46 onshore and offshore.

47 The main repositories of Himalayan detritus are preserved in the orogen's suture zone , peripheral and
48 axial foreland basins onshore (e.g. Hodges 2000, Najman 2006, Shah 2009), and the Indus and Bengal Fans
49 offshore, which are the world's largest sediment fans (Nyberg et al., 2018). Detritus from all these basins
50 has been studied to document hinterland evolution, using a variety of bulk rock and single grain analytical
51 techniques. For example, studies of the Indus River's sedimentary repository include detrital feldspar Pb
52 isotopic analyses applied to the Indus suture zone molasse (Clift et al., 2001b), detrital zircon fission track
53 and Sm-Nd bulk analyses have been applied to the peripheral foreland basin sedimentary rocks (e.g.,
54 Chirouze et al., 2015), detrital zircon U-Pb analyses have been applied to the axial foreland basin (e.g.,
55 Zhuang et al., 2015) and heavy mineral and petrography (Andò et al., 2020; Garzanti et al., 2020) data
56 applied to the Indus Fan. However, in order for the sediment archives to be robustly interpreted, a
57 knowledge of the river's palaeo-drainage evolution must be known, since significant drainage changes will
58 affect the sediment archive. Reconstruction of the lower Indus palaeo-drainage is the focus of this paper.

59 Today, the Indus River flows west along the Indus suture zone which separates the Indian and Asian plates,
60 before turning south across the Himalayas to flow eventually into the northern Indian Ocean, giving rise
61 to the Indus Fan (Fig. 1). Here we define the Lower Indus as that part of the Indus River downstream
62 (south) of the Himalayan mountain front, flowing axially, southward along the Indus Basin. We define the
63 Upper Indus as that part of the Indus River which flows through the mountains, sub-divided into the "west-
64 flowing axial Upper Indus" which flows from its headwaters, west, axially along the Indus suture zone,
65 and, further downstream, the "south-flowing transverse Upper Indus", which cuts south across the
66 mountain range (Fig 1A). The main tributaries to the Indus are the Punjabi or Himalayan tributaries of the
67 Jhelum, Chenab, Ravi, Beas and Sutlej Rivers, herein called the Punjabi tributary system, which drain
68 predominantly the Indian plate (Fig 1C).

69 Mid Eocene Owen Ridge sediments are considered to be early Indus Fan material derived from north of
70 the Indian plate (Clift et al., 2001a; Clift et al., 2002a). This places a lower bound on the timing of initiation
71 of the Indus River, although its upstream configuration is debated. For the Upper Indus, the proposed
72 time of its initiation as a river flowing westward along the suture zone ranges from early Eocene through
73 Miocene (Bhattacharya et al., 2020; Clift et al., 2001b; Henderson et al., 2011; Najman, 2006; Sinclair and
74 Jaffey, 2001).

75 This paper focusses on the palaeodrainage evolution of the Lower Indus River. Various suggestions have
76 been made regarding whether western Himalayan rivers switched between flowing east to the Ganges
77 and Bengal Fan catchment and west to the Indus River and Fan catchment. Whilst a number of authors
78 based their interpretations on palaeocurrent data from the peripheral foreland basin deposits (e.g. see
79 review in Burbank et al, 1996), Clift and Blusztajn (2005) used geochemical data from the Indus Fan. They
80 considered that changes in the geochemical signature of the Indus Fan sediment archive after 5 Ma
81 represented a major drainage change in the Lower Indus at this time, when the Punjab tributaries to the
82 Lower Indus River (Jhelum, Chenab, Ravi and Sutlej Rivers, Figs. 1A and 1C) switched from flowing east to
83 the Ganges and Bengal Fan, to west to the Indus River and Indus Fan. However, Chirouze et al. (2015),
84 considered that this geochemical change could be better interpreted as the result of variations in upland
85 exhumation. This suggestion was later agreed upon by the original proponents of the drainage diversion
86 hypothesis (Clift et al. (2019) and Zhou et al. (2021)) and thus the timing when the Indus River took on its
87 current configuration with respect to the Punjab tributary system remains unknown.

88 Using a similar rationale to Chirouze et al (2015), we compare provenance indicators from upstream and
89 downstream of the confluence of the Punjab tributary system with the Indus River, and from this
90 comparison we determine when a provenance change in the downstream repository is detected and thus
91 when the tributary system joined the trunk Indus. We extend the peripheral foreland basin Sr-Nd dataset

92 of Chirouze et al (2015) from the mid Miocene, down section as far as the Eocene in order to determine
93 when the provenance change occurred, and additionally we apply new provenance indicators, namely
94 detrital zircon U-Pb ages and detrital white mica Ar-Ar ages.

95

96 **2. Background:**

97 ***2.1 Himalayan geology***

98 *2.1.1 Tectonic units*

99 The Indus suture zone separates the Asian Lhasa Terrane to the north from the Indian plate to the south.
100 In the west, the Kohistan-Ladakh intra-oceanic island arc (KLIA) is sandwiched between the Indian and
101 Asian plates, with the southern margin of the Asian plate at this location comprised of the Karakoram and
102 Hindu Kush). The northern suture separating the Asian plate and KLIA is termed the Shyok Suture Zone
103 and the southern suture separating the KLIA and Indian plate is termed the Indus Suture zone, also known
104 as the Main Mantle Thrust in this region (Fig. 1C). The Indus River flows west along the Indus suture zone
105 before turning south to cross the Himalaya and foreland basin before debouching into the Arabian Sea.

106 The Lhasa terrane comprises Phanerozoic low grade metamorphic and sedimentary cover overlying
107 Precambrian-Cambrian basement (e.g., Leier et al., 2007). Along its southern rampart are intruded the
108 Gangdese continental arc batholiths of the Transhimalaya which represent the Andean-type southern
109 margin of Asia prior to consumption of the intervening Neo-Tethys ocean (Schärer et al., 1984). Whilst
110 Gangdese intrusions are Mesozoic-Paleogene aged, post-collisional igneous activity continued into
111 Miocene times (Hodges, 2000). To the west, the Lhasa terrane terminates against the Karakoram Fault.
112 West of the fault, the southern margin of Asia is represented by the Karakoram (Fig 1C). The Karakoram
113 terrane is divided into three units (Hildebrand et al., 1998; Hildebrand et al., 2001; Searle et al., 1999): the

114 Northern Karakoram Sedimentary Unit, the Southern Karakoram Metamorphic Belt and the intervening
115 Karakoram Batholith. The Northern Karakoram Sedimentary Unit comprises pre-Ordovician crystalline
116 basement covered by an Ordovician to Cretaceous sedimentary succession (Gaetani and Garzanti, 1991;
117 Gaetani et al., 1993; Zanchi and Gaetani, 2011). The Karakoram Batholith includes pre-India–Asia collision,
118 Andean-type subduction-related granitoids and post-India–Asia collision leucogranites. Age of
119 metamorphism of the Southern Karakoram Metamorphic Belt ranges from Late Cretaceous to late
120 Miocene (Fraser et al., 2001; Palin et al., 2012; Searle et al., 2010).

121 The Kohistan-Ladakh Island arc (KLIA) separates the Indian and Asian plates in the west of the orogen. It
122 consists of Late Cretaceous and Eocene plutonic belts, and pyroxene granulites, calc-alkaline volcanics,
123 amphibolites, and minor metasediments (Coward et al., 1984; Schaltegger et al., 2002), fringed by
124 ophiolitic melange in the southern suture (DiPietro et al., 2000; DiPietro and Pogue, 2004).

125 The Indian plate lies to the south of the KLIA. As summarised in Hodges (2000), in the central and eastern
126 part of the orogen, the Indian plate Himalaya is divided, from north to south, into the Tethyan Himalaya,
127 the Greater Himalaya, Lesser Himalaya and Cenozoic foreland basin sedimentary rocks of the Sub-
128 Himalaya (Fig 1C). Typically, the Tethyan Himalaya, separated from the Greater Himalaya to the south by
129 the South Tibetan Detachment System, consists of Paleozoic–Mesozoic sedimentary and low-grade
130 metasedimentary rocks which were deposited on the Tethyan ocean passive margin; the Greater
131 Himalaya, separated from the Lesser Himalaya to the south by the Main Central Thrust, consists predomi-
132 nantly of medium- to high-grade Neoproterozoic–Ordovician metamorphic rocks that were subjected to
133 metamorphism and anatexis during the Cenozoic Himalayan orogeny when they were intruded by
134 Neogene leucogranites; and the Lesser Himalaya, separated from the Cenozoic Sub-Himalaya foreland
135 basin sedimentary rocks to the south by the Main Boundary Thrust, consisting of Paleoproterozoic meta-
136 morphosed and unmetamorphosed Indian plate rocks. These lithologies also broadly constitute the Indian

137 plate Himalaya to the west in Pakistan. However, exact correlation is uncertain, the degree of
138 metamorphism differs, and the lithologies are not structurally imbricated in the same way (Treloar et al.,
139 2019). According to DiPietro and Pogue (2004), north of the Khairabad Thrust (MCT equivalent in
140 Pakistan) metamorphosed rocks of ages equivalent to the Tethyan, Greater and Lesser Himalaya are
141 found, whilst both Lesser and Tethyan equivalents are found between the Khairabad Thrust and the MBT.
142 The Nanga Parbat syntaxis is considered to be of Lesser, Greater and Tethyan Himalayan affinity (Argles
143 et al., 2003). In this paper, we refer to the Neoproterozoic-Ordovician rocks as *Greater Himalayan*
144 *lithological correlatives*, the Paleoproterozoic rocks as *Lesser Himalayan lithological correlatives*, and the
145 Paleozoic-Mesozoic rocks as *Tethyan Himalaya lithological correlatives*. Such terms do not reflect the
146 location of the rocks within the various thrust-bound terranes, as they do further east.

147 The units described above have distinct zircon U-Pb ages and Nd isotope signatures associated with
148 different crustal evolution histories (e.g., Argles et al., 2003; Clift et al., 2019; DeCelles et al., 2004;
149 DeCelles et al., 2016a; Gehrels et al., 2011; Najman, 2006). These differences, (Table 1), allow for use of
150 these techniques as provenance indicators in the detrital record downstream (e.g., Clift et al., 2019;
151 DeCelles et al., 2004; DeCelles et al., 2016a; Gehrels et al., 2011; Najman, 2006).

152 The overwhelming majority of zircons from the Indian plate have U-Pb ages >400 Ma (DeCelles et al.,
153 2004; Gehrels et al., 2011), with the minor exception of grains dated ~130 Ma from the Tethyan Himalaya
154 (e.g. Clift et al., 2014) and Neogene grains eroded from leucogranites (e.g. Hodges, 2000 and references
155 therein). Within the Indian plate, grains 1500-2300 Ma are characteristic of the Lesser Himalaya, and 300-
156 1250 Ma characteristic of the Greater and Tethyan Himalaya, although not uniquely so (Clift et al., 2019).
157 By contrast, zircons from the KLIA are exclusively aged 40-200 Ma, whilst the southern Asian margin (the
158 Karakoram and to the east the Lhasa Block) also have a high proportion of grains of such age, but also

159 with some Neogene grains, and older grains stretching to the Precambrian, derived from the substrate
160 into which the Mesozoic-Paleogene plutons intruded (e.g. Zhuang et al., 2018 and references therein).

161 The old continental crust of the Indian plate has a mean ϵ_{Nd} value of -15 for the Greater Himalaya, -22 for
162 the Lesser Himalaya, and -11 for the Tethyan Himalaya (Ahmad et al., 2000; Deniel et al., 1987; Richards
163 et al., 2005; Robinson et al., 2001; Zhang et al., 2004). By contrast, the Asian and intra-oceanic arc terranes
164 have more positive values, reflecting the dominance of Mesozoic-Paleogene plutons: the KLIA has values
165 around +5 (Bignold and Treloar, 2003; Khan et al., 1997; Khan et al., 2004; Khan et al., 2009), whilst the
166 Karakoram, which consists of both old sedimentary and metamorphic rocks as well as younger plutons,
167 has an average value around -9.6 (Mahéo et al., 2009; Miller et al., 1999). Data from the Lhasa Block are
168 mainly from the central and eastern part of the orogen: the Gangdese / Transhimalaya have values ranging
169 from +0.9 to 5.5 for the Mesozoic granitoids and +2.4-8.5 for the Paleocene-Eocene granitoids in contrast
170 to the Oligocene-Miocene granitoids with values of -9.4 to 5.5 (Ji et al., 2009; Pan et al., 2014), whilst the
171 continental substrate into which these plutons intruded have an average recorded ϵ_{Nd} value of -9 (Pan et
172 al., 2014; Zhu et al., 2009; Zhu et al., 2012).

173 *2.1.2. Tectonic evolution.*

174 Prior to India-Asia collision, India was subducting beneath Asia as Neo-Tethys closed, with the KLIA located
175 between the two continents in the west. The timing of India-Asia collision, and whether the island arc
176 collided with India or Asia first, is disputed; a majority of researchers consider India-Asia collision occurred
177 around 55-60 Ma (see review in Hu et al., 2016 and references therein) with other estimates extending to
178 c. 35 Ma or 25-20 Ma (Aitchison et al., 2007; Bouilhol et al., 2013; van Hinsbergen et al., 2012).

179 The west differs from the better studied east and central part of the orogen in both the presence of the
180 KLIA, and in the timing of exhumation of the Indian plate. In the west, a tectonic wedge consisting of the
181 KLIA, ophiolitic melange and thrust slices of Lesser Himalayan and Tethyan correlatives of the Indian plate

182 was in position and thrust over the Indian plate foreland prior to 47 Ma. Thereafter, Indian plate Lesser-,
183 Greater- and Tethyan Himalayan correlatives were exhumed from beneath the wedge (DiPietro et al.,
184 2008), predominantly during the Paleogene with a pulse of deformation also in the earliest Miocene, ~20
185 Ma (Argles et al., 2003, and references therein; DiPietro et al., 2021). Substantial rapid exhumation of the
186 Indian plate hinterland is not recorded after this time, except in the Nanga Parbat region (Fig 1C), a
187 syntaxis of Lesser, Greater and Tethyan Himalayan lithological correlatives, where rapidly accelerating
188 exhumation is recorded over the Pliocene (e.g., Schneider et al., 2001). Thrusting and exhumation
189 propagated south towards the foreland in the mid or late Miocene, continuing into the Pliocene (Burbank
190 and Tahirkheli, 1985; Yeats and Hussain, 1987).

191 To the north of the Indian plate, moderate exhumation is recorded from Eocene times in the Kohistan
192 Island arc (Van Der Beek et al., 2009) and by contrast, the Karakoram of the Asian plate records periods
193 of rapid exhumation around 27-35 Ma, 13-17 Ma, 7-8 Ma and 3.3-7.4 Ma (Dunlap et al., 1998; Wallis et
194 al., 2016; Zhuang et al., 2018).

195 **2.2. Foreland basin geology**

196 In Pakistan, current basinal environments along which the modern Indus River flows, consist of (1) the
197 peripheral foreland basin that strikes east-west along the southern margin of the orogen, and (2) the
198 north-south striking Lower Indus axial foreland basin along which the Lower Indus River debouches into
199 its delta in the Arabian Sea (Fig. 1A).

200 **2.2.1 Peripheral foreland basin**

201 Foreland basin stratigraphy is for the most part invariant along strike in the orogen, with local minor facies
202 variation, although formation names differ. In Pakistan, the Paleogene has a number of formation names
203 for equivalent units, in different areas (Pivnik and Wells, 1996). We adopt the formation names in our

204 area of study, which for our Paleogene samples is the Hazara-Kashmir syntaxis (HKS) (Figs. 1A and B), with
205 the stratigraphy as recorded in Table 1. At this location, the Paleocene Lockhart Limestone is overlain
206 successively by the latest Paleocene (57-55 Ma) Patala Formation, the early Eocene (55-53 Ma) Margala
207 Hill and Chorgali Formations, and the early-Mid Eocene (53-43 Ma) Kuldana Formation (Baig and Munir,
208 2007; Bossart and Ottiger, 1989; Ding et al., 2016b; Qasim et al., 2018). These formations, which stretch
209 from marine facies to the transitional Kuldana Formation, are separated from the overlying continental
210 alluvial facies by a late Eocene-Oligocene unconformity. Above the unconformity, there is the Murree
211 Formation, also called the Balakot Formation in the HKS. In this syntaxis, the Murree Formation has a
212 latest Oligocene maximum depositional age (MDA) as determined by the two youngest zircons within
213 error, with a weighted mean U-Pb age of 22.6 ± 1.0 Ma (this study, section 4.2) from a sample collected
214 near Paras, north of Balakot (Fig. 1B), supported by a grain dated at 22.7 ± 0.4 Ma (Ding et al 2016b)
215 from a section 15 kms south at Muzaffarabad (Fig. 1B). South-west of the Hazara-Kashmir syntaxis, at
216 Murree hill station (MHS) (Fig. 1B), detrital mica Ar-Ar ages indicate an MDA of <24 Ma (this study, section
217 4.3). These MDAs are in agreement with the early Miocene dating of the Murree Formation to the south,
218 based on mammal fossils (Shah, 2009). Further south, in the Kohat and Potwar Plateaus (Fig. 1A) are the
219 alluvial Kamlial Formation and overlying Siwalik Group, subdivided into the Chinji, Nagri and Dhok Pathan
220 Formations (see Table 1 for stratigraphy). These formations are dated by magnetostratigraphy (Johnson
221 et al., 1985), at 18-14 Ma, 14-11 Ma, 11-8.5 Ma, and <8.5 Ma, respectively.

222 *2.2.2 Lower Indus axial Basin*

223 The stratigraphy of the Lower Indus Basin in the Sulaiman and Kirthar regions are broadly correlative
224 (Shah, 2009). It encompasses the early Eocene Ghazij Formation, the middle-late Eocene Kirthar Group,
225 the Oligocene-early Miocene Chitarwata Formation, the late Early to middle Miocene Vihowa Formation,
226 and the middle Miocene-Pliocene rocks of the Siwalik Group (Roddaz et al., 2011; Shah, 2009; Zhuang et

227 al., 2015), as denoted in Table 1. Facies are predominantly marine until the Chitarwata Formation which
228 transitions up from deltaic to fluvial facies. Fluvial facies then persist until the top of the section.

229 **2.3 Paleodrainage models**

230 *2.3.1 The early drainage configuration of the palaeo-Indus: evidence from the Indus Fan sedimentary* 231 *archive*

232 The oldest eastern Indus Fan sample (IODP 355, U1456 and 1457, Fig 1A) to have been subject to detrital
233 zircon U-Pb analyses is 15 Ma. This shows evidence of input from the Karakoram (Zhou et al., 2022),
234 indicating the drainage basin of the palaeo-Indus stretched as far back as the Shyok Suture Zone by this
235 time (Fig 1C). The oldest sample subjected to detrital zircon U-Pb dating in the western part of the Indus
236 Fan (ODP 731, Fig 1A) is ~ 30 Ma. This sample shows evidence of input from the KLIA/Asian plate
237 (undifferentiated), indicating that the river stretched back at least beyond the Indus Suture Zone (Fig 1C)
238 by that time (Feng et al., 2021). Likewise, Mid Eocene Owen Ridge sediments from DSDP 224 (Fig 1A),
239 considered to be early Indus Fan deposits (Clift et al., 2001a; Clift et al., 2002a), show bulk rock ϵ_{Nd}
240 signatures and K-feldspars with Pb isotope compositions indicative of derivation from north of the Indian
241 plate (Clift et al., 2001a). This indicates that the river's drainage basin stretched back as least as far as the
242 Indus Suture Zone and KLIA at this time.

243 *2.3.2. The early drainage configuration of the upper axial palaeo-Indus: evidence from the Indus Suture* 244 *Zone molasse.*

245 Clift et al. (2001b) considered that various isotopic provenance datasets and palaeocurrents in Indus
246 suture zone sedimentary rocks of early Eocene age indicated contribution from the Lhasa Block to the
247 east, requiring along strike east to west flow along the suture zone at that time. However, Najman (2006)
248 argued that an alternative source with a suitable signature could potentially be that of the Karakoram,

249 located north of the suture zone sediments under discussion, and therefore not requiring along-strike
250 transport and axial flow. Sinclair and Jaffey (2001) considered their facies analyses of the suture zone
251 sediments indicated internal rather than through-flowing drainage until at least the early Miocene. Later,
252 Henderson et al. (2010) reported that white micas, interpreted as Indian-plate derived, first occurred in
253 the same suture sedimentary rocks as Asian-derived zircons, in suture zone sedimentary rocks dated <23
254 Ma. From these mixed source sedimentary rocks, and accompanying facies analysis, they considered the
255 Indus River was flowing in the suture zone at that time. However, it should be noted that (1) micas were
256 also recorded in older suture zone sedimentary rocks but they were of too small grain size to analyse, (2)
257 Indian plate material with low muscovite fertility such as from the Tethyan Himalaya may well have
258 contributed to the suture zone rocks earlier and (3) an open question remains as to why the first
259 appearance of micas interpreted as Indian-derived, was not also accompanied by an influx of Paleozoic
260 and older zircons, also typical of the Indian plate. Whilst subsequently, such old zircons, interpreted as
261 Indian rather than Asian-derived, have been documented in suture zone sediments as old as ca. 50 Ma
262 (Bhattacharya et al., 2020), nevertheless they are not present in the samples analysed for white mica Ar-
263 Ar analyses by Henderson et al (2010). Whilst mineral sorting due to different hydraulic regimes of zircon
264 versus mica (Malusà et al., 2016) might explain the difference, we suggest that, with the benefit of
265 subsequent better characterisation of the ages of micas from the southern margin of the Asian plate
266 (Zhuang et al 2018), an Asian Karakoram provenance might provide an alternative provenance for these
267 micas. Regardless, mixed Indian-Asian provenance, unaccompanied by facies data indicating deposition
268 in a major river, does not indicate east-west through-flow of drainage. Bhattacharya et al. (2020)
269 demonstrated from provenance data that detritus from the east was transported west by ca 27 Ma. Thus
270 we may conclude that an axial upper Indus flowed west by Oligocene times. Prior to that the suture zone
271 was a depocentre, but it may have been externally or internally drained.

272 *2.3.3. Early drainage configuration of the upper transverse palaeo-Indus River: evidence from the*
273 *peripheral foreland basin deposits*

274 In the peripheral foreland basin, detrital blue-green hornblende considered to be derived from the KLIA,
275 is first recorded in the Kohat and Potwar plateaus from 11 Ma (Nagri Formation), interpreted as palaeo-
276 Indus deposits (Abbasi and Friend, 1989; Cervený and Johnson, 1989). Ullah et al. (2015) applied
277 geochemistry and petrography to the Chinji Formation (14-11 Ma) to record material from the KLIA and
278 Indus suture zone. Based on petrography, Najman et al. (2003) recorded arc-derived detritus in the
279 Potwar plateau from the start of their studied section at 18 Ma, from which they interpreted that this
280 time represented the first arrival of sediment from the Upper Indus River to the foreland basin in this
281 region. Still later work (Ding et al., 2016b; Qasim et al., 2018) recorded arc-derived zircons in the foreland
282 basin latest Paleocene to Early Eocene Margala Hill and uppermost Patala Formations, indicating
283 derivation from north of the Indus Suture Zone / Main Mantle Thrust since at least 55 Ma.

284 Whilst the above provenance data indicates derivation from material as far north as the KLIA since Eocene
285 times, whether these rocks represent the deposits of the palaeo-Indus is debated (Cervený et al., 1989;
286 Willis, 1993; Zaleha, 1997). Chirouze et al (2015) proposed a Lhasa Block origin for detrital zircons with
287 old fission track ages in the Chinji Formation. This would indicate that the contributing drainage basin
288 stretched into the Shyok Suture Zone and Asian plate by this time, and was therefore likely the palaeo-
289 Indus. However, we suggest that such grains may also be derived from the Indian Himalayan units south
290 of the KLIA, as arguable by their occurrence in the Siwalik foreland basin sedimentary rocks of Nepal, that
291 were deposited by rivers which did not stretch back to Asia (Bernet et al., 2006).

292 However, more definitive evidence of deposition from the palaeo-Indus comes from detrital mica Ar-Ar
293 data. Lag times of detrital mica Ar-Ar ages from Kamliyal Formation Potwar Plateau sedimentary rocks
294 indicate rapid exhumation of the upland source region from 16-14 Ma (Najman et al., 2003). The exhuming

295 source area was interpreted by those authors to be the Karakoram and/or Nanga Parbat region, consistent
296 with both bedrock data from those regions (Treloar et al., 2000; Zhuang et al., 2018 and references
297 therein). Due to their locations, derivation of micas from either location strongly suggests transport by a
298 palaeo-Indus. Furthermore, detritus delivered by possible ancient smaller tributaries draining only the
299 Indian plate and arc would have had a distinct and different signature, with a higher proportion of Indian
300 plate detritus, for example the Mid Miocene Kamli Formation sample CP96-6A from Najman et al.
301 (2003), and presumably those samples from the Eocene Kuldana Formation with a high proportion of old
302 zircons at Muzaffarabad (Ding et al., 2016b) (see section 5.2 for further discussion).

303 *2.3.4. Evolution of the Lower Indus palaeodrainage*

304 Within the basin, the position of the Ganges-Indus drainage divide over time is long debated, with various
305 authors proposing that parts of the current Gangetic catchment used to flow into the Indus Fan (e.g.
306 DeCelles et al., 1998), and the current Indus River catchment into the Bengal Fan (e.g. Burbank et al.,
307 1996) at various times. Clift and Blusztajn (2005) noted a change to more negative ϵ_{Nd} values in the Indus
308 Fan at 5 Ma, which they interpreted as the drainage diversion of the major Indian-plate draining Punjabi
309 Indus River tributary system of the Jhelum, Chenab, Ravi and Sutlej rivers (Figs. 1A and C) from a previous
310 routing towards the Ganges and the Bengal Fan to the east.

311 However, the above argument was countered by Chirouze et al. (2015) who looked at both spatial and
312 temporal trends at the range front and Indus Fan. They considered that the change in the signal was due
313 to differential exhumation in the hinterland rather than drainage re-organization. They compared ϵ_{Nd} data
314 between the range front and Indus Fan for both the present day and the Miocene (using Chinji Formation
315 foreland basin data for the Miocene range front). They recorded a spatial variation of four ϵ_{Nd} units
316 between the range front and the Indus Fan for both mid-late Miocene times and modern day (Miocene
317 range front and Indus Fan values at -6 and -10 respectively; modern day range front and Indus Fan values

318 at -10 and -14 respectively. This suggests a stable drainage pattern for the lower Indus since at least the
319 mid-late Miocene. From the above data they noted a negative shift of $\sim 3 \epsilon_{Nd}$ units between Miocene and
320 the modern day at both the range front (comparison of Miocene foreland basin sedimentary rocks with
321 modern day Upper Indus values) and a similar shift in the Indus Fan. From this temporal shift they
322 therefore concluded that the variation over time was due to the changing exhumation rates of the
323 contributing source regions, with the exhumation and thus contribution of the Karakoram / Indian plate
324 syntaxial Himalaya increasing at the expense of the more positive KLIA (Table 1) to explain the shift in ϵ_{Nd}
325 values in the Indus Fan at 5 Ma. They supported their proposal of variations in exhumation using detrital
326 zircon fission track (ZFT) data, interpreting a decrease in older ZFT ages after 12 Ma as due to decreased
327 input from the KLIA. Later, the original proponents of the drainage capture hypothesis (Clift and Blusztajn,
328 2005) concurred with the view of Chirouze et al. (2015) that changes in the tectonics of the hinterland
329 was the more likely cause of the geochemical change in the Indus Fan at 6 Ma (Clift et al., 2019; Zhou et
330 al., 2022) thus the time when the Punjab tributary system joined the trunk Indus remains unknown. It is
331 towards this question, namely the evolution of the downstream Indus, that this paper focusses.

332 The location of the exit of the Indus River to the ocean in the past retains a level of uncertainty. Today the
333 Indus River debouches to the Arabian sea at the south of the Lower Indus axial Basin. These deposits are
334 recorded in eastern Sulaiman and Kirthar regions of the Lower Indus Axial Basin (Welcomme et al., 2001)
335 (Fig 1A). Zhuang et al. (2015) show that zircons from the KLIA are recorded in these sediments from at
336 least early Oligocene times; they considered that detrital zircon U-Pb data indicate input from the
337 Karakoram from at least Mid Miocene times, and that Sr-Nd data indicate a palaeo-Indus origin from 50
338 Ma. Roddaz et al. (2011) carried out mixture modelling on their Sr-Nd data and concluded that there was
339 an appreciable input from the Karakoram since 50 Ma.

340 However, Palaeogene deltaic facies have also been identified in the Katawaz remnant ocean Basin (Fig
341 1A) to the west (Qayyum et al., 2001). In view of the differing compositions and provenance between
342 these two deltaic systems, Roddaz et al. (2011) proposed two river-delta-fan systems, with the Katawaz
343 system debouching into the Khojak submarine fan and the sediments of the Lower Indus Axial Basin
344 debouching into the Indus Fan. Provenance data from the Katawaz rocks show that that drainage basin
345 stretched back at least as far as the KLIA by Miocene times (Carter et al, 2010) with a paucity of data
346 currently precluding earlier documentation. For a full evaluation of the Indus river-delta-fan system and
347 the spatial evolution, more data are needed from the Katawaz basin; data presented in this paper provide
348 a direct comparison between peripheral foreland basin records and terminal sinks in the delta and ocean.

349

350 **3. Methods**

351 ***3.1 Rationale and approach***

352 To determine when the Punjab tributary system joined the trunk Indus, we leverage that fact that the
353 tributaries have a very different drainage basin lithology to the trunk Indus; the former includes only
354 Himalayan units, whilst the drainage basin of the latter includes also the KLIA and Asian plate (Fig. 1C),
355 which have very different isotopic and geochemical signatures to the Indian plate (Table 1). This difference
356 is clearly reflected in both the Sm-Nd and zircon U-Pb characteristics of the trunk Indus river versus the
357 Punjabi tributary system: Figs 2 and 3A (inset) shows that, compared to the modern Indus trunk river, the
358 Punjabi tributaries have a more negative ϵ_{Nd} value and a much lower proportion of young arc-aged grains
359 (Alizai et al. 2011, Chirouze et al 2015), a signature which extended back into the ancient sedimentary
360 record (Exnicios et al., 2022; Najman et al., 2009).

361 We took a similar approach to Chirouze et al (2015) in hypothesising that prior to the time when the
362 Punjab tributary system joined the trunk Indus River, the sedimentary repositories upstream and

363 downstream of the confluence should look similar in terms of provenance. After the time when the
364 tributary system joined the Indus River, the repository upstream of the confluence should remain similar
365 (unless synchronously affected by a tectonic-induced change in the hinterland), but the downstream
366 repository should show increased input from Himalayan Indian plate units.

367 We therefore made comparison between data upstream (our new foreland basin data) and published
368 data downstream of the Punjab tributary system. Previous work used the Indus Fan as the downstream
369 comparative repository. We use both the deltaic record in the Sulaiman and Kirthar region, and the Indus
370 Fan archive, since onshore sedimentary archives are typically more prone to diagenetic alteration
371 compared to marine records, whilst distal deposits are more prone to the effects of hydraulic sorting (e.g.,
372 Garzanti et al., 2020) and contain evidence of subordinate extraneous (non-Indus River) sources to the
373 Himalayan orogen, such as the Deccan Traps of peninsular India input to the Indus Fan (Clift et al., 2019;
374 Garzanti et al., 2020; Yu et al., 2019). The Indus Fan record is a composite repository of material recovered
375 from the Owen Ridge and Western Fan from DSDP 224 (Eocene-Miocene) and ODP 720, 722 and 731
376 (Eocene-Pleistocene) sites, and IODP 355 sites of U1456 and 1457 of the Eastern Fan (Neogene only) (Clift
377 et al., 2019; Feng et al., 2021; Zhou et al., 2022) (Fig. 1A).

378 Our dataset builds on the previous work of Chirouze et al. (2015) in two ways. Firstly, it expands the time
379 range from the previous mid Miocene study of the Pakistan peripheral foreland basin to now include
380 foreland basin rocks from Eocene to late Miocene. This allows a more complete assessment of the
381 evolution of the lower Indus to be determined. Secondly, we incorporate not only ϵ_{Nd} data from
382 mudstones, but also new and previously published zircon U-Pb data to assess provenance, and mica
383 $^{40}Ar/^{39}Ar$ data to assess exhumation. Therefore, in addition to using both onshore and offshore
384 repositories to limit the potential effects of fertility, diagenetic, and hydraulic sorting biases, our multi-
385 proxy approach provides additional mitigation since: 1) zircons are resistant to diagenesis; 2) we assess

386 evidence from both the mud and sand grain size fractions with the use of both bulk and single grain
387 approaches and 3) we obtain data from both zircon and mica grains which respond differently to the
388 hydraulic regime (e.g., Garzanti and Andò, 2019; Garzanti et al., 2009; Malusà et al., 2016). Furthermore,
389 since white mica is rare in the KLIA, exhumation patterns of the Karakoram and Indian plate Himalaya can
390 be considered in isolation using this technique, unbiased by potential issues surrounding dilution and
391 fertility.

392

393 ***3.2. Samples and analyses***

394 *3.2.1 Samples*

395 We analysed 5 sandstones for detrital zircon, 10 mudstones for Sr-Nd isotopes, and 3 sandstones for mica
396 $^{40}\text{Ar}/^{39}\text{Ar}$. The locations of analysed samples (Figs. 1A and B) are from the Kuldana Formation in the HKS
397 at Paras north of Balakot, the Murree Formation in both the HKS and at Murree Hill Station (MHS), and
398 the Kamliyal, Chinji and Nagri Formations from the Chinji section on the Potwar Plateau, the latter being
399 the same location from which Chirouze et al (2015) took their samples. A summary of our sample
400 information is tabulated in S1. Our samples from the Kuldana Formation are structurally imbricated within
401 the Murree Formation (Najman et al., 2002). Originally, Najman et al (2002) considered these structural
402 imbrications to be Patala Formation, based on the work of Bossart and Ottiger (1989) who did not
403 recognise the Kuldana Formation. However, more recent detailed mapping (Ding et al., 2016b) and the
404 better agreement of biostratigraphic ages from the structural imbricates (early-mid Eocene; Bossart and
405 Ottiger (1989)) with the Kuldana Formation rather than Patala Formation (section 2.2.1), suggests
406 reassignment of these imbricates from the Patala to the Kuldana Formation.

407 *3.2.2 Sr-Nd bulk analyses*

408 Sr and Nd isotope analyses on bulk mudstones were carried out at the NERC Isotope Geosciences
409 Laboratory, Keyworth, Nottingham. Samples were leached in dilute acetic acid in order to remove
410 carbonate material, then dissolved using HF-HNO₃ and converted to chloride form. Sr and a bulk REE
411 fraction were separated using AG50x8 cation columns, and Nd was separated from the bulk REE using LN-
412 SPEC columns. Sr and Nd were analysed on a Thermo Scientific Triton mass spectrometer.

413 *3.2.3 Zircon U-Pb analyses*

414 Detrital zircon U-Pb ages were acquired using laser ablation ICPMS, at the London Geochronology Centre,
415 University College London. To avoid bias, polished grain mounts were made, without hand picking, directly
416 from Diidomethane sink fractions with a grain size $\leq 300 \mu\text{m}$. Each laser spot (25 μm) was placed on the
417 outermost parts of each grain to target the youngest growth stage. Between 150-320 grains were analysed
418 for each sample, providing statistical confidence of detecting all component ages. Data were processed
419 using GLITTER v4.4 data reduction software using age standard bracketing to correct for mass
420 fractionation. Between 8 and 15% of ages were rejected, due to high discordance from lead loss, zoning
421 or mixing of growth zones. One exception was the Chinji Formation that contained an unusually high
422 number (60%) of discordant grains. Most of these discordant grains are associated with ages between 75-
423 120 Ma and consistent with lead loss, likely due to source weathering.

424 *3.2.4 Muscovite Ar-Ar analyses*

425 Muscovite Ar-Ar ages were analysed at the Argon Geochronology Laboratory at VU University Amsterdam,
426 Netherlands. Individual grains ranging from 125-1000 μm were handpicked under a binocular microscope
427 to avoid obvious weathering or inclusions. After irradiation at the Oregon State University TRIGA nuclear
428 reactor, total fusion analyses were carried out with a ThermoFisher Scientific Helix MC plus multi-collector
429 mass spectrometer, fitted with 10^{13} Ohm amplifiers. Data reduction was done using ArArCALC2.5
430 (Koppers, 2002).

431 Detailed methodologies are provided in SI 1, and results are reported in Tables S1 (Sr-Nd data), S2 (zircon
432 U-Pb data) and S3 (mica Ar-Ar data).

433

434 **4. Results and integration with published data**

435 **4.1 Sr/Nd bulk (Figs. 2 and SI Fig S1, Table S1)**

436 There is little significant variation in ϵ_{Nd} values from the Eocene Kuldana Formation through to the late
437 Miocene Nagri Formation, with values ranging between -7.0 to -9.2 (Fig. 2A). The exception to this overall
438 similarity is the Murree Formation at MHS, with a value of -13.8. We note that previous work for the Chinji
439 Formation records values of -3.8 to -7.7 (Chirouze et al., 2015); this difference could perhaps reflect the
440 previous use of sand compared to analysis of muds in the current research (see Jonell et al., 2018 for
441 further discussion). There are no modern-day data available for the range front. The Upper Indus has a
442 value of -10.8 at Besham (Clift et al., 2002b) located just downstream of the Kohistan arc (Fig. 1B and C)
443 and we can extrapolate that values should be more negative than this at the range front, after the river
444 has passed over the Greater and Lesser Himalaya. Values at the delta front at Thatta are -14.9 (Clift et al.,
445 2002b).

446 We carried out mixture modelling on the foreland basin material (SI Fig. S1). The mixture modelling is
447 complicated by the number of end member contributors; today sediment in the Upper Indus River
448 contains material from the Lhasa Block, Karakoram, KLIA, suture zone, and the Indian plate units of the
449 Greater-, Lesser- and Tethyan Himalayan correlatives. Overlapping signatures of some units (e.g. between
450 the Karakoram and Tethyan Himalaya, and between the KLIA and ophiolitic melange of the suture zone)
451 also adds uncertainty. We started with the premise that, from the zircon data we are confident that the
452 foreland basin contains material from the KLIA (section 4.2.) from the oldest sediments studied, namely
453 the early-Mid Eocene Kuldana Formation. That therefore forms the apex of our model, and various
454 mixture couplings are calculated with this apex and other potential end members. The modelling shows

455 that all data can be explained by a mix of Indian plate and KLIA inputs, and contribution from the
456 Karakoram and Lhasa Block is equivocal. The Murree Formation sample from MHS requires considerable
457 input from Greater Himalayan lithological correlatives.

458 The Sr-Nd compositions of the samples plot on trends that are consistent with simple mixing between
459 mafic and more evolved sources. There is some scatter in the data towards high Sr^{87}/Sr^{86} values that may
460 result from weathering or diagenesis. However, we are confident that the dominant trends reflect
461 changes in provenance, as described above.

462 ***4. 2 Detrital zircon U-Pb analyses (Fig 3, SI Figs S2 and S3, Table S2)***

463 We compile our new data from the Murree Formation at Paras north of Balakot in the HKS and at MHS (<
464 24 Ma), and from the Kamlial (18-14 Ma), Chinji (14-11 Ma) and Nagri (11-8.5 Ma) Formations in the
465 Potwar Plateau, with previously published data from the Kuldana and Murree Formation rocks at Balakot,
466 Muzaffarabad and Kotli in the HKS and at MHS (Awais et al., 2021; Ding et al., 2016a; Qasim et al., 2018)
467 and modern river data collected at the MCT-correlative (Khairabad Thrust) at the range front at Attock
468 (Alizai et al 2011, Clift et al 2022) (Fig. 1B). We keep our observations of comparisons broad and
469 conservative in nature, since different approaches to both mineral separation and data processing
470 procedures by different labs can cause variation in proportions of different populations. We begin our
471 summary at the marine to continental transition (the Kuldana Formation, section 2.2.1). We focus on the
472 40-200 Ma “arc-aged” population characteristic of the KLIA and Karakoram, and the older grains typical
473 of the Indian plate and Karakoram, with emphasis on the 1500-2300 Ma population typical of the Lesser
474 Himalayan lithological correlatives and 300-1250 Ma population typical of the Greater Himalayan
475 lithological correlatives (section 2.1.1, Table 1).

476 With the exception of the Murree Formation (which we portray separately in Fig 3B and discuss separately
477 in section 5.2), the proportions of the 40-200 Ma “arc-aged” populations remain approaching or above

478 50% throughout the Neogene to present day. There is much variation within the Eocene Kuldana
479 Formation, with nevertheless a number of samples also showing a majority of grains to be arc aged (Fig.
480 3A, Table 1). By contrast, the Murree Formation has a very low proportion of grains in the 40-200 Ma
481 range in all samples analysed from MHS, Muzaffarabad and Balakot, although not at Paras north of Balakot
482 in the HKS (Fig 1B, Fig 3B and SI Fig S2). Instead, these Murree Formation samples from MHS,
483 Muazaffarabad and Balakot have a high proportion of grains with ages typical of the Greater Himalaya.
484 In contrast to the modern-day river sample at Attock (Fig. 3C, Fig 1B), there is no 1500-2300 Ma population
485 typical of the Lesser Himalayan lithological correlatives, in any of the formations.

486 **4.3 Mica Ar-Ar (Fig 4, SI Fig. S4, Table S3)**

487 We have integrated our new data from the Murree Formation at MHS, Chinji and Nagri Formations with
488 previous data from the Murree Formation in the HKS at Paras north of Balakot (Najman et al., 2001) and
489 Kamlial Formation (Najman et al., 2003) (SI Fig. SI4). We note the following, bearing in mind that the
490 number of grains analysed for the Murree Formation at Paras north of Balakot (n=257) and the Kamlial
491 Formation (n=277) are considerably higher than for the Murree Formation at MHS, Chinji and Nagri
492 samples (n=59, 94 and 43 respectively), resulting in more confidence that the Balakot and Kamlial
493 Formation datasets more completely capture the complete spectrum of ages:

494 The youngest grain in the Murree Formation at Paras in the HKS is 24.6 +/- 0.7 Ma, the weighted mean of
495 the youngest two grains overlapping within error at two sigmas is 24.8 Ma +/- 1.4 Ma and the youngest
496 peak population is 37 Ma. Pre-Cenozoic ages extend to >1500 Ma. Further south, the youngest grain in
497 the Murree Formation at MHS is 23.7 +/- 0.1 Ma, which also forms one of the two youngest grains
498 overlapping within error at 2 sigmas (weighted mean 23.85 +/- 0.12 Ma. The youngest peak population is
499 24-28 Ma. Pre-Cenozoic ages extend to ca 450 Ma. The youngest grain for the Kamlial Formation is 14.5
500 +/- 0.7 Ma, and weighted mean of the youngest two grains within error at 2 sigmas is 15.00 +/- 1.10a. The

501 youngest peak population is 18 Ma and Pre-Cenozoic ages extend to ca 450 Ma. The lowest Chinji
502 Formation sample (CP96-7A, Najman et al 2003, dated at 13.9 Ma) has a youngest grain at 14.1 +/- 0.7 Ma
503 and this also forms one of the two youngest grains within error at 2 sigmas (weighted mean 14.43 +/- 0.81
504 Ma). Pre-Cenozoic grains extend to 400 Ma. Our new sample from the Chinji Formation has a youngest
505 grain of 16.74 +/- 0.1 Ma, the weighted mean of the two youngest grains overlapping within error at 2
506 sigmas is 25.95 +/- 0.10 Ma, the youngest peak population is 28-29 Ma, and Pre-Cenozoic ages extend to
507 ca 450 Ma. The youngest grain in the Nagri Formation is 17.9 +/- 0.14 Ma, the weighted mean of the two
508 youngest two grains overlapping within error at 2 sigmas is 19.69 +/- 0.12 Ma the youngest peak
509 population is 21 Ma and Pre-Cenozoic ages extend to ca 200 Ma.

510 Rapid exhumation determined from short lag times was determined for the Kamliyal and lowest Chinji
511 Formation, between 16-14 Ma (Najman et al, 2003) (Fig. 4). Lack of independent depositional age
512 constraints precludes calculation of lag times for the newly analysed Murree, Chinji and Nagri Formation
513 samples. Up section from the Kamliyal Formation, there is no evidence of grain ages approaching
514 depositional age, until the modern river sample at Thatta, although the number of grains analysed is
515 relatively small.

516

517 **5. Interpretations of the evolution of the Lower Indus drainage**

518 ***5.1. When did the Punjabi tributary system join the paleo-Indus trunk river?***

519 As outlined in our rationale and approach (section 3.1), we determine when the Punjab tributary system
520 joined the main trunk river, by a comparison of provenance data from upstream and downstream of the
521 present day confluence, leveraging the fact that unlike the palaeo-Indus trunk River, the tributaries drain
522 only the Indian plate terranes (Fig 1C), and thus have a different provenance signature (section 3.1, Figs 2
523 and 3A inset).

524 As schematically presented in Fig 5, the following evidence should be met, at the time the tributary system
525 joined the trunk Indus:

526 (1) Prior to the time that the Punjab tributary system joined the Indus catchment, the proportion of
527 Indian plate detritus delivered to the Indus River should be comparable at the range front and at
528 the river mouth, i.e. upstream and downstream of where the Punjab tributary system now joins
529 the modern Indus.

530 (2) After the time when the Punjab tributary system joined the Indus River, the proportion of Indian
531 plate material in the Indus River downstream of the confluence with the Punjab Rivers should a)
532 increase relative to the downstream's previous pre-reorganisation proportion and b) be greater
533 than coeval sediments upstream. However, the proportion of Indian plate material in the
534 upstream should remain constant, pre and post the proposed drainage reorganisation.

535 For the above predictions to be explored, Indian plate, versus Karakoram, versus KLIA must be
536 differentiable in the foreland basin detritus. Table 1 provides the typical zircon U-Pb and ϵ_{Nd} signatures of
537 these units, alongside a summary of equivalent data from the peripheral foreland basin, and downstream
538 in both the Sulaiman-Kirthar region and Indus Fan. Figs 2 and 3A inset show the difference between the
539 modern trunk Indus which drains the Asian plate, arc and Indian plate, versus the modern Punjabi
540 tributary system which drains, for the most part, only the Indian plate.

541 For the interpretations made from this upstream-downstream comparison to be valid, the rocks at the
542 evaluated locations must be the products of the palaeo-Indus. Whilst all three repositories studied, the
543 peripheral foreland basin, the Lower Indus Axial Basin, and the Indus Fan, show evidence of derivation
544 from at least as far north as the KLIA since Eocene times, we acknowledge evidence for input from north
545 of the Shyok suture zone can be equivocal (see sections 2.3.1, 2.3.3 and 2.3.4).

546 Below, we summarise the salient points regarding the upstream and downstream repositories that are
547 relevant to the characteristics required to document the timing of conjoinment of the Punjabi tributary
548 system with the trunk Indus River as described above. We discuss the Murree Formation which is
549 anomalous at MHS, Muzaffarabad and Balakot, but not at Paras, separately in section 5.2.

550 *5.1.1. Comparison of the upstream peripheral foreland basin material with the downstream repositories*
551 *in terms of Sr-Nd data*

552 Our data from the upstream (peripheral foreland basin) show that values have remained broadly constant
553 from the start of our studied record in the early-Mid Eocene Kuldana Formation (Fig. 2A), until the late
554 Miocene Nagri Formation, when values become a little more negative. $\epsilon\text{Nd}(0)$ values in the downstream
555 repositories are similar to the upstream in the early Eocene. However, values in the downstream
556 repositories become more negative compared to the upstream, by Mid Eocene in the Indus Fan and
557 around the Eocene-Oligocene boundary in the Lower Indus Axial Basin (Fig. 2B). This shift indicates a
558 greater input of material from the Indian plate Himalayan terrane at this time.

559 From the more negative $\epsilon\text{Nd}(0)$ values recorded below compared to above the confluence throughout
560 the Neogene, we interpret that the Punjabi tributary system has drained into the palaeo-Indus throughout
561 the Neogene, and that the present drainage configuration was therefore established during the
562 Paleogene.

563 The consistency of ϵNd values from the Eocene to the Neogene in the upstream repository, in contrast to
564 the shift to more negative values in the downstream repositories should reflect the time when the Punjabi
565 tributary system joined the trunk Indus River. However, the difference in the time of the downstream
566 shift, at the Eocene-Oligocene time in the Lower Indus axial basin delta deposits and in the mid Eocene in
567 the Indus Fan indicates that more research is required before we can pinpoint the exact time that the

568 tributary system joined the trunk Indus. Nevertheless, with available data we can conclude that the
569 tributaries joined in the trunk Indus at or before the start of the Oligocene (Fig 5).

570 *5.1.2. Comparison of the upstream peripheral foreland basin material with the downstream repositories*
571 *in terms of detrital zircon U-Pb data*

572 Although intraformational variability, lack of data from the Oligocene in the peripheral foreland basin, and
573 lack of data from the Eocene in the downstream repositories limits the comparison, the data are
574 consistent with the interpretations determined the Sm-Nd data, that the Punjabi tributary system joined
575 the trunk River Indus by Oligocene times (section 5.1.1, Fig 5): The proportion of 40-200 Ma arc-aged
576 grains remains high throughout the Miocene in the peripheral foreland basin, and these values are higher
577 compared to Oligocene-Pliocene values in both downstream repositories (Fig. 3A and C, Table 1). Data
578 from the Eocene peripheral foreland basin is highly variable. However, at least some samples have a
579 proportion of arc-aged grains similar to the proportions of the Neogene peripheral foreland basin,
580 consistent with the pattern shown in the Sm-Nd data.

581 SI Figure SI 3 illustrates the river's evolution well, particularly by comparison to the Lower Indus Axial
582 Basin. Downstream samples have a greater affinity to Indian plate rocks and the modern Indus at its mouth
583 at Thatta, compared to the upstream peripheral foreland basin rocks which have greater affinity to the
584 arc and Asian plate, and the modern day Indus at the range front at Attock.

585 The variation in zircon U-Pb age spectra, and also in ϵ_{Nd} values, between the onshore and offshore
586 downstream palaeo-Indus, and between the Eastern and Western Indus Fan (SI Figs. S1 and S3) is
587 intriguing. It could be the result of a number of factors, for example differences in sample preparation
588 procedures between operators, downstream influence of hydraulics, or additional material contributing
589 downstream, for example.

590 **5.2. Interpretation of the Murree Formation**

591 Compared to the other peripheral foreland basin sediments sampled, the Zircon U-Pb data show
592 significantly higher proportions of old grains in the Murree Formation at MHS, Kotli, and at Balakot and
593 Muzaffarabad in the HKS, but not at Paras north of Balakot (Fig. 3B, SI Fig S2, Table 1). Where
594 accompanying Sr-Nd data are available (MHS and Paras only), there is a corresponding change to more
595 negative ϵ_{Nd} values at MHS (Fig 2A), mirroring the change noted in the zircon data. This signature indicates
596 a higher proportion of material derived from the Indian plate (see also SI Fig S3). These deposits may be
597 interpreted as the palaeo-Jhelum Punjab tributary, which has a similar zircon U-Pb spectrum to the
598 Murree Formation (Fig. 3B), and a drainage basin consisting predominantly of the Indian plate (Fig 1C).
599 The spatial distribution of our analysed samples is consistent with this interpretation: a Himalayan-derived
600 palaeo-Jhelum type signature is prevalent in Murree Formation samples at Muzaffarabad (Fig 1B) located
601 on the modern day Jhelum River, at MHS downstream and ca 10 miles to the west of the modern Jhelum
602 River, and at Kotli, downstream and 20 kms east of the modern Jhelum river. It is also prevalent at Balakot,
603 ca 15 miles upstream of the modern Jhelum River, which we suggest could have been in the flood plain of
604 the palaeo-Jhelum. 5 miles further north still, near Paras, the signature is more arc-like and in this palaeo-
605 drainage scenario, we propose lies outwith the floodplain of the palaeo-Jhelum. We note that at
606 Muzaffarabad only, through which the modern Jhelum River flows, a palaeo-Jhelum type signature is also
607 recorded, in some samples, in the underlying Eocene Kuldana Formation. This may reflect the early
608 initiation of this river, insufficiently large in its early evolution to affect the downstream.

609 Alternatively, the anomalous signature from the Murree Formation compared to the rest of the Cenozoic
610 sediments in the peripheral foreland basin may reflect increased input from the Himalaya attributable to
611 a pulse of exhumation recorded in the Himalaya in the early Miocene (section 2.1.2). A coeval change to
612 greater input from the Indian plate is also recorded in the Indus Fan (Feng et al., 2021) and Kirthar Ranges

613 (Zhuang et al., 2015), supporting this interpretation. Further analyses from Murree Formation samples
614 distal to the Jhelum River should distinguish between these two alternative hypotheses.

615 The difference in Murree signature compared to the rest of the foreland basin cannot be ascribed to bias
616 associated with grain size variation since the difference is reflected in both bulk rock Sr-Nd and zircon
617 proxies. Nor is there any reason to consider that a potential difference in the degree of diagenesis caused
618 the difference, since zircons are largely unaffected by this process.

619 ***5.3 What caused the change in the geochemical signature of the Indus Fan at 5-6 Ma?***

620 The more recently proposed alternatives to drainage reorganisation (Clift and Blusztajn, 2005) to explain
621 the geochemical shift in the Indus Fan at 5-6 Ma all involve tectonic explanations, namely variations in
622 exhumation of the hinterland terranes, although the extent to which increased exhumation of the Lesser
623 Himalaya versus Greater Himalaya versus Karakoram is responsible, is debated (Chirouze et al 2015, Clift
624 et al 2019, Zhou et al. 2022). Changes in monsoonal intensification are not thought to have been a major
625 influence (Clift et al., 2019, Zhou et al 2022).

626 To what extent do our data support a tectonic explanation? We focus on the peripheral foreland, which
627 should provide the most tectonically- influenced archive, above any downstream influence from the
628 Punjabi tributary system. We compare our data from the Nagri Formation (11-8.5 Ma; Table 1) to modern
629 day Indus data at the range front, this time period encompassing the 5-6 Ma date over which the
630 geochemical shift in the Indus Fan occurred.

631 The average ϵ_{Nd} value of the two samples from the Nagri Formation is -9.65. No data are available for the
632 modern Indus at the range front. The spatially closest sample is from Besham, just south of the MCT (Fig.
633 1C). This sample has a value of -10.7, and we would expect a more negative value by the time the river
634 had crossed to the range front, having flowed over more of the Greater Himalaya and most negative Lesser

635 Himalaya (Table 1). Thus, the shift to more negative ϵNd values between the Nagri Formation and the
636 estimated value for the range front in modern times I shows that variation in upland tectonics over this
637 time period could have resulted in the shift to more negative ϵNd values seen in the Indus Fan over this
638 time period.

639 Assignment of zircon U-Pb age populations to distinct provenances is challenging with respect to overlap
640 of the older Karakoram and Indian plate grains. Nevertheless, the 1500-2300 Ma population is typical of
641 the Lesser Himalaya. This population makes up 3% of the Nagri sample. There is no sample from the
642 modern Indus River at the range front. However, there is a sample from upstream at Attock (Fig. 1C). This
643 sample has an 11% contribution from the 1500-2300 Ma population, and we would predict a higher
644 proportion of that population after the river has flowed over a greater proportion of Indian plate material.
645 The shift to a higher proportion of zircons with ages indicative of Lesser Himalayan input between the
646 Nagri Formation and the modern day (Fig 3), therefore supports our observations from the Sm-Nd data,
647 that upstream variations in tectonics could have resulted in the geochemical shift in the Indus Fan.

648 There are no modern river mica $^{40}\text{Ar}/^{39}\text{Ar}$ data from the range front. Modern river $^{40}\text{Ar}/^{39}\text{Ar}$ mica data
649 from the trunk Indus at its river mouth at Thatta shows Plio-Pleistocene grains (1-5 Ma) indicative of rapid
650 exhumation (Clift et al., 2004). Recording of these young grains in the trunk river but not in the tributaries
651 draining only the Indian plate or Indian plate plus Hindu Kush (Clift et al., 2004; Najman et al., 2009;
652 Zhuang et al., 2018) is consistent with the viewpoints of, for example, Chirouze et al. (2015) and Clift et
653 al. (2022) Clift et al (2022), that the Karakoram and/or the Nanga Parbat syntaxis supplied this young
654 material. Lag times determined from mica data from the Neogene peripheral foreland basin sedimentary
655 rocks show no clear indication of rapid exhumation of the micas' source region after 14-16 Ma (Fig. 4)
656 although n values are small and therefore populations may have been missed. Therefore a period of rapid
657 exhumation occurred sometime between Nagri Formation times and present day, consistent with the

658 view that changing exhumation in the hinterland was responsible for the geochemical shift at 5 Ma in the
659 Indus Fan.

660

661 **6. Conclusions**

662 When the lower Indus River broadly attained its current drainage configuration, in particular when the
663 Punjab tributary system joined the main trunk river, is undocumented. Comparison of ϵ_{Nd} bulk rock data
664 and detrital zircon U-Pb data from Cenozoic paleo-Indus sedimentary rocks both upstream and
665 downstream of the confluence of the Indus with the Punjab tributary system shows that throughout the
666 Neogene, greater proportions of Indian plate material are recorded in the downstream compared to the
667 upstream repositories. We therefore conclude that the Punjabi tributary system, which transports
668 predominantly Indian plate detritus, had joined the trunk Indus River prior to the Neogene.

669 Whilst provenance indicators show that the proportion of Indian plate material remains constant from
670 Eocene to Neogene in the palaeo-Indus repository upstream of the confluence, the proportion of Indian
671 plate material increases in the downstream repositories, at the Eocene-Oligocene boundary in the palaeo-
672 delta, and in the mid Eocene in the Indus Fan. More research is required to understand the reasons for
673 this discrepancy in timing of the shift in the downstream repositories, but nevertheless we can conclude
674 that the Punjabi tributary system joined the palaeo-Indus trunk river at or before the start of the
675 Oligocene.

676 **Acknowledgements**

677 G.Z. acknowledges financial support of a Marie Curie Postdoctoral Fellowship under the Initial Training
678 Network iTECC funded by the EU REA under the FP7 implementation of the Marie Curie Action, under
679 grant agreement # 316966. Imran Khan helped with the sample collection. We thank Joe DiPietro for his

680 detailed insights into the geology of Pakistan and Peng Zhou for sharing the geology map shown in Fig.
681 1C. This paper benefitted from input from two anonymous reviewers.

682 **Figures**

683 **Figure 1: A:** Map showing modern drainage of the Indus River with the Punjab tributary system, and the
684 Indus Fan (black dotted line). Also shown are the onshore lower Indus (Kirthar and Sulaiman) and offshore
685 IODP, ODP and DSDP locations of previously published data (Roddaz et al. 2011, Zhuang et al 2015, Clift
686 et al. (2001), Clift and Blusztajn (2005), Clift et al. (2019) and Feng et al (2021). with which we compare
687 our new upstream data. Black rectangle shows the location of Fig 1B. **B:** locations of new data (this study)
688 and various towns and published sample sites discussed in text. MHS = Murree Hill Station; HKS = Hazara-
689 Kashmir Syntaxis. Samples with prefix KG96 or 99 are from Paras in the HKS, prefix MU96 are from Murree
690 Hill Station, and prefix KMSr, CHSr, and NgSr are from Chinji village area. **C:** Drainage superimposed on
691 regional geology (from Clift et al., 2019). ISZ = Indus Suture zone, SSZ = Shyok Suture Zone, MCT =Main
692 Central Thrust, MBT=Main Boundary Thrust, MFT = Main Frontal Thrust.

693 **Figure 2:** ϵ_{Nd} values from the upstream peripheral foreland basin in Pakistan (A), and downstream Lower
694 Indus axial basin and Indus Fan (B) through time. In A, numbers adjacent to squares refer to sample
695 numbers to left. Asterisks indicate new data. Open squares for modern Indus River at Besham and Thatta;
696 hexagons are from the modern Punjabi tributaries (data from Clift et al 2002, Alizai et al. 2011, Chirouze
697 et al 2015). In B: diamonds – data from the Sulaiman and Kirthar regions of the Lower Indus axial basin
698 (Roddaz et al, 2011, Zhuang et al 2015); circles – data from the Indus Fan from Clift et al. (2001), Clift and
699 Blusztajn (2005), Clift et al. (2019), Zhou et al 2021, and Feng et al (2021). Questions marks next to three
700 Mid Eocene samples represent uncertainties in age for those samples, as depicted in the original
701 publication of Clift et al (2001). HKS = Hazara-Kashmir Syntaxis; MHS = Murree Hill Station. MDA =

702 maximum depositional age as determined from detrital grain ages (sections 2.2.1, 4.2 and 4.3). Grey
703 horizontal shading between plots A and B denote roughly equivalent time periods.

704 **Figure 3:** Detrital zircon U-Pb data shown as cumulative age distribution plots. **A:** Pakistan peripheral
705 foreland basin data excluding Murree Formation data except our new data. Kuldana Fm samples are Early-
706 Mid Eocene, Murree Formation samples are Early Miocene, Kamlial Fm is Early-Mid Miocene, Chinji Fm is
707 Mid Miocene, Nagri Fm is Late Miocene (Table 1). A inset: modern river data comparing the Indus at the
708 range front at Attock, with rivers of the Punjabi tributary system. **B:** all Murree Formation data, both new
709 and published, with comparison to the Jhelum modern river data. Murree Formation is Early Miocene. **C:**
710 comparison between peripheral foreland basin data and downstream Lower Indus axial basin data and
711 (inset) Indus Fan data. Eocene peripheral foreland basin data are omitted from the figure as there are no
712 comparative data from the downstream repositories. HKS = Hazara-Kashmir Syntaxis, MHS = Murree Hill
713 Station. All new data are asterisked. Samples with superscripts are published data, as follows: ¹Ding et al
714 (2016), ²Qasim et al (2018), ³Awais et al (2021), ⁴Zhuang et al (2015), ⁵Clift et al (2002), ⁶Clift et al (2004),
715 ⁷Clift et al (2019), ⁸ (Zhou et al, 2021), ⁹Feng et al (2021), ¹⁰Alizai et al (2011).

716 **Figure 4:** Ar-Ar mica data plotted against depositional age for new (asterisked) and published samples.
717 Note: ¹Published data from ¹Clift et al (2004) for modern Indus River data at Thatta, ²published data from
718 Najman et al (2003) for lower Chinji Formation, ³published data from Najman et al (2003) for the Kamlial
719 Fm, and ⁴published data from Najman et al (2001) for the Murree Formation at HKS. Apart from the lower
720 Chinji Formation sample, Chinji and Nagri Formation samples are not tied to the magnetostratigraphically
721 dated section (Johnson et al 1985, section 2.2.1), and therefore the depositional age range of these
722 samples is shown by the grey bars. Note that Murree Formation samples are plotted on the y axis at the
723 age of their MDAs.

724 **Figure 5:** schematic figure showing expected and actual changes in provenance characteristics of
725 sedimentary archives upstream and downstream of the confluence, at the time when the Punjabi
726 tributary system joins the palaeo-Indus trunk river, superimposed on the modern geology. More detail on
727 analytical values summarised in this figure can be found in Table 1. Abbreviations: Av – average, Z – zircon,
728 Eoc – Eocene, KLIA – Kohistan-Ladakh Island Arc.

729 **Tables**

730 **Table 1:** comparison of provenance data from the peripheral foreland basin with those from the Lower
731 Indus Axial Basin and Indus Fan. Source region signatures also provided. Note that three “Mid Eocene”
732 data points from the Indus Fan are omitted as the age was noted as questionable in the original publication
733 of Clift et al (2001).

734 *Compiled source region data from Ahmad et al. (2000); Bignold and Treloar (2003); Clift et al. (2019);
735 DeCelles et al. (2004); DeCelles et al. (2016b); Deniel et al. (1987); Gehrels et al. (2011); Ji et al. (2009);
736 Khan et al. (1997); Khan et al. (2004); Khan et al. (2009); Mahéo et al. (2009); Miller et al. (1999); Najman
737 (2006); Pan et al. (2014); Richards et al. (2005); Robinson et al. (2001); Whittington et al. (1999); Zhang et
738 al. (2004); Zhu et al. (2012); Zhuang et al. (2018), and additional references as listed in Fig S3b.

739 **Supplementary information**

740 **Text S1:** detailed analytical methodologies and sample information

741 **Table S1:** Sr-Nd bulk mudstone data.

742 **Table S2:** detrital zircon U-Pb data.

743 **Table S3:** White mica Ar-Ar analyses.

744 **Figure SI 1:** Sr-Nd mixture modelling of end members (A), and new (asterisk) and previously published
745 bulk rock data from the peripheral and axial foreland basins in Pakistan, and the Indus Fan, plotted on to
746 a sub-region of Fig A (B). Downstream published data: ¹Roddaz et al (2011), ²Zhuang et al (2015), ³ Clift et
747 al. (2001), Clift and Blusztajn (2005), Clift et al. (2019), Zhou et al 2021, and ⁴Feng et al (2021). Means and
748 one standard errors are calculated from compiled data points (same symbols with smaller sizes and
749 transparency) (Zhuang et al., 2015 and references therein). HKS = Hazara-Kashmir Syntaxis; MHS = Murree
750 Hill Station.

751 **Figure SI 2a and b** – KDEs for new and published zircon U-Pb data, at two different scales, 0-400 Ma and
752 0-500 Ma. New data are shown by asterisks. Published data: ¹Alizai et al (2011) , ²Clift et al (2022), ³Ding
753 et al (2016), ⁴Qasim et al (2018), ⁵Awais et al (2021).

754 **Figure SI 3:** MDS plot showing zircon U-Pb data for our new samples (asterisk) from the peripheral
755 foreland basin (this study, red crosses), compared to downstream published data from the Kirthar and
756 Sulaiman ranges (purple crosses, sample prefixes SR and Z, data from Roddaz et al, 2011 and Zhuang et
757 al. 2015) and Indus Fan (data from Clift et al. 2019, Zhou et al 2021, Feng et al, 2022), grey crosses for
758 Western Fan, black crosses for Eastern Fan. Also shown are published data from the modern Indus River
759 upstream at Attock and downstream at Thatta (blue crosses, see Fig 1 for location, from Alizai et al 2011
760 and Clift et al. 2022), and end member source signatures (black hexagons for Asian plate and arc , KLA =
761 Kohistan-Ladakh Island arc, KK = Karakoram, HK = Hindu Kush, and red squares for Indian plate
762 TH=Tethyan Himalaya, GH = Greater Himalaya, LH = Lesser Himalaya. References for compiled end
763 member data are listed in Fig 3I 3b.

764 **Fig SI 4a and b** – KDEs for new and published mica Ar-Ar data from the peripheral foreland basin, Pakistan,
765 at two different scales, 0-500 Ma and 0-100 Ma. New data are asterisk, published data is referenced on
766 the plot.

768 **References**

- 769 Abbasi, I.A., Friend, P.F., 1989. Uplift and evolution of the Himalayan orogenic belts, as recorded in the
770 foredeep molasse sediments. *Zeitschrift für Geomorphologie NF Supplementband 76*, 75-88.
- 771 Ahmad, T., Harris, N., Bickle, M., Chapman, H., Bunbury, J., Prince, C., 2000. Isotopic constraints on the
772 structural relationships between the lesser Himalayan series and the high Himalayan crystalline series,
773 Garhwal Himalaya. *Geological Society of America Bulletin* 112, 467-477.
- 774 Aitchison, J.C., Ali, J.R., Davis, A.M., 2007. When and where did India and Asia collide? *Journal of*
775 *Geophysical Research: Solid Earth (1978–2012)* 112.
- 776 Alizai, A., Carter, A., Clift, P.D., VanLaningham, S., Williams, J.C., Kumar, R., 2011. Sediment provenance,
777 reworking and transport processes in the Indus River by U–Pb dating of detrital zircon grains. *Global and*
778 *Planetary Change* 76, 33-55.
- 779 Andò, S., Aharonovich, S., Hahn, A., George, S., Clift, P., Garzanti, E., 2020. Integrating heavy-mineral,
780 geochemical and biomarker analyses of Plio-Pleistocene sandy and silty turbidites: a novel approach for
781 provenance studies (Indus Fan, IODP Expedition 355). *Geological Magazine* 157, 929-938.
- 782 Argles, T., Foster, G., Whittington, A., Harris, N., George, M., 2003. Isotope studies reveal a complete
783 Himalayan section in the Nanga Parbat syntaxis. *Geology* 31, 1109-1112.
- 784 Awais, M., Qasim, M., Tanoli, J.I., Ding, L., Sattar, M., Baig, M.S., Pervaiz, S., 2021. Detrital Zircon
785 Provenance of the Cenozoic Sequence, Kotli, Northwestern Himalaya, Pakistan; Implications for India–
786 Asia Collision. *Minerals* 11, 1399.
- 787 Baig, M.S., Munir, M.-u.-H., 2007. Foraminiferal biostratigraphy of Yadgar area, Muzaffarabad Azad
788 Kashmir, Pakistan. *Journal of Himalayan Earth Sciences* 40.
- 789 Bernet, M., van der Beek, P., Pik, R., Huyghe, P., Mugnier, J.L., Labrin, E., Szulc, A., 2006. Miocene to Recent
790 exhumation of the central Himalaya determined from combined detrital zircon fission - track and U/Pb
791 analysis of Siwalik sediments, western Nepal. *Basin Research* 18, 393-412.
- 792 Bhattacharya, G., Robinson, D.M., Wielicki, M.M., 2020. Detrital zircon provenance of the Indus Group,
793 Ladakh, NW India: Implications for the timing of the India-Asia collision and other syn-orogenic processes.
794 *GSA Bulletin*.
- 795 Bignold, S., Treloar, P., 2003. Northward subduction of the Indian Plate beneath the Kohistan island arc,
796 Pakistan Himalaya: new evidence from isotopic data. *Journal of the Geological Society* 160, 377-384.
- 797 Bossart, P., Ottiger, R., 1989. Rocks of the Murree Formation in northern Pakistan: indicators of a
798 descending foreland basin of late Paleocene to middle Eocene age. *Eclogae Geologicae Helvetiae* 82, 133-
799 165.
- 800 Bouilhol, P., Jagoutz, O., Hanchar, J.M., Dudas, F.O., 2013. Dating the India–Eurasia collision through arc
801 magmatic records. *Earth and Planetary Science Letters* 366, 163-175.
- 802 Burbank, D.W., Beck, R.A., Mulder, T., 1996. The Himalayan foreland basin. *The Tectonic Evolution of Asia*,
803 Cambridge University Press, eds An Yin and Mark Harrison., 149-190.
- 804 Burbank, D.W., Tahirkheli, R.K., 1985. The magnetostratigraphy, fission-track dating, and stratigraphic
805 evolution of the Peshawar intermontane basin, northern Pakistan. *Geological Society of America Bulletin*
806 96, 539-552.
- 807 Cervený, P., Johnson, N., 1989. Tectonic and geomorphic implications of Siwalik Group. *Tectonics of the*
808 *Western Himalayas* 232, 129.
- 809 Cervený, P., Johnson, N., Tahirkheli, R., Bonis, N., 1989. Tectonic and geomorphic implications of Siwalik
810 Group heavy minerals, *Tectonics of western Himalayas*. *Geological Society of America Special Paper* 232,
811 129-136.

812 Chirouze, F., Huyghe, P., Chauvel, C., van der Beek, P., Bernet, M., Mugnier, J.-L., 2015. Stable Drainage
813 Pattern and Variable Exhumation in the Western Himalaya since the Middle Miocene. *The Journal of*
814 *Geology* 123, 1-20.

815 Clift, P., Shimizu, N., Layne, G., Blusztajn, J., Gaedicke, C., Schlüter, H.-U., Clark, M., Amjad, S., 2001a.
816 Development of the Indus Fan and its significance for the erosional history of the Western Himalaya and
817 Karakoram. *Geological Society of America Bulletin* 113, 1039-1051.

818 Clift, P.D., Blusztajn, J., 2005. Reorganization of the western Himalayan river system after five million years
819 ago. *Nature* 438, 1001-1003.

820 Clift, P.D., Campbell, I.H., Pringle, M.S., Carter, A., Zhang, X., Hodges, K.V., Khan, A.A., Allen, C.M., 2004.
821 Thermochronology of the modern Indus River bedload: New insight into the controls on the marine
822 stratigraphic record. *Tectonics* 23.

823 Clift, P.D., Carter, A., Jonell, T.N., 2014. U–Pb dating of detrital zircon grains in the Paleocene Stumpata
824 Formation, Tethyan Himalaya, Zaskar, India. *Journal of Asian Earth Sciences* 82, 80-89.

825 Clift, P.D., Carter, A., Krol, M., Kirby, E., 2002a. Constraints on India-Eurasia collision in the Arabian Sea
826 region taken from the Indus Group, Ladakh Himalaya, India. *The tectonic and climatic evolution of the*
827 *Arabian Sea region*, 97.

828 Clift, P.D., Lee, J.I., Hildebrand, P., Shimizu, N., Layne, G.D., Blusztajn, J., Blum, J.D., Garzanti, E., Khan, A.A.,
829 2002b. Nd and Pb isotope variability in the Indus River System: implications for sediment provenance and
830 crustal heterogeneity in the Western Himalaya. *Earth and Planetary Science Letters* 200, 91-106.

831 Clift, P.D., Mark, C., Alizai, A., Khan, H., Jan, M.Q., 2022. Detrital U–Pb rutile and zircon data show Indus
832 River sediment dominantly eroded from East Karakoram, not Nanga Parbat. *Earth and Planetary Science*
833 *Letters* 600, 117873.

834 Clift, P.D., Shimizu, N., Layne, G., Blusztajn, J., 2001b. Tracing patterns of unroofing in the Early Himalaya
835 through microprobe Pb isotope analysis of detrital K-feldspars in the Indus Molasse, India. *Earth and*
836 *Planetary Science Letters* 188, 475-491.

837 Clift, P.D., Zhou, P., Stockli, D.F., Blusztajn, J., 2019. Regional Pliocene exhumation of the Lesser Himalaya
838 in the Indus drainage. *Solid Earth* 10, 647-661.

839 Coward, M., Jan, Q., Rex, D., Tarney, J., Thirwall, M., 1984. Geology of the south Central Karakoram and
840 Kohistan, The international Karakoram project. *International conference*, pp. 71-85.

841 DeCelles, P., Gehrels, G., Najman, Y., Martin, A., Carter, A., Garzanti, E., 2004. Detrital geochronology and
842 geochemistry of Cretaceous–Early Miocene strata of Nepal: implications for timing and diachroneity of
843 initial Himalayan orogenesis. *Earth and Planetary Science Letters* 227, 313-330.

844 DeCelles, P.G., Carrapa, B., Gehrels, G.E., Chakraborty, T., Ghosh, P., 2016a. Along - strike continuity of
845 structure, stratigraphy, and kinematic history in the Himalayan thrust belt: The view from Northeastern
846 India. *Tectonics* 35, 2995-3027.

847 DeCelles, P.G., Castañeda, I.S., Carrapa, B., Liu, J., Quade, J., Leary, R., Zhang, L., 2016b. Oligocene -
848 Miocene Great Lakes in the India - Asia Collision Zone. *Basin Research*.

849 DeCelles, P.G., Gehrels, G.E., Quade, J., Ojha, T., 1998. Eocene - early Miocene foreland basin
850 development and the history of Himalayan thrusting, western and central Nepal. *Tectonics* 17, 741-765.

851 Deniel, C., Vidal, P., Fernandez, A., Le Fort, P., Peucat, J.-J., 1987. Isotopic study of the Manaslu granite
852 (Himalaya, Nepal): inferences on the age and source of Himalayan leucogranites. *Contributions to*
853 *Mineralogy and Petrology* 96, 78-92.

854 Ding, H., Zhang, Z., Dong, X., Tian, Z., Xiang, H., Mu, H., Gou, Z., Shui, X., Li, W., Mao, L., 2016a. Early Eocene
855 (c. 50 Ma) collision of the Indian and Asian continents: Constraints from the North Himalayan
856 metamorphic rocks, southeastern Tibet. *Earth and Planetary Science Letters* 435, 64-73.

857 Ding, L., Qasim, M., Jadoon, I.A., Khan, M.A., Xu, Q., Cai, F., Wang, H., Baral, U., Yue, Y., 2016b. The India–
858 Asia collision in north Pakistan: Insight from the U–Pb detrital zircon provenance of Cenozoic foreland
859 basin. *Earth and Planetary Science Letters* 455, 49-61.

860 DiPietro, J., Hussain, A., Ahmad, I., Khan, M.A., 2000. The main mantle thrust in Pakistan: its character and
861 extent. Geological Society, London, Special Publications 170, 375-393.

862 DiPietro, J.A., Ahmad, I., Hussain, A., 2008. Cenozoic kinematic history of the Kohistan fault in the Pakistan
863 Himalaya. Geological Society of America Bulletin 120, 1428-1440.

864 DiPietro, J.A., Pogue, K.R., 2004. Tectonostratigraphic subdivisions of the Himalaya: A view from the west.
865 Tectonics 23.

866 DiPietro, J.A., Pullen, A., Krol, M.A., 2021. Geologic history and thermal evolution in the hinterland region,
867 western Himalaya, Pakistan. Earth-Science Reviews 223, 103817.

868 Dunlap, W.J., Weinberg, R.F., Searle, M.P., 1998. Karakoram fault zone rocks cool in two phases. Journal
869 of the Geological Society 155, 903-912.

870 Exnicios, E.M., Carter, A., Najman, Y., Clift, P.D., 2022. Late Miocene unroofing of the Inner Lesser
871 Himalaya recorded in the NW Himalaya foreland basin. Basin Research 34, 1894-1916.

872 Feng, H., Lu, H., Carrapa, B., Zhang, H., Chen, J., Wang, Y., Clift, P.D., 2021. Erosion of the Himalaya-
873 Karakoram recorded by Indus Fan deposits since the Oligocene. Geology.

874 Fraser, J.E., Searle, M.P., Parrish, R.R., Noble, S.R., 2001. Chronology of deformation, metamorphism, and
875 magmatism in the southern Karakoram Mountains. Geological Society of America Bulletin 113, 1443-1455.

876 Gaetani, M., Garzanti, E., 1991. Multicyclic history of the Northern India continental margin (Northwestern
877 Himalaya)(1). AAPG Bulletin 75, 1427-1446.

878 Gaetani, M., Jadoul, F., Erba, E., Garzanti, E., 1993. Jurassic and Cretaceous orogenic events in the North
879 Karakoram: age constraints from sedimentary rocks. Geological Society, London, Special Publications 74,
880 39-52.

881 Garzanti, E., Andò, S., 2019. Heavy minerals for junior woodchucks. Minerals 9, 148.

882 Garzanti, E., Andò, S., Vezzoli, G., 2009. Grain-size dependence of sediment composition and
883 environmental bias in provenance studies. Earth and Planetary Science Letters 277, 422-432.

884 Garzanti, E., Andò, S., Vezzoli, G., 2020. Provenance of cenozoic indus fan sediments (IODP Sites U1456
885 and U1457). Journal of Sedimentary Research 90, 1114-1127.

886 Gehrels, G., Kapp, P., DeCelles, P., Pullen, A., Blakey, R., Weislogel, A., Ding, L., Guynn, J., Martin, A.,
887 McQuarrie, N., 2011. Detrital zircon geochronology of pre - Tertiary strata in the Tibetan - Himalayan
888 orogen. Tectonics 30.

889 Henderson, A.L., Najman, Y., Parrish, R., BouDagher - Fadel, M., Barford, D., Garzanti, E., Andò, S., 2010.
890 Geology of the Cenozoic Indus Basin sedimentary rocks: Paleoenvironmental interpretation of
891 sedimentation from the western Himalaya during the early phases of India - Eurasia collision. Tectonics
892 29.

893 Henderson, A.L., Najman, Y., Parrish, R., Mark, D.F., Foster, G.L., 2011. Constraints to the timing of India-
894 Eurasia collision; a re-evaluation of evidence from the Indus Basin sedimentary rocks of the Indus-
895 Tsangpo Suture Zone, Ladakh, India. Earth-Science Reviews 106, 265-292.

896 Hildebrand, P., Noble, S., Searle, M., Parrish, R., 1998. Tectonic significance of 24 Ma crustal melting in the
897 eastern Hindu Kush, Pakistan. Geology 26, 871-874.

898 Hildebrand, P., Noble, S., Searle, M., Waters, D., Parrish, R., 2001. Old origin for an active mountain range:
899 Geology and geochronology of the eastern Hindu Kush, Pakistan. Geological Society of America Bulletin
900 113, 625-639.

901 Hodges, K.V., 2000. Tectonics of the Himalaya and southern Tibet from two perspectives. Geological
902 Society of America Bulletin 112, 324-350.

903 Hu, X., Garzanti, E., Wang, J., Huang, W., An, W., Webb, A., 2016. The timing of India-Asia collision onset-
904 Facts, theories, controversies. Earth-Science Reviews 160, 264-299.

905 Ji, W., Wu, F., Liu, C., Chung, S., 2009. Geochronology and petrogenesis of granitic rocks in Gangdese
906 batholith, southern Tibet. Science in China Series D: Earth Sciences 52, 1240-1261.

907 Johnson, N.M., Stix, J., Tauxe, L., Cervený, P.F., Tahirkheli, R.A., 1985. Paleomagnetic chronology, fluvial
908 processes, and tectonic implications of the Siwalik deposits near Chinji Village, Pakistan. *The Journal of*
909 *Geology*, 27-40.

910 Jonell, T.N., Li, Y., Blusztajn, J., Giosan, L., Clift, P.D., 2018. Signal or noise? Isolating grain size effects on
911 Nd and Sr isotope variability in Indus delta sediment provenance. *Chemical Geology* 485, 56-73.

912 Khan, M.A., STERN, R.J., GRIBBLE, R.F., WINDLEY, B.F., 1997. Geochemical and isotopic constraints on
913 subduction polarity, magma sources, and palaeogeography of the Kohistan intra-oceanic arc, northern
914 Pakistan Himalaya. *Journal of the Geological Society* 154, 935-946.

915 Khan, S., Stern, R., Manton, M., Copeland, P., Kimura, J., Khan, M., 2004. Age, geochemical and Sr–Nd–Pb
916 isotopic constraints for mantle source characteristics and petrogenesis of Teru Volcanics, Northern
917 Kohistan Terrane, Pakistan. *Tectonophysics* 393, 263-280.

918 Khan, S.D., Walker, D.J., Hall, S.A., Burke, K.C., Shah, M.T., Stockli, L., 2009. Did the Kohistan-Ladakh island
919 arc collide first with India? *Geological Society of America Bulletin* 121, 366-384.

920 Koppers, A.A., 2002. ArArCALC—software for 40 Ar/39 Ar age calculations. *Computers & Geosciences* 28,
921 605-619.

922 Leier, A.L., Kapp, P., Gehrels, G.E., DeCelles, P.G., 2007. Detrital zircon geochronology of Carboniferous–
923 Cretaceous strata in the Lhasa terrane, Southern Tibet. *Basin Research* 19, 361-378.

924 Mahéo, G., Blichert-Toft, J., Pin, C., Guillot, S., Pêcher, A., 2009. Partial melting of mantle and crustal
925 sources beneath South Karakorum, Pakistan: implications for the Miocene geodynamic evolution of the
926 India–Asia convergence zone. *Journal of Petrology* 50, 427-449.

927 Malusà, M.G., Resentini, A., Garzanti, E., 2016. Hydraulic sorting and mineral fertility bias in detrital
928 geochronology. *Gondwana Research* 31, 1-19.

929 Miller, C., Schuster, R., Klötzli, U., Frank, W., Purtscheller, F., 1999. Post-collisional potassic and
930 ultrapotassic magmatism in SW Tibet: geochemical and Sr–Nd–Pb–O isotopic constraints for mantle
931 source characteristics and petrogenesis. *Journal of Petrology* 40, 1399-1424.

932 Najman, Y., 2006. The detrital record of orogenesis: A review of approaches and techniques used in the
933 Himalayan sedimentary basins. *Earth-Science Reviews* 74, 1-72.

934 Najman, Y., Bickle, M., Garzanti, E., Pringle, M., Barfod, D., Brozovic, N., Burbank, D., Ando, S., 2009.
935 Reconstructing the exhumation history of the Lesser Himalaya, NW India, from a multitechnique
936 provenance study of the foreland basin Siwalik Group. *Tectonics* 28.

937 Najman, Y., Garzanti, E., Pringle, M., Bickle, M., Stix, J., Khan, I., 2003. Early-Middle Miocene paleodrainage
938 and tectonics in the Pakistan Himalaya. *Geological Society of America Bulletin* 115, 1265-1277.

939 Najman, Y., Pringle, M., Godin, L., Oliver, G., 2001. Dating of the oldest continental sediments from the
940 Himalayan foreland basin. *Nature* 410, 194-197.

941 Najman, Y., Pringle, M., Godin, L., Oliver, G., 2002. A reinterpretation of the Balakot Formation:
942 Implications for the tectonics of the NW Himalaya, Pakistan. *Tectonics* 21, 9-1-9-18.

943 Nyberg, B., Helland-Hansen, W., Gawthorpe, R.L., Sandbakken, P., Eide, C.H., Sømme, T., Hadler-Jacobsen,
944 F., Leiknes, S., 2018. Revisiting morphological relationships of modern source-to-sink segments as a first-
945 order approach to scale ancient sedimentary systems. *Sedimentary Geology* 373, 111-133.

946 Palin, R., Searle, M., Waters, D., Horstwood, M., Parrish, R., 2012. Combined thermobarometry and
947 geochronology of peraluminous metapelites from the Karakoram metamorphic complex, North Pakistan;
948 New insight into the tectonothermal evolution of the Baltoro and Hunza Valley regions. *Journal of*
949 *Metamorphic Geology* 30, 793-820.

950 Pan, F.-B., Zhang, H.-F., Xu, W.-C., Guo, L., Wang, S., Luo, B.-j., 2014. U–Pb zircon chronology, geochemical
951 and Sr–Nd isotopic composition of Mesozoic–Cenozoic granitoids in the SE Lhasa terrane: Petrogenesis
952 and tectonic implications. *Lithos* 192, 142-157.

953 Pivnik, D.A., Wells, N.A., 1996. The transition from Tethys to the Himalaya as recorded in northwest
954 Pakistan. *Geological Society of America Bulletin* 108, 1295-1313.

955 Qasim, M., Ding, L., Khan, M.A., Jadoon, I.A., Haneef, M., Baral, U., Cai, F., Wang, H., Yue, Y., 2018. Tectonic
956 implications of detrital zircon ages from lesser Himalayan Mesozoic - Cenozoic strata, Pakistan.
957 *Geochemistry, Geophysics, Geosystems* 19, 1636-1659.

958 Qayyum, M., Niem, A.R., Lawrence, R.D., 2001. Detrital modes and provenance of the Paleogene Khojak
959 Formation in Pakistan: Implications for early Himalayan orogeny and unroofing. *Geological Society of
960 America Bulletin* 113, 320-332.

961 Richards, A., Argles, T., Harris, N., Parrish, R., Ahmad, T., Darbyshire, F., Draganits, E., 2005. Himalayan
962 architecture constrained by isotopic tracers from clastic sediments. *Earth and Planetary Science Letters*
963 236, 773-796.

964 Robinson, D.M., DeCelles, P.G., Patchett, P.J., Garzione, C.N., 2001. The kinematic evolution of the
965 Nepalese Himalaya interpreted from Nd isotopes. *Earth and Planetary Science Letters* 192, 507-521.

966 Roddaz, M., Said, A., Guillot, S., Antoine, P.-O., Montel, J.-M., Martin, F., Darrozes, J., 2011. Provenance of
967 Cenozoic sedimentary rocks from the Sulaiman fold and thrust belt, Pakistan: implications for the
968 palaeogeography of the Indus drainage system. *Journal of the Geological Society* 168, 499-516.

969 Schaltegger, U., Zeilinger, G., Frank, M., Burg, J.P., 2002. Multiple mantle sources during island arc
970 magmatism: U–Pb and Hf isotopic evidence from the Kohistan arc complex, Pakistan. *Terra Nova* 14, 461-
971 468.

972 Schärer, U., Hamet, J., Allègre, C.J., 1984. The Transhimalaya (Gangdese) plutonism in the Ladakh region:
973 a U Pb and Rb Sr study. *Earth and Planetary Science Letters* 67, 327-339.

974 Schneider, D., Zeitler, P., Kidd, W., Edwards, M., 2001. Geochronologic constraints on the tectonic
975 evolution and exhumation of Nanga Parbat, western Himalaya syntaxis, revisited. *The Journal of Geology*
976 109, 563-583.

977 Searle, M., Khan, M.A., Fraser, J., Gough, S., Jan, M.Q., 1999. The tectonic evolution of the Kohistan -
978 Karakoram collision belt along the Karakoram Highway transect, north Pakistan. *Tectonics* 18, 929-949.

979 Searle, M., Parrish, R.R., Thow, A., Noble, S., Phillips, R., Waters, D., 2010. Anatomy, age and evolution of
980 a collisional mountain belt: the Baltoro granite batholith and Karakoram Metamorphic Complex, Pakistani
981 Karakoram. *Journal of the Geological Society* 167, 183-202.

982 Shah, S.I., 2009. Stratigraphy of Pakistan. Government of Pakistan Ministry of Petroleum & Natural
983 Resources Geological Survey of Pakistan.

984 Sinclair, H., Jaffey, N., 2001. Sedimentology of the Indus Group, Ladakh, northern India: implications for
985 the timing of initiation of the palaeo-Indus River. *Journal of the Geological Society* 158, 151-162.

986 Treloar, P., Rex, D., Guise, P., Wheeler, J., Hurford, A., Carter, A., 2000. Geochronological constraints on
987 the evolution of the Nanga Parbat syntaxis, Pakistan Himalaya. Geological Society, London, Special
988 Publications 170, 137-162.

989 Treloar, P.J., Palin, R.M., Searle, M.P., 2019. Towards resolving the metamorphic enigma of the Indian
990 Plate in the NW Himalaya of Pakistan. Geological Society, London, Special Publications 483, 255-279.

991 Ullah, K., Arif, M., Shah, M.T., 2015. Geochemistry and provenance of the Lower Siwaliks from
992 southwestern Kohat, western Himalayan Foreland Basin, NW Pakistan. *Geologica Acta: an international
993 earth science journal* 13, 45-61.

994 Van Der Beek, P., Van Melle, J., Guillot, S., Pêcher, A., Reiners, P.W., Nicolescu, S., Latif, M., 2009. Eocene
995 Tibetan plateau remnants preserved in the northwest Himalaya. *Nature Geoscience* 2, 364-368.

996 van Hinsbergen, D.J., Lippert, P.C., Dupont-Nivet, G., McQuarrie, N., Doubrovine, P.V., Spakman, W.,
997 Torsvik, T.H., 2012. Greater India Basin hypothesis and a two-stage Cenozoic collision between India and
998 Asia. *Proceedings of the National Academy of Sciences* 109, 7659-7664.

999 Wallis, D., Carter, A., Phillips, R.J., Parsons, A.J., Searle, M.P., 2016. Spatial variation in exhumation rates
1000 across Ladakh and the Karakoram: New apatite fission track data from the Eastern Karakoram, NW India.
1001 *Tectonics*.

1002 Welcomme, J.-L., Benammi, M., Crochet, J.-Y., Marivaux, L., Métais, G., Antoine, P.-O., Baloch, I., 2001.
1003 Himalayan Forelands: palaeontological evidence for Oligocene detrital deposits in the Bugti Hills
1004 (Balochistan, Pakistan). *Geological Magazine* 138, 397-405.

1005 Whittington, A., Foster, G., Harris, N., Vance, D., Ayres, M., 1999. Lithostratigraphic correlations in the
1006 western Himalaya—an isotopic approach. *Geology* 27, 585-588.

1007 Willis, B., 1993. Evolution of Miocene fluvial systems in the Himalayan foredeep through a two kilometer-
1008 thick succession in northern Pakistan. *Sedimentary geology* 88, 77-121.

1009 Yeats, R.S., Hussain, A., 1987. Timing of structural events in the Himalayan foothills of northwestern
1010 Pakistan. *Geological society of America bulletin* 99, 161-176.

1011 Yu, Z., Colin, C., Wan, S., Saraswat, R., Song, L., Xu, Z., Clift, P., Lu, H., Lyle, M., Kulhanek, D., 2019. Sea
1012 level-controlled sediment transport to the eastern Arabian Sea over the past 600 kyr: Clay minerals and
1013 SrNd isotopic evidence from IODP site U1457. *Quaternary Science Reviews* 205, 22-34.

1014 Zaleha, M.J., 1997. Intra - and extrabasinal controls on fluvial deposition in the Miocene Indo - Gangetic
1015 foreland basin, northern Pakistan. *Sedimentology* 44, 369-390.

1016 Zanchi, A., Gaetani, M., 2011. The geology of the Karakoram range, Pakistan: the new 1: 100,000
1017 geological map of Central-Western Karakoram. *Italian journal of geosciences* 130, 161-262.

1018 Zhang, H., Harris, N., Parrish, R., Kelley, S., Zhang, L., Rogers, N., Argles, T., King, J., 2004. Causes and
1019 consequences of protracted melting of the mid-crust exposed in the North Himalayan antiform. *Earth and
1020 Planetary Science Letters* 228, 195-212.

1021 Zhou, P., Stockli, D.F., Ireland, T., Murray, R.W., Clift, P.D., 2021. Zircon U - Pb Age Constraints on NW
1022 Himalayan Exhumation from the Laxmi Basin, Arabian Sea. *Geochemistry, Geophysics, Geosystems*,
1023 e2021GC010158.

1024 Zhou, P., Stockli, D.F., Ireland, T., Murray, R.W., Clift, P.D., 2022. Zircon U - Pb Age Constraints on NW
1025 Himalayan Exhumation from the Laxmi Basin, Arabian Sea. *Geochemistry, Geophysics, Geosystems*,
1026 e2021GC010158.

1027 Zhu, D.-C., Mo, X.-X., Niu, Y., Zhao, Z.-D., Wang, L.-Q., Liu, Y.-S., Wu, F.-Y., 2009. Geochemical investigation
1028 of Early Cretaceous igneous rocks along an east–west traverse throughout the central Lhasa Terrane,
1029 Tibet. *Chemical Geology* 268, 298-312.

1030 Zhu, D.-C., Zhao, Z.-D., Niu, Y., Dilek, Y., Wang, Q., Ji, W.-H., Dong, G.-C., Sui, Q.-L., Liu, Y.-S., Yuan, H.-L.,
1031 2012. Cambrian bimodal volcanism in the Lhasa Terrane, southern Tibet: record of an early Paleozoic
1032 Andean-type magmatic arc in the Australian proto-Tethyan margin. *Chemical Geology* 328, 290-308.

1033 Zhuang, G., Najman, Y., Guillot, S., Roddaz, M., Antoine, P.-O., Métais, G., Carter, A., Marivaux, L., Solangi,
1034 S.H., 2015. Constraints on the collision and the pre-collision tectonic configuration between India and Asia
1035 from detrital geochronology, thermochronology, and geochemistry studies in the lower Indus basin,
1036 Pakistan. *Earth and Planetary Science Letters* 432, 363-373.

1037 Zhuang, G., Najman, Y., Tian, Y., Carter, A., Gemignani, L., Wijbrans, J., Jan, M.Q., Khan, M.A., 2018. Insights
1038 into the evolution of The Hindu Kush-Kohistan-Karakoram from modern river sand detrital geo- and
1039 thermochronological studies. *Journal of the Geological Society*, jgs2018-2007.

1040