

1 **SOIL BIOLOGY AND BIOCHEMISTRY**

2

3 **Impact of single and binary mixtures of phenanthrene and N-PAHs on microbial utilisation of ¹⁴C-glucose**
4 **in soil**

5

6 **Ihuoma N. Anyanwu^{a, b} and Kirk T. Semple^{a,*}**

7

8 ^a Lancaster Environment Centre, Lancaster University, Lancaster, LA1 4YQ, United Kingdom.

9

10 ^b Department of Biological Sciences, Federal University Ndufu-Alike Ikwo, P.M.B 1010, Abakaliki, Ebonyi
11 State, Nigeria.

12

13 Corresponding author: k.semple@lancs.ac.uk

14

15 **Abstract**

16 Microbes are susceptible to contaminant effects, and high concentration of chemicals in soil can impact on
17 microbial growth, density, viability and development. The impact of single and binary mixtures of phenanthrene
18 and its nitrogen-containing polycyclic aromatic hydrocarbon analogues (N-PAHs) on microbial metabolism of
19 ^{14}C -glucose in soil was measured over a 90 d soil-contact time. Impacts were assessed by measuring the rates and
20 mean overall extents of mineralisation (%), as well as the incorporation of ^{14}C -glucose into the microbial biomass.
21 The result revealed that the extents of ^{14}C -glucose mineralisation were consistently greater in N-PAH amended
22 soils than the control and phenanthrene soils with increased incubations. This indicates a trend of increasing
23 diversion of C from biosynthesis to maintenance requirement by soil microorganisms. Furthermore, biomass
24 uptake in the amended soils showed reduced substrate utilization (fixed- k_{EC}), suggesting that N-PAHs decreased
25 the amount of substrate-C that was incorporated into the microbial biomass. This however, signifies that N-PAHs
26 imposes oxidative stress on soil microbial community.

27

28 **Key words:** N-PAHs, phenanthrene, ^{14}C -biomass uptake, mineralisation, k_{EC} coefficient.

29

30

31 1. Introduction

32 The importance of microbial activity in the cycling of organic matter and regulating active nutrient pools suggests
33 that the effects of stress on microbial community will fundamentally impact on crops, natural vegetation and
34 ecosystem productivity (Killham, 1985; Anyanwu and Semple, 2016a; Siles and Margesin, 2017). Soil
35 microorganisms are very sensitive to environmental stress or change, and this often results in the diversion of
36 carbon from biosynthesis to maintenance of cells (Bargett and Saggar, 1994; Anyanwu and Semple, 2016a). Thus,
37 soil microbial biomass measurements are important in ascertaining the extent of chemical stress and/or
38 disturbance on soil ecosystem and the time dependence of microbial recovery. Most studies have used respiration
39 rate (Fournier et al., 1992; Nakamoto and Wakahara, 2004; Anyanwu and Semple, 2016a; Sun et al., 2017; Xu et
40 al., 2017) and changes in biomass (Anyanwu and Semple, 2016a; Mehnaz et al., 2017; Siles and Margesin, 2017).
41 Using a ^{14}C -substrate, the influence of synthetic and organophosphate sheep dip formulations (Boucard et al.,
42 2008), pesticides (Fournier et al., 1992), heavy metals (Bargett and Saggar, 1994; Bogomolov et al., 1996), sewage
43 sludge (Fließbach et al., 1994; Witter and Dahlin, 1995) and the ratio of ^{14}C -biomass-incorporated with ^{14}C -
44 respired (Sparling and West, 1989; Sparling et al., 1990; Gunina et al., 2017), have been determined on soil
45 microbial activity. The approach of using ^{14}C -glucose as a substrate to determine the ratio of respired-C, to
46 biomass-incorporated C, has shown that microorganisms in contaminated soils are less efficient in the utilization
47 of substrates for biomass synthesis and spend more energy in the maintenance requirements (Bargett and Saggar,
48 1994; Witter and Dahlin, 1995; Anyanwu and Semple, 2016a; Gunina et al., 2017). Thus, leading to a decrease in
49 the ratio, increases in stress, faster respiration, reduced efficiency of fresh substrate incorporation into new soil
50 microbial biomass and increased microbial turnover in contaminated soils (Fließbach et al., 1994; Bargett and
51 Saggar, 1994; Witter and Dahlin, 1995; Boucard et al., 2008; Gunina et al., 2017; Bore et al., 2017). These studies
52 have revealed that the growth, activity and physiological conditions of soil microbial community may be altered
53 and/or destroyed by the presence of contaminants.

54 Persistent contaminants are of particular concern due to their toxicity and widespread pollution that has occurred
55 during production, spills, combustion and disposition (Beelen and Doelman, 1997; Anyanwu and Semple, 2015a);
56 examples include metals, pesticides and polycyclic aromatic hydrocarbons (PAHs). However, for sustainable
57 environmental policies and regulations, risk assessment of other persistent contaminants such as, the nitrogen-
58 containing polycyclic aromatic hydrocarbons (N-PAHs) in the environment is of great importance. N-PAHs are
59 chemicals present in most contaminated sites worldwide and represent two-thirds of known organic xenobiotic

60 chemically synthesized (Rajasekhar et al., 2000; Anyanwu and Semple, 2015a). For example, they are used as
61 industrial solvents, dyes, explosives, pharmaceuticals and pesticides (Kaiser et al., 1996). The US Environmental
62 Protection Agency (USEPA) and International Agency for Research on Cancer (IARC) classified N-PAHs as
63 probable human carcinogens (IARC, 2012). Furthermore, many of these N-PAHs are antimicrobial (Vance et al.,
64 1986; Ferraz et al., 2017); therefore, their accumulation is a major threat to microbes because they have the
65 potency of inducing oxidative stress to soil microorganisms and other biotas.

66 Despite the widespread uses of N-PAHs, and previous N-PAHs studies in literature (Anyanwu and Semple, 2015a;
67 2015b; 2016a; Anyanwu et al., 2017), there has not been information of their impacts on microbial utilization of
68 ^{14}C -glucose and/or synthesis of cell biomass in soil. Functionally, microbes can act as relevant indicators of
69 environmental pollution; as a result, there is great need to assess the impact of N-PAHs on soil microbial
70 metabolism and biosynthesis of cell biomass. In this study therefore, the impact of single and binary mixtures of
71 phenanthrene and its nitrogen-containing analogues on microbial utilization of ^{14}C -glucose was investigated over
72 a 90 d incubation period in soil using respirometric assays.

73

74 **2. Materials and Methods**

75 *2.1 Chemicals*

76 Phenanthrene (Phen), 1,10-phenanthroline (1,10-Phen), 1,7-phenanthroline (1,7-Phen), 4,7-phenanthroline (4,7-
77 Phen) and benzo[h]quinoline (B[h]Q) and radiolabelled ^{14}C -glucose were obtained from Sigma-Aldrich, UK.
78 Goldstar liquid scintillation cocktails were supplied by Meridian Biotechnologies Ltd, UK.

79

80 *2.2 Soil preparation*

81 A pristine agricultural soil from Myerscough, UK, collected from the top layer of field under pasture, from a depth
82 of approximately 5-20 cm was prepared for the study (n = 3). The soil texture was sandy-loam (19.5% clay, 60.4%
83 sand, 20.0% silt), with organic matter content of 2.7%; total nitrogen of 0.14%; total organic carbon of 1.6% and
84 pH 6.5. The soil was thoroughly homogenized, air dried at room temperature and sieved with 2 mm mesh size.
85 The soil was rehydrated with deionised water back to 45% water holding capacity (WHC) and amended with
86 phenanthrene and the N-PAH analogues as described in Doick et al. (2003). Soil samples were placed in bowls:

87 $\frac{1}{3}$ (100 g; n = 3) were amended with phenanthrene and four N-PAH standards (benzo[h]quinoline, 1,10-
88 phenanthroline, 1,7-phenanthroline or 4,7-phenanthroline) dissolved in acetone to give concentration of 100 mg
89 kg^{-1} . The amended soils were kept in the fume hood for 3 h to allow the carrier solvent volatilize, after which the
90 soils were mixed with the remaining $\frac{2}{3}$ (200 g). Blanks were prepared using un-amended soils. Soils amended
91 with acetone only were also prepared to serve as a control. The amended soils were kept in amber glass jars and
92 aged in the dark at $21 \pm 1^\circ\text{C}$ for 1, 30, 60 and 90 d. Soil moisture content was checked regularly and lost water
93 was replenished with deionized water. After each ageing time (30 d interval), soils were analysed for microbial-
94 substrate-mineralisation and biomass uptake. Extractability of phenanthrene and the N-PAH analogues from soil
95 over time, and their percentage recoveries has been reported by Anyanwu and Semple (2015b, 2016a) (Table 1).

96

97 *2.3 Mineralisation of ^{14}C -glucose in soil.*

98 The ability of indigenous soil microorganisms to mineralise ^{14}C -glucose to $^{14}\text{CO}_2$ was assessed at 1, 30, 60 and 90
99 d contact time. Respirometric assays were carried out in modified 250 ml Schott bottles incorporating a Teflon-
100 lined screw cap containing 1 M NaOH to trap any $^{14}\text{CO}_2$ (Reid et al., 2001). A slurry system with a solid: liquid
101 ratio of 2:1 (20 g soil: 10 ml sterile water) was used to ensure complete $^{12/14}\text{C}$ -glucose distribution. Standards were
102 prepared in sterilized deionised water and delivered to give a ^{12}C -glucose concentration of 3 mM glucose solution
103 with an associated ^{14}C -activity of 800 Bq per respirometer. Controls were also prepared. Respirometers were
104 shaken at 100 rpm on an orbital shaker (Janke and Kunkel, IKA[®]-Labortechnik KS 510D), in the dark at $21 \pm 1^\circ\text{C}$.
105 Sampling was carried out every 1, 2, 4, 6, 8, 12, 24 h and 2, 3, 4, 5 d with the vials containing trapped $^{14}\text{CO}_2$.
106 Goldstar liquid scintillation cocktail was added to the vials. The vials were stored in the dark for 24 h before
107 sample quantification was carried out by liquid scintillation counting (LSC) using standard calibration and quench
108 correction techniques (Reid et al., 2001).

109

110 *2.4 Uptake of ^{14}C glucose into microbial biomass*

111 After each 5 d incubation, soil samples from respirometers were divided into three portions and analysed as
112 follows:

113 a) *Sample oxidation:* The first sample was oven dried at 30°C and combusted in a sample oxidizer (Packard
114 307) to determine the level of ^{14}C -activity remaining (i.e. residual ^{14}C -activity in soil). Soil (1 g), plus

115 200 µl of combustAid was combusted for 3 min. Carbon-sorb-E (10 ml) and Permaflour-E (10 ml) was
116 used as CO₂ trap and scintillation fluid, respectively. Sample quantification was carried out using LSC.

117 b) *Un-fumigated extraction:* The second sample (~4 g) was immediately extracted with 0.5 M K₂SO₄ (50
118 ml, pH 7) by shaking on an orbital shaker at 100 rpm for 30 min. The soil solutions were filtered using
119 Whatman No 1 filter papers and an aliquot of 5 ml supernatant was added to 15 ml scintillation cocktail.
120 The quantification of ¹⁴C-activity was carried out using the LSC.

121 c) *Fumigated extraction:* The third sample (~4 g) was placed in a desiccator and fumigated with ethanol-
122 free chloroform for 24 h to measure the ¹⁴C-activity within microbial biomass. After fumigation, the
123 samples were vented to remove chloroform residuals in the soil. After venting, samples were extracted
124 with 0.5 M K₂SO₄, filtered (using Whatman No 1 filter papers) and analysed as per the un-fumigated
125 extract.

126

127 2.5 Statistical analysis

128 The proportion of ¹⁴C-glucose incorporated into the microbial biomass was calculated as in Sparling et al. (1990)
129 and Boucard et al. (2008).

130 $^{14}\text{C-flush} = ^{14}\text{C-activity in fumigated soil} - ^{14}\text{C-activity in un-fumigated soil}.$

131 $^{14}\text{C-microbial biomass} = ^{14}\text{C-flush} \div k_{EC}.$

132 1. A fixed k_{EC} coefficient (0.35) was used to convert C-flush into microbial biomass Sparling et al., 1990;
133 Boucard et al., 2008).

134 2. Variable k_{EC} coefficients were also calculated from each amendment, at all the ageing times, and the ¹⁴C-
135 microbial biomass was re-calculated with the new coefficient. This process is based on the assumption
136 that; the calculated ¹⁴C-labelled microbial-C is a representative of the total microbial biomass and that
137 all the ¹⁴C-activity not taken into account by mineralisation and un-fumigated soil extraction has been
138 incorporated into the microbial biomass with negligible amount of extracellular metabolite (Sparling et
139 al., 1990; Boucard et al., 2008).

140 $k_{EC} = (^{14}\text{C-flush}) \div (^{14}\text{C}_{\text{init.}} - ^{14}\text{C-respired} - ^{14}\text{C-activity in un-fumigated soil}).$

141 ¹⁴C-flush and ¹⁴C-microbial biomass were later on expressed as percentages of the initial ¹⁴C-activity
142 ($^{14}\text{C}_{\text{init.}}$).

143 3. Biophysical quotients (BQ) were calculated as:

144
$$\text{BQ} = {}^{14}\text{CO}_2 \text{ respired} \div {}^{14}\text{C-microbial biomass (calculated from either fixed or variable } k_{EC}\text{)}.$$

145 Following blank corrections, data was statistically analysed using SigmaStat 3.5. Statistical significant differences
146 between the impacts of phenanthrene, N-PAHs, and soil contact time on soil microbial activity following addition
147 of ${}^{14}\text{C}$ -glucose was determined using analysis of variance (ANOVA). The statistical difference between the
148 biomass calculated with fixed and variable k_{EC} was also determined. Results are statistically significant when
149 $p < 0.05$. Data was presented as mean \pm SE and graphs were plotted using Sigma-Plot 10.0 version.

150

151 3. Results

152 3.1 Mineralisation of ${}^{14}\text{C}$ -glucose to ${}^{14}\text{CO}_2$ by soil microorganisms

153 The mineralisation of ${}^{14}\text{C}$ -glucose in the presence of 100 mg kg^{-1} phenanthrene and its N-PAH analogues was
154 measured (Fig. 1 and 2). Upon the addition of glucose, there was a considerable increase in % mineralisation in
155 the presence of the amended chemicals. However, the mineralisation of the ${}^{14}\text{C}$ -substrate (glucose) in the presence
156 of benzo[h]quinoline (B[h]Q) soil was reduced at 1 d compared to the control soils (Fig. 1 and 2).

157 The fastest rates of mineralisation were determined (Table 2), and the fastest rates ($\% {}^{14}\text{CO}_2 \text{ h}^{-1}$) recorded
158 maximum values after 24 h following addition of ${}^{14}\text{C}$ -glucose in all the amendments at all of the time points with
159 the exception of 4,7-Phen, B[h]Q and Phen, which recorded their fastest rates 48 h after addition of ${}^{14}\text{C}$ -glucose
160 (30 d) (Table 2). Furthermore, 1,10-Phen (single amendment) and 1,10-Phen + Phen (binary mixtures) recorded
161 maximum fastest rates at 6 h (90 d). From the data, the fastest rates followed a trend of decreased values with
162 increases in the soil-contact time. However, 1,7-Phen and B[h]Q amendments showed a dramatic rise of 50% and
163 70%, respectively, after 90 d (Table 2).

164 The extents of mineralisation (total ${}^{14}\text{CO}_2$ -respired (%)) were determined (Table 2). The results revealed that the
165 extents of mineralisation of ${}^{14}\text{C}$ -glucose appeared to be consistently greater in amended soils than the control soils
166 with increase in ageing time; with the exception of 1,10-Phen, 4,7-Phen and Phen amended soils (90 d). The
167 overall extent of mineralisation followed a trend of increased ${}^{14}\text{C}$ -glucose mineralisation at 1 d in all the
168 amendments. Among the N-PAHs, however, B[h]Q soils recorded increased mineralisation with increase in soil-
169 contact time, but, this declined a little at 90 d (Table 2).

170 While the extents of mineralisation in the single amendments displayed decreased and increased values, a
171 consistent decrease in ¹⁴C-glucose mineralisation was observed in the binary mixtures over time (Table 2).
172 Analysis of data among the treatment groups showed no statistically significant differences between the mean
173 values at 1 d ($p>0.05$); however, statistically significant differences was observed after 30 d ($p<0.05$) (Table 2).
174 Furthermore, statistical analysis of data showed statistically significance differences between phenanthrene and
175 N-PAH amended soils over time ($p<0.05$). In addition, incubation times were observed to affect the %
176 mineralisation of ¹⁴C-glucose in all the amendments at all the time points ($p<0.001$).

177

178 *3.2 Impact of phenanthrene and N-PAHs on the k_{EC} coefficients*

179 The impact of 100 mg/kg phenanthrene and its nitrogen-containing analogues on the k_{EC} coefficients was
180 calculated (Table 2). It was noted that the fumigation-extraction released 0.3–15% of the incorporated microbial-
181 C giving the calculated k_{EC} ranges of 0.003–0.149. However, variation among the chemical amendments was
182 observed in the calculated k_{EC} values obtained after the fumigation-extraction (Table 2). Furthermore, the
183 calculated k_{EC} values (0.003–0.149) were lower than the fixed k_{EC} value (0.35) in all the amendments (Table 2).
184 Although the data showed a disparaging statistical difference, all the amendments showed a similar trend of low
185 k_{EC} values at 1 d and 60 d and high k_{EC} values at 30 d and 90 d (with the exception of Phen and 1,7-Phen + Phen
186 chemicals). Also, the presence of 1,10-Phen and 1,7-Phen in soil recorded lower k_{EC} coefficients (30 d and 60 d)
187 compared to the control soil values (Table 2).

188 Soils amended with binary mixtures of phenanthrene and N-PAHs recorded low, but varying k_{EC} values compared
189 to the control soils at all the time points. For example, while control fixed k_{EC} values ranged from $3.06 \pm 0.87 -$
190 19.92 ± 3.65 , values of $0.66 \pm 0.13 - 15.08 \pm 3.47$ and $1.12 \pm 0.11 - 9.93 \pm 1.22$ (fixed k_{EC}) were recorded in the
191 single amendments and binary mixtures, respectively. Also, while control values for variable $k_{EC} = 46.67 \pm 8.55$
192 $- 63.45 \pm 18.1$, values range of $34.03 \pm 3.11 - 62.45 \pm 11.22$ and $41.45 \pm 1.93 - 59.37 \pm 9.45$ (variable k_{EC}) were
193 measured (single amendments and binary mixtures, respectively) (Table 2). In addition, the calculated k_{EC} values
194 of $0.003 \pm 0.00 - 0.140 \pm 0.03$ (single amendments), and $0.007 \pm 0.00 - 0.013 \pm 0.00$ (binary mixtures) was
195 obtained, while, control values = $0.016 \pm 0.00 - 0.149 \pm 0.02$ (Table 2). Furthermore, incubation time was noted
196 to have statistically significant effect on the k_{EC} values in binary mixtures ($p<0.001$). Although not consistent,
197 there was a trend of increases in the extraction efficiency of K_2SO_4 at 30 d and 90 d.

198

199 *3.3 Uptake of ¹⁴C-glucose into microbial biomass*

200 The incorporation of the ¹⁴C-substrate into the microbial biomass in soils amended with 100 mg/kg single and
201 binary mixtures of phenanthrene and the N-PAH analogues were calculated using fixed and variable k_{ECs} ,
202 respectively (Table 2). Although not showing a consistent trend, the results showed that increase in fixed k_{EC}
203 values, lead to decrease in variable k_{EC} values and vice versa in all the amendments, with the exception of B[h]Q
204 amendment (Table 2). Statistical analysis of data showed statistically significant differences in the calculated data
205 for both the fixed and variable k_{EC} values obtained after 1 d – 90 d soil-contact time ($p < 0.001$). Also, there was a
206 consistent trend of higher values in the C-flush resulting in higher biomass values (fixed k_{EC}) and lower biomass
207 values (variable k_{EC}) (Table 2).

208 The BQs (using biomass values calculated with fixed and variable k_{ECs}) was determined, and the results differed
209 significantly in all the amendments at all the time points (Table 2). For example, BQs calculated with the fixed
210 k_{EC} varied widely compared to that of the variable k_{EC} . Furthermore, the amended soils recorded significant BQ
211 values at 1 d compared with the control soils ($p < 0.05$); and Phen amendment recorded the highest BQ value of
212 54.76% (Table 2). A trend of high BQ values at 1 d and 60 d (fixed k_{EC}), and low values at 30 d and 90 d (variable
213 k_{EC}) was observed in all of the amendments (Table 2). Among the chemical amendments, B[h]Q showed a
214 consistent increase in BQ value with increased ageing, recording values > 1 ($p < 0.05$). Although showing high
215 variability, the calculated BQs recorded high values with increased ageing in the binary mixtures.

216

217 **4. Discussion**

218 *4.1 Mineralisation of ¹⁴C-glucose to ¹⁴CO₂ by soil microorganisms*

219 The impact of single and binary mixtures of phenanthrene and its nitrogen-containing analogues on microbial
220 utilisation of ¹⁴C-glucose in soil was studied over a 90-d incubation. Loss of phenanthrene, benzo[h]quinoline,
221 1,10-phenanthroline, 1,7-phenanthroline or 4,7-phenanthroline through volatilisation was considered minimal due
222 to the sealed nature of the incubations (Hofman et al., 2008; Towell et al., 2011). From the results, mineralisation
223 of ¹⁴C-glucose was greater in the N-PAHs amended soils than the control soil after 1 d; indicating that
224 microorganisms utilized energy for cell maintenance rather than biosynthesis of new cells. This phenomenon

225 agrees with the observations of Bargett and Saggar (1994), Witter and Dahlin (1995), Chnader and Joergensen
226 (2001), Boucard et al. (2008), and Bore et al. (2017). In support, Fließbach et al. (1994) reported that in heavily
227 contaminated sites, soil respiration increased substantially compared to the corresponding low contaminated soils.
228 Respiration has been linked as a process and microbial biomass as a pool to metabolic quotient for CO₂ (qCO_2)
229 by Anderson and Domsch (1986). Thus, it is widely accepted that a high qCO_2 is a surprisingly common
230 characteristic of soil microbial biomass in chronically contaminated soils (Fließbach et al., 1994). This has been
231 suggested to be a useful indicator of oxidative stress in soils (Brookes, 1993; Mooshammer et al., 2017). In
232 addition, Gunina et al. (2017) reported that an increased qCO_2 indicates stress to the soil microbial community.

233 In this study, N-PAHs (B[h]Q amendment) recorded low mineralisation at 1 d; this is an evidence of reduced
234 microbial substrate utilization efficiency under chemical stress. Hattori (1992) and Molaei et al. (2017)
235 documented that initial microbial respiratory responses are the most sensitive in quantifying the impact of
236 contaminants following their introduction into soil. However, the consistent increase in B[h]Q mineralisation after
237 30 d could be attributed to oxidative stress and/or chemical bioavailability, due to its lower K_{ow} (Anyanwu and
238 Semple, 2015b, 2016a); since the total concentration did not exceed that of other amended soils.

239 A decline in mineralisation (%) was observed over time. The notable decline may be as a result of chemical
240 sequestration (into soil organic matter) there by rendering the contaminants less available to microorganisms
241 (Semple et al., 2007), microbial degradation (Anyanwu and Semple, 2015b; 2016a) and/or adaptation to toxicity
242 (Granato et al., 2017; Anyanwu and Semple, 2017b). Organic contaminants are known to be retained within the
243 soil through chemical or physical sequestration processes, such as binding, sorption to clay and/or soil organic
244 matter as well as occlusion within the 3-dimensional structure of the soil (Semple et al., 2007). Furthermore,
245 factors which include, soil organic matter content and physico-chemical properties of the chemical (aqueous
246 solubility, polarity, hydrophobicity, molecular structure, K_{ow} , and lipophilicity) are known to control the fate and
247 behaviour of organic contaminants (N-PAHs) in soil (Anyanwu and Semple, 2015b; 2017b; Zhu et al., 2017;
248 Doley et al., 2017).

249

250 *4.2 Uptake of ¹⁴C-glucose into soil microbial biomass*

251 The study revealed that microbial uptake in the chemically amended soils did, however, show reduced substrate
252 utilization. Thus, the amount of glucose incorporated was lower in amended soils. Studies have shown that

253 microorganisms subjected to stress exhibit a higher ratio of respired-C to biomass-incorporated-C; indicating a
254 reduced microbial substrate utilization efficiency under chemical stress and a change in community structure
255 following substrate addition (Bargett and Saggarr, 1994; Witter and Dahlin, 1995; Frostegård et al., 1996; Knight
256 et al., 1997; Boucard et al., 2008; Gunina et al., 2017). Killham (1985) recorded that increasing stress often causes
257 a reduction in soil respiration, soil dehydrogenase activity and an increase in the ratio of respired-C to biomass
258 incorporated-C. Furthermore, the decreased biomass uptake observed with N-PAHs (B[h]Q) over time may be
259 attributed to microbial toxicity and/or oxidative stress as shown by the consistent increase in BQ to >1. (It should
260 be noted that BQ values >1 signifies oxidative stress to microbial community). McGrath et al. (1995) observed
261 that long-term exposure results in decreased soil microbial biomass. In this study, it may be because
262 microorganisms differ in their sensitivity to chemicals and prolonged N-PAHs exposure may have increased the
263 mortality of cells due to disturbance in the normal functioning, and/or gradually changed the community sizes due
264 to alterations in viability or competence (Van Beelen and Doelman, 1997; Giller et al., 1998; Anyanwu and
265 Semple 2016a; Molaei et al., 2017; Siles and Margesin, 2017).

266 Biomass uptake varied significantly over time; and the variations among chemicals were observed to be
267 consistent. This confirms the findings of Chander and Brookes (1991); Bardgett and Sagger (1994); Boucard et
268 al. (2008). Despite the variations, however, it could be concluded that soil microorganisms subjected to long term
269 N-PAH exposure, may not be able to maintain the same overall biomass as in un-contaminated soil.

270

271 *4.3 Impact of phenanthrene and its nitrogen-containing analogues on the k_{EC} coefficient*

272 The K_{EC} coefficient is related to the extractability from the soil of the microbial-C after it has been released from
273 dead fumigated cells. The k_{EC} coefficient, which is used to convert the C-flush of oxidizable organic-C to
274 microbial-C, allows for the incomplete release and extraction of microbial-C, and was obtained by calibrating
275 against alternative methods to estimate the microbial-C (Sparling et al., 1990).

276 Variation in k_{EC} coefficients (fixed and variable) was observed. The observed variations in k_{EC} coefficients after
277 fumigation are consistent with the findings in water content (Sparling and West, 1989; Ross, 1990 b) and sheep
278 dip formulation (Boucard et al., 2008). The cause is not known, however, difference in chemical amendments
279 may be attributable. This portrays the impact of contaminants on soil microbial uptake and further showed that
280 the fixed k_{EC} coefficient (0.35), fails to consider the impact of contaminated sites on soil microorganism; thus

281 overestimating (and/or underestimating) the biomass uptakes in contaminated soils (if the biomass uptakes
282 calculated with the variable k_{EC} values are considered to be more accurate). In addition, the calculated k_{EC}
283 coefficient showed that PAH and N-PAH contaminants can greatly affect the amount of substrate-C extracted by
284 0.5 M K_2SO_4 after fumigation.

285 In this present study, it could be that: (1) N-PAHs may have impacted the k_{EC} coefficients by influencing the
286 factors that modify the toxicity of contaminants in soil; such as, physico-chemical properties and/or the
287 physiological state of the microbes (Boucard et al., 2008; Siles and Margesin, 2017); or (2) The impact of N-
288 PAHs may have resulted in a possible reduction in the efficiency of chloroform disintegration of the microbial
289 cell membrane (lysis) or interference with the K_2SO_4 extraction (Sparling et al., 1990; Joergensen et al., 1995;
290 Badalocco et al., 1997; Boucard et al., 2008). However, N-PAHs bioavailability and/or differences in microbial
291 community structure between soils that vary in their sensitivity to chemical toxicity (Butler et al., 2011), could be
292 an important factor in explaining the variability in k_{EC} coefficients.

293

294 **5. Conclusions**

295 In this current study, the presence of N-PAHs resulted in alterations to soil microbial activity and functions. It
296 could be that the increased energy requirement for repair and maintenance probably was the main reason for the
297 increased respiration, but synergistic process cannot be neglected. However, the study was unable to ascertain if
298 the biomass uptakes in the chemically amended soils were characterized by either a low substrate utilization
299 efficiency or death rate; if stress increased the burden of the microbial community. Nevertheless, B[h]Q, may
300 have persistent deleterious impacts on soil microorganisms. From an ecotoxicity perspective, future investigations
301 should consider the impact of these contaminants on changes in the soil microbial community structure. Further
302 studies could also investigate the development of bacterial and fungal degrading populations within the microbial
303 community which may be able to exploit the C and N for their metabolic needs.

304

305 **Acknowledgment**

306 This work was funded by Petroleum Technology Development Fund (PTDF), Nigeria.

307

308 **References**

- 309 Anderson, T.H., Domsch, K.H., 1986. Carbon assimilation and microbial activity in soil. *Zeitschrift für*
310 *Pflanzenernährung und Bodenkunde* 149, 457-468.
- 311 Anyanwu, I.N., Semple, K.T., 2015a. Fate and behaviour of nitrogen-containing polycyclic aromatic
312 hydrocarbons in soil. *Environmental Technology and Innovation* 3, 108-120.
- 313 Anyanwu, I.N., Semple, K.T., 2015b. Biodegradation of phenanthrene-nitrogen-containing analogues in soil.
314 *Water, Air, Soil Pollution* 226, 252.
- 315 Anyanwu, I.N., Semple, K.T., 2016a. Assessment of the effects of phenanthrene and its nitrogen heterocyclic
316 analogues on microbial activity in soil. *SpringerPlus* 5, 279.
- 317 Anyanwu, I.N., Semple, K.T., 2016b. Effect of phenanthrene and its nitrogen-heterocyclic analogues aged in
318 soil on earthworm *Eisenia fetida*. *Applied Soil Ecology* 105, 151-159.
- 319 Anyanwu, I.N., Clifford, O.I., Semple, K.T., 2017. Impact of nitrogen-containing polycyclic aromatic
320 hydrocarbons on phenanthrene and benzo[a]pyrene mineralisation in soil. *Ecotoxicology and*
321 *Environmental Safety* 147C, 594–601.
- 322 Badalucco, L., De Cesare, F., Grego, S., Landi, L., Nannipieri, P., 1997. Do physical properties of soil affect
323 chloroform efficiency in lysing microbial biomass? *Soil Biology and Biochemistry* 29, 1135-1142.
- 324 Bardgett, R.D., Sgar, S., 1994. Effects of heavy metal contamination on the short-term decomposition of
325 labelled [¹⁴C] glucose in a pasture soil. *Soil Biology and Biochemistry* 26, 727-733.
- 326 Bogomolov, D.M., Chen, S.K., Parmelee, R.W., Subler, S., Edwards, C.A., 1996. An ecosystem approach to
327 soil toxicity testing: a study of copper contamination in laboratory soil microcosms. *Applied Soil*
328 *Ecology* 4, 95-105.
- 329 Bore, E.K., Apostel, C., Halicki, S., Kuzyakov, Y., Dippold, M.A., 2017. Microbial Metabolism in Soil at
330 Subzero Temperatures: Adaptation Mechanisms Revealed by Position-Specific ¹³C Labeling. *Frontiers*
331 *of Microbiology* 8, 946.
- 332 Boucard, T.K., McNeill, C., Bardgett, R.D., Paynter, C.D., Semple, K.T., 2008. The impact of synthetic
333 pyrethroid and organophosphate sheep dip formulations on microbial activity in soil. *Environmental*
334 *Pollution* 153, 207-214.

335 Brookes, P.C., 1995. The potential of microbiological properties as indicators in soil pollution monitoring. In
336 soil monitoring. Early detection and surveying of soil contamination and degradation, eds. R.
337 Schulin, A. Desaulles, R. Webster and B. von Steiger, 229-254.

338 Butler, E., Whelan, M.J., Ritz, K., Sarkrabani, R., van Egmond, R., 2011. Effects of triclosan on soil respiration.
339 Environmental Toxicology and Chemistry 30, 360-366.

340 Chander, K., Brookes P.C., 1991a. Microbial biomass dynamics during the decomposition of glucose and maize
341 in metal-contaminated soils. Soil Biology and Biochemistry 23, 917-925.

342 Chander, K., Joergensen, R.G., 2001. Decomposition of ¹⁴C glucose in two soils with different amounts of
343 heavy metal contamination. Soil Biology and Biochemistry 33, 1811-1816.

344 Dioc, K.J., Lee, P.H., Semple, K.T., 2003. Assessment of spiking procedures for the introduction of a
345 phenanthrene-LNAPL mixture into field-wet soil. Environmental Pollution 126, 399-406.

346 Doley, R., Barthakur, M., Goswami, B.S., 2017. Microbial Degradation of Aromatic hydrocarbon: Naphthalene
347 through *Nocardia* sp. RD3. International Journal of Current Microbiology and Applied Science 6,
348 1174–1181.

349 Ferraz, M.C., Mano, R.A., Oliveira, D.H., Maia, D.S.V., Silva, W.P., Savegnago, L., Lenardão, E.J., Jacob,
350 R.G., 2017. Synthesis, Antimicrobial, and Antioxidant Activities of Chalcogen-Containing Nitro-
351 Derivatives from (R)-citronellal. Medicines 4, 39.

352 Fließbach, A., Martens, R., Reber, H.H., 1994. Soil microbial biomass and microbial activity in soils treated
353 with heavy metal contaminated sewage sludge. Soil Biology and Biochemistry 26, 1201-1205.

354 Fournier, J.C., Horatallah, A., Collu, T., Froncek, B., 1992. Labelling of microbial biomass with radioactive as
355 a means to estimate pesticide effects in soil. Science of the Total Environment 123/124, 325-332.

356 Frostergård, Å., Tunlid, A., Bååth, E., 1993. Changes in microbial community structure during long-term
357 incubation in two soils experimentally contaminated with metals. Soil Biology and Biochemistry 28,
358 55-63.

359 Giller, K.E., Witter, E., McGrath, S.P., 1998. Toxicity of heavy metals to microorganisms and microbial
360 processes in agricultural soils: a review. Soil Biology and Biochemistry 30, 1389-1414.

361 Granato, M.Q., Gonçalves, Dd.S., Seabra, S.H., McCann, M., Devereux, M., dos Santos, A.L.S., Kneipp,
362 L.F., 2017. 1,10-Phenanthroline-5,6-Dione– Based Compounds Are Effective in Disturbing Crucial
363 Physiological Events of *Phialophora verrucosa*. *Frontiers of Microbiology* 8, 76.

364 Gunina, A., Dippold, M., Glaser, B., Kuzyakov, Y., 2017. Turnover of microbial groups and cell components in
365 soil: ¹³C analysis of cellular biomarkers. *Biogeosciences* 14, 271-283.

366 Hattori, H., 1992. Influence of heavy metals on soil microbial activities. *Soil Science and Plant Nutrition* 38,
367 93-100.

368 IARC, 2012. In: IARC monographs on the evaluation of carcinogenic risk of chemicals to humans. Agents
369 classified by IARC Monographs. World Health Organization, 1-105.

370 Joergensen, R.G., Schmaedeke, F., Windhorst, K., Meyer, B., 1995. Biomass and activity of microorganisms in
371 a fuel oil contaminated soil. *Soil Biology and Biochemistry* 27, 1137-1143.

372 Kaiser, J.P., Feng, Y., Bollag, J.M., 1996. Microbial metabolism of pyridine, quinoline, acridine and their
373 derivatives under aerobic and anaerobic conditions. *Microbiological Reviews* 60, 483-498.

374 Killham, K., 1985. A physiological determination of the impact of environmental stress on the activity of
375 microbial biomass. *Environmental Pollution* 38, 283-294.

376 Knight, B., McGrath, S.P., Chaudri, A.M., 1997. Biomass carbon measurements and substrate utilization
377 patterns of microbial populations from soil amended with cadmium, copper or zinc. *Applied and*
378 *Environmental Microbiology* 63, 39-43

379 Leite, M.F.A., Pan, Y., Bloem, J., ten Berge, H., Kuramae, E.E., 2017. Organic nitrogen rearranges both
380 structure and activity of the soil-borne microbial seedbank. *Scientific Report* 7, 42634.

381 McGrath, S.P., Chaudri, A.M., Giller, K.E., 1995. Long-term effects of land application of sewage sludge: soils,
382 microorganisms and plants. *Journal of Industrial Microbiology* 14, 94-104.

383 Mehnaz, K., Keitel, C., Dijkstra, F., 2017. Effect of phosphorus and carbon addition on gross nitrogen
384 mineralization, microbial respiration and N₂O emission in a grassland soil. *Geophysical Research*
385 *Abstracts* 19, EGU 2017-6614, EGU General Assembly.

386 Molaei, A., Lakzian, A., Haghnia, G., Astaraei, A., Rasouli-Sadaghiani, M., Ceccherini, T.M., Datta, R., 2017.
387 Assessment of some cultural experimental methods to study the effects of antibiotics on microbial
388 activities in a soil: An incubation study. *PLoS ONE* 12 (7), e0180663.

389 Mooshammer, M., Hofhansl, F., Frank, A.H., Wanek, W., Hämmerle, I., Leitner, S., Schenecker, J., Wild, B.,
390 Watzka, M., Keiblinger, K.M., Zechmeister-Boltenstern, S., Richter, A., 2017. Decoupling of microbial
391 carbon, nitrogen, and phosphorus cycling in response to extreme temperature events. *Science Advances*
392 3, e1602781.

393 Nakamoto, T., Wakahara, S., 2004. Development of substrate induced respiration (SIR) method combined with
394 selective inhibition for estimating fungal and bacterial biomass in humic andosols. *Plant Production*
395 *Science* 7, 70-76.

396 Rajasekhar, N., Sasikala, C., Ramana, C.V., 2000. Toxicity of N-containing heterocyclic aromatic compounds
397 and their utilization for growth by few purple non-sulfur bacteria. *Bulletin of Environmental*
398 *Contamination and Toxicology* 65, 375 – 382.

399 Reid, B.J., MacLeod, C.J.A., Lee, P.L., Morriss, A.W.J., Stokes, J.D., Semple, K.T., 2001. A simple ¹⁴C-
400 respirometric method for assessing microbial catabolic potential and contaminant bioavailability.
401 *FEMS Microbiology Letters* 196, 141-146.

402 Ross, D.J., 1990b. Estimation of soil microbial C by a fumigation-extraction method: influence of seasons, soils
403 and calibration with the fumigation-incubation procedure. *Soil Biology and Biochemistry* 22, 295-300.

404 Semple, K.T., Doick, K.J., Wick, L.Y., Harms, H., 2007. Microbial interactions with organic contaminants in
405 soil: definitions, processes and measurement. *Environmental Pollution* 150, 166-176

406 Siles, J.A., Margesin, R., 2017. Seasonal soil microbial responses are limited to changes in functionality at two
407 Alpine forest sites differing in altitude and vegetation. *Scientific Reports* 7, 2204.

408 Sparling, G.P., West, A.E., 1988. Modifications to the fumigation-extraction of soil microbial C and N.
409 *Communications in Soil Science and Plant Analyses* 19, 327-344.

410 Sparling, G.P., West, A.E., Feltham, C.W., Reynolds, J., 1990. Estimation of soil microbial C by a fumigation-
411 extraction method: use on soils of high organic matter content, and a reassessment of the k_{ec} -factor.
412 *Soil Biology and Biochemistry* 22, 301-307.

413 Sun, Q., Meyer, W.S., Koerber, G.R., Marschner, P., 2017. Response of microbial activity to labile C addition
414 in sandy soil from semi-arid woodland is influenced by vegetation patch and wildfire. *Journal of Soil*
415 *Science and Plant Nutrition* 17, 62-73.

416 Van Beelen, P., Doelman, P., 1997. Significance and application of microbial toxicity tests in assessing
417 ecotoxicological risks of contaminants in soils and sediments. *Chemosphere* 34, 455-499.

418 Vance, E.D., Brookes, P.C., Jenkinson, D.S., 1987. An extraction method for measuring soil microbial biomass
419 in soil. *Soil Biology and Biochemistry* 19, 703-707.

420 Witter, E., Dahlin, S., 1995. Microbial utilization of [U-¹⁴C]-labelled straw and [U-¹³C]-labelled glucose in soils
421 of contrasting pH and metal status. *Soil Biology and Biochemistry* 27, 1507-1516.

422 Xu, W., Cai, Y.P., Yang, Z.F., Yin, X.A., Tan, Q., 2017. Microbial nitrification, denitrification and respiration
423 in the leached cinnamon soil of the upper basin of Miyun Reservoir. *Scientific Report* 7, 42032.

424 Zhu, F., Storey, S., Ashaari, M. M., Clipson, N., Doyle, E., 2017. Benzo(a)pyrene degradation and microbial
425 community responses in composted soil. *Environmental Science and Pollution Research* 24, 5404–
426 5414.

427

428

429

430

431

432

433

434

435

436

437

438

439

440

441

442

443

444 **Table 1.** Extractability (%) of phenanthrene and its N-PAHs from soil over time

Chemical	Initial chemical conc (mg/kg)	Mean chemicals extracted (mg/kg)			
		Time (d)			
		0	30	60	90
Phen	100	78.00± 8.00	42.10±5.70	11.30 ± 3.10	3.10 ± 0.30
1,7-Phen	100	59.30± 9.00	54.40±12.50	49.60 ± 4.00	38.00 ±8.00
B[h]Q	100	59.20± 9.00	89.40 ± 7.00	64.10 ± 6.00	58.70 ± 4.20
4,7-Phen	100	91.90 ± 4.20	41.70 ± 6.40	29.30 ± 4.00	29.60 ± 3.90

445 Source: Anyanwu and Semple (2015b).

Table 2. Distribution of ¹⁴C-glucose in soils amended with 100 mg/kg single and binary mixtures of phenanthrene and its N-PAH analogues after 5 d

Chemical	Time (d)	Fastest rates (% h ⁻¹)	Respired ¹⁴ CO ₂ (%)	C-flush ^a	Biomass uptake ^b (fixed <i>k_{EC}</i> = 0.35)	Biophysical quotient ^c	Biomass uptake ^d (variable <i>k_{EC}</i>)	Biophysical quotient ^e	Calculated <i>k_{EC}</i> ^f
Control	1	0.46 ± 0.08	42.62 ± 2.20	2.59 ± 0.33	7.41 ± 0.94**	5.75 ± 2.32*	53.42 ± 6.81**	0.79 ± 0.32	0.048 ± 0.00
	30	0.25 ± 0.01	35.07 ± 0.45*	3.63 ± 0.57	10.36 ± 1.63**	3.38 ± 0.27	54.02 ± 8.52**	0.64 ± 0.05	0.067 ± 0.01
	60	0.18 ± 0.03	31.07 ± 2.00*	1.07 ± 0.31	3.06 ± 0.87**	10.15 ± 2.28	63.45 ± 18.1**	0.48 ± 0.11	0.016 ± 0.00
	90	0.18 ± 0.00	28.42 ± 1.26*	6.97 ± 1.28	19.92 ± 3.65**	1.42 ± 0.34	46.67 ± 8.55**	0.60 ± 0.14	0.149 ± 0.02
1,10-Phen	1	0.43 ± 0.02	38.81 ± 2.84	1.49 ± 0.27	4.24 ± 0.77**	9.14 ± 0.00*	58.91 ± 10.78**	0.65 ± 0.26	0.025 ± 0.00
	30	0.29 ± 0.02	36.98 ± 0.81*	2.96 ± 0.02	8.46 ± 0.06**	4.72 ± 0.03	51.90 ± 0.39**	0.71 ± 2.02	0.057 ± 0.00
	60	0.23 ± 0.03	31.83 ± 1.55*	0.77 ± 0.36	2.21 ± 1.04**	14.42 ± 0.00	61.00 ± 23.82**	0.52 ± 0.05	0.012 ± 0.00
	90	0.15 ± 0.03	23.13 ± 1.70*	2.35 ± 0.04	6.72 ± 0.10**	3.44 ± 0.08	56.01 ± 0.85**	0.41 ± 1.98	0.042 ± 0.00
1,7-Phen	1	0.43 ± 0.02	41.77 ± 1.99	2.27 ± 0.08	6.49 ± 0.22**	6.43 ± 0.01*	54.21 ± 1.87**	0.76 ± 1.06	0.041 ± 0.00
	30	0.30 ± 0.03	36.90 ± 1.27*	2.82 ± 0.32	8.05 ± 0.90**	4.58 ± 0.00	51.35 ± 5.79**	0.71 ± 0.21	0.054 ± 0.00
	60	0.21 ± 0.02	36.83 ± 2.06*	1.67 ± 0.74	4.77 ± 2.11**	7.72 ± 0.00	53.72 ± 23.83**	0.68 ± 0.08	0.031 ± 0.01
	90	0.50 ± 0.07	34.74 ± 2.43*	5.28 ± 1.22	15.08 ± 3.47**	2.30 ± 0.00	37.49 ± 8.63**	0.92 ± 0.28	0.140 ± 0.03
4,7-Phen	1	0.26 ± 0.03	33.51 ± 2.93	0.31 ± 0.06	0.90 ± 0.16**	37.24 ± 0.03*	62.45 ± 11.22**	0.53 ± 0.26	0.005 ± 0.00
	30	0.22 ± 0.03	35.31 ± 1.78*	3.03 ± 0.44	8.65 ± 1.25**	4.08 ± 0.00	56.11 ± 8.16**	0.62 ± 0.21	0.053 ± 0.01
	60	0.20 ± 0.00	33.57 ± 0.27*	1.27 ± 0.16	3.64 ± 0.44**	9.22 ± 0.00	57.71 ± 7.10**	0.58 ± 0.03	0.022 ± 0.00
	90	0.17 ± 0.04	27.31 ± 2.35*	2.21 ± 0.62	6.31 ± 1.77**	4.32 ± 0.00	39.41 ± 11.08**	0.69 ± 0.21	0.056 ± 0.01
B[h]Q	1	0.37 ± 0.05	39.61 ± 1.83	0.53 ± 0.09	1.52 ± 0.25**	26.01 ± 0.01*	56.72 ± 9.43**	0.69 ± 0.19	0.009 ± 0.00
	30	0.28 ± 0.03	42.52 ± 2.73*	3.06 ± 0.61	8.75 ± 1.72**	4.85 ± 0.00	46.12 ± 9.11**	0.92 ± 0.29	0.066 ± 0.01
	60	0.34 ± 0.05	44.62 ± 2.30*	1.13 ± 0.47	3.22 ± 1.33**	13.86 ± 0.00	41.51 ± 17.26**	1.07 ± 0.13	0.027 ± 0.01
	90	0.70 ± 0.05	37.01 ± 2.25*	2.50 ± 0.23	7.16 ± 0.65**	5.17 ± 0.01	34.03 ± 3.11**	1.08 ± 0.72	0.073 ± 0.00
Phen	1	0.35 ± 0.02	36.42 ± 1.21	0.23 ± 0.00	0.66 ± 0.13**	54.76 ± 0.17*	60.99 ± 1.27**	0.59 ± 0.95	0.003 ± 0.00
	30	0.27 ± 0.01	37.17 ± 0.92*	3.32 ± 0.18	9.48 ± 0.51**	3.91 ± 0.00	42.74 ± 2.31**	0.86 ± 0.39	0.077 ± 0.00
	60	0.23 ± 0.01	32.01 ± 2.47*	1.39 ± 1.03	3.96 ± 2.94**	8.07 ± 0.00	58.41 ± 43.33**	0.54 ± 0.05	0.023 ± 0.01
	90	0.19 ± 0.01	26.78 ± 0.21*	2.19 ± 0.35	6.27 ± 1.00**	4.27 ± 0.00	51.48 ± 8.24**	0.52 ± 0.02	0.042 ± 0.00
1,10-Phen+Phen	1	0.41 ± 0.07	41.82 ± 3.49	0.64 ± 0.08	1.82 ± 0.21**	22.02 ± 0.03*	50.59 ± 6.09**	0.82 ± 0.57	0.012 ± 0.00
	30	0.26 ± 0.04	36.20 ± 2.80*	1.53 ± 0.56	4.39 ± 1.58**	8.25 ± 0.00	55.02 ± 19.92**	0.65 ± 0.14	0.027 ± 0.01

	60	0.25 ± 0.02	36.33 ± 1.44*	1.13 ± 1.16	3.73 ± 3.31**	9.74 ± 0.00	53.70 ± 47.77**	0.67 ± 0.03	0.024 ± 0.02
	90	0.15 ± 0.03	23.87 ± 0.87*	2.47 ± 0.42	7.05 ± 1.20**	3.38 ± 0.00	50.07 ± 8.55**	0.47 ± 0.10	0.049 ± 0.00
1,7- Phen+Phen	1	0.44 ± 0.10	42.72 ± 2.11	0.77 ± 0.11	2.19 ± 0.31**	19.51 ± 0.01*	55.99 ± 8.12**	0.79 ± 0.25	0.013 ± 0.00
	30	0.26 ± 0.04	38.02 ± 1.10*	2.56 ± 0.18	7.32 ± 0.51**	5.19 ± 0.00	52.74 ± 3.70**	0.72 ± 0.29	0.048 ± 0.00
	60	0.25 ± 0.03	36.41 ± 2.38*	1.44 ± 0.61	4.10 ± 1.74**	8.87 ± 0.00	53.44 ± 22.77**	0.68 ± 0.10	0.026 ± 0.01
	90	0.40 ± 0.04	31.99 ± 2.22*	2.50 ± 0.12	7.16 ± 0.33**	4.47 ± 0.03	41.45 ± 1.93**	0.77 ± 1.14	0.060 ± 0.00
4,7- Phen+Phen	1	0.53 ± 0.02	44.20 ± 0.72	0.52 ± 0.08	1.47 ± 0.23**	29.98 ± 0.00*	59.37 ± 9.45**	0.74 ± 0.07	0.008 ± 0.00
	30	0.27 ± 0.01	39.88 ± 1.00*	3.30 ± 0.28	9.44 ± 0.79**	4.22 ± 0.00	53.39 ± 4.49**	0.74 ± 0.22	0.061 ± 0.00
	60	0.21 ± 0.03	35.89 ± 1.93*	1.89 ± 1.19	5.41 ± 3.39**	6.63 ± 0.00	57.95 ± 36.37**	0.61 ± 0.05	0.032 ± 0.02
	90	0.28 ± 0.05	31.21 ± 1.62*	3.02 ± 0.61	8.63 ± 1.73**	3.61 ± 0.00	52.73 ± 10.57**	0.59 ± 0.15	0.057 ± 0.01
BhQ+Phen	1	0.43 ± 0.04	43.89 ± 1.91	0.39 ± 0.04	1.12 ± 0.11**	39.21 ± 0.03*	54.64 ± 5.66**	0.80 ± 0.33	0.007 ± 0.00
	30	0.31 ± 0.04	42.91 ± 1.73*	2.38 ± 0.22	6.79 ± 0.61**	6.31 ± 0.00	49.32 ± 4.48**	0.86 ± 0.38	0.048 ± 0.00
	60	0.37 ± 0.03	43.00 ± 2.00*	2.42 ± 0.14	6.92 ± 0.39**	6.21 ± 0.01	50.66 ± 2.92**	0.84 ± 0.68	0.047 ± 0.00
	90	0.34 ± 0.05	32.63 ± 2.34*	3.48 ± 0.43	9.93 ± 1.22**	3.28 ± 0.00	44.93 ± 5.55**	0.72 ± 0.24	0.077 ± 0.00

^a C-flush = ¹⁴C-activity in fumigated soil – ¹⁴C- activity in un-fumigated soil

^b ¹⁴C-microbial biomass = ¹⁴C-flush/fixed k_{EC} 0.35

^c BQ = ¹⁴C respired/¹⁴C in biomass (using fixed k_{EC} 0.35)

^d ¹⁴C-microbial biomass = ¹⁴C-flush/variable k_{EC}

^e BQ = ¹⁴C respired/¹⁴C in biomass (using variable k_{EC})

^f k_{EC} = (¹⁴C-flush)/(initial ¹⁴C-activity added – ¹⁴C respired – ¹⁴C activity in un-fumigated soil)

Conc = 100 mg/kg

n = 3

* = p < 0.05

** = p < 0.001




Fig. 1. Microbial mineralisation of ¹⁴C-glucose in soils amended with phenanthrene and its N-PAH analogues (single amendments). The 1–90 d incubation graphs shows: control (●), 1,10-Phen (○), 1,7-Phen (▼), 4,7-Phen (△), B[h]Q (■) and Phen (□). Conc = 100 mg/kg.




Fig. 2. Microbial mineralisation of ^{14}C -glucose in soils amended with phenanthrene and its N-PAH analogues (binary mixtures). The 1–90 d incubation graphs shows: control (●), 1,10-Phen + Phen (○), 1,7-Phen + Phen (▼), 4,7-Phen + Phen (Δ) and B[h]Q + Phen (■). Conc = 100 mg/kg.