

# 1 Genomic dissection of bipolar disorder and schizophrenia including 28 subphenotypes

## 3 Primary Author List

4 Bipolar Disorder and Schizophrenia Working Group of the Psychiatric Genomics Consortium

## 6 Secondary Author List

7 Douglas M Ruderfer, Stephan Ripke, Andrew McQuillin, James Boocock, Eli A Stahl, Jennifer M Whitehead  
8 Pavlides, Niamh Mullins, Alexander W Charney, Anil P S Ori, Loes M Olde Loohuis, Enrico Domenici, Arianna Di  
9 Florio, Sergi Papiol, Janos L. Kalman, Vassily Trubetskoy, Rolf Adolfsson, Ingrid Agartz, Esben Agerbo,  
10 Huda Akil, Diego Albani, Margot Albus, Martin Alda, Madeline Alexander, Ney Alliey-Rodriguez, Thomas D Als,  
11 Farooq Amin, Adebayo Anjorin, Maria J Arranz, Swapnil Awasthi, Silviu A Bacanu, Judith A Badner, Marie  
12 Baekvad-Hansen, Steven Bakker, Gavin Band, Jack D Barchas, Ines Barroso, Nicholas Bass, Michael Bauer,  
13 Bernhard T Baune, Martin Begemann, Celine Bellenguez, Richard A Belliveau Jr, Frank Bellivier, Stephan Bender,  
14 Judit Bene, Sarah E Bergen, Wade H Berrettini, Elizabeth Bevilacqua, Joanna M Biernacka, Tim B Bigdeli, Donald  
15 W Black, Hannah Blackburn, Jenefer M Blackwell, Douglas HR Blackwood, Carsten Bocker Pedersen, Michael  
16 Boehnke, Marco Boks, Anders D Borglum, Elvira Bramon, Gerome Breen, Matthew A Brown, Richard Bruggeman,  
17 Nancy G Buccola, Randy L Buckner, Monika Budde, Brendan Bulik-Sullivan, Suzannah J Bumpstead, William  
18 Bunney, Margit Burmeister, Joseph D Buxbaum, Jonas Bybjerg-Grauholm, William Byerley, Wiepke Cahn,  
19 Guiqing Cai, Murray J Cairns, Dominique Campion, Rita M Cantor, Vaughan J Carr, Noa Carrera, Juan P Casas,  
20 Miquel Casas, Stanley V Catts, Pablo Cervantes, Kimberley D Chambert, Raymond CK Chan, Eric YH Chen,  
21 Ronald YL Chen, Wei Cheng, Eric FC Cheung, Siow Ann Chong, Toni-Kim Clarke, C Robert Cloninger, David  
22 Cohen, Nadine Cohen, Jonathan R I Coleman, David A Collier, Paul Cormican, William Coryell, Nicholas  
23 Craddock, David W Craig, Benedicto Crespo-Facorro, James J Crowley, Cristiana Cruceanu, David Curtis, Piotr M  
24 Czerski, Anders M Dale, Mark J Daly, Udo Dannlowski, Ariel Darvasi, Michael Davidson, Kenneth L Davis,  
25 Christiaan A de Leeuw, Franziska Degenhardt, Jurgen Del Favero, Lynn E DeLisi, Panos Deloukas, Ditte Demontis,  
26 J Raymond DePaulo, Marta di Forti, Dimitris Dikeos, Timothy Dinan, Srdjan Djurovic, Amanda L Dobbyn, Peter  
27 Donnelly, Gary Donohoe, Elodie Drapeau, Serge Dronov, Jubao Duan, Frank Dudbridge, Audrey Duncanson,  
28 Howard Edenberg, Sarah Edkins, Hannelore Ehrenreich, Peter Eichhammer, Torbjorn Elvsashagen, Johan Eriksson,  
29 Valentina Escott-Price, Tonu Esko, Laurent Essioux, Bruno Etain, Chun Chieh Fan, Kai-How Farh, Marttilias S  
30 Farrell, Matthew Flickinger, Tatiana M Foroud, Liz Forty, Josef Frank, Lude Franke, Christine Fraser, Robert  
31 Freedman, Colin Freeman, Nelson B Freimer, Joseph I Friedman, Menachem Fromer, Mark A Frye, Janice M  
32 Fullerton, Katrin Gade, Julie Garnham, Helena A Gaspar, Pablo V Gejman, Giulio Genovese, Lyudmila Georgieva,  
33 Claudia Giambartolomei, Eleni Giannoulatou, Ina Giegling, Michael Gill, Matthew Gillman, Marianne Giortz  
34 Pedersen, Paola Giusti-Rodriguez, Stephanie Godard, Fernando Goes, Jacqueline I Goldstein, Srihari Gopal, Scott D  
35 Gordon, Katherine Gordon-Smith, Jacob Gratten, Emma Gray, Elaine K Green, Melissa J Green, Tiffany A  
36 Greenwood, Maria Grigoriou-Serbanescu, Jakob Grove, Weihua Guan, Hugh Gurling, Jose Guzman Parra, Rhian  
37 Gwilliam, Lieuwe de Haan, Jeremy Hall, Mei-Hua Hall, Christian Hammer, Naomi Hammond, Marian L Hamshere,  
38 Mark Hansen, Thomas Hansen, Vahram Haroutunian, Annette M Hartmann, Joanna Hauser, Martin Hautzinger, Urs  
39 Heilbronner, Garrett Hellenthal, Frans A Henskens, Stefan Herms, Maria Hipolito, Joel N Hirschhorn, Per  
40 Hoffmann, Mads V Hollegaard, David M Hougaard, Hailiang Huang, Laura Huckins, Christina M Hultman, Sarah E  
41 Hunt, Masashi Ikeda, Nakao Iwata, Conrad Iyegbe, Assen V Jablensky, Stephane Jamain, Janusz Jankowski,  
42 Alagurevathi Jayakumar, Inge Joa, Ian Jones, Lisa A Jones, Erik G Jonsson, Antonio Julia, Anders Jureus, Anna K  
43 Kahler, Rene S Kahn, Luba Kalaydjieva, Radhika Kandaswamy, Sena Karachanak-Yankova, Juha Karjalainen,  
44 Robert Karlsson, David Kavanagh, Matthew C Keller, Brian J Kelly, John Kelsoe, James L Kennedy, Andrey  
45 Khrunin, Yunjung Kim, George Kirov, Sarah Kittel-Schneider, Janis Klovins, Jo Knight, Sarah V Knott, James A  
46 Knowles, Manolis Kogevinas, Bettina Konte, Eugenia Kravariti, Vaidutis Kucinskas, Zita Ausrele Kucinskiene,  
47 Ralph Kupka, Hana Kuzelova-Ptackova, Mikael Landen, Cordelia Langford, Claudine Laurent, Jacob Lawrence,  
48 Stephen Lawrie, William B Lawson, Markus Leber, Marion Leboyer, Phil H Lee, Jimmy Lee Chee Keong, Sophie E  
49 Legge, Todd Lencz, Bernard Lerer, Douglas F Levinson, Shawn E Levy, Cathryn M Lewis, Jun Z Li, Miaoxin Li,  
50 Qingqin S Li, Tao Li, Kung-Yee Liang, Jennifer Liddle, Jeffrey Lieberman, Svetlana Limborska, Kuang Lin, Don H  
51 Linszen, Jolanta Lissowska, Chunyu Liu, Jianjun Liu, Jouko Lonnqvist, Carmel M Loughland, Jan Lubinski,  
52 Susanne Lucae, Milan Macek Jr, Donald J MacIntyre, Patrik KE Magnusson, Brion S Maher, Pamela B Mahon,  
53 Wolfgang Maier, Anil K Malhotra, Jacques Mallet, Ulrik F Malt, Hugh S Markus, Sara Marsal, Nicholas G Martin,  
54 Ignacio Mata, Christopher G Mathew, Manuel Mattheisen, Morten Mattingsdal, Fermin Mayoral, Owen T McCann,  
55 Robert W McCarley, Steven A McCarroll, Mark I McCarthy, Colm McDonald, Susan L McElroy, Peter McGuffin,

56 Melvin G McInnis, Andrew M McIntosh, James D McKay, Francis J McMahon, Helena Medeiros, Sarah E  
57 Medland, Sandra Meier, Carin J Meijer, Bela Melegh, Ingrid Melle, Fan Meng, Raquelle I Mesholam-Gately,  
58 Andres Metspalu, Patricia T Michie, Lili Milani, Vihra Milanova, Philip B Mitchell, Younes Mokrab, Grant W  
59 Montgomery, Jennifer L Moran, Gunnar Morken, Derek W Morris, Ole Mors, Preben B Mortensen, Bryan J Mowry,  
60 Thomas W Mühleisen, Bertram Müller-Myhsok, Kieran C Murphy, Robin M Murray, Richard M Myers, Inez Myin-  
61 Germeys, Benjamin M Neale, Mari Nelis, Igor Nenadic, Deborah A Nertney, Gerald Nestadt, Kristin K Nicodemus,  
62 Caroline M Nievergelt, Liene Nikitina-Zake, Vishwajit Nimgaonkar, Laura Nisenbaum, Merete Nordentoft, Annelie  
63 Nordin, Markus M Nöthen, Evaristus A Nwulia, Eadbhard O'Callaghan, Claire O'Donovan, Colm O'Dushlaine, F  
64 Anthony O'Neill, Ketil J Oedegaard, Sang-Yun Oh, Ann Olincy, Line Olsen, Lilijana Oruc, Jim Van Os, Michael J  
65 Owen, Sara A Paciga, Colin N A Palmer, Aarno Palotie, Christos Pantelis, George N Papadimitriou, Elena  
66 Parkhomenko, Carlos Pato, Michele T Pato, Tiina Paunio, Richard Pearson, Psychosis Endophenotypes International  
67 Consortium, Diana O Perkins, Roy H Perlis, Amy Perry, Tune H Pers, Tracey L Petryshen, Andrea Pfenning, Marco  
68 Picchioni, Olli Pietilainen, Jonathan Pimm, Matti Pirinen, Robert Plomin, Andrew J Pocklington, Danielle  
69 Posthuma, James B Potash, Simon C Potter, John Powell, Alkes Price, Ann E Pulver, Shaun M Purcell, Digby  
70 Queded, Josep Antoni Ramos-Quiroga, Henrik B Rasmussen, Anna Rautanen, Radhi Ravindrarajah, Eline J Regeer,  
71 Abraham Reichenberg, Andreas Reif, Mark A Reimers, Marta Ribases, John P Rice, Alexander L Richards,  
72 Michelle Ricketts, Brien P Riley, Fabio Rivas, Margarita Rivera, Joshua L Roffman, Guy A Rouleau, Panos  
73 Roussos, Dan Rujescu, Veikko Salomaa, Cristina Sanchez-Mora, Alan R Sanders, Stephen J Sawcer, Ulrich Schall,  
74 Alan F Schatzberg, William A Scheftner, Peter R Schofield, Nicholas J Schork, Sibylle G Schwab, Edward M  
75 Scolnick, Laura J Scott, Rodney J Scott, Larry J Seidman, Alessandro Serretti, Pak C Sham, Cynthia Shannon  
76 Weickert, Tatyana Shehktman, Jianxin Shi, Paul D Shilling, Engilbert Sigurdsson, Jeremy M Silverman, Kang Sim,  
77 Claire Slaney, Petr Slominsky, Olav B Smeland, Jordan W Smoller, Hon-Cheong So, Janet L Sobell, Erik  
78 Soderman, Christine Soholm Hansen, Chris C A Spencer, Anne T Spijker, David St Clair, Hreinn Stefansson, Kari  
79 Stefansson, Stacy Steinberg, Elisabeth Stogmann, Eystein Stordal, Amy Strange, Richard E Straub, John S Strauss,  
80 Fabian Streit, Eric Strengman, Jana Strohmaier, T Scott Stroup, Zhan Su, Mythily Subramaniam, Jaana Suvisaari,  
81 Dragan M Svrakic, Jin P Szatkiewicz, Szabolcs Szelinger, Avazeh Tashakkori-Ghanbaria, Srinivas Thirumalai,  
82 Robert C Thompson, Thorgeir E Thorgeirsson, Draga Toncheva, Paul A Tooney, Sarah Tosato, Timothea  
83 Toulopoulou, Richard C Trembath, Jens Treutlein, Vassily Trubetsky, Gustavo Turecki, Arne E Vaaler, Helmut  
84 Vedder, Eduard Vieta, John Vincent, Peter M Visscher, Ananth C Viswanathan, Damjan Vukcevic, John  
85 Waddington, Matthew Waller, Dermot Walsh, Muriel Walshe, James TR Walters, Dai Wang, Qiang Wang, Weiqing  
86 Wang, Yunpeng Wang, Stanley J Watson, Bradley T Webb, Thomas W Weickert, Daniel R Weinberger, Matthias  
87 Weisbrod, Mark Weiser, Thomas Werge, Paul Weston, Pamela Whittaker, Sara Widaa, Durk Wiersma, Dieter B  
88 Wildenauer, Nigel M Williams, Stephanie Williams, Stephanie H Witt, Aaron R Wolen, Emily HM Wong, Nicholas  
89 W Wood, Brandon K Wormley, Wellcome Trust Case-Control Consortium , Jing Qin Wu, Simon Xi, Wei Xu, Allan  
90 H Young, Clement C Zai, Peter Zandi, Peng Zhang, Xuebin Zheng, Fritz Zimprich, Sebastian Zollner, Aiden  
91 Corvin, Ayman H Fanous, Sven Cichon, Marcella Rietschel, Elliot S Gershon, Thomas G Schulze, Alfredo B  
92 Cuellar-Barboza, Andreas J Forstner, Peter A Holmans, John I Nurnberger, Ole A Andreassen, S Hong Lee, Michael  
93 C O'Donovan, Patrick F Sullivan, Roel A Ophoff, Naomi R Wray, Pamela Sklar\*#, Kenneth S Kendler\*

94  
95 \*Authors contributed equally to this work

96 #Deceased

97

98 **Corresponding author and lead contact:** Douglas M. Ruderfer ([douglas.ruderfer@vanderbilt.edu](mailto:douglas.ruderfer@vanderbilt.edu))

99

100

101 **Summary**

102 Schizophrenia and bipolar disorder are two distinct diagnoses that share symptomology.  
103 Understanding the genetic factors contributing to the shared and disorder-specific symptoms will  
104 be crucial for improving diagnosis and treatment. In genetic data consisting of 53,555 cases  
105 (20,129 BD, 33,426 SCZ) and 54,065 controls, we identified 114 genome-wide significant loci  
106 implicating synaptic and neuronal pathways shared between disorders. Comparing SCZ to BD  
107 (23,585 SCZ, 15,270 BD) identified four genomic regions including one with disorder-  
108 independent causal variants and potassium ion response genes as contributing to differences in  
109 biology between the disorders. Polygenic risk score (PRS) analyses identified several significant  
110 correlations within case-only phenotypes including SCZ PRS with psychotic features and age of  
111 onset in BD. For the first time, we discover specific loci that distinguish between BD and SCZ  
112 and identify polygenic components underlying multiple symptom dimensions. These results  
113 point to the utility of genetics to inform symptomology and potentially treatment.

114

115

116 **Introduction**

117 Bipolar disorder (BD) and schizophrenia (SCZ) are severe psychiatric disorders and among the  
118 leading causes of disability worldwide(Whiteford et al., 2013). Both disorders have significant  
119 genetic components with heritability estimates ranging from 60-80%(Nöthen et al., 2010).  
120 Recent genetic and epidemiological studies have demonstrated substantial overlap between these  
121 two disorders with a genetic correlation from common variation near 0.6-0.7(Cross-Disorder  
122 Group of the Psychiatric Genomics Consortium, 2013) and high relative risks (RR) among  
123 relatives of both BD and SCZ patients (RRs for parent/offspring: BD/BD: 6.4, BD/SCZ: 2.4;

124 SCZ/BD: 5.2, SCZ/SCZ: 9.9)(Lichtenstein et al., 2009). Despite shared genetics and  
125 symptomology, the current diagnostic systems(“Diagnostic and Statistical Manual of Mental  
126 Disorders | DSM Library,” n.d.)(“WHO | International Classification of Diseases,” n.d.) adhere  
127 to historical distinctions from the late 19<sup>th</sup> century and represent BD and SCZ as independent  
128 categorical entities differentiated on the basis of their clinical presentation, with BD  
129 characterized by predominant mood symptoms, mood-congruent delusions and an episodic  
130 disease course and SCZ considered a prototypical psychotic disorder. Identifying genetic  
131 components contributing to both disorders provides insight into the biology underlying the  
132 shared symptoms of the disorders.

133 While the shared genetic component is substantial, studies to date have also implicated genetic  
134 architecture differences between these two disorders(Curtis et al., 2011; Ruderfer et al., 2014). A  
135 polygenic risk score created from a case only SCZ vs BD genome-wide association study  
136 (GWAS) significantly correlated with SCZ or BD diagnosis in an independent sample(Ruderfer  
137 et al., 2014), providing the first evidence that differences between the disorders also have a  
138 genetic basis. An enrichment of rare, moderate to highly penetrant copy number variants (CNVs)  
139 and *de novo* CNVs are seen in SCZ patients(CNV and Schizophrenia Working Groups of the  
140 Psychiatric Genomics Consortium, 2017; Gulsuner and McClellan, 2015; Kirov et al., 2012;  
141 Stone et al., 2008; Szatkiewicz et al., 2014), while, the involvement of CNVs in BD is less  
142 clear(Green et al., 2016). Although the role of *de novo* single nucleotide variants in BD and SCZ  
143 has been investigated in only a handful of studies, enrichment in pathways associated with the  
144 postsynaptic density has been reported for SCZ, but not BD(Fromer et al., 2014; Kataoka et al.,  
145 2016). Identifying disorder-specific variants and quantifying the contribution of genetic variation  
146 to specific symptom dimensions remain important open questions. Characterizing these genetic

147 differences will facilitate an understanding of the dimensions of the disorders instead of the  
148 dichotomous diagnosis. For example, we have shown that SCZ patients with greater manic  
149 symptoms have higher polygenic risk for BD(Ruderfer et al., 2014). These findings demonstrate  
150 shared genetic underpinnings for symptoms across disorders and may enable us to characterize  
151 patients by genetic liability to symptom dimensions thereby informing disease course and  
152 treatment.

153 Here, we utilize large collections of genotyped samples for BD and SCZ along with clinically-  
154 relevant measures identifying 28 subphenotypes to address three questions: 1) Are there specific  
155 variants, genes or pathways that are either shared by, or differentiate BD and SCZ? 2) Are the  
156 shared symptoms between these disorders driven by the same underlying genetic profiles? and 3)  
157 Can we demonstrate independent genetic signatures for subphenotypes within these disorders?

158

## 159 **Results**

160

### 161 **Shared genetic contribution to BD and SCZ**

162 We performed association analysis of BD and SCZ combined into a single phenotype, totaling  
163 53,555 cases (20,129 BD, 33,426 SCZ) and 54,065 controls on 15.5 million SNP allele dosages  
164 imputed from 1000 genomes phase 3(The 1000 Genomes Project Consortium, 2015). Logistic  
165 regression was performed controlling for 13 principal components of ancestry, study sites and  
166 genotyping platform. We identified 11,231 SNPs with p-value below our genome-wide  
167 significance (GWS) threshold of  $5 \times 10^{-8}$ . After grouping SNPs in linkage disequilibrium with  
168 each other ( $r^2 > 0.2$ ), 114 genomic risk loci remained. For the most significant variant in each of  
169 the 114 GWS loci, we performed conditional analysis with any GWS hit within 1Mb of the

170 extent of the locus from the previously performed single disease GWAS of SCZ(Schizophrenia  
171 Working Group of the Psychiatric Genomics Consortium, 2014) and BD(Stahl et al., 2017) and  
172 identified 32 loci that were independently significant defined strictly as no single disease locus  
173 within 1Mb or a GWS p-value after conditional analysis (Supplementary Table 1). We further  
174 performed gene-set based tests using MAGMA(Leeuw et al., 2015) across 10,891 curated  
175 pathways(Watanabe et al., 2017) and identified 8 pathways surpassing Bonferroni correction ( $p <$   
176  $4.6 \times 10^{-6}$ ) with all but one pathway implicating synaptic and neuronal biology (Supplementary  
177 Table 2a). Establishing independent controls (see Methods) allowed us to perform disorder-  
178 specific GWAS in 20,129 BD cases vs 21,524 BD controls and 33,426 SCZ cases and 32,541  
179 SCZ controls. Using these results, we compared effect sizes of these 114 loci across each  
180 disorder independently showing the subsets of variants that had larger effects in SCZ compared  
181 to BD and vice versa (Figure 1a).

182

### 183 **Differentiating genetic contribution to BD and SCZ**

184 To identify loci with divergent effects on BD and SCZ, we performed an association analysis  
185 comparing 23,585 SCZ cases with 15,270 BD cases matched for shared ancestry and genotyping  
186 platform (see Methods, Figure 1b, Table 1). Two genome-wide significant loci were identified,  
187 the most significant of which was rs56355601 located on chromosome 1 at position 173,811,455  
188 within an intron of *DARS2* (Supplementary Figure 1). The second most significant locus was  
189 rs200005157, a four base-pair insertion/deletion, on chromosome 20 at position 47638976 in an  
190 intron of *ARFGEF2* (Supplementary Figure 2). For both variants, the minor allele frequency was  
191 higher in BD cases than SCZ cases and disease-specific GWAS showed opposite directions of  
192 effect when compared to controls. We sought to identify additional disease-specific loci by

193 comprehensively incorporating expression information with association results to perform fine-  
194 mapping and identify novel variants(Gamazon et al., 2015; Giambartolomei et al., 2014; Gusev  
195 et al., 2016; He et al., 2013). Here, we applied the summary-data-based Mendelian  
196 randomization (SMR) method(Zhu et al., 2016) (see Methods) utilizing the cis-QTLs derived  
197 from peripheral blood(Westra et al., 2013), human dorsolateral prefrontal cortex  
198 (DLPFC)(Fromer et al., 2016) from the Common Mind Consortium and 11 brain regions from  
199 the GTEx consortium(Consortium, 2015). We identified one SNP-probe combination that  
200 surpassed the threshold for genome-wide significance in blood but was also the most significant  
201 finding in brain. We found that SNP rs4793172 in gene *DCAKD* is associated with SCZ vs BD  
202 analysis ( $p_{\text{GWAS}} = 2.8 \times 10^{-6}$ ) and is an eQTL for probe ILMN 1811648 ( $p_{\text{eQTL}} = 2.9 \times 10^{-168}$ ),  
203 resulting in  $p_{\text{SMR}} = 4.1 \times 10^{-6}$  in blood ( $p_{\text{eQTL}} = 2.9 \times 10^{-25}$ ,  $p_{\text{SMR}} = 2.0 \times 10^{-5}$  in DLFC, and  $p_{\text{eQTL}} =$   
204  $4.6 \times 10^{-15}$ ,  $p_{\text{SMR}} = 6.0 \times 10^{-5}$  in GTEx cerebellar hemisphere) (Supplementary Table 3,  
205 Supplementary Figure 3) and shows no evidence of heterogeneity ( $p_{\text{HET}} = 0.66$ ) which implies  
206 only a single causal variant in the locus.

207 In an effort to prioritize genes for the two GWS loci from the GWAS, we performed fine-  
208 mapping(Benner et al., 2016) using an LD map derived from a majority of the control samples.  
209 We then performed SMR on each of the variants with causal probability greater than 1% using  
210 all eQTLs from the CommonMind Consortium DLPFC reference. All the most likely causal  
211 variants were shown to most significantly regulate the same gene suggesting *CSEIL* is the most  
212 likely relevant gene on chromosome 20 (rs200005157: causal probability=0.21,  $p_{\text{GWAS}}=2.4 \times 10^{-8}$ ,  
213  $p_{\text{eQTL}} 3 \times 10^{-8}$ ,  $p_{\text{SMR}}=8.5 \times 10^{-5}$ ,  $p_{\text{HET}}=0.34$ ). For the locus on chromosome 1, *SLC9C2* is the most  
214 significantly regulated gene. However, a highly significant heterogeneity test indicates a  
215 complex genetic architecture making it difficult to infer a causal role for the associated SNP.

216 Therefore, *DARS2* presents as the most likely relevant gene on chromosome 1 (rs56355601:  
217  $p_{\text{GWAS}}=5.6 \times 10^{-9}$ , causal probability=0.079,  $p_{\text{eQTL}} 7.4 \times 10^{-13}$ ,  $p_{\text{SMR}}=6.17 \times 10^{-6}$ ,  $p_{\text{HET}}=0.03$ ). We note  
218 however, that in both cases there are less associated variants that are stronger eQTLs for these  
219 genes complicating a straightforward causal interpretation. Finally, using the same gene-set test  
220 used for the combined analysis GO biological process “response to potassium ion” ( $p=1.6 \times 10^{-6}$ )  
221 was the only pathway surpassing our Bonferroni corrected significance threshold  
222 (Supplementary Table 2b).

223

### 224 **Regional joint association**

225 We expanded our efforts to identify disorder-specific genomic regions by jointly analyzing  
226 independent GWAS results from BD and SCZ (Pickrell et al., 2016). The genome was split into  
227 1,703 previously defined approximately LD independent regions (Berisa and Pickrell, 2015).  
228 Thirteen percent, or 223 regions, had a posterior probability greater than 0.5 of having a causal  
229 variant for at least one disorder. Of these, 132 best fit the model of a shared causal variant  
230 influencing both BD and SCZ, 88 were most likely specific to SCZ, 3 demonstrated evidence of  
231 two independent variants (with one impacting each of the two disorders) and none were BD-  
232 specific. Of note, this approach calculates a prior probability that any given region is disease-  
233 specific and from these data the probability of having a BD specific region was 0.1% compared  
234 to 15% for SCZ, likely a result of increased power from the larger SCZ sample size and/or a  
235 difference in genetic architecture between these disorders.

236 The 114 GWS SNPs from the combined BD and SCZ GWAS localized into 99 independent  
237 regions (13 regions had multiple GWS SNPs), of which 78 (79%) were shared with a posterior  
238 probability of greater than 0.5. Sixty regions had at least one GWS SNP in the independent SCZ



239 GWAS, of which 30 (50%) are shared and 8 regions contained a GWS SNP in the independent  
240 BD GWAS, of which 6 (75%) are shared using the same definition. For the three regions  
241 showing evidence for independent variants, two had highly non-overlapping association signals  
242 in the same region stemming from independent variants. The third, on chromosome 19 presented  
243 a different scenario where association signals were overlapping. The most significant variant in  
244 BD was rs111444407 (chr19:19358207,  $p = 8.67 \times 10^{-10}$ ) and for SCZ was rs2315283  
245 (chr19:19480575,  $p=4.41 \times 10^{-7}$ ). After conditioning on the most significant variant in the other  
246 disorder, the association signals of the most significant variant in BD and SCZ were largely  
247 unchanged (BD rs111444407  $=1.3 \times 10^{-9}$ , SCZ rs2315283  $p=6.7 \times 10^{-5}$ ). We further calculated the  
248 probability of each variant in the region being causal for both BD and SCZ (Benner et al., 2016)  
249 and found no correlation ( $r= -0.00016$ ). The most significant variants had the highest posterior  
250 probability of being causal (SCZ: rs2315283, prob = 0.02, BD: rs111444407, prob = 0.16). Both  
251 variants most significantly regulate the expression of *GATAD2A* in brain (Fromer et al., 2016) but  
252 in opposite directions (rs111444407  $p_{eQTL} = 6 \times 10^{-15}$ , beta = 0.105; rs2315283  $p_{eQTL} = 1.5 \times 10^{-28}$ ,  
253 beta = -0.11).

254

### 255 **Regional SNP-heritability estimation**

256 Across the genome, regional SNP-heritabilities ( $h^2_{\text{snp}}$ ) were estimated separately for SCZ and  
257 BD (Shi et al., 2016) and were found to be moderately correlated ( $r=0.25$ ). We next defined risk  
258 regions as those containing the most associated SNP for each GWS locus. In total, there were  
259 101 SCZ risk regions from the 105 autosomal GWS loci reported previously (Schizophrenia  
260 Working Group of the Psychiatric Genomics Consortium, 2014) and 29 BD risk regions from 30  
261 GWS loci reported previously (Stahl et al., 2017). Ten regions were risk regions for both BD and

262 SCZ comprising 33% of BD risk regions and 10% of SCZ risk regions. We further stratified  
263 regional  $h^2_{\text{snp}}$  by whether a region was a risk region in one disorder, none or both (Supplementary  
264 Figure 4). Since the discovery data for the regions overlapped with the data used for the  
265 heritability estimation, we expected within-disorder analyses to show significant results. In risk  
266 regions specific to SCZ (n=91) there was a significant increase in regional  $h^2_{\text{snp}}$  in SCZ, as  
267 expected ( $p = 1.1 \times 10^{-22}$ ), but also in BD ( $p = 1.2 \times 10^{-6}$ ). In risk regions specific to BD (n=19),  
268 significantly increased regional  $h^2_{\text{snp}}$  was observed in BD, as expected ( $p = 0.0007$ ), but not in  
269 SCZ ( $p = 0.89$ ). Risk regions shared by both disorders had significantly higher  $h^2_{\text{snp}}$  in both  
270 disorders, as expected (BD  $p = 5.3 \times 10^{-5}$ , SCZ  $p = 0.006$ ), compared to non-risk regions.  
271 However, we observed a significant increase in BD  $h^2_{\text{snp}}$  in shared risk regions compared to BD  
272 risk regions (BD  $p = 0.003$ ) but not SCZ  $h^2_{\text{snp}}$  for shared risk regions compared to SCZ risk  
273 regions ( $p = 0.62$ ). Using a less stringent p-value threshold for defining risk regions ( $p < 5 \times 10^{-6}$ ),  
274 thereby substantially increasing the number of regions, resulted in similar results. Seven regions  
275 contributed to substantially higher  $h^2_{\text{snp}}$  in SCZ compared to BD but no region showed the  
276 inverse pattern. Of these regions, all but one was in the major histocompatibility region (MHC),  
277 the sole novel region was chr10:104380410-106695047 with regional  $h^2_{\text{snp}} = 0.0019$  in SCZ and  
278  $h^2_{\text{snp}} = 0.00063$  in BD.

279

## 280 **Polygenic dissection of subphenotypes**

281 Subphenotypes were collected for a subset of patients with either BD or SCZ (see Methods). For  
282 SCZ, we had clinical quantitative measurements of manic, depressive, positive and negative  
283 symptoms generated from factor analysis of multiple instruments as described  
284 previously (Ruderfer et al., 2014) but in larger sample sizes (n=6908, 6907, 8259, 8355

285 respectively). For BD, 24 subphenotypes were collected among nearly 13,000 cases in distinct  
286 categories including comorbidities, clinical information such as rapid cycling and psychotic  
287 features as well as additional disease course data such as age of onset and number of  
288 hospitalizations. For each BD or SCZ patient, we calculated a polygenic risk score (PRS) using  
289 all SNPs, from each of the four main GWAS analyses (BD+SCZ, BD, SCZ and SCZvsBD). We  
290 then used regression analysis including principal components and site to assess the relationship  
291 between each subphenotype and the 4 PRS. Specifically, we tested whether polygenic risk scores  
292 of BD+SCZ, BD, SCZ or SCZvsBD were correlated with each of these subphenotypes separately  
293 within BD and SCZ cases. When testing if the variance explained by the PRS was different from  
294 zero, we applied a significance cutoff of  $p < 0.0004$  based on Bonferroni correction for 112 tests.  
295 In total, we identified 6 significant results after correction (Figure 2, Table 2).

296  
297 A significant positive correlation existed between BD PRS and manic symptoms in SCZ cases as  
298 seen previously (Ruderfer et al., 2014) ( $p=2 \times 10^{-5}$ ,  $t=4.26$ ) and BD PRS and psychotic features in  
299 BD patients ( $p=5.3 \times 10^{-5}$ ,  $t=4.04$ ). A significant increase in SCZ PRS was seen for BD cases with  
300 versus without psychotic features ( $p=1.2 \times 10^{-10}$ ,  $t=6.45$ ) and patients with increased negative  
301 symptoms in SCZ patients ( $p=3.60 \times 10^{-6}$ ,  $t=4.64$ ). The BD+SCZ vs controls PRS was  
302 significantly associated with psychotic features in BD ( $p=7.9 \times 10^{-13}$ ,  $t=7.17$ ) and negative  
303 symptoms in SCZ ( $p=1.5 \times 10^{-5}$ ,  $t=4.33$ ). The next two most significant results which did not  
304 survive our conservative correction were both indicative of a more severe course in BD:  
305 increased BD+SCZ PRS with increased numbers of hospitalizations in BD cases ( $p=4.2 \times 10^{-4}$ ,  
306  $t=3.53$ ) and increased SCZ PRS with earlier onset of BD ( $p=7.9 \times 10^{-4}$ ,  $t=-3.36$ ). We assessed the  
307 role of BD subtype on the correlation between SCZ PRS and psychotic features and identified a

308 significant correlation when restricted to only BD type I cases indicating the result was not likely  
309 driven by BD patients with a schizoaffective subtype (BDI: 3,763 with psychosis, 2,629 without,  
310  $p=1.55 \times 10^{-5}$ , Supplementary Table 4).

311  
312 We performed a GWAS for all 8 quantitative subphenotypes and 9 binary subphenotypes with at  
313 least 1,000 cases and calculated heritability and genetic correlation with BD and SCZ. Only two  
314 subphenotypes had significant  $h^2_{\text{snp}}$  estimates using LD-score regression (Bulik-Sullivan et al.,  
315 2015) both in BD: psychotic features in BD ( $h^2_{\text{snp}}=0.15$ ,  $SE=0.06$ ) and suicide attempt  
316 ( $h^2_{\text{snp}}=0.25$ ,  $SE=0.1$ ). Only psychotic features demonstrated a significant genetic correlation with  
317 SCZ ( $r_g=0.34$ ,  $SE=0.13$ ,  $p=0.009$ ). The significant genetic correlation demonstrates a genome-  
318 wide relationship between common variants contributing to SCZ risk and those contributing to  
319 psychotic features in BD cases. We tested whether the most significantly associated SCZ loci  
320 contributed directly to psychotic features in BD. One hundred of the 105 autosomal genome-  
321 wide significant SCZ SNPs previously published (Schizophrenia Working Group of the  
322 Psychiatric Genomics Consortium, 2014) were in our dataset after QC and 60 were in the same  
323 direction of effect for risk of psychotic features in BD ( $p=0.028$ , one-sided binomial-test).

324  
325  
326 **Discussion**

327 Here we present a genetic dissection of bipolar disorder and schizophrenia from over 100,000  
328 genotyped subjects. Consistent with earlier results (Cross-Disorder Group of the Psychiatric  
329 Genomics Consortium, 2013), we found extensive genetic sharing between these two disorders,  
330 identifying 114 genome-wide significant loci contributing to both disorders of which 32 are  
331 novel. These findings point to the relevance of neuronal and synaptic biology for the shared

332 genetic substrate of these disorders. However, despite this degree of sharing, we identified  
333 several loci that significantly differentiated between the two disorders, having opposite directions  
334 of effect. We also found polygenic components that significantly correlated from one disorder to  
335 symptoms of the other.

336

337 Two GWS loci were identified from the case only SCZ versus BD analysis providing  
338 opportunities to inform the underlying biological distinctions between BD and SCZ. The most  
339 significant locus implicates *DARS2* (coding for the mitochondrial Aspartate-tRNA ligase) which  
340 is highly expressed in the brain and significantly regulated by the most significant SNP  
341 rs56355601 ( $p_{eQTL}=2.5 \times 10^{-11}$ ). Homozygous mutations in *DARS2* are responsible for  
342 leukoencephalopathy with brainstem and spinal cord involvement and lactate elevation (LBSL),  
343 which was characterized by neurological symptoms such as psychomotor developmental delay,  
344 cerebellar ataxia and delayed mental development(Yamashita et al., 2013, p. 2). Based on  
345 methylation analysis from the prefrontal cortex of stress models (rats and monkeys) and from  
346 peripheral samples (in monkeys and human newborns), *DARS2*, among others, has been  
347 suggested as a potential molecular marker of early-life stress and vulnerability to psychiatric  
348 disorders(Luoni et al., 2016). The second most significant locus implicates *CSE1L*, a nuclear  
349 transport factor that plays a role in cellular proliferation as well as in apoptosis(Bera et al., 2001).  
350 Intronic SNPs in *CSE1L* have been associated with subjective well-being(Okbay et al., 2016)  
351 and, nominally to antidepressant response(Li et al., 2016). More interestingly, *CSE1L* is a  
352 potential target gene of miR-137, one of the well-known schizophrenia risk loci(Schizophrenia  
353 Working Group of the Psychiatric Genomics Consortium, 2014), which is able to negatively  
354 regulate CSE1L by interacting with complementary sequences in the 3' UTR of *CSE1L*(Li et al.,

355 2013). Although falling short of genome-wide significance, the third most significant locus  
356 implicates *ARNTL* (Aryl Hydrocarbon Receptor Nuclear Translocator Like), which is a core  
357 component of the circadian clock. *ARNTL* has been previously hypothesized for relevance in  
358 bipolar disorder,(Yang et al., 2008) although human genetic evidence is currently limited(Byrne  
359 et al., 2014).

360

361 The ability to generate transcriptional data on multiple tissues across many individuals using  
362 RNA-sequencing has provided detailed information on the role common variants play in  
363 regulating expression of specific genes in specific tissues. These eQTLs can be integrated with  
364 the genetic association data from GWAS to inform on the relationship between variant  
365 association and variant regulation of expression for each gene. Performing this integration, we  
366 identified a third genome-wide significant finding in *DCAKD*. The gene codes for Dephospho-  
367 CoA Kinase Domain Containing protein, a member of the human postsynaptic density proteome  
368 from human neocortex(Bayés et al., 2011). In the mouse cortical synaptoproteome *DCAKD* is  
369 among the proteins with the highest changes between juvenile postnatal days and adult stage,  
370 suggesting a putative role in brain development(Gonzalez-Lozano et al., 2016; Moczulska et al.,  
371 2014). Discerning between pleiotropy (variant independently regulates expression and alters risk  
372 to disease) from causality (variant regulates expression which thereby alters risk to disease)  
373 through statistical analysis alone is difficult, this analytical approach is stringent in excluding  
374 loci where colocalised SNP-phenotype and SNP-expression associations may reflect  
375 confounding driven by linkage disequilibrium (LD) (one variant regulates expression and a  
376 different variant alters risk but the variants in the region are in LD). Hence, this approach utilizes  
377 currently available data to prioritize genes, including direction of effect, for functional follow-up.

378 These analyses will become more powered with increased sample sizes for both phenotype and  
379 eQTL data sets.

380

381 Performing pathway analysis based on the full association results shows enrichment of genes  
382 involved in response to potassium ions, including potassium voltage-gated channel subfamily  
383 members and a number of genes regulated by cellular potassium concentration. This is in line  
384 with previous genetic evidence pointing to a key etiologic role of potassium channels, in  
385 particular, in BD(Judy and Zandi, 2013), which could be explained by their role in multiple  
386 neurobiological mechanisms involved in the development of psychiatric disorders such as  
387 regulation of the dopaminergic circuits, synaptic plasticity, and myelination(Balaraman et al.,  
388 2015).

389

390 We further assessed the contribution of regions of the genome to each disorder through joint  
391 regional association and heritability estimation. These results point to an additional locus that  
392 may contribute differentially to liability to BD and SCZ. The region on chr19 shows overlapping  
393 association peaks that are driven by independent causal variants for each disorder. Both variants  
394 significantly regulate the same gene *GATAD2A* but in opposite directions. *GATAD2A* is a  
395 transcriptional repressor, which is targeted by *MBD2* and is involved in methylation-dependent  
396 gene silencing. The protein is part of the large NuRD (nucleosome remodeling and deacetylase)  
397 complex, for which also HDAC1/2 are essential components. NurD complex proteins have been  
398 associated with autism(Li et al., 2015). Their members, including *GATAD2A*, display preferential  
399 expression in fetal brain development(Li et al., 2015) and in recent work has been implicated in  
400 SCZ through open chromatin(Fullard et al., n.d.). Further, p66 $\alpha$  (mouse *GATAD2A*) was recently

401 shown to participate in memory preservation through long-lasting histone modification in  
402 hippocampal memory-activated neurons(Ding et al., 2017). SNP-heritability appears to be  
403 consistently shared across regions and chromosomes between these two disorders. Regions with  
404 GWS loci often explain higher proportions of heritability as expected. When looking at the effect  
405 on heritability of the presence of a GWS locus in the other disorder, we identified a significant  
406 increase in BD heritability for regions containing a GWS locus for SCZ but no significant  
407 increase in SCZ heritability in regions having a BD one. This result suggests a directionality to  
408 the genetic sharing of these disorders with a larger proportion of BD loci being specific to BD.  
409 However, we cannot exclude that the asymmetry of results may reflect less power of discovery  
410 for BD than SCZ. The degree to which power and subphenotypes contribute to this result  
411 requires further examination.

412

413 We note that as with nearly all GWAS findings, the calculated population-based effect sizes of  
414 the variants identified here are small and independently explain only a modest fraction to the  
415 heritability of these disorders. The identification of these variants is dependent on the ability to  
416 have highly accurate allele frequency estimates that can only be ascertained from large sample  
417 sizes. As sample sizes get larger the power to identify variants of smaller effect increases  
418 meaning that increasing sample size results in the identification of variants of smaller effect.  
419 However, a small population effect size does not exclude the possibility of a substantially larger  
420 effect on molecular phenotypes nor does it preclude the utility of association regions in  
421 understanding biology or having a clinical impact. Efforts following up GWAS results to date  
422 have demonstrated the value of these findings in pointing to genes that can aid in understanding  
423 the underlying biology of the trait(Claussnitzer et al., 2015; Mohanan et al., 2018; Sekar et al.,



424 2016). Further, there is a clear relationship between GWAS results of a phenotype and gene  
425 targets of drugs that treat that phenotype pointing to the potential for improved therapeutic  
426 understanding(Nelson et al., 2015; Ruderfer et al., 2016). A major challenge of GWAS is the  
427 sheer number of findings and the substantial time/cost required for functional follow up of these  
428 findings in the classical paradigms used for genes causal for monogenic disorders. In silico  
429 bioinformatic analyses (such as SMR used here) that integrate GWAS results with ‘omics data  
430 (transcription, protein, epigenetic, etc.) have the potential to put a clearer biological focus on  
431 GWAS results. Such analyses can become more complex as more reference omics data sets (with  
432 genome-wide genotyping) become available. Additional analytical efforts will be required to  
433 facilitate the transition from GWAS to biology but substantial data has shown there is much to be  
434 learned from these variants despite their small effects(Visscher et al., 2017).

435

436 We have now identified multiple genomic signatures that correlate between one disorder and a  
437 clinical symptom in the other disorder, illustrating genetic components underlying particular  
438 symptom dimensions within these disorders. Medical symptoms, including those seen in  
439 psychiatric disorders, can manifest through a multitude of causes. The classic example often used  
440 is headache for which many different paths lead to the same symptom. Psychiatric symptoms  
441 also have many potential causes. For example, symptoms of psychosis can be the result of highly  
442 heritable diseases such as BD and SCZ but also infectious and neurodegenerative diseases,  
443 sleep/sensory deprivation or psychedelic drugs. Demonstrating a shared biological underpinning  
444 to these symptoms suggests they could be treated through modulating the same pathway. As  
445 previously shown, we find a significant positive correlation between the PRS of BD and manic  
446 symptoms in SCZ. We also demonstrate that BD cases with psychotic features carry a

447 significantly higher SCZ PRS than BD cases without psychotic features and this result is not  
448 driven by the schizoaffective BD subtype. Further, we show that increased PRS is associated  
449 with more severe illness. This is true for BD with psychotic features having increased SCZ PRS,  
450 earlier onset BD having higher SCZ PRS and cases with higher BD+SCZ PRS having a larger  
451 number of hospitalizations. We demonstrated that psychotic features within BD is a heritable  
452 trait and GWS loci for SCZ have a consistent direction of effect in psychotic features in BD,  
453 demonstrating the potential to study psychosis more directly to identify variants contributing to  
454 that symptom dimension.

455

456 This work illustrates the utility of genetic data, in aggregate, at dissecting symptom  
457 heterogeneity among related disorders and suggests that further work could aid in characterizing  
458 patients for more personalized treatment. Genetic risk scores have demonstrated their ability to  
459 inform and predict pathology (Cleyne et al., 2016) and more recently have been shown to be  
460 able to identify patients with risk equivalent to monogenic variants (Khera et al., 2017). In  
461 psychiatry, we lack objective biological measurements (biomarkers) with which to assess the  
462 ability of a genetic signature to predict or inform. Lacking diagnostic pathology for psychiatric  
463 disorders leaves a genuine opportunity for the genetics to drive diagnosis and treatment to a  
464 much larger degree than in other domains. One potential model assumes that each individual has  
465 a quantitative loading of a series of symptom dimensions (i.e. manic, psychotic, cognitive, etc.)  
466 and that these symptoms can be assessed at the genetic level to characterize a patient's  
467 dysfunction and used to inform disease course and optimal treatment. Making this a reality will  
468 require more detailed information on disease course and outcomes. For example, if treatment  
469 response data existed for these samples one could ask whether a genetic loading for psychosis

470 was correlated with response to treatment. Initial work has already shown the potential of this  
471 approach using a SCZ PRS to inform lithium response in BD(Amare et al., 2018). Ultimately,  
472 the goal will be to quantify multiple genetic loadings of each individual's illness and use those  
473 measures to inform treatment based on the outcomes of previous individuals with similar  
474 profiles.

475  
476 In conclusion, we present a detailed genetic dissection of BD and SCZ pointing to substantial  
477 shared genetic risk but also demonstrating that specific loci contribute to the phenotypic  
478 differences of these disorders. We show that genetic risk scores can correspond to symptoms  
479 within and across disorders. Finally, we present data that points to these disorders being neither  
480 independent nor the same but sharing particular symptom dimensions that can be captured from  
481 the genetics and used to characterize patients to ultimately inform diagnosis and treatment.

482

483 **Author Contributions:**

484 DMR, PS and KSK managed and organized the group. DMR, SR, JB, EAS, JMWP, NM, AWC,  
485 APSO, LMOL and VT contributed to analyses. Subphenotype collection and organization was  
486 led by AM and AHF. Initial manuscript was drafted by DMR, ED, ADF, SP, JLK. Manuscript  
487 contributions and interpretation of results was provided by DMR, ED, SHL, MCO, PFS, RAO,  
488 NRW, PS and KSK. The remaining authors contributed to the recruitment, genotyping, or data  
489 processing for the contributing components of the study. All other authors saw, had the  
490 opportunity to comment on, and approved the final draft.

491

492

494 **Bipolar Disorder and Schizophrenia Working Group of the Psychiatric Genomics**  
495 **Consortium**

496 Douglas M Ruderfer, Stephan Ripke, Andrew McQuillin, James Boocock, Eli A Stahl, Jennifer M Whitehead  
 497 Pavlides, Niamh Mullins, Alexander W Charney, Anil P S Ori, Loes M Olde Loohuis, Enrico Domenici, Arianna Di  
 498 Florio, Sergi Papiol, Janos L. Kalman, Vassily Trubetskoy, Rolf Adolfsson, Ingrid Agartz, Esben Agerbo,  
 499 Huda Akil, Diego Albani, Margot Albus, Martin Alda, Madeline Alexander, Ney Alliey-Rodriguez, Thomas D Als,  
 500 Farooq Amin, Adebayo Anjorin, Maria J Arranz, Swapnil Awasthi, Silviu A Bacanu, Judith A Badner, Marie  
 501 Baekvad-Hansen, Steven Bakker, Gavin Band, Jack D Barchas, Ines Barroso, Nicholas Bass, Michael Bauer,  
 502 Bernhard T Baune, Martin Begemann, Celine Bellenguez, Richard A Belliveau Jr, Frank Bellivier, Stephan Bender,  
 503 Judit Bene, Sarah E Bergen, Wade H Berrettini, Elizabeth Bevilacqua, Joanna M Biernacka, Tim B Bigdeli, Donald  
 504 W Black, Hannah Blackburn, Jenefer M Blackwell, Douglas HR Blackwood, Carsten Bocker Pedersen, Michael  
 505 Boehnke, Marco Boks, Anders D Borglum, Elvira Bramon, Gerome Breen, Matthew A Brown, Richard Bruggeman,  
 506 Nancy G Buccola, Randy L Buckner, Monika Budde, Brendan Bulik-Sullivan, Suzannah J Bumpstead, William  
 507 Bunney, Margit Burmeister, Joseph D Buxbaum, Jonas Bybjerg-Grauholm, William Byerley, Wiekpe Cahn,  
 508 Guiqing Cai, Murray J Cairns, Dominique Champion, Rita M Cantor, Vaughan J Carr, Noa Carrera, Juan P Casas,  
 509 Miquel Casas, Stanley V Catts, Pablo Cervantes, Kimberley D Chambert, Raymond CK Chan, Eric YH Chen,  
 510 Ronald YL Chen, Wei Cheng, Eric FC Cheung, Siow Ann Chong, Toni-Kim Clarke, C Robert Cloninger, David  
 511 Cohen, Nadine Cohen, Jonathan R I Coleman, David A Collier, Paul Cormican, William Coryell, Nicholas  
 512 Craddock, David W Craig, Benedicto Crespo-Facorro, James J Crowley, Cristiana Cruceanu, David Curtis, Piotr M  
 513 Czerski, Anders M Dale, Mark J Daly, Udo Dannlowski, Ariel Darvasi, Michael Davidson, Kenneth L Davis,  
 514 Christiaan A de Leeuw, Franziska Degenhardt, Jurgen Del Favero, Lynn E DeLisi, Panos Deloukas, Ditte Demontis,  
 515 J Raymond DePaulo, Marta di Forti, Dimitris Dikeos, Timothy Dinan, Srdjan Djurovic, Amanda L Dobbyn, Peter  
 516 Donnelly, Gary Donohoe, Elodie Drapeau, Serge Dronov, Jubao Duan, Frank Dudbridge, Audrey Duncanson,  
 517 Howard Edenberg, Sarah Edkins, Hannelore Ehrenreich, Peter Eichhammer, Torbjorn Elvsashagen, Johan Eriksson,  
 518 Valentina Escott-Price, Tonu Esko, Laurent Essioux, Bruno Etain, Chun Chieh Fan, Kai-How Farh, Martilias S  
 519 Farrell, Matthew Flickinger, Tatiana M Foroud, Liz Forty, Josef Frank, Lude Franke, Christine Fraser, Robert  
 520 Freedman, Colin Freeman, Nelson B Freimer, Joseph I Friedman, Menachem Fromer, Mark A Frye, Janice M  
 521 Fullerton, Katrin Gade, Julie Garnham, Helena A Gaspar, Pablo V Gejman, Giulio Genovese, Lyudmila Georgieva,  
 522 Claudia Giambartolomei, Eleni Giannoulatou, Ina Giegling, Michael Gill, Matthew Gillman, Marianne Giortz  
 523 Pedersen, Paola Giusti-Rodriguez, Stephanie Godard, Fernando Goes, Jacqueline I Goldstein, Srihari Gopal, Scott D  
 524 Gordon, Katherine Gordon-Smith, Jacob Gratten, Emma Gray, Elaine K Green, Melissa J Green, Tiffany A  
 525 Greenwood, Maria Grigoriou-Serbanescu, Jakob Grove, Weihua Guan, Hugh Gurling, Jose Guzman Parra, Rhian  
 526 Gwilliam, Lieuwe de Haan, Jeremy Hall, Mei-Hua Hall, Christian Hammer, Naomi Hammond, Marian L Hamshere,  
 527 Mark Hansen, Thomas Hansen, Vahram Haroutunian, Annette M Hartmann, Joanna Hauser, Martin Hautzinger, Urs  
 528 Heilbronner, Garrett Hellenthal, Frans A Henskens, Stefan Herms, Maria Hipolito, Joel N Hirschhorn, Per  
 529 Hoffmann, Mads V Hollegaard, David M Hougaard, Hailiang Huang, Laura Huckins, Christina M Hultman, Sarah E  
 530 Hunt, Masashi Ikeda, Nakao Iwata, Conrad Iyegbe, Assen V Jablensky, Stephane Jamain, Janusz Jankowski,  
 531 Alagurevathi Jayakumar, Inge Joa, Ian Jones, Lisa A Jones, Erik G Jonsson, Antonio Julia, Anders Jureus, Anna K  
 532 Kahler, Rene S Kahn, Luba Kalaydjieva, Radhika Kandaswamy, Sena Karachanak-Yankova, Juha Karjalainen,  
 533 Robert Karlsson, David Kavanagh, Matthew C Keller, Brian J Kelly, John Kelsoe, James L Kennedy, Andrey  
 534 Khrunin, Yunjung Kim, George Kirov, Sarah Kittel-Schneider, Janis Klovins, Jo Knight, Sarah V Knott, James A  
 535 Knowles, Manolis Kogevinas, Bettina Konte, Eugenia Kravariti, Vaidutis Kucinskas, Zita Ausrele Kucinskiene,  
 536 Ralph Kupka, Hana Kuzelova-Ptackova, Mikael Landen, Cordelia Langford, Claudine Laurent, Jacob Lawrence,  
 537 Stephen Lawrie, William B Lawson, Markus Leber, Marion Leboyer, Phil H Lee, Jimmy Lee Chee Keong, Sophie E  
 538 Legge, Todd Lencz, Bernard Lerer, Douglas F Levinson, Shawn E Levy, Cathryn M Lewis, Jun Z Li, Miaoxin Li,  
 539 Qingqin S Li, Tao Li, Kung-Yee Liang, Jennifer Liddle, Jeffrey Lieberman, Svetlana Limborska, Kuang Lin, Don H  
 540 Linszen, Jolanta Lissowska, Chunyu Liu, Jianjun Liu, Jouko Lonqvist, Carmel M Loughland, Jan Lubinski,  
 541 Susanne Lucae, Milan Macek Jr, Donald J MacIntyre, Patrik KE Magnusson, Brion S Maher, Pamela B Mahon,  
 542 Wolfgang Maier, Anil K Malhotra, Jacques Mallet, Ulrik F Malt, Hugh S Markus, Sara Marsal, Nicholas G Martin,  
 543 Ignacio Mata, Christopher G Mathew, Manuel Mattheisen, Morten Mattingsdal, Fermin Mayoral, Owen T McCann,  
 544 Robert W McCarley, Steven A McCarroll, Mark I McCarthy, Colm McDonald, Susan L McElroy, Peter McGuffin,  
 545 Melvin G McInnis, Andrew M McIntosh, James D McKay, Francis J McMahon, Helena Medeiros, Sarah E  
 546 Medland, Sandra Meier, Carin J Meijer, Bela Melegh, Ingrid Melle, Fan Meng, Raquelle I Meshulam-Gately,

547 Andres Metspalu, Patricia T Michie, Lili Milani, Vihra Milanova, Philip B Mitchell, Younes Mokrab, Grant W  
548 Montgomery, Jennifer L Moran, Gunnar Morken, Derek W Morris, Ole Mors, Preben B Mortensen, Bryan J Mowry,  
549 Thomas W Mühleisen, Bertram Müller-Myhsok, Kieran C Murphy, Robin M Murray, Richard M Myers, Inez Myin-  
550 Germeys, Benjamin M Neale, Mari Nelis, Igor Nenadic, Deborah A Nertney, Gerald Nestadt, Kristin K Nicodemus,  
551 Caroline M Nievergelt, Liene Nikitina-Zake, Vishwajit Nimgaonkar, Laura Nisenbaum, Merete Nordentoft, Annelie  
552 Nordin, Markus M Nöthen, Evaristus A Nwulia, Eadhbhard O'Callaghan, Claire O'Donovan, Colm O'Dushlaine, F  
553 Anthony O'Neill, Ketil J Oedegaard, Sang-Yun Oh, Ann Olincy, Line Olsen, Lilijana Oruc, Jim Van Os, Michael J  
554 Owen, Sara A Paciga, Colin N A Palmer, Aarno Palotie, Christos Pantelis, George N Papadimitriou, Elena  
555 Parkhomenko, Carlos Pato, Michele T Pato, Tiina Paunio, Richard Pearson, Psychosis Endophenotypes International  
556 Consortium, Diana O Perkins, Roy H Perlis, Amy Perry, Tune H Pers, Tracey L Petryshen, Andrea Pfennig, Marco  
557 Picchioni, Olli Pietilainen, Jonathan Pimm, Matti Pirinen, Robert Plomin, Andrew J Pocklington, Danielle  
558 Posthuma, James B Potash, Simon C Potter, John Powell, Alkes Price, Ann E Pulver, Shaun M Purcell, Digby  
559 Quested, Josep Antoni Ramos-Quiroga, Henrik B Rasmussen, Anna Rautanen, Radhi Ravindrarajah, Eline J Regeer,  
560 Abraham Reichenberg, Andreas Reif, Mark A Reimers, Marta Ribases, John P Rice, Alexander L Richards,  
561 Michelle Ricketts, Brien P Riley, Fabio Rivas, Margarita Rivera, Joshua L Roffman, Guy A Rouleau, Panos  
562 Roussos, Dan Rujescu, Veikko Salomaa, Cristina Sanchez-Mora, Alan R Sanders, Stephen J Sawcer, Ulrich Schall,  
563 Alan F Schatzberg, William A Scheftner, Peter R Schofield, Nicholas J Schork, Sibylle G Schwab, Edward M  
564 Scolnick, Laura J Scott, Rodney J Scott, Larry J Seidman, Alessandro Serretti, Pak C Sham, Cynthia Shannon  
565 Weickert, Tatyana Shehktman, Jianxin Shi, Paul D Shilling, Engilbert Sigurdsson, Jeremy M Silverman, Kang Sim,  
566 Claire Slaney, Petr Slominsky, Olav B Smeland, Jordan W Smoller, Hon-Cheong So, Janet L Sobell, Erik  
567 Soderman, Christine Soholm Hansen, Chris C A Spencer, Anne T Spijker, David St Clair, Hreinn Stefansson, Kari  
568 Stefansson, Stacy Steinberg, Elisabeth Stogmann, Eystein Stordal, Amy Strange, Richard E Straub, John S Strauss,  
569 Fabian Streit, Eric Strengman, Jana Strohmaier, T Scott Stroup, Zhan Su, Mythily Subramaniam, Jaana Suvisaari,  
570 Dragan M Svrakic, Jin P Szatkiewicz, Szabolcs Szelinger, Avazeh Tashakkori-Ghanbaria, Srinivas Thirumalai,  
571 Robert C Thompson, Thorgeir E Thorgeirsson, Draga Toncheva, Paul A Tooney, Sarah Tosato, Timothea  
572 Touloupoulou, Richard C Trembath, Jens Treutlein, Vassily Trubetsky, Gustavo Turecki, Arne E Vaaler, Helmut  
573 Vedder, Eduard Vieta, John Vincent, Peter M Visscher, Ananth C Viswanathan, Damjan Vukcevic, John  
574 Waddington, Matthew Waller, Dermot Walsh, Muriel Walshe, James TR Walters, Dai Wang, Qiang Wang, Weiqing  
575 Wang, Yunpeng Wang, Stanley J Watson, Bradley T Webb, Thomas W Weickert, Daniel R Weinberger, Matthias  
576 Weisbrod, Mark Weiser, Thomas Werge, Paul Weston, Pamela Whittaker, Sara Widaa, Durk Wiersma, Dieter B  
577 Wildenauer, Nigel M Williams, Stephanie Williams, Stephanie H Witt, Aaron R Wolen, Emily HM Wong, Nicholas  
578 W Wood, Brandon K Wormley, Wellcome Trust Case-Control Consortium , Jing Qin Wu, Simon Xi, Wei Xu, Allan  
579 H Young, Clement C Zai, Peter Zandi, Peng Zhang, Xuebin Zheng, Fritz Zimprich, Sebastian Zollner, Aiden  
580 Corvin, Ayman H Fanous, Sven Cichon, Marcella Rietschel, Elliot S Gershon, Thomas G Schulze, Alfredo B  
581 Cuellar-Barboza, Andreas J Forstner, Peter A Holmans, John I Nurnberger, Ole A Andreassen, S Hong Lee, Michael  
582 C O'Donovan, Patrick F Sullivan, Roel A Ophoff, Naomi R Wray, Pamela Sklar, Kenneth S Kendler  
583  
584

## 585 **Acknowledgements**

586 The work of the contributing groups was supported by numerous grants from governmental and  
587 charitable bodies as well as philanthropic donation. Specifically, DMR (R01MH111776), NRW  
588 (NHMRC 1078901, 1087889).  
589

## 590 **Declaration of Interests**

591 The authors declare no competing interests.  
592

## 593 **References**

594 Amare, A.T., Schubert, K.O., Hou, L., Clark, S.R., Papiol, S., Heilbronner, U., Degenhardt, F.,  
 595 Tekola-Ayele, F., Hsu, Y.-H., Shekhtman, T., Adli, M., Akula, N., Akiyama, K., Arda, R.,  
 596 Arias, B., Aubry, J.-M., Backlund, L., Bhattacharjee, A.K., Bellivier, F., Benabarre, A.,  
 597 Bengesser, S., Biernacka, J.M., Birner, A., Brichant-Petitjean, C., Cervantes, P., Chen,  
 598 H.-C., Chillotti, C., Cichon, S., Cruceanu, C., Czerski, P.M., Dalkner, N., Dayer, A., Zompo,  
 599 M.D., DePaulo, J.R., Étain, B., Falkai, P., Forstner, A.J., Frisen, L., Frye, M.A., Fullerton,  
 600 J.M., Gard, S., Garnham, J.S., Goes, F.S., Grigoriu-Serbanescu, M., Grof, P., Hashimoto,  
 601 R., Hauser, J., Herms, S., Hoffmann, P., Hofmann, A., Jamain, S., Jiménez, E., Kahn, J.-P.,  
 602 Kassem, L., Kuo, P.-H., Kato, T., Kelsoe, J., Kittel-Schneider, S., Kliwicki, S., König, B.,  
 603 Kusumi, I., Laje, G., Landén, M., Lavebratt, C., Leboyer, M., Leckband, S.G., Tortorella,  
 604 A., Manchia, M., Martinsson, L., McCarthy, M.J., McElroy, S., Colom, F., Mitjans, M.,  
 605 Mondimore, F.M., Monteleone, P., Nievergelt, C.M., Nöthen, M.M., Novák, T.,  
 606 O'Donovan, C., Ozaki, N., Ösby, U., Pfennig, A., Potash, J.B., Reif, A., Reininghaus, E.,  
 607 Rouleau, G.A., Rybakowski, J.K., Schalling, M., Schofield, P.R., Schweizer, B.W.,  
 608 Severino, G., Shilling, P.D., Shimoda, K., Simhandl, C., Slaney, C.M., Squassina, A.,  
 609 Stamm, T., Stopkova, P., Maj, M., Turecki, G., Vieta, E., Volkert, J., Witt, S., Wright, A.,  
 610 Zandi, P.P., Mitchell, P.B., Bauer, M., Alda, M., Rietschel, M., McMahon, F.J., Schulze,  
 611 T.G., Baune, B.T., 2018. Association of Polygenic Score for Schizophrenia and HLA  
 612 Antigen and Inflammation Genes With Response to Lithium in Bipolar Affective  
 613 Disorder: A Genome-Wide Association Study. *JAMA Psychiatry* 75, 65–74.  
 614 <https://doi.org/10.1001/jamapsychiatry.2017.3433>  
 615 Balaraman, Y., Lahiri, D.K., Nurnberger, J.I., 2015. Variants in Ion Channel Genes Link  
 616 Phenotypic Features of Bipolar Illness to Specific Neurobiological Process Domains.  
 617 *Mol. Neuropsychiatry* 1, 23–35. <https://doi.org/10.1159/000371886>  
 618 Bayés, À., van de Lagemaat, L.N., Collins, M.O., Croning, M.D.R., Whittle, I.R., Choudhary, J.S.,  
 619 Grant, S.G.N., 2011. Characterization of the proteome, diseases and evolution of the  
 620 human postsynaptic density. *Nat. Neurosci.* 14, 19–21.  
 621 <https://doi.org/10.1038/nn.2719>  
 622 Benner, C., Spencer, C.C.A., Havulinna, A.S., Salomaa, V., Ripatti, S., Pirinen, M., 2016.  
 623 FINEMAP: efficient variable selection using summary data from genome-wide  
 624 association studies. *Bioinformatics* 32, 1493–1501.  
 625 <https://doi.org/10.1093/bioinformatics/btw018>  
 626 Bera, T.K., Bera, J., Brinkmann, U., Tessarollo, L., Pastan, I., 2001. Cse1l Is Essential for Early  
 627 Embryonic Growth and Development. *Mol. Cell. Biol.* 21, 7020–7024.  
 628 <https://doi.org/10.1128/MCB.21.20.7020-7024.2001>  
 629 Berisa, T., Pickrell, J.K., 2015. Approximately independent linkage disequilibrium blocks in  
 630 human populations. *Bioinformatics* 32, 283–285.  
 631 <https://doi.org/10.1093/bioinformatics/btv546>  
 632 Bulik-Sullivan, B.K., Loh, P.-R., Finucane, H.K., Ripke, S., Yang, J., Schizophrenia Working  
 633 Group of the Psychiatric Genomics Consortium, Patterson, N., Daly, M.J., Price, A.L.,  
 634 Neale, B.M., 2015. LD Score regression distinguishes confounding from polygenicity  
 635 in genome-wide association studies. *Nat. Genet.* 47, 291–295.  
 636 <https://doi.org/10.1038/ng.3211>  
 637 Byrne, E.M., Heath, A.C., Madden, P.A., Pergadia, M.L., Hickie, I.B., Montgomery, G.W., Martin,  
 638 N.G., Wray, N.R., 2014. Testing the role of circadian genes in conferring risk for

639 psychiatric disorders. *Am. J. Med. Genet. Part B Neuropsychiatr. Genet. Off. Publ. Int.*  
640 *Soc. Psychiatr. Genet.* 0, 254–260. <https://doi.org/10.1002/ajmg.b.32230>

641 Claussnitzer, M., Dankel, S.N., Kim, K.-H., Quon, G., Meuleman, W., Haugen, C., Glunk, V.,  
642 Sousa, I.S., Beaudry, J.L., Puvindran, V., Abdennur, N.A., Liu, J., Svensson, P.-A., Hsu,  
643 Y.-H., Drucker, D.J., Mellgren, G., Hui, C.-C., Hauner, H., Kellis, M., 2015. FTO Obesity  
644 Variant Circuitry and Adipocyte Browning in Humans. *N. Engl. J. Med.* 373, 895–907.  
645 <https://doi.org/10.1056/NEJMoa1502214>

646 Cleynen, I., Boucher, G., Jostins, L., Schumm, L.P., Zeissig, S., Ahmad, T., Andersen, V.,  
647 Andrews, J.M., Annese, V., Brand, S., Brant, S.R., Cho, J.H., Daly, M.J., Dubinsky, M.,  
648 Duerr, R.H., Ferguson, L.R., Franke, A., Gearry, R.B., Goyette, P., Hakonarson, H.,  
649 Halfvarson, J., Hov, J.R., Huang, H., Kennedy, N.A., Kupcinskis, L., Lawrance, I.C., Lee,  
650 J.C., Satsangi, J., Schreiber, S., Théâtre, E., van der Meulen-de Jong, A.E., Weersma,  
651 R.K., Wilson, D.C., Parkes, M., Vermeire, S., Rioux, J.D., Mansfield, J., Silverberg, M.S.,  
652 Radford-Smith, G., McGovern, D.P.B., Barrett, J.C., Lees, C.W., 2016. Inherited  
653 determinants of Crohn’s disease and ulcerative colitis phenotypes: a genetic  
654 association study. *Lancet Lond. Engl.* 387, 156–167.  
655 [https://doi.org/10.1016/S0140-6736\(15\)00465-1](https://doi.org/10.1016/S0140-6736(15)00465-1)

656 CNV and Schizophrenia Working Groups of the Psychiatric Genomics Consortium, 2017.  
657 Contribution of copy number variants to schizophrenia from a genome-wide study  
658 of 41,321 subjects. *Nat. Genet.* 49, 27–35. <https://doi.org/10.1038/ng.3725>

659 Consortium, T.Gte., 2015. The Genotype-Tissue Expression (GTEx) pilot analysis:  
660 Multitissue gene regulation in humans. *Science* 348, 648–660.  
661 <https://doi.org/10.1126/science.1262110>

662 Cross-Disorder Group of the Psychiatric Genomics Consortium, 2013. Genetic relationship  
663 between five psychiatric disorders estimated from genome-wide SNPs. *Nat. Genet.*  
664 45, 984–994. <https://doi.org/10.1038/ng.2711>

665 Curtis, D., Vine, A., McQuillin, A., Bass, N., Pereira, A., Kandaswamy, R., Lawrence, J., Anjorin,  
666 A., Choudhury, K., Datta, S., Puri, V., Krasucki, R., Pimm, J., Thirumalai, S., Quedsted, D.,  
667 Gurling, H., 2011. Case-case genome wide association analysis reveals markers  
668 differentially associated with schizophrenia and bipolar disorder and implicates  
669 calcium channel genes. *Psychiatr. Genet.* 21, 1–4.  
670 <https://doi.org/10.1097/YPG.0b013e3283413382>

671 Delaneau, O., Zagury, J.-F., Marchini, J., 2013. Improved whole-chromosome phasing for  
672 disease and population genetic studies. *Nat. Methods* 10, 5–6.  
673 <https://doi.org/10.1038/nmeth.2307>

674 Diagnostic and Statistical Manual of Mental Disorders | DSM Library [WWW Document],  
675 n.d. URL  
676 <http://dsm.psychiatryonline.org/doi/book/10.1176/appi.books.9780890425596>  
677 (accessed 3.14.17).

678 Ding, X., Liu, S., Tian, M., Zhang, W., Zhu, T., Li, D., Wu, J., Deng, H., Jia, Y., Xie, W., Xie, H.,  
679 Guan, J.-S., 2017. Activity-induced histone modifications govern Neurexin-1 mRNA  
680 splicing and memory preservation. *Nat. Neurosci.* advance online publication.  
681 <https://doi.org/10.1038/nn.4536>

682 Fromer, M., Pocklington, A.J., Kavanagh, D.H., Williams, H.J., Dwyer, S., Gormley, P.,  
683 Georgieva, L., Rees, E., Palta, P., Ruderfer, D.M., Carrera, N., Humphreys, I., Johnson,  
684 J.S., Roussos, P., Barker, D.D., Banks, E., Milanova, V., Grant, S.G., Hannon, E., Rose,

685 S.A., Chambert, K., Mahajan, M., Scolnick, E.M., Moran, J.L., Kirov, G., Palotie, A.,  
686 McCarroll, S.A., Holmans, P., Sklar, P., Owen, M.J., Purcell, S.M., O'Donovan, M.C.,  
687 2014. De novo mutations in schizophrenia implicate synaptic networks. *Nature* 506,  
688 179–184. <https://doi.org/10.1038/nature12929>

689 Fromer, M., Roussos, P., Sieberts, S.K., Johnson, J.S., Kavanagh, D.H., Perumal, T.M., Ruderfer,  
690 D.M., Oh, E.C., Topol, A., Shah, H.R., Klei, L.L., Kramer, R., Pinto, D., Gümüş, Z.H., Cicek,  
691 A.E., Dang, K.K., Browne, A., Lu, C., Xie, L., Readhead, B., Stahl, E.A., Xiao, J., Parvizi, M.,  
692 Hamamsy, T., Fullard, J.F., Wang, Y.-C., Mahajan, M.C., Derry, J.M.J., Dudley, J.T.,  
693 Hemby, S.E., Logsdon, B.A., Talbot, K., Raj, T., Bennett, D.A., De Jager, P.L., Zhu, J.,  
694 Zhang, B., Sullivan, P.F., Chess, A., Purcell, S.M., Shinobu, L.A., Mangravite, L.M.,  
695 Toyoshiba, H., Gur, R.E., Hahn, C.-G., Lewis, D.A., Haroutunian, V., Peters, M.A., Lipska,  
696 B.K., Buxbaum, J.D., Schadt, E.E., Hirai, K., Roeder, K., Brennand, K.J., Katsanis, N.,  
697 Domenici, E., Devlin, B., Sklar, P., 2016. Gene expression elucidates functional impact  
698 of polygenic risk for schizophrenia. *Nat. Neurosci.* 19, 1442–1453.  
699 <https://doi.org/10.1038/nn.4399>

700 Fullard, J.F., Giambartolomei, C., Hauberg, M.E., Xu, K., Voloudakis, G., Shao, Z., Bare, C.,  
701 Dudley, J.T., Mattheisen, M., Robakis, N.K., Haroutunian, V., Roussos, P., n.d. Open  
702 chromatin profiling of human postmortem brain infers functional roles for non-  
703 coding schizophrenia loci. *Hum. Mol. Genet.* <https://doi.org/10.1093/hmg/ddx103>

704 Gamazon, E.R., Wheeler, H.E., Shah, K.P., Mozaffari, S.V., Aquino-Michaels, K., Carroll, R.J.,  
705 Eyler, A.E., Denny, J.C., GTEx Consortium, Nicolae, D.L., Cox, N.J., Im, H.K., 2015. A  
706 gene-based association method for mapping traits using reference transcriptome  
707 data. *Nat. Genet.* 47, 1091–1098. <https://doi.org/10.1038/ng.3367>

708 Giambartolomei, C., Vukcevic, D., Schadt, E.E., Franke, L., Hingorani, A.D., Wallace, C.,  
709 Plagnol, V., 2014. Bayesian Test for Colocalisation between Pairs of Genetic  
710 Association Studies Using Summary Statistics. *PLOS Genet.* 10, e1004383.  
711 <https://doi.org/10.1371/journal.pgen.1004383>

712 Gonzalez-Lozano, M.A., Klemmer, P., Gebuis, T., Hassan, C., van Nierop, P., van Kesteren, R.E.,  
713 Smit, A.B., Li, K.W., 2016. Dynamics of the mouse brain cortical synaptic proteome  
714 during postnatal brain development. *Sci. Rep.* 6.  
715 <https://doi.org/10.1038/srep35456>

716 Green, E.K., Rees, E., Walters, J.T.R., Smith, K.-G., Forty, L., Grozeva, D., Moran, J.L., Sklar, P.,  
717 Ripke, S., Chambert, K.D., Genovese, G., McCarroll, S.A., Jones, I., Jones, L., Owen, M.J.,  
718 O'Donovan, M.C., Craddock, N., Kirov, G., 2016. Copy number variation in bipolar  
719 disorder. *Mol. Psychiatry* 21, 89–93. <https://doi.org/10.1038/mp.2014.174>

720 Gulsuner, S., McClellan, J.M., 2015. Copy Number Variation in Schizophrenia.  
721 *Neuropsychopharmacology* 40, 252–254. <https://doi.org/10.1038/npp.2014.216>

722 Gusev, A., Ko, A., Shi, H., Bhatia, G., Chung, W., Penninx, B.W.J.H., Jansen, R., de Geus, E.J.C.,  
723 Boomsma, D.I., Wright, F.A., Sullivan, P.F., Nikkola, E., Alvarez, M., Civelek, M., Lusi,  
724 A.J., Lehtimäki, T., Raitoharju, E., Kähönen, M., Seppälä, I., Raitakari, O.T., Kuusisto, J.,  
725 Laakso, M., Price, A.L., Pajukanta, P., Pasaniuc, B., 2016. Integrative approaches for  
726 large-scale transcriptome-wide association studies. *Nat. Genet.* 48, 245–252.  
727 <https://doi.org/10.1038/ng.3506>

728 He, X., Fuller, C.K., Song, Y., Meng, Q., Zhang, B., Yang, X., Li, H., 2013. Sherlock: Detecting  
729 Gene-Disease Associations by Matching Patterns of Expression QTL and GWAS. *Am.*  
730 *J. Hum. Genet.* 92, 667–680. <https://doi.org/10.1016/j.ajhg.2013.03.022>



731 Howie, B., Marchini, J., Stephens, M., 2011. Genotype Imputation with Thousands of  
732 Genomes. *G3 GenesGenomesGenetics* 1, 457–470.  
733 <https://doi.org/10.1534/g3.111.001198>

734 Judy, J.T., Zandi, P.P., 2013. A review of potassium channels in bipolar disorder. *Front.*  
735 *Genet.* 4. <https://doi.org/10.3389/fgene.2013.00105>

736 Kataoka, M., Matoba, N., Sawada, T., Kazuno, A.-A., Ishiwata, M., Fujii, K., Matsuo, K., Takata,  
737 A., Kato, T., 2016. Exome sequencing for bipolar disorder points to roles of de novo  
738 loss-of-function and protein-altering mutations. *Mol. Psychiatry* 21, 885–893.  
739 <https://doi.org/10.1038/mp.2016.69>

740 Khera, A.V., Chaffin, M., Aragam, K., Emdin, C.A., Klarin, D., Haas, M., Roselli, C., Natarajan, P.,  
741 Kathiresan, S., 2017. Genome-wide polygenic score to identify a monogenic risk-  
742 equivalent for coronary disease. *bioRxiv* 218388. <https://doi.org/10.1101/218388>

743 Kirov, G., Pocklington, A.J., Holmans, P., Ivanov, D., Ikeda, M., Ruderfer, D., Moran, J.,  
744 Chambert, K., Toncheva, D., Georgieva, L., Grozeva, D., Fjodorova, M., Wollerton, R.,  
745 Rees, E., Nikolov, I., van de Lagemat, L.N., Bayés, À., Fernandez, E., Olason, P.I.,  
746 Böttcher, Y., Komiyama, N.H., Collins, M.O., Choudhary, J., Stefansson, K., Stefansson,  
747 H., Grant, S.G.N., Purcell, S., Sklar, P., O'Donovan, M.C., Owen, M.J., 2012. De novo CNV  
748 analysis implicates specific abnormalities of postsynaptic signalling complexes in  
749 the pathogenesis of schizophrenia. *Mol. Psychiatry* 17, 142–153.  
750 <https://doi.org/10.1038/mp.2011.154>

751 Leeuw, C.A. de, Mooij, J.M., Heskes, T., Posthuma, D., 2015. MAGMA: Generalized Gene-Set  
752 Analysis of GWAS Data. *PLOS Comput. Biol.* 11, e1004219.  
753 <https://doi.org/10.1371/journal.pcbi.1004219>

754 Li, J., Ma, Z., Shi, M., Maly, R.H., Aoki, H., Minic, Z., Phanse, S., Jin, K., Wall, D.P., Zhang, Z.,  
755 Urban, A.E., Hallmayer, J., Babu, M., Snyder, M., 2015. Identification of Human  
756 Neuronal Protein Complexes Reveals Biochemical Activities and Convergent  
757 Mechanisms of Action in Autism Spectrum Disorders. *Cell Syst.* 1, 361–374.  
758 <https://doi.org/10.1016/j.cels.2015.11.002>

759 Li, K.K.-W., Yang, L., Pang, J.C.-S., Chan, A.K.-Y., Zhou, L., Mao, Y., Wang, Y., Lau, K.-M., Poon,  
760 W.S., Shi, Z., Ng, H.-K., 2013. MIR-137 Suppresses Growth and Invasion, is  
761 Downregulated in Oligodendroglial Tumors and Targets CSE1L. *Brain Pathol.* 23,  
762 426–439. <https://doi.org/10.1111/bpa.12015>

763 Li, Q.S., Tian, C., Seabrook, G.R., Drevets, W.C., Narayan, V.A., 2016. Analysis of 23andMe  
764 antidepressant efficacy survey data: implication of circadian rhythm and  
765 neuroplasticity in bupropion response. *Transl. Psychiatry* 6, e889.  
766 <https://doi.org/10.1038/tp.2016.171>

767 Lichtenstein, P., Yip, B.H., Björk, C., Pawitan, Y., Cannon, T.D., Sullivan, P.F., Hultman, C.M.,  
768 2009. Common genetic determinants of schizophrenia and bipolar disorder in  
769 Swedish families: a population-based study. *The Lancet* 373, 234–239.  
770 [https://doi.org/10.1016/S0140-6736\(09\)60072-6](https://doi.org/10.1016/S0140-6736(09)60072-6)

771 Luoni, A., Massart, R., Nieratschker, V., Nemoda, Z., Blasi, G., Gilles, M., Witt, S.H., Suderman,  
772 M.J., Suomi, S.J., Porcelli, A., Rizzo, G., Fazio, L., Torretta, S., Rampino, A., Berry, A.,  
773 Gass, P., Cirulli, F., Rietschel, M., Bertolino, A., Deuschle, M., Szyf, M., Riva, M.A., 2016.  
774 Ankyrin-3 as a molecular marker of early-life stress and vulnerability to psychiatric  
775 disorders. *Transl. Psychiatry* 6, e943. <https://doi.org/10.1038/tp.2016.211>

776 Moczulska, K.E., Pichler, P., Schutzbier, M., Schleiffer, A., Rumpel, S., Mechtler, K., 2014. Deep  
777 and Precise Quantification of the Mouse Synaptosomal Proteome Reveals  
778 Substantial Remodeling during Postnatal Maturation. *J. Proteome Res.* 13, 4310–  
779 4324. <https://doi.org/10.1021/pr500456t>

780 Mohanan, V., Nakata, T., Desch, A.N., Lévesque, C., Boroughs, A., Guzman, G., Cao, Z., Creasey,  
781 E., Yao, J., Boucher, G., Charron, G., Bhan, A.K., Schenone, M., Carr, S.A., Reinecker, H.-  
782 C., Daly, M.J., Rioux, J.D., Lassen, K.G., Xavier, R.J., 2018. C1orf106 is a colitis risk gene  
783 that regulates stability of epithelial adherens junctions. *Science* eaan0814.  
784 <https://doi.org/10.1126/science.aan0814>

785 Nelson, M.R., Tipney, H., Painter, J.L., Shen, J., Nicoletti, P., Shen, Y., Floratos, A., Sham, P.C.,  
786 Li, M.J., Wang, J., Cardon, L.R., Whittaker, J.C., Sanséau, P., 2015. The support of  
787 human genetic evidence for approved drug indications. *Nat. Genet.* advance online  
788 publication. <https://doi.org/10.1038/ng.3314>

789 Nöthen, M.M., Nieratschker, V., Cichon, S., Rietschel, M., 2010. New findings in the genetics  
790 of major psychoses. *Dialogues Clin. Neurosci.* 12, 85–93.

791 Okbay, A., Baselmans, B.M.L., De Neve, J.-E., Turley, P., Nivard, M.G., Fontana, M.A., Meddens,  
792 S.F.W., Linnér, R.K., Rietveld, C.A., Derringer, J., Gratten, J., Lee, J.J., Liu, J.Z., de  
793 Vlaming, R., Ahluwalia, T.S., Buchwald, J., Cavadino, A., Frazier-Wood, A.C., Furlotte,  
794 N.A., Garfield, V., Geisel, M.H., Gonzalez, J.R., Haitjema, S., Karlsson, R., van der Laan,  
795 S.W., Ladwig, K.-H., Lahti, J., van der Lee, S.J., Lind, P.A., Liu, T., Matteson, L., Mihailov,  
796 E., Miller, M.B., Minica, C.C., Nolte, I.M., Mook-Kanamori, D., van der Most, P.J.,  
797 Oldmeadow, C., Qian, Y., Raitakari, O., Rawal, R., Realo, A., Rueedi, R., Schmidt, B.,  
798 Smith, A.V., Stergiakouli, E., Tanaka, T., Taylor, K., Wedenoja, J., Wellmann, J., Westra,  
799 H.-J., Willems, S.M., Zhao, W., LifeLines Cohort Study, Amin, N., Bakshi, A., Boyle, P.A.,  
800 Cherney, S., Cox, S.R., Davies, G., Davis, O.S.P., Ding, J., Direk, N., Eibich, P., Emeny,  
801 R.T., Fatemifar, G., Faul, J.D., Ferrucci, L., Forstner, A., Gieger, C., Gupta, R., Harris,  
802 T.B., Harris, J.M., Holliday, E.G., Hottenga, J.-J., De Jager, P.L., Kaakinen, M.A., Kajantie,  
803 E., Karhunen, V., Kolcic, I., Kumari, M., Launer, L.J., Franke, L., Li-Gao, R., Koini, M.,  
804 Loukola, A., Marques-Vidal, P., Montgomery, G.W., Mosing, M.A., Paternoster, L.,  
805 Pattie, A., Petrovic, K.E., Pulkki-Råback, L., Quaye, L., Räikkönen, K., Rudan, I., Scott,  
806 R.J., Smith, J.A., Sutin, A.R., Trzaskowski, M., Vinkhuyzen, A.E., Yu, L., Zabaneh, D.,  
807 Attia, J.R., Bennett, D.A., Berger, K., Bertram, L., Boomsma, D.I., Snieder, H., Chang, S.-  
808 C., Cucca, F., Deary, I.J., van Duijn, C.M., Eriksson, J.G., Bültmann, U., de Geus, E.J.C.,  
809 Groenen, P.J.F., Gudnason, V., Hansen, T., Hartman, C.A., Haworth, C.M.A., Hayward,  
810 C., Heath, A.C., Hinds, D.A., Hyppönen, E., Iacono, W.G., Järvelin, M.-R., Jöckel, K.-H.,  
811 Kaprio, J., Kardia, S.L.R., Keltikangas-Järvinen, L., Kraft, P., Kubzansky, L.D.,  
812 Lehtimäki, T., Magnusson, P.K.E., Martin, N.G., McGue, M., Metspalu, A., Mills, M., de  
813 Mutsert, R., Oldehinkel, A.J., Pasterkamp, G., Pedersen, N.L., Plomin, R., Polasek, O.,  
814 Power, C., Rich, S.S., Rosendaal, F.R., den Ruijter, H.M., Schlessinger, D., Schmidt, H.,  
815 Svento, R., Schmidt, R., Alizadeh, B.Z., Sørensen, T.I.A., Spector, T.D., Steptoe, A.,  
816 Terracciano, A., Thurik, A.R., Timpson, N.J., Tiemeier, H., Uitterlinden, A.G.,  
817 Vollenweider, P., Wagner, G.G., Weir, D.R., Yang, J., Conley, D.C., Smith, G.D., Hofman,  
818 A., Johannesson, M., Laibson, D.I., Medland, S.E., Meyer, M.N., Pickrell, J.K., Esko, T.,  
819 Krueger, R.F., Beauchamp, J.P., Koellinger, P.D., Benjamin, D.J., Bartels, M., Cesarini,  
820 D., 2016. Genetic variants associated with subjective well-being, depressive

821 symptoms, and neuroticism identified through genome-wide analyses. *Nat. Genet.*  
822 advance online publication. <https://doi.org/10.1038/ng.3552>

823 Pickrell, J.K., Berisa, T., Liu, J.Z., Séguirel, L., Tung, J.Y., Hinds, D.A., 2016. Detection and  
824 interpretation of shared genetic influences on 42 human traits. *Nat. Genet.* 48, 709–  
825 717. <https://doi.org/10.1038/ng.3570>

826 Ruderfer, D.M., Charney, A.W., Readhead, B., Kidd, B.A., Kähler, A.K., Kenny, P.J., Keiser, M.J.,  
827 Moran, J.L., Hultman, C.M., Scott, S.A., Sullivan, P.F., Purcell, S.M., Dudley, J.T., Sklar, P.,  
828 2016. Polygenic overlap between schizophrenia risk and antipsychotic response: a  
829 genomic medicine approach. *Lancet Psychiatry* 3, 350–357.  
830 [https://doi.org/10.1016/S2215-0366\(15\)00553-2](https://doi.org/10.1016/S2215-0366(15)00553-2)

831 Ruderfer, D.M., Fanous, A.H., Ripke, S., McQuillin, A., Amdur, R.L., Gejman, P.V., O'Donovan,  
832 M.C., Andreassen, O.A., Djurovic, S., Hultman, C.M., Kelsoe, J.R., Jamain, S., Landén, M.,  
833 Leboyer, M., Nimgaonkar, V., Nurnberger, J., Smoller, J.W., Craddock, N., Corvin, A.,  
834 Sullivan, P.F., Holmans, P., Sklar, P., Kendler, K.S., 2014. Polygenic dissection of  
835 diagnosis and clinical dimensions of bipolar disorder and schizophrenia. *Mol.*  
836 *Psychiatry* 19, 1017–1024. <https://doi.org/10.1038/mp.2013.138>

837 Schizophrenia Working Group of the Psychiatric Genomics Consortium, 2014. Biological  
838 insights from 108 schizophrenia-associated genetic loci. *Nature* 511, 421–427.  
839 <https://doi.org/10.1038/nature13595>

840 Sekar, A., Bialas, A.R., de Rivera, H., Davis, A., Hammond, T.R., Kamitaki, N., Tooley, K.,  
841 Presumey, J., Baum, M., Van Doren, V., Genovese, G., Rose, S.A., Handsaker, R.E.,  
842 Schizophrenia Working Group of the Psychiatric Genomics Consortium, Daly, M.J.,  
843 Carroll, M.C., Stevens, B., McCarroll, S.A., 2016. Schizophrenia risk from complex  
844 variation of complement component 4. *Nature* 530, 177–183.  
845 <https://doi.org/10.1038/nature16549>

846 Shi, H., Kichaev, G., Pasaniuc, B., 2016. Contrasting the Genetic Architecture of 30 Complex  
847 Traits from Summary Association Data. *Am. J. Hum. Genet.* 99, 139–153.  
848 <https://doi.org/10.1016/j.ajhg.2016.05.013>

849 Stahl, E., Forstner, A., McQuillin, A., Ripke, S., Pgc, B.D.W.G. of the, Ophoff, R., Scott, L.,  
850 Cichon, S., Andreassen, O.A., Sklar, P., Kelsoe, J., Breen, G., 2017. Genomewide  
851 association study identifies 30 loci associated with bipolar disorder. *bioRxiv*  
852 173062. <https://doi.org/10.1101/173062>

853 Stone, J.L., O'Donovan, M.C., Gurling, H., Kirov, G.K., Blackwood, D.H.R., Corvin, A., Craddock,  
854 N.J., Gill, M., Hultman, C.M., Lichtenstein, P., McQuillin, A., Pato, C.N., Ruderfer, D.M.,  
855 Owen, M.J., Clair, D.S., Sullivan, P.F., Sklar, P., (Leader), S.M.P., Korn, J., Macgregor, S.,  
856 Morris, D.W., O'Dushlaine, C.T., Daly, M.J., Visscher, P.M., Holmans, P.A., Purcell, S.M.,  
857 Scolnick, E.M., (Leader), P.S., Williams, N.M., Georgieva, L., Nikolov, I., Norton, N.,  
858 Williams, H., Toncheva, D., Milanova, V., Thelander, E.F., Sullivan, P., Kenny, E.,  
859 Waddington, J.L., Choudhury, K., Datta, S., Pimm, J., Thirumalai, S., Puri, V., Krasucki,  
860 R., Lawrence, J., Quedsted, D., Bass, N., Curtis, D., Crombie, C., Fraser, G., Kwan, S.L.,  
861 Walker, N., Muir, W.J., McGhee, K.A., Pickard, B., Malloy, P., Maclean, A.W., Beck, M.V.,  
862 Pato, M.T., Medeiros, H., Middleton, F., Carvalho, C., Morley, C., Fanous, A., Conti, D.,  
863 Knowles, J.A., Ferreira, C.P., Macedo, A., Azevedo, M.H., McCarroll, S.A., Daly, M.,  
864 Chambert, K., Gates, C., Gabriel, S.B., Mahon, S., Ardlie, K., 2008. Rare chromosomal  
865 deletions and duplications increase risk of schizophrenia. *Nature* 455, 237–241.  
866 <https://doi.org/10.1038/nature07239>

867 Szatkiewicz, J.P., O'Dushlaine, C., Chen, G., Chambert, K., Moran, J.L., Neale, B.M., Fromer, M.,  
868 Ruderfer, D., Akterin, S., Bergen, S.E., Kähler, A., Magnusson, P.K.E., Kim, Y., Crowley,  
869 J.J., Rees, E., Kirov, G., O'Donovan, M.C., Owen, M.J., Walters, J., Scolnick, E., Sklar, P.,  
870 Purcell, S., Hultman, C.M., McCarroll, S.A., Sullivan, P.F., 2014. Copy number variation  
871 in schizophrenia in Sweden. *Mol. Psychiatry* 19, 762–773.  
872 <https://doi.org/10.1038/mp.2014.40>

873 The 1000 Genomes Project Consortium, 2015. A global reference for human genetic  
874 variation. *Nature* 526, 68–74. <https://doi.org/10.1038/nature15393>

875 Visscher, P.M., Wray, N.R., Zhang, Q., Sklar, P., McCarthy, M.I., Brown, M.A., Yang, J., 2017. 10  
876 Years of GWAS Discovery: Biology, Function, and Translation. *Am. J. Hum. Genet.*  
877 101, 5–22. <https://doi.org/10.1016/j.ajhg.2017.06.005>

878 Watanabe, K., Taskesen, E., Bochoven, A. van, Posthuma, D., 2017. FUMA: Functional  
879 mapping and annotation of genetic associations. *bioRxiv* 110023.  
880 <https://doi.org/10.1101/110023>

881 Westra, H.-J., Peters, M.J., Esko, T., Yaghootkar, H., Schurmann, C., Kettunen, J., Christiansen,  
882 M.W., Fairfax, B.P., Schramm, K., Powell, J.E., Zhernakova, A., Zhernakova, D.V.,  
883 Veldink, J.H., Van den Berg, L.H., Karjalainen, J., Withoff, S., Uitterlinden, A.G.,  
884 Hofman, A., Rivadeneira, F., 't Hoen, P.A.C., Reinmaa, E., Fischer, K., Nelis, M., Milani,  
885 L., Melzer, D., Ferrucci, L., Singleton, A.B., Hernandez, D.G., Nalls, M.A., Homuth, G.,  
886 Nauck, M., Radke, D., Völker, U., Perola, M., Salomaa, V., Brody, J., Suchy-Dicey, A.,  
887 Gharib, S.A., Enquobahrie, D.A., Lumley, T., Montgomery, G.W., Makino, S., Prokisch,  
888 H., Herder, C., Roden, M., Grallert, H., Meitinger, T., Strauch, K., Li, Y., Jansen, R.C.,  
889 Visscher, P.M., Knight, J.C., Psaty, B.M., Ripatti, S., Teumer, A., Frayling, T.M.,  
890 Metspalu, A., van Meurs, J.B.J., Franke, L., 2013. Systematic identification of trans  
891 eQTLs as putative drivers of known disease associations. *Nat. Genet.* 45, 1238–1243.  
892 <https://doi.org/10.1038/ng.2756>

893 Whiteford, H.A., Degenhardt, L., Rehm, J., Baxter, A.J., Ferrari, A.J., Erskine, H.E., Charlson,  
894 F.J., Norman, R.E., Flaxman, A.D., Johns, N., Burstein, R., Murray, C.J., Vos, T., 2013.  
895 Global burden of disease attributable to mental and substance use disorders:  
896 findings from the Global Burden of Disease Study 2010. *The Lancet* 382, 1575–1586.  
897 [https://doi.org/10.1016/S0140-6736\(13\)61611-6](https://doi.org/10.1016/S0140-6736(13)61611-6)

898 WHO | International Classification of Diseases [WWW Document], n.d. . WHO. URL  
899 <http://www.who.int/classifications/icd/en/> (accessed 3.14.17).

900 Yamashita, S., Miyake, N., Matsumoto, N., Osaka, H., Iai, M., Aida, N., Tanaka, Y., 2013.  
901 Neuropathology of leukoencephalopathy with brainstem and spinal cord  
902 involvement and high lactate caused by a homozygous mutation of DARS2. *Brain*  
903 *Dev.* 35, 312–316. <https://doi.org/10.1016/j.braindev.2012.05.007>

904 Yang, S., Van Dongen, H.P.A., Wang, K., Berrettini, W., Bućan, M., 2008. Assessment of  
905 circadian function in fibroblasts of patients with bipolar disorder. *Mol. Psychiatry*  
906 14, 143–155. <https://doi.org/10.1038/mp.2008.10>

907 Zhu, Z., Zhang, F., Hu, H., Bakshi, A., Robinson, M.R., Powell, J.E., Montgomery, G.W.,  
908 Goddard, M.E., Wray, N.R., Visscher, P.M., Yang, J., 2016. Integration of summary data  
909 from GWAS and eQTL studies predicts complex trait gene targets. *Nat. Genet.* 48,  
910 481–487. <https://doi.org/10.1038/ng.3538>

911

912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934

**Figure Legends**

**Figure 1. Associated Genomic Loci Shared and Divergent Between BD and SCZ**

a) Odds ratios (OR) from independent data sets of BD (blue) and SCZ (red) for each of the 114 genome-wide significant variants in the BD and SCZ vs controls GWAS. b) Manhattan plot for SCZ vs BD GWAS.

**Figure 2. Polygenic Risk Score Dissection of Clinical Symptom Dimensions**

Effect size (calculated by dividing regression estimate by standard error) from regression analysis including ancestry covariates for each subphenotype and PRS for BD (x-axis) and SCZ (y-axis). Point size represents  $-\log_{10}(\text{p-value})$  with SCZ (red) and BD (blue). Numbered subphenotypes are 1) comorbid migraine, 2) panic attacks 3) suicide attempt 4) mixed states 5) rapid cycling 6) comorbid eating disorder 7) comorbid OCD 8) year of birth 9) suicide ideation 10) panic disorder 11) number of suicide attempts 12) depressive symptoms (SCZ) 13) episodes depressive 14) episodes total 15) positive symptoms (SCZ) 16) irritable mania 17) age of onset depression 18) family history 19) episodes mixed mania 20) unipolar mania 21) alcohol substance dependence 22) age of onset mania 23) age at interview 24) number of hospitalizations. All subphenotypes are in BD except those labeled (SCZ).

935 **Table Legends**

936

937 **Table 1. Most Significant Associated Loci from SCZ vs BD GWAS**

938 Association results for the five most significant variants in the SCZ vs BD GWAS with the top  
939 two being genome-wide significant. Each variant includes results from the independent BD vs  
940 controls and SCZ vs controls GWAS and the comparable p-value from a heterogeneity test when  
941 performing a two cohort meta-analysis of SCZ and BD.

942

943 **Table 2. Complete Results of Polygenic Risk Score Dissection Analysis**

944 Polygenic scoring results of all four GWAS phenotypes (BD+SCZ vs controls, BD vs controls,  
945 SCZ vs controls and SCZ vs BD) and 24 subphenotypes from BD and 4 subphenotypes from  
946 SCZ, rows without case/control counts are quantitative measures. Significance and effects are  
947 from regression analysis of subphenotype on PRS including principal components of ancestry  
948 and site as covariates. Effect is the regression estimate divided by the standard error.

949

950 **Supplementary Figure Legends**

951

952 **Figure S1. Related to Figure 1b. Regional Association Plot and Forest Plot for the First**  
953 **Genome-wide Significant Hit in the SCZ vs BD GWAS.**

954 **Figure S2. Related to Figure 1b. Regional Association Plot and Forest Plot for the Second**  
955 **Genome-wide Significant Hit in the SCZ vs BD GWAS.**

956

957

958 **Figure S3. Related to Summary-data-based Mendelian Randomization. Detailed**  
959 **Association of DCAKD from SMR.**

960 Results at the *DCAKD* locus from SMR analysis of SCZ vs BD. Top plot, brown dots represent  
961 the *P* values for SNPs from SCZ vs BD GWAS, diamonds represent the *P* values for probes from  
962 the SMR test. Bottom plot, the eQTL *P* values of SNPs from the Westra study for the  
963 ILMN\_1811648 probe tagging *DCAKD*. The top and bottom plots include all the SNPs available  
964 in the region in the GWAS and eQTL summary data, respectively, rather than only the SNPs  
965 common to both data sets. Highlighted in red is the gene (*DCAKD*) that passed the SMR and  
966 HEIDI tests.

967

968 **Figure S4. Related to Regional SNP-heritability estimation. Heritability Estimates for BD**  
969 **and SCZ in Genome-wide Significant Regions of BD and SCZ.**

970 Regional SNP-heritability estimates for SCZ and BD stratified by whether the region contains  
971 the most significant variant in a genome-wide significant locus in BD, SCZ, neither or both.

972

973

974 **STAR Methods**

975 **CONTACT FOR REAGENT AND RESOURCE SHARING**

976 Genotype and phenotype data use is restricted and governed by the Psychiatric Genetics  
977 Consortium. Further information and requests for analytical results or additional information  
978 should be directed to and will be fulfilled by the Lead Contact, Douglas Ruderfer  
979 ([douglas.ruderfer@vanderbilt.edu](mailto:douglas.ruderfer@vanderbilt.edu)).

980

981 **SUBJECT DETAILS**

982 **Genotyped Sample Description**

983 SCZ samples are a substantial subset of those analyzed previously(Schizophrenia Working  
984 Group of the Psychiatric Genomics Consortium, 2014). BD samples are the newest collection  
985 from Psychiatric Genomics Consortium Bipolar Disorder Working Group(Stahl et al., 2017).

986 Below we provide information on the individual samples used here as provided by the original  
987 PGC disorder publications. Additionally, most studies have been described in detail in the  
988 citations provided. The boldfaced first line for each sample is study PI, PubMed ID, country  
989 (study name), and the PGC internal tag or study identifier.

990

991 **European ancestry, case-control design**

992 *Schizophrenia*

993 **Adolfsson, R | NP | Umeå, Sweden | scz\_umeb\_eur**

994 **Adolfsson, R | NP | Umeå, Sweden | scz\_umes\_eur**

995 Cases of European ancestry were ascertained from multiple different studies of schizophrenia  
996 (1992-2009). The diagnostic processes were similar between studies, and the final diagnosis is a  
997 best-estimate consensus lifetime diagnosis based on multiple sources of information such as  
998 clinical evaluation by research psychiatrists, different types of semi-structured interviews made  
999 by trained research nurses and research psychiatrists, medical records, course of the disease and  
1000 data from multiple informants. Diagnosis was made in accordance with the Diagnostic and  
1001 Statistical Manual of Mental Disorders-Version IV (DSM-IV) or International Classification of  
1002 Diseases, 10th Revision (ICD-10) criteria. Controls were recruited from the Betula study, an  
1003 ongoing longitudinal, prospective, population-based study from the same geographic area (North



1004 Sweden) that is studying aging, health, and cognition in adults. All subjects (cases and controls)  
1005 participated after giving written informed consent and the regional Ethical Review Board at the  
1006 University of Umeå approved all original studies and participation in the PGC. GWAS  
1007 genotyping was performed at Broad Institute.

1008 **Andreassen, O | 19571808 | Norway (TOP) | scz\_top8\_eur**

1009 In the TOP study (Tematisk område psykoser), cases of European ancestry, born in Norway,  
1010 were recruited from psychiatric hospitals in the Oslo region. Patients were diagnosed according  
1011 to SCID and further ascertainment details have been reported. Healthy control subjects were  
1012 randomly selected from statistical records of persons from the same catchment area as the patient  
1013 groups. All participants provided written informed consent and the human subjects protocol was  
1014 approved by the Norwegian Scientific-Ethical Committee and the Norwegian Data Protection  
1015 Agency.

1016 **Blackwood, D | 19571811 | Edinburgh, UK | scz\_edin\_eur**

1017 Cases and controls were recruited from the southeast of Scotland, and ascertainment has been  
1018 previously described as part of the International Schizophrenia Consortium studies. All  
1019 participating subjects gave written, informed consent and the human subjects protocol was  
1020 approved by the Scotland A Research Ethics Committee. DNA samples were genotyped at the  
1021 Broad Institute.

1022 **Børglum, A | 19571808 | Denmark | scz\_aarh\_eur**

1023 DNA samples for all subjects were collected from blood spots systematically collected by the  
1024 Danish Newborn Screening Biobank), with case/control status established using the Danish  
1025 Psychiatric Central Register. Cases were diagnosed clinically according to ICD-10 criteria.

1026 Controls were selected to match the cases by birth cohort. The Danish Data Protection Agency  
1027 and the ethics committees in Denmark approved the human subjects protocol.

1028 **Bramon | 23871474 | Seven countries (PEIC, WTCCC2) | scz\_pewb\_eur**

1029 **Bramon | 23871474 | Spain (PEIC, WTCCC2) | scz\_pewb\_eur**

1030 The Psychosis Endophenotypes International Consortium (PEIC) was part of WTCCC2. Samples  
1031 were collected through seven centers in Europe and Australia (the Institute of Psychiatry, King's  
1032 College London, London; GROUP (consisting of the University of Amsterdam, Amsterdam; the  
1033 University of Groningen, Groningen; Maastricht University Medical Centre, Maastricht; and the  
1034 University of Utrecht, Utrecht); the University of Western Australia, Perth; the Universidad de  
1035 Cantabria, Santander; the University of Edinburgh, Edinburgh; Heidelberg University,  
1036 Heidelberg and Ludwig-Maximilians-Universität München, Munich). To allow for a DSM-IV  
1037 diagnosis to be ascertained or ruled out, all participants (including controls and unaffected family  
1038 members) underwent a structured clinical interview with the Schedule for Affective Disorders  
1039 and Schizophrenia (SADS), the Structured Clinical Interview for DSM Disorders (SCID), or the  
1040 Schedules for Clinical Assessment in Neuropsychiatry (SCAN). We included cases with  
1041 schizophrenia and schizoaffective disorder. Participants in all groups were excluded if they had a  
1042 history of neurological disease or head injury resulting in loss of consciousness.

1043 **Buxbaum, J | 20489179 | New York, US & Israel | scz\_msaf\_eur**

1044 Samples contributed by Mount Sinai were derived from three cohorts. In all cohorts, ethical  
1045 approval was obtained from all participating sites, and all subjects provided informed consent.  
1046 Two of the cohorts were in a prior paper on copy number variation. One of the cohorts was from  
1047 the Mount Sinai brain bank, where DNA was extracted from postmortem samples, and another

1048 comprised of patients ascertained in Israel. The third cohort included subjects more recently  
1049 recruited through the Mount Sinai Conte Center.

1050 **Corvin, A | 19571811 | Ireland | scz\_dubl\_eur**

1051 The case sample was collected primarily in the Dublin area and the ascertainment procedure has  
1052 been previously described. The controls were recruited, from the same region through the Irish  
1053 Blood Transfusion Services. All participants gave written, informed consent and the collections  
1054 were approved through the Federated Dublin Hospitals and Irish Blood Transfusion Services  
1055 Research Ethics Committees, respectively. DNA samples were genotyped at the Broad Institute.

1056 **Corvin, A; Riley, B | 22883433 | Ireland (WTCCC2) | scz\_irwt\_eur**

1057 The case sample was recruited from the Republic of Ireland and Northern Ireland. All cases had  
1058 four Irish grandparents and ascertainment details have been reported elsewhere. Ethics approval  
1059 was obtained from all participating hospitals and centers. Controls were blood donors from the  
1060 Irish Blood Transfusion Service, whose Ethics Committee approved the human subjects  
1061 protocol. All participants gave written informed consent. Samples were genotyped at Affymetrix  
1062 (Santa Clara, California, US) laboratory as part of the WTCCC2 genotyping pipeline.

1063 **Ehrenreich, H | 20819981 | Germany (GRAS) | scz\_gras**

1064 The Gottingen Research Association for Schizophrenia (GRAS) collection included cases  
1065 recruited across 23 German hospitals. Controls were unscreened blood donors recruited at the  
1066 Georg-August-University according to national blood donation guidelines. Cases completed a  
1067 structured clinical interview and were diagnosed with DSM-IV schizophrenia or schizoaffective  
1068 disorder. The study was approved by the Georg-August-University ethics committee and local  
1069 internal review boards of the participating centers. All participants gave written informed  
1070 consent.

1071 **Esko, T | 15133739 | Estonia (EGCUT) | scz\_egcu\_eur**

1072 The Estonian cohort comes from the population-based biobank of the Estonian Genome Project

1073 of University of Tartu (EGCUT). The project was conducted according to the Estonian Gene

1074 Research Act and all participants provided informed consent (www.biobank.ee). In total, 52,000

1075 individuals aged 18 years or older participated in this cohort (33% men, 67% women). The

1076 population distributions of the cohort reflect those of the Estonian population (83% Estonians,

1077 14% Russians and 3% other). General practitioners (GP) and physicians in the hospitals

1078 randomly recruited the participants. A Computer-Assisted Personal interview was conducted

1079 over 1-2 ours at doctors' offices. Data on demographics, genealogy, educational and

1080 occupational history, lifestyle and anthropometric and physiological data were assessed.

1081 Schizophrenia was diagnosed prior to the recruitment by a psychiatrist according to ICD-10

1082 criteria and identified from the Estonian Biobank phenotype database. Controls were drawn from

1083 a larger pool of genotyped biobank samples by matching on gender, age and genetic ancestry.

1084 All the controls were population-based and have not been sampled for any specific disease.

1085 **Esko, T; Li, Q; Dominici E | 15133739, 24166486 | J&J and Roche cases, EGCUT controls |**

1086 **scz\_jr3a\_eur**

1087 **Esko, T; Li, Q; Domenici E | 15133739, 24166486 | J&J and Roche cases, EGCUT controls |**

1088 **scz\_jr3b\_eur**

1089 **Esko, T; Li, Q; Domenici E | 15133739, 24166486 | J&J and Roche cases, EGCUT controls |**

1090 **scz\_jri6\_eur**

1091 **Esko, T; Li, Q; Dominici E | 15133739, 24166486 | J&J and Roche cases cases, EGCUT**

1092 **controls | scz\_jrsa\_eur**

1093 Cases were collected by Johnson and Johnson (J&J) and Roche as part of clinical collaborations  
1094 with hospitals and outpatient centers. Cases were diagnosed according to DSMIV criteria, with  
1095 medical record review by a trained psychiatrist. There were reliability trials across centers for the  
1096 J&J studies. The J& J cases were mostly collected in Eastern Europe, with most coming from  
1097 Estonian and Russia (>100); intermediate numbers from Austria, the Czech Republic, Latvia,  
1098 Lithuania, and Spain (50-100); and smaller collections from Bulgaria, Hungary, and Poland  
1099 (<50). The Roche cases were assessed with a structured psychiatric assessment by trained  
1100 interviewers. Most of the Eastern European controls were from the Estonian Biobank project  
1101 (EGCUT) and were ancestrally matched with cases from the J&J sample.

1102 **Gejman, P | 19571809 | US, Australia (MGS) | scz\_mgs2\_eur**

1103 European ancestry case samples were collected by the Molecular Genetics of Schizophrenia  
1104 (MGS) collaboration across multiple sites in the USA and Australia as described in detail  
1105 elsewhere. Cases gave written informed consent, and IRBs at each collecting site approved the  
1106 human subjects protocol. A survey company (Knowledge Networks, under MGS guidance)  
1107 collected the European ancestry control sample and ascertainment is described in detail  
1108 elsewhere. DNA samples were genotyped at the Broad Institute.

1109 **Gurling, H | 19571811 | London, UK | scz\_uclo\_eur**

1110 All cases and controls were collected by University College London and had both parents from  
1111 England, Scotland or Wales. All participants gave written informed consent and the U.K.  
1112 National Health Service multicenter and local research ethics committee approved the human  
1113 subjects protocol. Further details on ascertainment are available elsewhere. The samples were  
1114 genotyped at the Broad Institute.

1115 **Jönsson, E | 19571808 | Sweden (Hubin) | scz\_ersw\_eur**

1116 Cases were recruited from northwestern Stockholm County and ascertainment has been  
1117 described previously. Cases gave informed consent and the human subjects protocol was  
1118 approved by the ethical committees of the Karolinska Hospital and the Stockholm Regional  
1119 Ethical Committee. Controls were recruited either among subjects previously participating in  
1120 biological research at the Karolinska Institute or drawn from a representative register of the  
1121 population of Stockholm County. All participants provided informed consent.

1122 **Kirov, G | Not published | Bulgaria | scz\_buls\_eur**

1123 All cases were recruited from Bulgaria and had a history of hospitalization for treatment of  
1124 schizophrenia. Controls were recruited from the two largest cities in Bulgaria as previously  
1125 described. All participants gave written informed consent and the study was approved by local  
1126 ethics committees at the participating centers.

1127 **Knight, J; Collier DA; Nisenbaum L| Not published | Canada (Toronto) -US(Lilly)-US  
1128 (MIGen)| scz\_lktu\_eur**

1129 Toronto cases were recruited by referral and advertisement. Diagnoses were made according to  
1130 DSM-III or DSM-IV criteria following interview and medical record review. US cases were  
1131 recruited from schizophrenia clinical trials in a range of settings as part of a trial with Eli Lilly.  
1132 Diagnoses were made according to DSM-III or DSM-IV criteria following interview by  
1133 psychiatrist and medical record review. No controls were sampled as part of the study, and  
1134 ancestrally-matched controls were chosen from the Myocardial Infarction Genetics Consortium  
1135 (MIGen, dbGaP ID phs000294.v1.p1) that was genotyped with the same SNP array.

1136 **Lencz, T; Darvasi A | 23325106 | Israel | scz\_ajsz\_eur**

1137 Cases and controls were sampled from an Ashkenazi Jewish repository (Hebrew University  
1138 Genetic Resource, <http://hugr.huji.ac.il>). Patients were recruited from hospitalized inpatients at 7

1139 medical centers in Israel and were diagnosed with DSM-IV schizophrenia or schizoaffective  
1140 disorder. Controls were sampled through the Israeli Blood Bank and did not report any chronic  
1141 disease or regularly prescribed medication at the time of assessment. Full ascertainment details  
1142 have previously been reported. Local ethics committees and the National Genetic Committee of  
1143 the Israeli Ministry of Health approved the studies and all participants gave informed, written  
1144 consent.

1145 **Levinson, D | 22885689 | Six countries, WTCCC controls | scz\_lacw\_eur**

1146 Cases collected as part of a larger pedigree-based study were partitioned into two subsamples.  
1147 Cases with two genotyped parents were analyzed as trios (see PI Levinson, ms.scz\_lemu\_eur in  
1148 the Trio section below). Unrelated cases who could not be used as part of a trio were included as  
1149 a separate case-control analysis, using independent controls, matched by ancestry and  
1150 genotyping array, from the Wellcome Trust Case Control Consortium. Cases were identified  
1151 from different clinical settings (e.g. inpatients, outpatients and community facilities) in six  
1152 countries (Australia, France, Germany, Ireland, UK, and the US). Diagnoses were established  
1153 using semi-structured interviews, psychiatric records and informant reports. Case subjects were  
1154 diagnosed with schizophrenia or schizoaffective disorder according to DSM-III-R criteria. All  
1155 protocols were approved by local IRBs, and all cases provided written informed consent.

1156 **Malhotra, A | 17522711 | New York, US | scz\_zhh1\_eur**

1157 The case and control subjects were recruited in the New York metropolitan area and  
1158 ascertainment methods have been described previously. All participants gave written, informed  
1159 consent and the IRB of the North Shore-Long Island Jewish Health System approved the human  
1160 subjects protocols. DNA was genotyped at Zucker Hillside.

1161 **Mowry, B | 21034186 | Australia | scz\_asrb\_eur**

1162 These subjects were part of the Australian Schizophrenia Research Bank. The case sample was  
1163 recruited in four Australian States (New South Wales, Queensland, Western Australia and  
1164 Victoria) through hospital inpatient units, community mental health services, outpatient clinics  
1165 and rehabilitation services, non-government mental illness support organizations, and, in the  
1166 initial stages, through a large-scale, national, multi-media advertising campaign. This sample is  
1167 comprised of 509 cases from larger metropolitan centers of Brisbane, Newcastle, Sydney,  
1168 Melbourne, and Perth. Cases gave written informed consent, and the human subjects protocol  
1169 was initially approved by the Hunter New England Area Health Research Committee and  
1170 subsequently approved by relevant Institutional Ethics Committees in Brisbane, Sydney,  
1171 Melbourne and Perth. Healthy controls were recruited through multi-media advertisements, and  
1172 other sources. Controls were from the metropolitan centers of Brisbane, Newcastle, Sydney,  
1173 Melbourne, and Perth. Controls gave written informed consent, and the human subjects protocol  
1174 was approved by the Hunter New England Area Health Research Committee and Institutional  
1175 Ethics Committees in Brisbane, Sydney, Melbourne and Perth. The samples were genotyped in  
1176 two stages at the Hunter Medical Research Institute, University of Newcastle, Newcastle,  
1177 Australia.

1178 **O'Donovan, M: Owen, M | 19571811 | Cardiff, UK | scz\_caws\_eur**

1179 The case sample included European ancestry schizophrenia cases recruited in the British Isles  
1180 and described previously. All cases gave written informed consent to. The study was approved  
1181 by the Multicentre Research Ethics Committee in Wales and Local Research Ethics Committees  
1182 from all participating sites. The control sample used the Wellcome Trust CaseControl  
1183 Consortium (WTCCC) sample described elsewhere, but included similar numbers of individuals



1184 from the 1958 British Birth Cohort and a panel of consenting blood donors (UK Blood Service).  
1185 Samples were genotyped at Affymetrix service lab (San Francisco, USA).

1186 **O'Donovan, M: Owen, M: Walters, J | 22614287 | UK (CLOZUK) | scz\_clm2\_eur**

1187 **O'Donovan, M: Owen, M: Walters, J | 22614287 | UK (CLOZUK) | scz\_clo3\_eur**

1188 CLOZUK cases were taking the antipsychotic clozapine and had received a clinical diagnosis of  
1189 treatment-resistant schizophrenia. Patients taking clozapine provide blood samples to allow  
1190 detection of adverse drug-effects. Through collaboration with Novartis (the manufacturer of a  
1191 proprietary form of clozapine, Clozaril), we acquired blood from people with treatment-resistant  
1192 schizophrenia according to the clozapine registration forms completed by treating psychiatrists  
1193 as previously reported. The samples were genotyped at the Broad Institute. The UK Multicentre  
1194 Research Ethics Committee (MREC) approved the study. The controls were drawn from the  
1195 WTCCC2 control samples (~3,000 from the 1958 British Birth Cohort and ~3,000 samples from  
1196 the UK Blood Service Control Group). An additional 900 controls, held by Cardiff University,  
1197 were recruited from the UK National Blood Transfusion Service. They were not specifically  
1198 screened for psychiatric illness. All control samples were from participants who provided  
1199 informed consent.

1200 **Ophoff, R | 19571808 | Netherlands | scz\_ucla\_eur**

1201 The case sample consisted of inpatients and outpatients recruited through psychiatric hospitals  
1202 and institutions throughout the Netherlands. Cases with DSM-IV schizophrenia were included in  
1203 the analysis. Further details on ascertainment are provided elsewhere. Controls came from the  
1204 University Medical Centre Utrecht and were volunteers with no psychiatric history. Ethical  
1205 approval was provided by local ethics committees and all participants gave written informed  
1206 consent.

1207 **Palotie, A | 19571808 | Finland | scz\_fi3m\_eur**

1208 **Palotie, A | Not published | Finnish | scz\_fii6\_eur**

1209 Finnish cases were drawn from a nationwide collection of families with schizophrenia spectrum  
1210 disorders. The control sample was derived from the Finnish Health 2000 survey. All participants  
1211 provided written informed consent and approval was obtained from the ethics committees at each  
1212 location.

1213 **Pato, C | 19571811 | Portugal | scz\_port\_eur**

1214 Cases and controls lived in Portugal, the Azorean and Madeiran islands, or were the direct  
1215 (first or second-generation) Portuguese immigrant population in the US, as previously described.  
1216 Controls were not biologically related to cases. All participants gave written informed consent  
1217 and the IRB of SUNY Upstate Medical University approved the protocol. The samples were  
1218 genotyped at the Broad Institute.

1219 **Petryshen, T | 24424392 | Boston, US (CIDAR) | scz\_cims\_eur**

1220 Cases were recruited from inpatient and outpatient settings in the Boston area by clinician  
1221 referral, through review of medical records, or through advertisements in local media. Cases  
1222 were diagnosed with DSM-IV schizophrenia through a structured clinical interview (SCID) by  
1223 trained interviewers with review of medical records and a best estimate diagnostic procedure  
1224 including reliability trials across interviewers. A psychiatrist or a PhD-level mental health  
1225 professional made the final diagnostic determination. Controls were ascertained through local  
1226 advertisements from the same geographical area. Ethical approval was provided by local ethics  
1227 committees and all participants gave written informed consent.

1228 **Rietschel/Rujescu/Nöthen | 19571808 | Bonn/Mannheim, Germany | scz\_boco\_eur**

1229 These German samples were collected by separate groups within the MooDS Consortium in  
1230 Mannheim, Bonn, Munich and Jena. For the PGC analyses, the samples were combined by chip  
1231 and ancestry. In Bonn/Mannheim, cases were ascertained as previously described. Controls were  
1232 drawn from three population-based epidemiological studies (PopGen), the Cooperative Health  
1233 Research in the Region of Augsburg (KORA) study, and the Heinz Nixdorf Recall (HNR) study.  
1234 All participants gave written informed consent and the local ethics committees approved the  
1235 human subjects protocols. Additional controls were randomly selected from a Munich-based  
1236 community sample and screened for the presence of anxiety and affective disorders using the  
1237 Composite International Diagnostic Screener. Only individuals negative for the above mentioned  
1238 disorders were included in the sample.

1239 **Rujescu, D | 19571808 | Munich, Germany | scz\_munc\_eur**

1240 For the Munich sample, cases were ascertained from the Munich area of Germany, as described  
1241 previously. The controls were unrelated volunteers randomly selected from the general  
1242 population of Munich. All were screened to exclude a history of psychosis/central neurological  
1243 disease either personally or in a first-degree relative. All participants gave written informed  
1244 consent and the local ethics committees approved the human subjects protocols.

1245 **St Clair, D | 19571811 | Aberdeen, UK | scz\_aber\_eur**

1246 Ascertainment and inclusion/exclusion criteria for cases and controls have been previously  
1247 described. All participating subjects were born in the UK (95% Scotland) and gave written  
1248 informed consent. Both local and multiregional academic ethical committee approved the human  
1249 subjects protocol. The samples were genotyped at the Broad Institute.

1250 **Sullivan, PF | 18347602 | US (CATIE) | scz\_cati\_eur**

1251 Cases were collected as part of the Clinical Antipsychotics Trials of Intervention Effectiveness  
1252 (CATIE) project and ascertainment was previously described. Participants were recruited from  
1253 multiple sites in the USA with informed written consent and approval from the IRBs at each  
1254 CATIE site and the University of North Carolina (Chapel Hill). The control subjects were  
1255 collected by MGS (described above) and gave online informed consent and were fully  
1256 anonymized. There was no overlap with controls included in the MGS collaboration sample.

1257 **Sullivan, PF; Sklar P; Hultman C | 23974872 | Sweden | scz\_swe1\_eur**

1258 **Sullivan, PF; Sklar P; Hultman C | 23974872 | Sweden | scz\_s234\_eur**

1259 **Sullivan, PF; Sklar P; Hultman C | 23974872 | Sweden | scz\_swe5\_eur**

1260 **Sullivan, PF; Sklar P; Hultman C | 23974872 | Sweden | scz\_swe6\_eur**

1261 Samples from the Swedish Schizophrenia Study were collected in a multi-year project and  
1262 genotypes in six batches (sw1-6). All procedures were approved by ethical committees at the  
1263 Karolinska Institutet and the University of North Carolina, and all subjects provided written  
1264 informed consent (or legal guardian consent and subject assent). All samples were genotyped at  
1265 the Broad Institute. Cases with schizophrenia were identified via the Swedish Hospital Discharge  
1266 Register which captures all public and private inpatient hospitalizations. The register is complete  
1267 from 1987 and is augmented by psychiatric data from 1973-1986. The register contains  
1268 International Classification of Disease discharge diagnoses made by attending physicians for  
1269 each hospitalization. Case inclusion criteria included  $\geq 2$  hospitalizations with a discharge  
1270 diagnosis of schizophrenia, both parents born in Scandinavia and age  $\geq 18$  years. Case exclusion  
1271 criteria included hospital register diagnosis of any medical or psychiatric disorder mitigating a  
1272 confident diagnosis of schizophrenia as determined by expert review. The validity of this case  
1273 definition of schizophrenia was strongly supported by clinical, epidemiological, genetic

1274 epidemiological and genetic evidence. Controls were selected at random from Swedish  
1275 population registers, with the goal of obtaining an appropriate control group and avoiding ‘super-  
1276 normal’ controls. Control inclusion criteria included never being hospitalized for schizophrenia  
1277 or bipolar disorder (given evidence of genetic overlap with schizophrenia), both parents born in  
1278 Scandinavia and age of  $\geq 18$  years.

1279 **Walters, J | 21850710 | Cardiff, UK (CogUK) | scz\_cou3\_eur**

1280 Cases were recruited from community mental health teams in Wales and England on the basis of  
1281 a clinical diagnosis of schizophrenia or schizoaffective disorder (depressed sub-type) as  
1282 described previously. 35 Diagnosis was confirmed following a SCAN interview and review of  
1283 case notes followed by consensus diagnosis according to DSM-IV criteria. The samples were  
1284 genotyped at the Broad Institute. The UK Multicentre Research Ethics Committee (MREC)  
1285 approved the study and all participants provided valid informed consent.

1286 **Weinberger, D | 11381111 | NIMH CBDB | scz\_lie2\_eur**

1287 **Weinberger, D | 11381111 | NIMH CBDB | scz\_lie5\_eur**

1288 Subjects were recruited from the Clinical Brain Disorders Branch of the NIMH ‘Sibling Study’  
1289 as previously described. In brief, cases and controls gave informed consent and only participants  
1290 of European ancestry were included in the current analysis. Cases completed a structured clinical  
1291 interview and were diagnosed with schizophrenia-spectrum disorders. Samples were genotyped  
1292 at the NIMH.

1293 **Wendland/Schubert | Pfizer | Not Published | Multiple countries | scz\_pfla\_eur**

1294 Pfizer contributed anonymized individual genotypes for cases from seven multi-center  
1295 randomized, double-blind efficacy and safety clinical trials (A1281063, A1281134, A1281148,  
1296 A245-102, NRA7500001, NRA7500002, NRA7500003, and NRA7500004) as well as a set of

1297 purchased samples (NRA9000099). Trial samples were collected for antipsychotic medications  
1298 across outpatient and inpatient treatment settings. All participating cases had a diagnosis of  
1299 schizophrenia and were assessed using a structural clinical interview by trained interviewers,  
1300 with systematic procedures to quality-control diagnostic accuracy and reliability trials across  
1301 participating sites in the United States and internationally. Purchased blood samples were  
1302 obtained from PrecisionMed International by Pharmacia and Upjohn Corporation, and were  
1303 collected from diagnosed subjects with schizophrenia and schizoaffective disorder. All studies  
1304 were reviewed by both central and local institutional review boards, depending on the study site,  
1305 before recruitment of subjects started. Protocol amendments were approved while the study was  
1306 in progress and before the data were unblinded. The studies were conducted in conformity with  
1307 the U.S. Food and Drug Administration Code of Federal Regulations (21CFR, Part 50) and the  
1308 Declaration of Helsinki and its amendments, and were consistent with Good Clinical Practice  
1309 and the applicable regulatory requirements. Participants provided written informed consent  
1310 before enrollment. An optional blood sample was collected from clinical trial subjects for  
1311 pharmacogenetic analysis to investigate potential associations between genetic variant drug  
1312 response and general characteristics of schizophrenia and related disorders. Sample collection  
1313 was not required for participation in the original clinical trials. The controls (A9011027) were  
1314 recruited in a multi-site, cross-sectional, non-treatment prospective trial to collect data, including  
1315 DNA, from cognitive normal and free of psychiatric diseases elderly subjects in the US. Subjects  
1316 were specifically recruited to match the gender, age, and ethnicity information from the LEADe  
1317 and UCSD MCI studies. The study described here is within the scope of patient consent.

1318 **Werge, T | 19571808 | Denmark | scz\_denm\_eur**

1319 Cases were ascertained through psychiatric departments and twin pair studies, and were of  
1320 Danish parentage for at least the prior three generations. The controls were collected at the  
1321 University of Aarhus, and included 500 medical students, all of Danish parentage for at least  
1322 three generations. All subjects gave written informed consent and the Danish Data Protection  
1323 Agency and the ethics committees of Denmark approved the human subjects protocol.

1324

1325 *Bipolar Disorder*

1326 **Adolfsson, R | Not published | Umeå, Sweden | bip\_ume4\_eur**

1327 Clinical characterization of the patients included the Mini-International Neuropsychiatric  
1328 Interview (MINI), the Diagnostic Interview for Genetic Studies (DIGS), the Family Interview for  
1329 Genetic Studies (FIGS) and the Schedules for Clinical Assessment in Neuropsychiatry (SCAN).  
1330 The final diagnoses were made according to the DSM-IV-TR and determined by consensus of 2  
1331 research psychiatrists. The unrelated Swedish control individuals, consisting of a large  
1332 population-based sample representative of the general population of the region, were randomly  
1333 selected from the ‘Betula study’.

1334 **Alda, M; Smoller, J | Not published | Nova Scotia, Canada; I2B2 controls | bip\_hal2\_eur**

1335 The case samples were recruited from patients longitudinally followed at specialty mood  
1336 disorders clinics in Halifax and Ottawa (Canada). Cases were interviewed in a blind fashion with  
1337 the Schedule of Affective Disorders and Schizophrenia-Lifetime version (SADS-L) and  
1338 consensus diagnoses were made according to DSM-IV and Research Diagnostic Criteria (RDC).  
1339 Protocols and procedures were approved by the local Ethics Committees and written informed  
1340 consent was obtained from all patients before participation in the study. Control subjects were  
1341 drawn from the I2B2 (Informatics for Integrating Biology and the Bedside) project. The study

1342 consists of de-identified healthy individuals recruited from a healthcare system in the Boston,  
1343 MA, US area. The de-identification process meant that the Massachusetts General Hospital  
1344 Institutional Review Board elected to waive the requirement of seeking informed consent as  
1345 detailed by US Code of Federal Regulations, Title 45, Part 46, Section 116 (46.116).

1346 **Andreassen, OA | PMID:21926972 [PGC1], PMID:20451256 | Norway (TOP) |**  
1347 **bip\_top7\_eur**

1348 In the TOP study (Tematisk område psykoser), cases of European ancestry, born in Norway,  
1349 were recruited from psychiatric hospitals in the Oslo region. Patients were diagnosed according  
1350 to the SCID and further ascertainment details have been reported. Healthy control subjects were  
1351 randomly selected from statistical records of persons from the same catchment area as the patient  
1352 groups. The control subjects were screened by interview and with the Primary Care Evaluation  
1353 of Mental Disorders (PRIME-MD). None of the control subjects had a history of  
1354 moderate/severe head injury, neurological disorder, mental retardation or an age outside the age  
1355 range of 18-60 years. Healthy subjects were excluded if they or any of their close relatives had a  
1356 lifetime history of a severe psychiatric disorder. All participants provided written informed  
1357 consent and the human subjects protocol was approved by the Norwegian Scientific-Ethical  
1358 Committee and the Norwegian Data Protection Agency.

1359 **Andreassen, OA | Not published | Norway (TOP) | bip\_top8\_eur**

1360 The TOP8 bipolar disorder cases and controls were ascertained in the same way as the  
1361 bip\_top7\_eur (TOP7) samples described above, and recruited from hospitals across Norway.

1362 **Biernacka, JM; Frye, MA | 27769005 | Mayo Clinic, USA | bip\_may1\_eur**

1363 Bipolar cases were drawn from the Mayo Clinic Bipolar Biobank. Enrolment sites included  
1364 Mayo Clinic, Rochester, Minnesota; Lindner Center of HOPE/University of Cincinnati College



1365 of Medicine, Cincinnati, Ohio; and the University of Minnesota, Minneapolis, Minnesota.  
1366 Enrolment at each site was approved by the local Institutional Review Board approval, and all  
1367 participants consented to use of their data for future genetic studies. Participants were identified  
1368 through routine clinical appointments, from in-patients admitted in mood disorder units, and  
1369 recruitment advertising. Participants were required to be between 18 and 80 years old and be able  
1370 to speak English, provide informed consent, and have DSM-IV-TR diagnostic confirmation of  
1371 type 1 or 2 bipolar disorder or schizoaffective bipolar disorder as determined using the SCID.  
1372 Controls were selected from the Mayo Clinic Biobank. Potential controls with ICD9 codes for  
1373 bipolar disorder, schizophrenia or related diagnoses in their electronic medical record were  
1374 excluded.

1375 **Blackwood, D | 18711365 [PGC1] | Edinburgh, UK | bip\_edi1\_eur**

1376 This sample comprised Caucasian individuals contacted through the inpatient and outpatient  
1377 services of hospitals in South East Scotland. A BD-I diagnosis was based on an interview with  
1378 the patient using the SADS-L supplemented by case note review and frequently by information  
1379 from medical staff, relatives and caregivers. Final diagnoses, based on DSM-IV criteria were  
1380 reached by consensus between two trained psychiatrists. Ethnically-matched controls from the  
1381 same region were recruited through the South of Scotland Blood Transfusion Service. Controls  
1382 were not directly screened to exclude those with a personal or family history of psychiatric  
1383 illness. The study was approved by the Multi-Centre Research Ethics Committee for Scotland  
1384 and patients gave written informed consent for the collection of DNA samples for use in genetic  
1385 studies.

1386 **Breen, G; Vincent, JB | 24387768; 19416921; 21926972 [PGC1] | London, UK; Toronto,**  
1387 **Canada [BACC] | bip\_bac1\_eur**

1388 The total case/control cohort (N=1922) includes 871 subjects from Toronto, Canada (N=431  
1389 cases (160 male; 271 female); N=440 controls (176 male; 264 female)), 1051 subjects from  
1390 London, UK (N=538 cases (180 male; 358 female); N=513 controls (192 male; 321 female)). A  
1391 summary of mean and median age at interview, age of onset (AOO), diagnostic subtypes (BD 1  
1392 versus BD 2), presence of psychotic symptoms, suicide attempt and family history of psychiatric  
1393 disorders has been provided previously for both the Toronto and London cohorts. From the  
1394 Toronto site (Centre for Addiction & Mental Health (CAMH)), BD individuals and unrelated  
1395 healthy controls matched for age, gender and ethnicity were recruited. Inclusion criteria for  
1396 patients: a) diagnosed with DSMIV/ICD 10 BD 1 or 2; b) 18 years old or over; c) Caucasian, of  
1397 Northern and Western European origin, and three out of four grandparents also N.W. European  
1398 Caucasian. Exclusion criteria include: a) Use of intravenous drugs; b) Evidence of intellectual  
1399 disability; c) Related to an individual already in the study; d) Manias that only ever occurred in  
1400 relation to or resulting from alcohol or substance abuse/dependence, or medical illness; e)  
1401 Manias resulting from non-psychotropic substance usage. The SCAN interview (Schedule for  
1402 Clinical Assessments in Neuropsychiatry) was used for subject assessment. Using the SCAN  
1403 interview along with case note review, each case was assigned DSM-IV and ICD 10 diagnoses  
1404 by two independent diagnosticians, according to lifetime consensus best-estimate diagnosis.  
1405 Lifetime occurrence of psychiatric symptoms was also recorded using the OPCRIT checklist,  
1406 modified for use with mood disorders. Similar methods and criteria were also used to collect a  
1407 sample of 538 BD cases and 513 controls for the London cohort (King's College London; KCL).  
1408 Both studies were approved by respective institutional research ethics committees (the CAMH  
1409 Research Ethics Board (REB) in Toronto, and the College Research Ethics Committee (CREC)  
1410 at KCL), and informed written consent was obtained from all participants. GWAS results have

1411 previously been published for the entire KCL/CAMH cohort.

1412 **Corvin, A | 18711365 [PGC1] | Ireland | bip\_dub1\_eur**

1413 Samples were collected as part of a larger study of the genetics of psychotic disorders in the  
1414 Republic of Ireland, under protocols approved by the relevant IRBs and with written informed  
1415 consent that permitted repository use. Cases were recruited from Hospitals and Community  
1416 psychiatric facilities in Ireland by a psychiatrist or psychiatric nurse trained to use the SCID.  
1417 Diagnosis was based on the structured interview supplemented by case note review and collateral  
1418 history where available. All diagnoses were reviewed by an independent reviewer. Controls were  
1419 ascertained with informed consent from the Irish GeneBank and represented blood donors who  
1420 met the same ethnicity criteria as cases. Controls were not specifically screened for psychiatric  
1421 illness.

1422 **Rietschel, M; Nöthen, MM, Cichon, S | 21926972 [PGC1] | BOMA-Germany I |**  
1423 **bip\_bonn\_eur**

1424 Cases for the BOMA-Bipolar Study were ascertained from consecutive admissions to the  
1425 inpatient units of the Department of Psychiatry and Psychotherapy at the University of Bonn and  
1426 at the Central Institute for Mental Health in Mannheim, University of Heidelberg, Germany.  
1427 DSM-IV lifetime diagnoses of bipolar I disorder were assigned using a consensus best-estimate  
1428 procedure, based on all available information, including a structured interview with the SCID  
1429 and SADS-L, medical records, and the family history method. In addition, the OPCRIT checklist  
1430 was used for the detailed polydiagnostic documentation of symptoms. Controls were ascertained  
1431 from three population-based studies in Germany (PopGen, KORA, and Heinz-Nixdorf-Recall  
1432 Study). The control subjects were not screened for mental illness. Study protocols were reviewed  
1433 and approved in advance by Institutional Review Boards of the participating institutions. All

1434 subjects provided written informed consent.

1435 **Rietschel, M; Nöthen, MM; Schulze, TG; Reif, A; Forstner, AJ | 24618891 | BOMA-**

1436 **Germany II | bip\_bmg2\_eur**

1437 Cases were recruited from consecutive admissions to psychiatric in-patient units at the

1438 University Hospital Würzburg. All cases received a lifetime diagnosis of BD according to the

1439 DSM-IV criteria using a consensus best-estimate procedure based on all available information,

1440 including semi-structured diagnostic interviews using the Association for Methodology and

1441 Documentation in Psychiatry, medical records and the family history method. In addition, the

1442 OPCRIT system was used for the detailed polydiagnostic documentation of symptoms.

1443 Control subjects were ascertained from the population-based Heinz Nixdorf Recall (HNR) Study.

1444 The controls were not screened for a history of mental illness. Study protocols were reviewed

1445 and approved in advance by Institutional Review Boards of the participating institutions. All

1446 subjects provided written informed consent.

1447 **Rietschel, M; Nöthen, MM; Schulze, TG; Bauer, M; Forstner, AJ; Müller-Myhsok, B |**

1448 **24618891 | BOMA-Germany III | bip\_bmg3\_eur**

1449 Cases were recruited at the Central Institute of Mental Health in Mannheim, University of

1450 Heidelberg, and other collaborating psychiatric hospitals in Germany. All cases received a

1451 lifetime diagnosis of BD according to the DSM-IV criteria using a consensus best-estimate

1452 procedure based on all available information including structured diagnostic interviews using the

1453 AMDP, Composite International Diagnostic Screener (CID-S), SADS-L and/or SCID, medical

1454 records, and the family history method. In addition, the OPCRIT system was used for the

1455 detailed polydiagnostic documentation of symptoms. Controls were selected randomly from a

1456 Munich-based community sample and recruited at the Max-Planck Institute of Psychiatry. They

1457 were screened for the presence of anxiety and mood disorders using the CID-S. Only individuals  
1458 without mood and anxiety disorders were collected as controls. Study protocols were reviewed  
1459 and approved in advance by Institutional Review Boards of the participating institutions. All  
1460 subjects provided written informed consent.

1461 **Hauser, J; Lissowska, J; Forstner, AJ | 24618891 | BOMA-Poland | bip\_bmpo\_eur**

1462 Cases were recruited at the Department of Psychiatry, Poznan University of Medical Sciences,  
1463 Poznan, Poland. All cases received a lifetime diagnosis of BD according to the DSM-IV criteria  
1464 on the basis of a consensus best-estimate procedure and structured diagnostic interviews using  
1465 the SCID. Controls were drawn from a population-based case-control sample recruited by the  
1466 Cancer-Center and Institute of Oncology, Warsaw, Poland and a hospital-based case-control  
1467 sample recruited by the Nofer Institute of Occupational Medicine, Lodz, Poland. The Polish  
1468 controls were produced by the International Agency for Research on Cancer (IARC) and the  
1469 Centre National de Génotypage (CNG) GWAS Initiative for a study of upper aerodigestive tract  
1470 cancers. The controls were not screened for a history of mental illness. Study protocols were  
1471 reviewed and approved in advance by Institutional Review Boards of the participating  
1472 institutions. All subjects provided written informed consent.

1473 **Rietschel, M; Nöthen, MM; Rivas, F; Mayoral, F; Kogevinas, M; others | 24618891 |**  
1474 **BOMA-Spain | bip\_bmsp\_eur**

1475 Cases were recruited at the mental health departments of the following five centers in Andalusia,  
1476 Spain: University Hospital Reina Sofia of Córdoba, Provincial Hospital of Jaen; Hospital of  
1477 Jerez de la Frontera (Cádiz); Hospital of Puerto Real (Cádiz); Hospital Punta Europa of  
1478 Algeciras (Cádiz); and Hospital Universitario San Cecilio (Granada). Diagnostic assessment was  
1479 performed using the SADS-L; the OPCRIT; a review of medical records; and interviews with

1480 first and/or second degree family members using the Family Informant Schedule and Criteria  
1481 (FISC). Consensus best estimate BD diagnoses were assigned by two or more independent senior  
1482 psychiatrists and/or psychologists, and according to the RDC, and the DSM-IV. Controls were  
1483 Spanish subjects drawn from a cohort of individuals recruited in the framework of the European  
1484 Community Respiratory Health Survey (ECRHS, <http://www.ecrhs.org/>). The controls were not  
1485 screened for a history of mental illness. Study protocols were reviewed and approved in advance  
1486 by Institutional Review Boards of the participating institutions. All subjects provided written  
1487 informed consent.

1488 **Fullerton, J.M.; Mitchell, P.B.; Schofield, P.R.; Martin N.G.; Cichon, S. | 24618891 |**  
1489 **BOMA-Australia | bip\_bmau\_eur**

1490 Cases were recruited at the Mood Disorder Unit, Prince of Wales Hospital in Sydney. All cases  
1491 received a lifetime diagnosis of BD according to the DSM-IV criteria on the basis of a consensus  
1492 best-estimate procedure and structured diagnostic interviews using the DIGS, FIGS, and the  
1493 SCID. Controls were parents of unselected adolescent twins from the Brisbane Longitudinal  
1494 Twin Study. The controls were not screened for a history of mental illness. Study protocols were  
1495 reviewed and approved in advance by Institutional Review Boards of the participating  
1496 institutions. All subjects provided written informed consent.

1497 **Grigoriu-Serbanescu, M; Nöthen, MM | 21353194 | BOMA-Romania | bip\_rom3\_eur**

1498 Cases were recruited from consecutive admissions to the Obregia Clinical Psychiatric Hospital,  
1499 Bucharest. Patients were administered the DIGS and FIGS interviews. Information was also  
1500 obtained from medical records and close relatives. The diagnosis of BP-I was assigned according  
1501 to DSM-IV criteria using the best estimate procedure. All patients had at least two hospitalized  
1502 illness episodes. Population-based controls were evaluated using the DIGS to exclude a lifetime

1503 history of major affective disorders, schizophrenia, schizoaffective disorders, and other  
1504 psychoses, obsessive-compulsive disorder, eating disorders, and alcohol or drug addiction.

1505 **Craddock, N, Jones, I, Jones, L | 17554300 | WTCCC | bip\_wtcc\_eur\_sr-qc**

1506 Cases were all over the age of 17 yr, living in the UK and of European descent. Recruitment was  
1507 undertaken throughout the UK and included individuals who had been in contact with mental  
1508 health services and had a lifetime history of high mood. After providing written informed  
1509 consent, participants were interviewed by a trained psychologist or psychiatrist using a semi-  
1510 structured lifetime diagnostic psychiatric interview (Schedules for Clinical Assessment in  
1511 Neuropsychiatry) and available psychiatric medical records were reviewed. Using all available  
1512 data, best-estimate life-time diagnoses were made according to the RDC. In the current study we  
1513 included cases with a lifetime diagnosis of RDC bipolar 1 disorder, bipolar 2 disorder or schizo-  
1514 affective disorder, bipolar type. Controls were recruited from two sources: the 1958 Birth Cohort  
1515 study and the UK Blood Service (blood donors) and were not screened for history of mental  
1516 illness. All cases and controls were recruited under protocols approved by the appropriate IRBs.  
1517 All subjects gave written informed consent.

1518 **Kelsoe, J | 21926972 [PGC1] | USA (GAIN) | bip\_gain\_eur**

1519 *Genetic Association Information Network (GAIN)/ The Bipolar Genome Study (BiGS)* The BD  
1520 sample was collected under the auspices of the NIMH Genetics Initiative for BD  
1521 (<http://zork.wustl.edu/nimh/>), genotyped as part of GAIN and analyzed as part of a larger GWAS  
1522 conducted by the BiGS consortium. Approximately half of the GAIN sample was collected as  
1523 multiplex families or sib pair families (waves 1-4), the remainder were collected as individual  
1524 cases (wave 5). Subjects were ascertained at 11 sites: Indiana University, John Hopkins  
1525 University, the NIMH Intramural Research Program, Washington University at St. Louis,

1526 University of Pennsylvania, University of Chicago, Rush Medical School, University of Iowa,  
1527 University of California, San Diego, University of California, San Francisco, and University of  
1528 Michigan. All investigations were carried out after the review of protocols by the IRB at each  
1529 participating institution. At all sites, potential cases were identified from screening admissions to  
1530 local treatment facilities and through publicity programs or advocacy groups. Potential cases  
1531 were evaluated using the DIGS, FIGS, and information from relatives and medical records. All  
1532 information was reviewed through a best estimate diagnostic procedure by two independent and  
1533 non-interviewing clinicians and a consensus best-estimate diagnosis was reached. In the event of  
1534 a disagreement, a third review was done to break the tie. Controls were from the NIMH Genetic  
1535 Repository sample obtained by Dr. P. Gejman through a contract to Knowledge Networks, Inc.  
1536 Only individuals with complete or near-complete psychiatric questionnaire data who did not  
1537 fulfill diagnostic criteria for major depression and denied a history of psychosis or BD were  
1538 included as controls for BiGS analyses. Controls were matched for gender and ethnicity to the  
1539 cases.

1540 **Kelsoe, J; Sklar, P; Smoller, J | [PGC1 Replication] | USA (FAT2; FaST, BiGS, TGEN) |**  
1541 **bip\_fat2\_eur**

1542 Cases were collected from individuals at the 11 U.S. sites described for the GAIN sample.  
1543 Eligible participants were age 18 or older meeting DSM-IV criteria for BD-I or BD-II by  
1544 consensus diagnosis based on interviews with the Affective Disorders Evaluation (ADE) and  
1545 MINI. All participants provided written informed consent and the study protocol was approved  
1546 by IRBs at each site. Collection of phenotypic data and DNA samples were supported by NIMH  
1547 grants MH063445 (JW Smoller); MH067288 (PI: P Sklar), and MH63420 (PI: V Nimgaonkar).  
1548 The control samples were NIMH controls that were using the methods described in that section.



1549 The case and control samples were independent of those included in the GAIN sample.

1550 **Kirov, G | 25055870 | Bulgarian trios | bip\_butr\_eur**

1551 All cases were recruited in Bulgaria from psychiatric inpatient and outpatient services. Each  
1552 proband had a history of hospitalisation and was interviewed with an abbreviated version of the  
1553 SCAN. Consensus best-estimate diagnoses were made according to DSM-IV criteria by two  
1554 researchers. All participants gave written informed consent and the study was approved by local  
1555 ethics committees at the participating centers.

1556 **Kirov, G | 25055870 | UK trios | bip\_uktr\_eur**

1557 The BD subjects were recruited from lithium clinics and interviewed in person by a senior  
1558 psychiatrist, using abbreviated version of the SCAN. Consensus best-estimate diagnoses were  
1559 made based on the interview and hospital notes. Ethics committee approval for the study was  
1560 obtained from the relevant research ethics committees and all individuals provided written  
1561 informed consent for participation.

1562 **Landén, M; Sullivan, PF; Sklar, P | [ICCBD] | Sweden (ICCBD) | bip\_swa2\_eur**

1563 The BD subjects were identified using the Swedish National Quality Register for Bipolar  
1564 Disorders (Bipolär) and the Swedish National Patient Register (using a validated algorithm  
1565 requiring at least two hospitalizations with a BD diagnosis). A confirmatory telephone interview  
1566 with a diagnostic review was conducted. Additional subjects were recruited from the St. Göran  
1567 Bipolar Project (Affective Center at Northern Stockholm Psychiatry Clinic, Sweden), enrolling  
1568 new and ongoing patients diagnosed with BD using structured clinical interviews. Diagnoses  
1569 were made according to the DSM-IV criteria (Bipolär and St. Göran Bipolar Project) and ICD-  
1570 10 (National Patient Register). The control subjects used were the same as for the SCZ analyses  
1571 described above. All ascertainment procedures were approved by the Regional Ethical

1572 Committees in Sweden.

1573 **Landén, M; Sullivan, PF; Sklar, P | [ICCBD] | Sweden (ICCBD) | bip\_swei\_eur**

1574 The cases and controls in the bip\_swei\_eur sample were recruited using the same ascertainment  
1575 methods described for the bip\_swa2\_eur sample.

1576 **Leboyer, M | [PGC1 replication] | France | bip\_fran\_eur**

1577 Cases with BD1 or BD2 and control samples were recruited as part of a large study of genetics of  
1578 BD in France (Paris-Creteil, Bordeaux, Nancy) with a protocol approved by relevant IRBs and  
1579 with written informed consent. Cases were of French descent for more than 3 generations were  
1580 assessed by a trained psychiatrist or psychologist using structured interviews supplemented by  
1581 medical case notes, mood scales and self-rating questionnaire assessing dimensions.

1582 **Li, Q | 24166486; 27769005 | USA (Janssen), SAGE controls | bip\_jst5\_eur**

1583 The study included unrelated patients with bipolar 1 disorder from 6 clinical trials (IDs:  
1584 NCT00253162, NCT00257075, NCT00076115, NCT00299715, NCT00309699, and  
1585 NCT00309686). Participant recruitment was conducted by Janssen Research & Development,  
1586 LLC (formerly known as Johnson & Johnson Pharmaceutical Research & Development, LLC) to  
1587 assess the efficacy and safety of risperidone. Bipolar cases were diagnosed according to DSM-  
1588 IV-TR criteria. The diagnosis of bipolar disorder was confirmed by the Schedule for Affective  
1589 Disorders and Schizophrenia for School-Age Children-Present and Lifetime Version (K-SADS-  
1590 PL) in NCT00076115, by the SCID in NCT00257075 and NCT00253162, or by the MINI in  
1591 NCT00299715 and NCT00309699, and NCT00309686, respectively. Additional detailed  
1592 descriptions of these clinical trials can be found at ClinicalTrials.gov. Only patients of European  
1593 ancestry with matching controls were included in the current analysis. Controls subjects were  
1594 drawn from the Study of Addiction: Genetics and Environment (SAGE, dbGaP Study Accession:

1595 phs000092.v1.p1). Control subjects did not have alcohol dependence or drug dependence  
1596 diagnoses; however, mood disorders were not an exclusion criterion.

1597 **McQuillin, A; Gurling, H | 18317468 [PGC1] | UCL (University College London), London,**  
1598 **UK | bip\_uclo\_eur**

1599 The UCL sample comprised Caucasian individuals who were ascertained and received clinical  
1600 diagnoses of bipolar 1 disorder according to UK National Health Service (NHS) psychiatrists at  
1601 interview using the categories of the International Classification of Disease version 10. In  
1602 addition bipolar subjects were included only if both parents were of English, Irish, Welsh or  
1603 Scottish descent and if three out of four grandparents were of the same descent. All volunteers  
1604 read an information sheet approved by the Metropolitan Medical Research Ethics Committee  
1605 who also approved the project for all NHS hospitals. Written informed consent was obtained  
1606 from each volunteer. The UCL control subjects were recruited from London branches of the  
1607 National Blood Service, from local NHS family doctor clinics and from university student  
1608 volunteers. All control subjects were interviewed with the SADS-L to exclude all psychiatric  
1609 disorders.

1610 **Craddock, N; Jones, I; Jones, L | [ICCBD] | Cardiff and Worcester, UK (ICCBD-BDRN) |**  
1611 **bip\_icuk\_eur**

1612 Cases were all over the age of 17 yr, living in the UK and of European descent. Cases were  
1613 recruited via systematic and not systematic methods as part of the Bipolar Disorder Research  
1614 Network project ([www.bdrn.org](http://www.bdrn.org)), provided written informed consent and were interviewed  
1615 using a semi-structured diagnostic interview, the Schedules for Clinical Assessment in  
1616 Neuropsychiatry. Based on the information gathered from the interview and case notes review,  
1617 best-estimate lifetime diagnosis was made according to DSM-IV. Inter-rater reliability was

1618 formally assessed using 20 randomly selected cases (mean  $\kappa$  Statistic = 0.85). In the current  
1619 study we included cases with a lifetime diagnosis of DSM-IV bipolar disorder or schizo-affective  
1620 disorder, bipolar type. The BDRN study has UK National Health Service (NHS) Research Ethics  
1621 Committee approval and local Research and Development approval in all participating NHS  
1622 Trusts/Health Boards. Controls were part of the Wellcome Trust Case Control Consortium  
1623 common control set, which comprised healthy blood donors recruited from the UK Blood  
1624 Service and samples from the 1958 British Birth Cohort. Controls were not screened for a history  
1625 of mental illness. All cases and controls were recruited under protocols approved by the  
1626 appropriate IRBs. All subjects gave written informed consent.

1627 **Ophoff, RA | Not Published | Netherlands | bip\_ucla\_eur**

1628 The case sample consisted of inpatients and outpatients recruited through psychiatric hospitals  
1629 and institutions throughout the Netherlands. Cases with DSM-IV bipolar disorder, determined  
1630 after interview with the SCID, were included in the analysis. Controls were collected in parallel  
1631 at different sites in the Netherlands and were volunteers with no psychiatric history after  
1632 screening with the (MINI). Ethical approval was provided by UCLA and local ethics committees  
1633 and all participants gave written informed consent.

1634 **Paciga, S | [PGC1] | USA (Pfizer) | bip\_pf1e\_eur**

1635 This sample comprised Caucasian individuals recruited into one of three Geodon (ziprasidone)  
1636 clinical trials (NCT00141271, NCT00282464, NCT00483548). Subjects were diagnosed by a  
1637 clinician with a primary diagnosis of Bipolar 1 Disorder, most recent episode depressed, with or  
1638 without rapid cycling, without psychotic features, as defined in the DSM-IV-TR (296.5x) and  
1639 confirmed by the MINI (version 5.0.0). Subjects also were assessed as having a HAM-D-17  
1640 total score of >20 at the screening visit. The trials were conducted in accordance with the

1641 protocols, International Conference on Harmonization of Good Clinical Practice Guidelines, and  
1642 applicable local regulatory requirements and laws. Patients gave written informed consent for the  
1643 collection of blood samples for DNA for use in genetic studies.

1644 **Pato, C | [ICCBD] | Los Angeles, USA (ICCBD-GPC)| bip\_usc2\_eur**

1645 Genomic Psychiatry Consortium (GPC) cases and controls were collected via the University of  
1646 Southern California healthcare system, as previously described. Using a combination of focused,  
1647 direct interviews and data extraction from medical records, diagnoses were established using the  
1648 OPCRIT and were based on DSM-IV-TR criteria. Age and gender-matched controls were  
1649 ascertained from the University of Southern California health system and assessed using a  
1650 validated screening instrument and medical records.

1651 **Scott, L; Myer, RM; Boehnke, M | 19416921 [PGC1] | Michigan, USA (Pritzker and**  
1652 **NIMH) | bip\_mich\_eur**

1653 The Pritzker Neuropsychiatric Disorders Research Consortium (NIMH/Pritzker) case and  
1654 controls samples were from the NIMH Genetics Initiative Genetics Initiative Repository. Cases  
1655 were diagnosed according to DMS-III or DSM-IV criteria using diagnostic interviews and/or  
1656 medical record review. Cases with low confidence diagnoses were excluded. From each wave 1-  
1657 5 available non-Ashkenazi European-origin family, two BD1 siblings were included when  
1658 possible and the proband was preferentially included if available (n=946 individuals in 473  
1659 sibling pairs); otherwise a single BD1 case was included (n=184). The bipolar sibling pairs were  
1660 retained within the NIMH/Pritzker sample when individuals in more than one study were  
1661 uniquely assigned to a study set. Controls had non-Ashkenazi European-origin, were aged 20-70  
1662 years and reported no diagnosis with or treatment for BD or schizophrenia, and that they had not  
1663 heard voices that others could not hear. Individuals with suspected major depression were

1664 excluded based on answers to questions related to depressive mood. NIMH controls were further  
1665 selected as the best match(es) to NIMH cases based on self-reported ancestry.

1666 **Sklar, P; Smoller, J | 18317468 [PGC1] | USA (STEP1) | bip\_stp1\_eur**

1667 The Systematic Treatment Enhancement Program for Bipolar Disorder (STEP-BD) was a seven-  
1668 site, national U.S., longitudinal cohort study designed to examine the effectiveness of treatments  
1669 and their impact on the course of BD that enrolled 4,361 participants who met DSM-IV criteria  
1670 for BD1, BD2, bipolar not otherwise specified (NOS), schizoaffective manic or bipolar type, or  
1671 cyclothymic disorder based on diagnostic interviews. From the parent study, 2,089 individuals  
1672 who were over 18 years of age with BD1 and BD2 diagnoses consented to the collection of  
1673 blood samples for DNA. BD samples with a consensus diagnosis of BD1 were selected for  
1674 inclusion in STEP1. Two groups of controls samples from the NIMH repository were used. One  
1675 comprised DNA samples derived from US Caucasian anonymous cord blood donors. The  
1676 second were controls who completed the online self-administered psychiatric screen and were  
1677 ascertained as described above, by Knowledge Networks Inc. For the second sample of controls  
1678 only those without history of schizophrenia, psychosis, BD or major depression with functional  
1679 impairment were used.

1680 **Sklar, P; Smoller, J | 18711365 [PGC1] | USA (STEP2) | bip\_stp2\_eur**

1681 The STEP2 sample included BD-1 and BD-2 samples from the STEP-BD study described above  
1682 along with BD-2 subjects from UCL study also described above. The controls samples for this  
1683 study were from the NIMH repository as described above for the STEP1 study.

1684

1685 **European ancestry, trio design**

1686 *Schizophrenia*

1687 **Kirov, G: Owen M | 22083728| Bulgaria | ms.scz\_butr\_eur**

1688 Families from Bulgaria were recruited if a proband had schizophrenia or schizoaffective  
1689 disorder, both parents were available, and all members of the trio agreed to participate in the  
1690 study. Recruitment took place between 1999 and 2004 in several psychiatric hospitals in  
1691 Bulgaria. Ethical Committee approval was obtained from each of these hospitals. All probands  
1692 and all parents received an Information Sheet and signed Informed Consent Forms. All  
1693 participants had attended mainstream schools, which at the time in Bulgaria, excluded people  
1694 with mental retardation. Probands were either in- or out-patients at the time of the study but each  
1695 had a history of hospitalization. A team of psychiatrists was trained in using the rating scales and  
1696 methods of the study. We used the SCAN instrument to perform an interview for psychotic and  
1697 mood symptoms. This instrument has been translated into Bulgarian and validated by one of its  
1698 authors (A. Jablensky). Consensus diagnoses were made according to DSM-IV criteria on the  
1699 basis of an interview and inspection of hospital notes by two clinicians. If consensus was not  
1700 attained, the patient was re-interviewed by a research interview trained clinician and was  
1701 excluded if consensus could still not be reached. In addition, approximately 23% of the sample  
1702 was selected at random and re-interviewed by a research interview trained clinician. Hospital  
1703 notes were also collected for affected relatives in order to confirm diagnoses.

1704 **Levinson, D | 22885689 | Six countries | ms.scz\_lemu\_eur**

1705 Schizophrenia cases were included from the family sample of European-ancestry pedigrees  
1706 described by Levinson et al. Participants and their families in this trio study, probands were  
1707 ascertained and recruited from different clinical settings (e.g. inpatients, outpatients and  
1708 community facilities) in six countries (Australia, France, Germany, Ireland, UK, and the US).  
1709 (Unrelated individuals were included as part of a case-control design, see Levinson, D,

1710 scz\_lacw\_eur above.) Diagnoses were established using semi-structured interviews, psychiatric  
1711 records and informant reports. Case probands were diagnosed with schizophrenia or  
1712 schizoaffective disorder according to DSM-III-R criteria. The trio-based analysis included  
1713 families where there was at least one affected proband and two available parents. Each affected  
1714 sibling in such families was included, with the parents, as an independent trio. All protocols were  
1715 approved by local IRBs, and all cases provided written informed consent.

1716 **Kirov, G: Owen, M | Not Published | Bulgaria | ms.scz\_uktr\_eur**

1717 All cases and parents were recruited from UK and had a history of hospitalization for treatment  
1718 of schizophrenia. Diagnosis was confirmed following a SCAN interview and review of case  
1719 notes followed by consensus diagnosis according to DSM-IV criteria. The samples were  
1720 genotyped at the Broad Institute. All participants gave written informed consent and the study  
1721 was approved by local ethics committees at the participating centers. The samples were  
1722 genotyped at the Broad Institute.

1723

#### 1724 **Genotype Quality Control**

1725 To ensure independence of the data sets, individuals were excluded until no individual showed a  
1726 relatedness ( $r_{\text{hat}}$ ) value greater than 0.2 to any other individual in the collection, while  
1727 preferentially keeping the case over the control for case-control related pairs. In total 1,795 BD  
1728 cases, 1,165 SCZ cases and 27,274 controls were removed (most of which were previously  
1729 known), leaving 20,129 BD cases 33,426 SCZ cases and 54,065 controls for the final meta-  
1730 analysis.

1731 For analyses directly comparing BD and SCZ, we matched cases from both phenotypes on  
1732 genotyping platform and ancestry, resulting in 15,270 BD cases versus 23,585 SCZ cases.



1733 Hence, we were able to match 76% of BD cases and 71% of SCZ cases for this case vs case  
1734 analysis.

1735 Among our entire dataset, 44% of the sample was female, 51% was male and 5% were  
1736 unreported by the collection site. This work focused explicitly on the autosomes and sought  
1737 maximal power across the analyses, sex was not used except for during quality control and sex-  
1738 specific analyses were not performed in this effort. Individual ages were not provided. For a  
1739 subset of cases, we had information for age of onset which were used in subphenotype specific  
1740 analyses only.

1741

#### 1742 **Sub-phenotype Description**

1743 BD sub-phenotypes were collected by each study site using a combination of diagnostic  
1744 instruments, case records and participant interviews. Ascertainment details for each study site are  
1745 described in the supplementary data of the PGC Bipolar Working Group paper(Stahl et al.,  
1746 2017). The selection of phenotypes for collection by this group was determined by literature  
1747 searches in order to determine phenotypes with prior evidence for heritability. It was further  
1748 refined dependent on the availability of phenotype data across a range of study sites and the  
1749 consistency by which the phenotypes were defined. Schizophrenia subphenotypes represent  
1750 quantitative traits extracted using factor analysis from a set of standard psychiatric assessments  
1751 and represent four symptom dimensions (manic, depressive, positive and negative). These  
1752 subphenotypes were used previously(Ruderfer et al., 2014) but in this work we have increased  
1753 the sample size with additional cohorts being added.

1754

#### 1755 **METHOD DETAILS**

1756

## 1757 **QUANTIFICATION AND STATISTICAL ANALYSIS**

1758

### 1759 **Quality Control, Imputation, Association Analysis and Polygenic Risk Score Testing**

1760 Quality control and imputation were performed on each of the study cohort datasets (n=81),  
1761 according to standards established by the Psychiatric Genomics Consortium (PGC). The quality  
1762 control parameters for retaining SNPs and subjects were: SNP missingness < 0.05 (before  
1763 sample removal); subject missingness ( $p < 0.02$ ); autosomal heterozygosity deviation ( $| F_{het} | <$   
1764  $0.2$ ); SNP missingness < 0.02 (after sample removal); difference in SNP missingness between  
1765 cases and controls < 0.02; and SNP Hardy-Weinberg equilibrium ( $p > 10^{-6}$  in controls or  $p >$   
1766  $10^{-10}$  in cases). Genotype imputation was performed using the pre-phasing/imputation stepwise  
1767 approach implemented in IMPUTE2(Howie et al., 2011) / SHAPEIT(Delaneau et al., 2013)  
1768 (chunk size of 3 Mb and default parameters). The imputation reference set consisted of 2,186  
1769 phased haplotypes from the full 1000 Genomes Project dataset (August 2012, 30,069,288  
1770 variants, release “v3.macGT1”), all variants align to human genome build 19 (hg19). After  
1771 imputation, we used the best guess genotypes (genotype probability > 0.8), for further robust  
1772 relatedness testing and population structure analysis. Here we required very high imputation  
1773 quality (INFO > 0.8) and low missingness (<1%) for further quality control. After linkage  
1774 disequilibrium (LD) pruning ( $r^2 < 0.02$ ) and frequency filtering (MAF > 0.05), there were 14,473  
1775 autosomal SNPs in the data set. Principal component estimation was done with the same  
1776 collection of autosomal SNPs. We tested the first 20 principal components for phenotype  
1777 association (using logistic regression with study indicator variables included as covariates) and  
1778 evaluated their impact on the genome-wide test statistics using  $\lambda$ . Thirteen principal components

1779 namely 1,2,3,4,5,6,7,8,10,12,15,18,20 were included in all association analyses  
1780 ( $\lambda=1.45$ ). Analytical steps were repeated for SCZ vs BD analysis.  
1781 We performed four main association analyses (Figure 1), i.e. (i) GWAS of BD and SCZ as a  
1782 single combined case phenotype, as well as disorder-specific GWAS using independent control  
1783 sets in (ii) BD cases vs BD controls and (iii) SCZ cases vs SCZ controls, and (iv) association  
1784 analysis of SCZ cases vs BD cases. For all GWS loci from the GWAS of BD and SCZ vs  
1785 controls we identified any GWS loci within 1Mb from the extent of the locus in the previously  
1786 published PGC SCZ vs controls(Schizophrenia Working Group of the Psychiatric Genomics  
1787 Consortium, 2014) and the most recent PGC GWAS of BD vs controls(Stahl et al., 2017) and  
1788 performed conditional analysis. Specifically, we transformed the genotype probabilities of the  
1789 disease variant into dosages and used it as an additional covariate for the association analysis for  
1790 the BD+SCZ vs controls index SNP. This was done within each cohort and an OR based inverse  
1791 SE weighted meta-analysis was performed for the final result. All datasets were included except  
1792 for those with trios.

1793

#### 1794 **Summary-data-based Mendelian Randomization (SMR)**

1795 SMR(Zhu et al., 2016) is a method that integrates summary level GWAS data with gene  
1796 expression quantitative trait loci (eQTL) identified in independent data sets. This integration  
1797 aims to identify variants that have pleiotropic effects on expression of a given gene and the  
1798 phenotype. While significant findings may indeed reflect a causal path from variant to phenotype  
1799 through expression, it is impossible to discern statistically between pleiotropy and causality.  
1800 However, the method can remove linkage as driving the result, and uses the data available to  
1801 prioritise amongst genes in genomic regions that show association with disease. We used SMR

1802 as a statistical fine-mapping tool applied to the SCZ vs BD GWAS results to identify loci with  
1803 strong evidence of causality via gene expression. SMR analysis is limited to significant (FDR <  
1804 0.05) cis SNP-expression quantitative trait loci (eQTLs) with MAF > 0.01. eQTLs passing these  
1805 thresholds were combined with GWAS results in the SMR test, with significance ( $p_{SMR}$ ) reported  
1806 at a Bonferroni-corrected threshold for each eQTL data set. The eQTL architecture may differ  
1807 between genes. For example, through LD, many SNPs can generate significant associations with  
1808 the same gene, but in some instances multiple SNPs may be independently associated with the  
1809 expression of a gene. After identification of significant SNP-expression-trait association through  
1810 the SMR test, a follow-up heterogeneity test aims to prioritize variants by excluding regions for  
1811 which there is conservative evidence for multiple causal loci ( $p_{HET} < 0.05$ ). SMR analyses were  
1812 conducted using eQTL data from whole peripheral blood(Westra et al., 2013), dorsolateral  
1813 prefrontal cortex generated by the CommonMind Consortium<sup>8</sup>, and 11 brain sub-regions from  
1814 the GTEx consortium(Consortium, 2015).

1815

### 1816 **Regional joint GWAS**

1817 Summary statistic Z-scores were calculated for each marker in each of the four main GWAS  
1818 results, using the logistic regression coefficient and its standard error. Rare SNPs (MAF < 0.01),  
1819 and SNPs with a low INFO score (< 0.3) in either dataset were removed. The causal variant  
1820 relationships between SCZ and BD were investigated using the Bayesian method software pw-  
1821 gwas (v0.2.1), with quasi-independent regions determined by estimate LD blocks in an analysis  
1822 of European individuals (n=1,703)(Berisa and Pickrell, 2015; Pickrell et al., 2016). Briefly, pw-  
1823 gwas takes a Bayesian approach to determine the probability of five independent models of  
1824 association. (1) There is no causal variant in BD or SCZ; (2) a causal variant in BD, but not SCZ

1825 (3); a causal variant in SCZ, but not BD; (4) a shared causal variant influencing both BD and  
1826 SCZ; (5) two causal variants where one influences BD, and one influences SCZ (Figure 2). The  
1827 posterior probability of each model is calculated using model priors, estimated empirically  
1828 within pw-gwas. Regions were considered to support a particular model when the posterior  
1829 probability of the model was greater than 0.5.

1830

### 1831 **Regional SNP-heritability estimation**

1832 We calculated local SNP-heritability independently for SCZ and BD using the Heritability  
1833 Estimator from Summary Statistics (HESS) software(Shi et al., 2016) for each of the  
1834 independent regions defined above. The sum of these regional estimates is the total SNP-  
1835 heritability of the trait. To calculate local SNP-heritability HESS requires reference LD matrices  
1836 representative of the population from which the GWAS samples were drawn. We utilized the  
1837 1000 genomes European individuals as the reference panel(The 1000 Genomes Project  
1838 Consortium, 2015). Unlike pw-gwas(Pickrell et al., 2016), HESS does not assume that only one  
1839 causal variant can be present in each region.

1840

### 1841 **DATA AND SOFTWARE AVAILABILITY**

1842 Summary statistics from GWAS are publically available at  
1843 <https://www.med.unc.edu/pgc/results-and-downloads/downloads>.

1844

1845

1846

1847