

## IMAGING THE ENVIRONMENT OF A $Z = 6.3$ SUBMILLIMETER GALAXY WITH SCUBA-2

E. I. ROBSON<sup>1</sup>, R. J. IVISON<sup>2,3</sup>, IAN SMAIL<sup>4</sup>, W. S. HOLLAND<sup>1,3</sup>, J. E. GEACH<sup>5</sup>, A. G. GIBB<sup>6</sup>, D. RIECHERS<sup>7</sup>, P. A. R. ADE<sup>8</sup>,  
D. BINTLEY<sup>9</sup>, J. BOCK<sup>10</sup>, E. L. CHAPIN<sup>11</sup>, S. C. CHAPMAN<sup>12</sup>, D. L. CLEMENTS<sup>13</sup>, A. CONLEY<sup>14</sup>, A. COORAY<sup>15</sup>, J. S. DUNLOP<sup>3</sup>,  
D. FARRAH<sup>16</sup>, M. FICH<sup>17</sup>, HAI FU<sup>15</sup>, T. JENNESS<sup>7</sup>, N. LAPORTE<sup>18</sup>, S. J. OLIVER<sup>19</sup>, A. OMONT<sup>20,21</sup>, I. PÉREZ-FOURNON<sup>18,22</sup>,  
DOUGLAS SCOTT<sup>6</sup>, A. M. SWINBANK<sup>4</sup>, AND J. WARDLOW<sup>23</sup>

*To be submitted to The Astrophysical Journal*

### ABSTRACT

We describe a search for submillimeter emission in the vicinity of one of the most distant, luminous galaxies known, HerMES FLS3 at  $z = 6.34$ , exploiting it as a signpost to a potentially biased region of the early Universe, as might be expected in hierarchical structure formation models. Imaging to the confusion limit with the innovative, wide-field submillimeter bolometer camera, SCUBA-2, we are sensitive to colder and/or less luminous galaxies in the surroundings of HFLS3. We use the Millennium Simulation to illustrate that HFLS3 may be expected to have companions if it is as massive as claimed, but find no significant evidence from the surface density of SCUBA-2 galaxies in its vicinity, or their colors, that HFLS3 marks an over-density of dusty, star-forming galaxies. We cannot rule out the presence of dusty neighbours with confidence, but deeper 450- $\mu\text{m}$  imaging has the potential to more tightly constrain the redshifts of nearby galaxies, at least one of which likely lies at  $z \gtrsim 5$ . If associations with HFLS3 can be ruled out, this could be taken as evidence that HFLS3 is less biased than a simple extrapolation of the Millennium Simulation may imply. This could suggest either that it represents a rare short-lived, but highly luminous, phase in the evolution of an otherwise typical galaxy, or that this system has suffered amplification due to a foreground gravitational lens and so is not as intrinsically luminous as claimed.

**Keywords:** galaxies: high-redshift — galaxies: starburst — submillimeter: galaxies — infrared: galaxies — radio continuum: galaxies

### 1. INTRODUCTION

Dust extinction and a profusion of less luminous foreground galaxies makes it difficult to select high-redshift ultra-luminous star-forming galaxies ( $L_{\text{IR}} \geq 10^{12} L_{\odot}$ ) at rest-frame ultraviolet/optical wavelengths. Although extinction is not an issue at radio wavelengths, an unfavourable  $K$ -correction works against detecting the highest redshift examples,  $z \gg 3$ . Since the advent of large-format submillimeter (submm) cameras such as the Submillimeter Common-User Bolometer Array (SCUBA – Holland et al. 1999), however, it has been possible to exploit the *negative*  $K$ -correction in the submm waveband to select dusty, star-forming galaxies (submm-selected galaxies, or SMGs) almost independently of their redshift (e.g. Franceschini et al. 1991; Blain & Longair 1993).

The scope of this field has been substantially expanded by *Herschel* (Pilbratt et al. 2010) which has surveyed approximately a hundred square degrees of extragalactic sky to the confusion limit at 500  $\mu\text{m}$  (as defined by Nguyen et al. 2010), with simultaneous imaging at 250 and 350  $\mu\text{m}$ , using the SPIRE instrument (Griffin et al. 2010). A SPIRE image of the *Spitzer* First Look Survey (FLS) field, obtained as part of the *Herschel* Multi-Tiered Extragalactic Survey (HerMES<sup>24</sup> – Oliver et al. 2012), led to the discovery of 1HERMES S350 J170647.8+584623 (hereafter HFLS3 – Riechers et al. 2013; Dowell et al. 2014) as an unusually red SPIRE source with  $S_{250} < S_{350} < S_{500}$ , i.e. with its thermal dust peak within or beyond the 500- $\mu\text{m}$  band (see also Cox et al. 2011; Combes et al. 2012; Rawle et al. 2014). Some of these “500- $\mu\text{m}$  risers” are in fact due to synchrotron emission from bright, flat-spectrum radio quasars (e.g. Jenness et al. 2010), but HFLS3 does not exhibit such

<sup>1</sup> UK ATC, Royal Observatory, Blackford Hill, Edinburgh EH9 3HJ, UK

<sup>2</sup> ESO, Karl Schwarzschild Strasse 2, D-85748 Garching, Germany

<sup>3</sup> Institute for Astronomy, University of Edinburgh, Royal Observatory, Blackford Hill, Edinburgh EH9 3HJ, UK

<sup>4</sup> Institute for Computational Cosmology, Durham University, South Road, Durham DH1 3LE, UK

<sup>5</sup> Centre for Astrophysics Research, University of Hertfordshire, Hatfield AL10 9AB, UK

<sup>6</sup> Dept of Physics & Astronomy, University of British Columbia, 6224 Agricultural Road, Vancouver, BC V6T 1Z1, Canada

<sup>7</sup> Astronomy Department, Cornell University, Ithaca, NY 14853

<sup>8</sup> Astronomy and Instrumentation Group, Cardiff University, Cardiff, Wales, UK

<sup>9</sup> Joint Astronomy Centre, 660 N. Ahoku Place, University Park, Hilo, Hawaii 96720

<sup>10</sup> Jet Propulsion Laboratory, National Aeronautics and Space Administration, Pasadena, CA 91109

<sup>11</sup> XMM SOC, ESAC, Apartado 79, 28691 Villanueva de la Canada, Madrid, Spain

<sup>12</sup> Dalhousie University, Department of Physics and Atmospheric Science, Coburg Road, Halifax B3H 1A6, Canada

<sup>13</sup> Astrophysics Group, Imperial College London, Blackett Laboratory, Prince Consort Road, London SW7 2AZ, UK

<sup>14</sup> Center for Astrophysics and Space Astronomy, 389 UCB, University of Colorado, Boulder, CO 80309

<sup>15</sup> Department of Physics and Astronomy, University of California, Irvine, CA 92697

<sup>16</sup> Department of Physics, Virginia Tech, Blacksburg, VA 24061

<sup>17</sup> Department of Physics and Astronomy, University of Waterloo, Waterloo, Ontario N2L 3G1, Canada

<sup>18</sup> IAC, E-38200 La Laguna, Tenerife, Spain

<sup>19</sup> Astronomy Centre, Department of Physics and Astronomy, University of Sussex, Brighton, BN1 9QH, UK


<sup>20</sup> UPMC Univ Paris 06, UMR 7095, IAP, 75014, Paris, France

<sup>21</sup> CNRS, UMR7095, IAP, F-75014, Paris, France

<sup>22</sup> Departamento de Astrofísica, Universidad de La Laguna, E-38205 La Laguna, Tenerife, Spain

<sup>23</sup> Dark Cosmology Centre, Niels Bohr Institute, University of Copenhagen, Denmark

<sup>24</sup> hermes.sussex.ac.uk



**Figure 1.** *Left:* SCUBA-2 imaging at  $850\ \mu\text{m}$ , with  $450\text{-}\mu\text{m}$  and  $850\text{-}\mu\text{m}$  sources marked by red squares and white circles, respectively, for the  $67.2\ \text{arcmin}^2$  where  $\sigma_{850} \leq 1.5\ \text{mJy beam}^{-1}$ . The negative bowl around HFLS3 is a typical artifact of the filtering procedures employed here. The FWHM of the SCUBA-2 beams are shown as solid ellipses. *Right:* Three-color representation of the data obtained using SPIRE at  $250\ \mu\text{m}$ ,  $350\ \mu\text{m}$  and  $500\ \mu\text{m}$  for the same field around HFLS3, superimposed with blue PACS  $160\text{-}\mu\text{m}$  contours. Several of the SCUBA-2  $850\text{-}\mu\text{m}$  sources are associated with green SPIRE sources – those with SEDs peaking at  $350\ \mu\text{m}$ , consistent with  $z \approx 2$ ; others have no obvious SPIRE counterparts and may lie at considerably higher redshifts. The region over which PACS sensitivity is better than half the best is outlined in blue. Positions of faint  $1.4\text{-GHz}$  sources from the  $\sim 1.3''$ -resolution,  $\sigma \sim 11\ \mu\text{Jy beam}^{-1}$  Karl G. Jansky Very Large Array imaging described in Riechers et al. (2013) are marked ‘+’ (the radio catalog covers only  $\approx 25\%$  of the region shown, hence the detection rate is unremarkable). N is up; E is left; offsets from  $\alpha_{2000} = 17:06:47.8$ ,  $\delta_{2000} = +58:46:23$  are marked.

powerful AGN-driven radio emission. Panchromatic spectral-line observations place HFLS3 at  $z = 6.34$  via the detection of  $\text{H}_2\text{O}$ ,  $\text{CO}$ ,  $\text{OH}$ ,  $\text{OH}^+$ ,  $\text{NH}_3$ ,  $[\text{C I}]$  and  $[\text{C II}]$  emission and absorption lines. Its continuum spectral energy distribution (SED) is consistent with a characteristic dust temperature,  $T_d = 56\ \text{K}$ , and a dust mass of  $1.1 \times 10^9 M_\odot$ . Its infrared luminosity,  $L_{\text{IR}} = 2.9 \times 10^{13} L_\odot$ , suggests a star-formation rate (SFR) of  $2900\ \mu\text{L}^{-1} M_\odot\ \text{yr}^{-1}$  for a Chabrier (2003) initial mass function, where the lensing magnification suffered by HFLS3 due to a foreground galaxy less than  $2''$  away has been estimated to be in the range  $\mu_L = 1.2\text{--}1.5$  (Riechers et al. 2013, although Cooray et al. 2014 estimate  $\mu_L = 2.2 \pm 0.3$ ).

It is expected that the most massive galaxies found at very high redshifts grew in (and thus signpost) the densest peaks in the early Universe, making them useful tracers of distant proto-clusters. Above  $z \sim 6$ , such sources may also contribute to the rapid evolution of the neutral fraction of the Universe, during the so-called ‘era of reionisation’, and to the earliest phase of enrichment of the interstellar medium in galaxies, less than  $1\ \text{Gyr}$  after the Big Bang. They may also host the highest redshift quasars. In the submm regime, to explore distant galaxies and their environments we have observed radio galaxies and quasars, typically detecting factor  $\sim 2\text{--}4\times$  over-densities of submm companions around these signposts (e.g. Ivison et al. 2000; Stevens et al. 2003, 2010; Robson et al. 2004; Priddey et al. 2008). Here, we continue this tradition, targeting the most distant known submm galaxy, HFLS3 at  $z = 6.34$ , with the  $10,000$ -pixel SCUBA-2 bolometer camera (Holland et al. 2013), which is more sensitive than *Herschel* to cold dust in high-redshift galaxies.

In §2 we describe our SCUBA-2 observations of the field surrounding HFLS3, after its discovery with SPIRE aboard

*Herschel*, and our reduction of those data. In §3 we analyze the surface density of SCUBA-2 galaxies in the field, and their color, and discuss whether there is any evidence that HFLS3 inhabits an over-dense region of the Universe, as might be expected in hierarchical structure-formation models (e.g. Kaufman et al. 1999; Springel et al. 2005). We finish with our conclusions in §4. Throughout, we adopt a cosmology with  $H_0 = 71\ \text{km s}^{-1} \text{Mpc}^{-1}$ ,  $\Omega_m = 0.27$  and  $\Omega_\Lambda = 0.73$ , so  $1''$  equates to  $5.7\ \text{kpc}$  at  $z = 6.34$ .

## 2. OBSERVATIONS AND DATA REDUCTION

### 2.1. SCUBA-2 imaging and catalogues

Data were obtained simultaneously at  $450$  and  $850\ \mu\text{m}$  in 2011 September 23-24 and 2012 March 14 (project M11BGT01) using SCUBA-2 on the  $15\text{-m}$  James Clerk Maxwell Telescope (JCMT). The observations were taken with the constant speed DAISY pattern, which provides uniform exposure-time coverage in the central  $3'$ -diameter region of a field, and useful coverage over  $12'$ . A total of  $6.9\ \text{hr}$  was spent integrating on HFLS3. Observing conditions were good or excellent, with precipitable-water-vapor levels typically  $1\ \text{mm}$  or less, corresponding to a  $225\text{-GHz}$  optical depth of  $0.05$ . The data were calibrated in flux density against the primary calibrators Uranus and Mars, and also secondary sources CRL 618 and CRL 2688 from the JCMT calibrator list (Dempsey et al. 2013), with estimated calibration uncertainties amounting to  $5\%$  at  $850\ \mu\text{m}$  and  $10\%$  at  $450\ \mu\text{m}$ .

The data were reduced using the Dynamic Iterative Map-Maker within the STARLINK SMURF package (Chapin et al. 2013) called from the ORAC-DR automated pipeline (Cavanagh et al. 2008). The chosen recipe accounted for attenuation of the signal as a result of time-series filtering

and removed residual low-frequency noise from the map using a jack-knife method. The maps were made using inverse-variance weighting, with  $1''$  pixels at both wavelengths, before application of a matched filter (e.g. Chapin et al. 2011).

The map-maker used a ‘blank field’ configuration file, optimized for faint, unresolved or compact sources. This applies a high-pass filter with a spatial cutoff of  $200''$ , corresponding to about 0.8 Hz for a typical scanning speed of  $155'' \text{ s}^{-1}$ . This removes the majority of low-frequency (large spatial-scale) noise, while the remainder is removed using a Fourier-space whitening filter. This is derived from the power spectrum of the central  $9'$  region of a jack-knife map, produced from two independent halves of the total dataset.

This filtering attenuates the peak signal of sources in the map. To estimate the magnitude of this effect, the pipeline re-makes each map with a fake 10-Jy Gaussian added to the raw data, offset from the nominal map centre by  $30''$  to avoid contamination by any target at the map centre. The amplitude of the fake Gaussian in the output map is measured to determine a correction factor. The standard flux conversion factor (FCF), as determined from observations of primary and secondary calibrators, is then multiplied by this factor (1.17 and 1.15 at 850 and  $450 \mu\text{m}$ , respectively) and applied to the final image to give a map calibrated in  $\text{Jy beam}^{-1}$ . The maps with the fake Gaussian are also used to form the point spread function (PSF) for the matched filter since they reflect the effective point-source transfer function of the map-maker.

The SCUBA-2 850- and  $450\text{-}\mu\text{m}$  images shown in Fig. 1 reach noise levels of 0.9 and  $5.0 \text{ mJy beam}^{-1}$  over the central  $3'$ -diameter regions, yielding detections of HFLS3 at approximately the 41- and  $7\text{-}\sigma$  levels, respectively. At  $850 \mu\text{m}$ , the central  $67.2 \text{ arcmin}^2$  of the map has a noise level of  $1.5 \text{ mJy beam}^{-1}$  or better. The astrometry of the SCUBA-2 images was found to be accurate to better than  $1''$  by stacking at the positions of  $3.6\text{-}\mu\text{m}$  and  $1.4\text{-GHz}$  sources in the field.

Following Geach et al. (2013), we create a catalogue of sources from the  $450\text{-}$  and  $850\text{-}\mu\text{m}$  images by searching for peaks in the beam-convolved signal-to-noise ratio maps, recording their coordinates, flux densities and local noise levels. We then mask a region  $1.5\times$  the beam size and then repeat the search. Above a signal-to-noise level of 3.75 the contamination rate due to false detections is below 5%. We adopt this as our detection threshold, listing the 26 sources with  $850\text{-}\mu\text{m}$  flux density uncertainties below  $1.5 \text{ mJy}$  in Table 1, alongside  $450\text{-}\mu\text{m}$  sources selected from the same area at the same significance threshold.

We calculate our completeness levels and flux boosting following Geach et al. (2013), who followed Weiß et al. (2009), injecting  $10^5$  artificial point sources into a map with the same noise properties as the real image. We correct for false positives using the jack-knife map.

## 2.2. Herschel imaging

The acquisition and reduction of 16.8 hrs of *Herschel* SPIRE and (shallow) PACS data for the FLS field (OD159, 164) as part of HerMES is described in detail by Oliver et al. (2012). The SPIRE data, which are confusion limited, are shown as a three-color image in Fig. 1.

We have obtained much deeper data from PACS (Poglitsch et al. 2010) via a 3.9-hr integration as part of programme OT2\_DRIECHER\_3 (OD1329) (see Riechers et al. 2013, for further details). Observations were carried out on 2013 January 01 in mini-scan mapping mode ( $4 \times 15$  re-

**Table 1**  
Sources detected at 850 and  $450 \mu\text{m}$  near HFLS3.

IAU name	$S / \text{mJy}^a$	S/N
850 $\mu\text{m}$ :		
S2FLS850 J170647.67+584623.0 <sup>b</sup>	$35.4 \pm 0.9$	40.9
S2FLS850 J170631.07+584812.9	$11.4 \pm 1.3$	8.6
S2FLS850 J170621.93+584826.8	$9.3 \pm 1.4$	6.5
S2FLS850 J170646.64+584816.0	$5.6 \pm 1.0$	5.5
S2FLS850 J170647.80+584735.0	$4.9 \pm 0.9$	5.3
S2FLS850 J170701.41+584318.0	$7.5 \pm 1.4$	5.3
S2FLS850 J170625.54+584725.9	$6.3 \pm 1.3$	4.8
S2FLS850 J170642.92+584456.0	$4.7 \pm 1.0$	4.6
S2FLS850 J170717.24+584535.8	$5.8 \pm 1.3$	4.6
S2FLS850 J170659.77+584859.0	$5.6 \pm 1.2$	4.6
S2FLS850 J170723.05+584719.7	$6.1 \pm 1.4$	4.5
S2FLS850 J170642.00+585004.0	$5.4 \pm 1.2$	4.5
S2FLS850 J170630.54+585015.9	$6.2 \pm 1.4$	4.4
S2FLS850 J170710.97+584811.9	$5.4 \pm 1.3$	4.2
S2FLS850 J170653.32+584246.0	$5.5 \pm 1.3$	4.2
S2FLS850 J170713.78+584618.8	$4.9 \pm 1.2$	4.2
S2FLS850 J170638.27+585009.0	$5.3 \pm 1.3$	4.2
S2FLS850 J170710.54+584403.9	$6.1 \pm 1.5$	4.1
S2FLS850 J170724.30+584514.7	$5.9 \pm 1.5$	4.0
S2FLS850 J170700.38+584225.0	$5.9 \pm 1.5$	4.0
S2FLS850 J170715.48+584859.8	$5.4 \pm 1.3$	4.0
S2FLS850 J170614.39+584458.7	$6.0 \pm 1.5$	4.0
S2FLS850 J170641.49+585039.0	$5.3 \pm 1.3$	4.0
S2FLS850 J170637.26+584616.0	$3.9 \pm 1.0$	3.9
S2FLS850 J170633.28+584430.0	$5.2 \pm 1.3$	3.8
S2FLS850 J170654.87+584614.0	$3.3 \pm 0.9$	3.8
450 $\mu\text{m}$ :		
S2FLS450 J170647.80+584620.0 <sup>b</sup>	$39.8 \pm 5.5$	7.3
S2FLS450 J170701.05+584715.0	$29.4 \pm 6.9$	4.3
S2FLS450 J170631.33+584813.9	$36.3 \pm 8.8$	4.1
S2FLS450 J170636.12+584224.0	$32.5 \pm 8.3$	3.9
S2FLS450 J170658.59+584419.0	$30.2 \pm 7.8$	3.9
S2FLS450 J170711.99+584734.9	$34.5 \pm 8.9$	3.9

<sup>a</sup> Deboosted flux densities; errors exclude 5 and 10% calibration uncertainties at 850 and  $450 \mu\text{m}$ , respectively.

<sup>b</sup> HFLS3.


peats), using the 70- plus  $160\text{-}\mu\text{m}$  parallel mode and the 110- plus  $160\text{-}\mu\text{m}$  parallel mode for one orthogonal cross-scan pair each. In the 70-, 110-, and  $160\text{-}\mu\text{m}$  bands, the r.m.s. sensitivities at the position of HFLS3 are 0.67, 0.73, and  $1.35 \text{ mJy beam}^{-1}$ , respectively. Data reduction and mosaicing were carried out using standard procedures. The absolute flux density scale is accurate to 5%. The  $160\text{-}\mu\text{m}$  PACS image, the only one potentially useful in the context of faint, distant galaxies, is shown in Fig. 1.

The 250-, 350- and  $500\text{-}\mu\text{m}$  flux densities,  $S_{250}$ ,  $S_{350}$  and  $S_{500}$ , at the positions<sup>25</sup> of the 19 SMGs discussed in §2.1 were determined using beam-convolved SPIRE maps. None of our SMGs lie near bright SPIRE sources so we expect the uncertainties associated with these flux densities should be close to the typical confusion levels,  $\approx 6 \text{ mJy}$  (where hereafter we adopt  $\sigma_{\text{conf}}$  in each SPIRE band from Nguyen et al. 2010).

## 3. RESULTS, ANALYSIS AND DISCUSSION

HFLS3 dominates the submm sky in the  $67.2\text{-arcmin}^2$  ( $8 \text{ Mpc}^2$ ) region we have mapped at  $850 \mu\text{m}$  with SCUBA-2, being three times brighter than the next-brightest submm emitter (Fig. 1; Table 1). At  $450 \mu\text{m}$ , HFLS3 is the brightest SMG in the region, despite the peak of its SED having moved beyond that filter; it is one of two sources detected formally

<sup>25</sup> Positions are known to  $\sigma_{\text{pos}} \approx 2.2''$  even for the least significant SMGs, a small fraction of the beam-convolved SPIRE point spread function.



**Figure 2.** Source counts in several flux density bins at 450 and 850  $\mu\text{m}$  in the 67.2-arcmin<sup>2</sup> region around HFLS3, excluding HFLS3 itself, relative to those found in typical blank fields by Coppin et al. (2006) and Geach et al. (2013). We see no indication of a strong excess of submm emitters across the field, compared to blank-field counts, which suggests there is no over-density of SMGs on  $\sim 1.5$ -Mpc scales around HFLS3. The raw counts, uncorrected for incompleteness, are shown as open symbols.

at both 450 and 850  $\mu\text{m}$ . Perhaps unsurprisingly, there are no sources in common between the SCUBA-2 and PACS images.


We see no evidence for an over-density of SMGs on  $< 1.5$ -Mpc scales around the position of HFLS3 in either our 450 or 850- $\mu\text{m}$  maps (Fig. 1).

### 3.1. Number counts relative to blank fields

Although no obvious cluster of submm emitters is visible near HFLS3 in Fig. 1, we must ask whether the entire 8-Mpc<sup>2</sup> field might be over-dense in SMGs? Fig. 2 shows the density of sources brighter than  $S_\nu$  at 450 and 850  $\mu\text{m}$  – extracted at the 3.75- $\sigma$  level and corrected for incompleteness using the analysis discussed in §2.1, excluding HFLS3 itself – relative to the source density seen in typical blank-field surveys, where the same techniques have been used to construct catalogues and correct for incompleteness (Coppin et al. 2006; Geach et al. 2013). The only hint of an over-density comes in the 850- $\mu\text{m}$  bin at 5 mJy, but a  $\approx 2$ - $\sigma$  deviation is not unusual when plotting seven points. Simplifying matters, then, by taking only one large bin above our detection threshold at 850  $\mu\text{m}$ , we find 26 sources (25 if we ignore HFLS3) against an expectation of 20 from Coppin et al. 2006 – on the face of it, a  $\approx 1$ - $\sigma$  over-density. When we take into account the errors associated with the Coppin et al. 2006 counts, we expect to see 26 (25) sources in a random field of the same size 13% (19%) of the time, so the result obtained intuitively from studying Fig. 2 is confirmed – any excess is statistically unconvincing. No significant over-density of SMGs is apparent in the vicinity of HFLS3, at least not on Mpc scales at flux densities above the JCMT confusion limit.

### 3.2. Redshift constraints for sources in the field

The  $K$ -correction in the submm waveband means our SCUBA-2 maps are sensitive to SMGs across a very wide



**Figure 3.** Submm color-color plots for HFLS3 and its neighbouring SMGs, adapted from Ivison et al. (2012) and Swinbank et al. (2014). For sources with flux densities below twice the confusion noise, ( $\sigma_{\text{conf}}$ ; Nguyen et al. 2010), we show limits based on the measured flux density (zero, if negative) plus  $\sigma_{\text{conf}}$ . The colored backgrounds indicate the typical redshift of the subset of  $10^7$  model SEDs that fall in each pixel, where in the *top* panel we have adopted the dust temperature of HFLS3 ( $T_d = 56$  K), a flat redshift distribution from  $z = 0$ –7, a flat  $\beta$  distribution from 1.5–2.0 (for HFLS3,  $\beta = 1.92 \pm 0.12$ ), and 10% flux density errors. HFLS3 is the star symbol. The error bar is typical for sources detected at 350, 500 and 850  $\mu\text{m}$ . Those sources without detections at 350 and 500  $\mu\text{m}$  are plotted arbitrarily at  $S_{350}/S_{500} = 1.5$ , inside a rectangular box; some could be considerably redder than this. Of those detected in both 350 and 500  $\mu\text{m}$ , S2FLS J170621.93+584826.8 is the reddest in  $S_{350}/S_{500}$ . A handful are as red as HFLS3 in  $S_{850}/S_{500}$ , e.g. S2FLS J170647.80+584735.0, which is a 5.3- $\sigma$  SCUBA-2 source with no SPIRE counterpart. Despite its relatively high  $T_d$ , HFLS3 is the reddest of the sources detected in the three bands, though we note that redder objects may exist amongst the SPIRE-undetected SMGs. The *lower* plot shows a flat  $\beta$  distribution from 1.5–2.0 and  $T_d = 35$  K, illustrating how  $T_d$  influences the colors of an SMG. Comparing the redshift constraints from both panels makes it clear that nothing can be usefully concluded about the likely redshifts from the far-infrared colors unless there are independent constraints on the dust temperature.

redshift range, reducing the contrast of any potential structure around HFLS3. However, with SCUBA-2, SPIRE and PACS photometry in hand, we are able to crudely constrain the likely redshifts of the galaxies detected in the field surrounding HFLS3, using their far-infrared/submm colors. Fig. 3 shows color-color plots for HFLS3 and its neighbouring SMGs, designed to exploit information from SCUBA-2 at  $850\ \mu\text{m}$  to constrain the redshifts of galaxies at  $z > 2$  (Ivison et al. 2012), probing their colors across the rest-frame  $\approx 100\text{-}\mu\text{m}$  bump seen in the SEDs of all dusty, star-forming galaxies. The colored backgrounds in the upper and lower panels of Fig. 3 indicate the typical redshift of the subset of  $10^7$  model SEDs that fall in each pixel, where we have adopted a flat redshift distribution ( $z = 0\text{--}7$ ), a flat distribution for the spectral dependence of the dust emissivity ( $\beta = 1.8\text{--}2.0$ , centered on the  $\beta$  measured for HFLS3) and 10% flux density uncertainties. For the upper panel of Fig. 3 we adopt the dust temperature of HFLS3 ( $T_d = 56\ \text{K}$ ).

We concentrate only on those galaxies detected by SCUBA-2, since SPIRE-detected galaxy with a typical SED in the vicinity of HFLS3 could not have evaded detection at  $850\ \mu\text{m}$ . Despite its relatively high  $T_d$ , HFLS3 is the reddest source detected in the three bands used to make Fig. 3.


For SMGs with SPIRE flux densities below  $2\sigma_{\text{conf}}$ , we plot limits based on the measured flux density (zero, if negative) plus  $\sigma_{\text{conf}}$ . We have placed those sources without detections at 350 and  $500\ \mu\text{m}$  at  $S_{350}/S_{500} = 2$ , arbitrarily; some could be considerably redder than this in  $S_{350}/S_{500}$ , but we cannot constrain this color reliably with the relatively shallow *Herschel* data at our disposal. Several SMGs may also be as red as HFLS3 in  $S_{850}/S_{500}$ , perhaps redder. One particularly interesting example is S2FLS850J170647.80+584735.0, a  $5.3\text{-}\sigma$  SCUBA-2 source with no significant SPIRE emission. With  $S_{850}/S_{500} > 0.9$ , this SMG likely lies at  $z > 5$ , with a lower  $T_d$  and luminosity than HFLS3.

The lower panel of Fig. 3 shows the effect of lowering  $T_d$ , illustrating an issue long-known to hamper studies of this kind: far-infrared/submm colors are sensitive only to  $(1+z)/T_d$  (Blain 1999), i.e. redshift and  $T_d$  are degenerate. As a result, our current data does not allow us to conclude with certainty that the environment surrounding HFLS3 contains other luminous, dusty starbursts; neither can we rule it out.

Single-dish imaging of this field at 450, 850, 1100 and  $2000\ \mu\text{m}$  is possible from the ground, reaching  $\sigma_{450} \sim 2.5\ \text{mJy}$  and  $\sigma_{2000} \sim 0.1\ \text{mJy}$  over tens of arcmin<sup>2</sup> with existing facilities in a few tens of hours. Would these data be capable of further constraining the redshifts of the SMGs discovered here? Fig. 4 shows  $S_{850}/S_{450}$  versus  $S_{2000}/S_{850}$  and we see that the latter color offers little insight. For  $S_{850}/S_{450} \lesssim 0.3$  we can rule out  $z \gtrsim 2$  for all but the warmest dust;  $S_{850}/S_{450} \gtrsim 1$  suggests  $z \gtrsim 5$ , even for  $T_d = 35\ \text{K}$ , with significantly cooler dust unlikely in this redshift regime. Deeper  $450\text{-}\mu\text{m}$  imaging would therefore be useful.

### 3.3. Predictions from the Millennium Simulation

Is the field surrounding HFLS3 less over-dense in submm sources than expected for such a massive galaxy living in a biased environment at high redshift, similar to those found around high-redshift radio galaxies and radio-loud quasars at  $z = 2\text{--}4$  (e.g. Stevens et al. 2003, 2004)? The answer to this question may have implications for the potential gravitational amplification suffered by HFLS3 (see §1) or for investigating the potential presence of a buried AGN and its role in supporting its high IR luminosity.



**Figure 4.**  $S_{850}/S_{450}$  versus  $S_{2000}/S_{850}$  ratios for which deep, relatively unconfused and unbiased data can be obtained from the ground, covering tens of square arcminutes. The observed colors of HFLS3 are shown. The colored background indicates the typical redshift of the subset of  $10^7$  model SEDs that fall in each pixel, where we have adopted the same dust parameters as Fig. 3. *Inset:* Same plot, having changed only  $T_d$ , to 35 K, illustrating the degeneracy between redshift and  $T_d$ .  $S_{850}/S_{450} \lesssim 0.3$  suggests  $z \lesssim 2$  for all but the warmest dust;  $S_{850}/S_{450} \gtrsim 1$  suggests  $z \gtrsim 5$ , with dust much cooler than 35 K unlikely at these redshifts;  $S_{2000}/S_{850}$  offers less insight.

By necessity this comparison will be crude. We therefore selected the implementation of the Bower et al. (2006) galaxy-formation recipe in the Millennium Simulation (Springel et al. 2005) and searched the  $z = 6.2$  output for galaxies with a total baryonic mass in excess of  $1.3 \times 10^{11} M_\odot$ , consistent with the combined mass of gas and stars estimated for HFLS3 (Riechers et al. 2013). By adopting a total baryonic mass cut we are less sensitive to details of the early star-formation histories of galaxies in the model.

We find just one galaxy in the  $3.2 \times 10^8\ \text{Mpc}^3$  volume at  $z = 6.2$  with a total baryonic mass above  $1.3 \times 10^{11} M_\odot$ . All its baryonic mass is in stars; it hosts a  $2 \times 10^8 M_\odot$  black hole and is the central galaxy of a  $6 \times 10^{12} M_\odot$  halo,  $4\times$  more massive than the next most massive galaxy's halo within the volume, and the optimal environment to find merging galaxies according to the simulations of Hopkins et al. (2008). Another 16 galaxies are spread across a  $\sim 0.7$ -comoving-Mpc-diameter region around the most massive galaxy, but most of these are dwarf galaxies with baryonic masses,  $\lesssim 10^9 M_\odot$ . Inside a sphere with an angular size of  $9'$  diameter, centered on the  $1.3 \times 10^{11} M_\odot$  galaxy, only two galaxies have total baryonic masses of  $\gtrsim 15\%$  of the mass of HFLS3 (we choose this limit as the faintest submm emitters in this field have  $850\text{-}\mu\text{m}$  flux densities of around 15% that of HFLS3); the next most massive galaxy has half this mass. The total masses of these two companion galaxies in stars and gas are approximately  $3$  and  $4 \times 10^{10} M_\odot$  and their predicted  $K_{\text{vega}}$  magnitudes are 23.5 and 25.0. The HFLS3 clone is predicted to have  $K_{\text{vega}} = 22.5$  for a distance modulus of 49.0, about a magnitude fainter than observed.

From this simple theoretical comparison, we thus expect  $2 \pm 2$  detectable galaxies in the vicinity of HFLS3 if, as ex-

pected, their starburst lifetimes are a significant fraction of the time available at this early epoch. This is consistent with the fact that some high-redshift SMGs do have submm-bright companions (e.g. GN 20 – Daddi et al. 2009) while others have Lyman-break galaxies nearby but no submm-bright companions (e.g. AzTEC-3 – Capak et al. 2011).

Having found no clear evidence for or against the level of over-density expected in simulations, we can draw no strong conclusions regarding the likely gravitational amplification suffered by HFLS3, or for the likely fraction of its luminosity provided by a buried AGN.

#### 4. CONCLUSIONS

We have detected the most distant, dusty starburst galaxy, HFLS3, at high significance with SCUBA-2. We detect another 29 dusty galaxies within an area of  $67.2 \text{ arcmin}^2$  surrounding HFLS3, most of them likely at lower redshift. We find no compelling evidence, from surface density or color, for an over-density of SMGs around HFLS3, although applying similar selection criteria to theoretical models suggests that a modest excess could be expected, as is found for some other high-redshift SMGs (e.g. GN 20 – Daddi et al. 2009). We can therefore draw no strong conclusions regarding the likely gravitational amplification suffered by HFLS3, or for the likely fraction of its luminosity provided by a buried AGN.

#### ACKNOWLEDGEMENTS

We thank John Helly for help with the Millennium Simulation. RJI and IRS acknowledge support from the European Research Council (ERC) in the form of Advanced Investigator programs, COSMICISM and DUSTYGAL, respectively. IRS also acknowledges support from the UK’s Science and Technology Facilities Council (STFC, ST/I001573/1), a Leverhulme Fellowship and a Royal Society/Wolfson Merit Award. JEG acknowledges the Royal Society for support. The Dark Cosmology Centre is funded by the Danish National Research Foundation. The JCMT is operated by the Joint Astronomy Centre on behalf of STFC, the National Research Council of Canada and (until 31 March 2013) the Netherlands Organisation for Scientific Research. Additional funds for the construction of SCUBA-2 were provided by the Canada Foundation for Innovation. *Herschel* is an ESA space observatory with science instruments provided by European-led Principal Investigator consortia and with important participation from NASA. SPIRE was developed by a consortium of institutes led by Cardiff Univ. (UK) and including: Univ. Lethbridge (Canada); NAOC (China); CEA, LAM (France); IFSI, Univ. Padua (Italy); IAC (Spain);

Stockholm Observatory (Sweden); Imperial College London, RAL, UCL-MSSL, UKATC, Univ. Sussex (UK); and Caltech, JPL, NHSC, Univ. Colorado (USA). This development has been supported by national funding agencies: CSA (Canada); NAOC (China); CEA, CNES, CNRS (France); ASI (Italy); MCINN (Spain); SNSB (Sweden); STFC, UKSA (UK); and NASA (USA). The JCMT and *Herschel* data used in this paper can be obtained from the JCMT archive ([www.jach.hawaii.edu/JCMT/archive](http://www.jach.hawaii.edu/JCMT/archive)) and the *Herschel* Database in Marseille, HeDaM, ([hedam.oamp.fr/HerMES](http://hedam.oamp.fr/HerMES)), respectively.

*Facilities:* JCMT, *Herschel*

#### REFERENCES

- Blain, A. W. 1999, MNRAS, 309, 955
- Blain, A. W., & Longair, M. S. 1993, MNRAS, 264, 509
- Bower, R. G., Benson, A. J., Malbon, R., et al. 2006, MNRAS, 370, 645
- Capak, P. L., Riechers, D., Scoville, N. Z., et al. 2011, Nature, 470, 233
- Cavanagh, B., Jenness, T., Economou, F., & Currie, M. J. 2008, Astronomische Nachrichten, 329, 295
- Chabrier, G. 2003, PASP, 115, 763
- Chapin, E. L., Berry, D. S., Gibb, A. G., et al. 2013, MNRAS, 430, 2545
- Chapin, E. L., Chapman, S. C., Coppin, K. E., et al. 2011, MNRAS, 411, 505
- Combes, F., Rex, M., Rawle, T. D., et al. 2012, A&A, 538, L4
- Cooray, A., Calanog, J., Wardlow, J. L., et al. 2014, ArXiv e-prints, arXiv:1404.1378
- Coppin, K., Chapin, E. L., Mortier, A. M. J., et al. 2006, MNRAS, 372, 1621
- Cox, P., Krips, M., Neri, R., et al. 2011, ApJ, 740, 63
- Daddi, E., Dannerbauer, H., Stern, D., et al. 2009, ApJ, 694, 1517
- Dempsey, J. T., Friberg, P., Jenness, T., et al. 2013, MNRAS, 430, 2534
- Dowell, C. D., Conley, A., Glenn, J., et al. 2014, ApJ, 780, 75
- Franceschini, A., Toffolatti, L., Mazzei, P., Danese, L., & de Zotti, G. 1991, AAPS, 89, 285
- Geach, J. E., Chapin, E. L., Coppin, K. E. K., et al. 2013, MNRAS, 432, 53
- Griffin, M. J., Abergel, A., Abreu, A., et al. 2010, A&A, 518, L3+
- Holland, W. S., Robson, E. I., Gear, W. K., et al. 1999, MNRAS, 303, 659
- Holland, W. S., Bintley, D., Chapin, E. L., et al. 2013, MNRAS, 430, 2513
- Hopkins, P. F., Hernquist, L., Cox, T. J., & Kereš, D. 2008, ApJS, 175, 356
- Iverson, R. J., Dunlop, J. S., Smail, I., et al. 2000, ApJ, 542, 27
- Iverson, R. J., Smail, I., Amblard, A., et al. 2012, MNRAS, 425, 1320
- Jenness, T., Robson, E. I., & Stevens, J. A. 2010, MNRAS, 401, 1240
- Kaufman, M. J., Wolfire, M. G., Hollenbach, D. J., & Luhman, M. L. 1999, ApJ, 527, 795
- Nguyen, H. T., Schulz, B., Levenson, L., et al. 2010, A&A, 518, L5+
- Oliver, S. J., Bock, J., Altieri, B., et al. 2012, MNRAS, 424, 1614
- Pilbratt, G. L., Riedinger, J. R., Passvogel, T., et al. 2010, A&A, 518, L1+
- Poglitsch, A., Waelkens, C., Geis, N., et al. 2010, A&A, 518, L2+
- Priddey, R. S., Iverson, R. J., & Isaak, K. G. 2008, MNRAS, 383, 289
- Rawle, T. D., Egami, E., Bussmann, R. S., et al. 2014, ApJ, 783, 59
- Riechers, D. A., Bradford, C. M., Clements, D. L., et al. 2013, Nature, 496, 329
- Robson, I., Priddey, R. S., Isaak, K. G., & McMahon, R. G. 2004, MNRAS, 351, L29
- Springel, V., White, S. D. M., Jenkins, A., et al. 2005, Nature, 435, 629
- Stevens, J. A., Jarvis, M. J., Coppin, K. E. K., et al. 2010, MNRAS, 405, 2623
- Stevens, J. A., Page, M. J., Iverson, R. J., Smail, I., & Carrera, F. J. 2004, ApJ, 604, L17
- Stevens, J. A., Iverson, R. J., Dunlop, J. S., et al. 2003, Nature, 425, 264
- Swinbank, A. M., Simpson, J. M., Smail, I., et al. 2014, MNRAS, 438, 1267
- Weiß, A., Kovács, A., Coppin, K., et al. 2009, ApJ, 707, 1201