



WILEY

## Physical and biological controls on fine sediment transport and storage in rivers

Journal:	<i>WIREs Water</i>
Manuscript ID	WATER-450.R1
Wiley - Manuscript type:	Advanced Review
Date Submitted by the Author:	n/a
Complete List of Authors:	<p>Wilkes, Martin; Coventry University, Centre for Agroecology Water and Resilience  Gittins, Joshua; Lancaster Environment Centre  Mathers, Kate; Eawag - Swiss Federal Institute of Aquatic Science and Technology  Mason, Richard; Loughborough University Department of Geography  Casas-Mulet, Roser; Technical University of Munich, Department of Ecology and Ecosystem Management; Cardiff University; University of Melbourne Department of Infrastructure Engineering  Vanzo, Davide; Eawag - Swiss Federal Institute of Aquatic Science and Technology  Mckenzie, Morwenna; Coventry University, Centre for Agroecology Water and Resilience  Murray-Bligh, John; Environment Agency  England, Judy; Environment Agency  Gurnell, Angela; Queen Mary, University of London, Geography  Jones, J. Iwan; Queen Mary University of London - Mile End Campus</p>
Choose 1-3 topics to categorize your article:	
Keywords:	River, Fine sediment, Ecosystem engineering, Geomorphology, Ecology

SCHOLARONE™  
Manuscripts

**PLEASE READ ALL 'COMMENT' TEXT BEFORE PREPARING YOUR ARTICLE.** If you do not see the Comments, select View > Print Layout. Please delete them before submitting (Review > Delete All Comments in Document) so that peer reviewers see a clean copy of the manuscript.

**Remember that you are writing for an interdisciplinary audience. Please be sure to discuss interdisciplinary themes, issues, debates, etc. where appropriate.** Note that the WIREs are forums for review articles, rather than primary literature describing the results of original research.

If you have any questions, [contact your editorial office](#).



**Article Title: Physical and biological controls on fine sediment transport and storage in rivers**

**Article Type:**

ADVANCED REVIEW

**Authors:**

[List each person's full name, [ORCID ID](#), affiliation, email address, and any conflicts of interest. Copy rows as necessary for additional authors. Please use an asterisk (\*) to indicate the corresponding author.]

**Martin A. Wilkes\***

orcid.org/0000-0002-2377-3124

martin.wilkes@coventry.ac.uk

Centre for Agroecology, Water and Resilience, Coventry University, United Kingdom

**Joshua Gittins**

orcid.org/0000-0002-6098-8917

j.gittins1@lancaster.ac.uk

Lancaster Environment Centre, Lancaster University, United Kingdom

**Kate L. Mathers**

orcid.org/0000-0003-3741-1439

kate.mathers@eawag.ch

Eawag - Swiss Federal Institute of Aquatic Science and Technology, Kastanienbaum, Switzerland

**Richard Mason**

R.J.Mason@lboro.ac.uk

Department of Geography, Loughborough University, United Kingdom


<p><b>Roser Casas-Mulet</b>  orcid.org/0000-0002-7139-8859  rosier.casas-mulet@tum.de  Aquatic Systems Biology Unit, Department of Ecology and Ecosystem Management,  Technical University of Munich, Bavaria, Germany; Water Research Institute, Cardiff  University, Wales, UK; Department of Infrastructure Engineering, The University of  Melbourne, Victoria, Australia</p>
<p><b>Davide Vanzo</b>  orcid.org/0000-0002-2033-9197  davide.vanzo@eawag.ch  Eawag - Swiss Federal Institute of Aquatic Science and Technology, Kastanienbaum,  Switzerland</p>
<p><b>Morwenna Mckenzie</b>  orcid.org/0000-0001-8906-4286  mckenz36@uni.coventry.ac.uk  Centre for Agroecology, Water and Resilience, Coventry University, United Kingdom</p>
<p><b>John Murray-Bligh</b>  john.murray-bligh@environment-agency.gov.uk  Environment Agency, Exeter, United Kingdom</p>
<p><b>Judy England</b>  orcid.org/0000-0001-5247-4812  judy.england@environment-agency.gov.uk  Environment Agency, Exeter, United Kingdom</p>
<p><b>Angela Gurnell</b>  orcid.org/0000-0002-7249-8202  a.m.gurnell@qmul.ac.uk  School of Geography, Queen Mary University of London, United Kingdom</p>
<p><b>J. Iwan Jones</b>  orcid.org/0000-0002-7238-2509  j.i.jones@qmul.ac.uk  School of Biological and Chemical Sciences, Queen Mary University of London, United  Kingdom</p>

## Abstract

Excess fine sediment, comprising particles <2 mm in diameter, is a major cause of ecological degradation in rivers. The erosion of fine sediment from terrestrial or aquatic sources, its delivery to the river, and its storage and transport in the fluvial environment are controlled by a complex interplay of physical, biological and anthropogenic factors. Whilst the physical controls exerted on fine sediment dynamics are relatively well-documented, the role of biological processes and their interactions with hydraulic and physico-chemical phenomena has been largely overlooked. The activities of biota, from primary producers to predators, exert strong controls on fine sediment deposition, infiltration and resuspension. For example, extracellular polymeric substances (EPS) associated with biofilms increase deposition and decrease resuspension. In lower energy rivers, aquatic macrophyte growth and senescence **is-are** intimately linked to sediment retention and loss, whereas riparian trees are dominant ecosystem

engineers in high energy systems. Fish and invertebrates also have profound effects on fine sediment dynamics through activities that drive both particle deposition and erosion depending on species composition and abiotic conditions. The functional traits of species present will determine not only these biotic *effects* but also the *responses* of river ecosystems to excess fine sediment. We discuss which traits are involved and put them into context with spatial processes that occur throughout the river network. Whilst strides towards better understanding of the impacts of excess fine sediment have been made, further progress to identify the most effective management approaches is urgently required through close communication between authorities and scientists.

### Graphical/Visual Abstract and Caption



Controls on the delivery, transport and storage of fine sediment and the resulting ecological responses in river networks. From the sediment sources in the landscape to the ecological impact in the river, the transport of particles and the responses of biota are subject to potential lags and delays as sediment is temporarily stored and time is taken for local populations to reach critical life-stages where fine sediment limits survival.

### Introduction

Excess fine sediment ('fines'), typically defined as organic and inorganic particles <2 mm in diameter, is one of the principal reasons for the failure of waterbodies to achieve good ecological status (GES) under the EU Water Framework Directive (WFD)<sup>1,2</sup>. In light of this, the recent finding that most lowland rivers in the UK are transport-limited, leading to saturation of river-bedsstreambeds with fine sediment<sup>3</sup>, is of critical importance. We may safely assume that this result can be generalised to other similar environments worldwide. It implies that in-channel transport and storage processes are primary controls on the volume of fines stored on or within the river-bedstreambed, whereas both exogenous (i.e. sediment supply) and endogenous processes (e.g. bioturbation) primarily influence the concentration of fine sediments carried in suspension and their flux through the river network. Greater understanding of these processes is urgently required. To facilitate progress in the primary research areas involved, we provide a synthesis of current knowledge on the physical and biological factors controlling the transport, storage and ecological impacts of fine sediment in river channels to complement recent reviews on catchment-scale, terrestrial processes (e.g.<sup>4</sup>).

In reviewing the factors that control fine sediment transport and storage in river channels, we emphasise biological controls as these have received far less attention than physical controls (e.g.<sup>5</sup>). Bottom-up, biologically-driven forces are increasingly recognised as important elements for understanding fluvial geomorphic processes<sup>6</sup>. We build upon this knowledge in the specific context of fines. Ecological impacts at organism- and community- levels, driven by top-down abiotic forces, have been reviewed in detail elsewhere<sup>7–10</sup> but how the ecological response to fine sediment is mediated by the structure and function of ecosystems across multiple levels of organisation is less well-understood. We therefore also include information on ecological resistance ([ability to withstand perturbation](#)) and resilience ([ability to recover from disturbance](#)) to fine sediment stress at the level of individual organisms to regional biogeographies. We mostly limit the discussion to processes occurring after the sediment has been delivered to the river channel but also present some exogenous processes insofar as they influence in-channel controls. Our discussion is structured according to the conceptual model in Figure 1, going beyond a recent review on infiltration ('colmation')<sup>11</sup> to provide a more holistic perspective on the fine sediment 'problem'.




Figure 1 – Controls on the delivery, transport and storage of fine sediment and the resulting ecological responses in river networks. Natural and anthropogenic factors control the quantity and quality of sediment potentially available for delivery to the river channel. Whether or not this potential is realised, and the quantity and composition of sediment conserved during the pluvial phase, depends on delivery pathways that connect the source to the river channel. Once delivered to the river channel, particles may be transported through suspension or saltation, temporarily stored as surface-deposited material, or become infiltrated in the river bedstreambed and stored on a longer-term basis. Which compartment particles are transported or stored in depends on a number of controls, including the inherent properties of the particles, hydraulics, and the activities of biota. The latter is in turn influenced by

1  
2  
3 physico-chemical factors (e.g. oxygen, nutrients) that work to limit or enhance biological processes. The  
4 hydraulic and biological controls determine the compartments within which particles are transported or  
5 stored, structuring the physical habitat, modifying available resources and potentially filtering out  
6 unsuitable species from the species pool on the basis of their traits. Such environmental filtering effects  
7 may be interfered with by biogeographical processes related to dispersal. The fate of fine sediment, and  
8 the extent and nature of the impact it causes is subject to potential lags as sediment is temporarily  
9 stored within the landscape. Ecological impacts may also be delayed as time is taken for local  
10 populations to reach critical life-stages where fine sediment limits survival.  
11  
12  
13  
14  
15  
16

### 17 **CONTROLS ON SEDIMENT YIELD AND DELIVERY TO THE RIVER CHANNEL**

18 The yield of fine sediment to the river channel from the catchment (expressed as specific yield,  $t\ km^{-2}\ yr^{-1}$ )  
19 is a consequence of the mobilisation of sediment from a variety of sources, the transport of that  
20 sediment through the landscape and any storage, either temporary or longer-term, within the  
21 landscape. Hence, many factors influence the quantity and composition (including *inter alia* grain size,  
22 organic content, and associated contaminants and nutrients) of fine sediment delivered to the channel.  
23 Whilst soil type, climate and geology have a major influence on the mobilisation of fine sediment, their  
24 effects are modified by local factors, including topography, land-use, vegetation and management. As  
25 the mobilised fine sediment is transported through the landscape from the source to the river, there is  
26 further potential for sorting and retention to influence the quantity and composition of the sediment  
27 ultimately delivered to the river channel. For this reason, well-connected sources tend to contribute  
28 most to the fine sediment yield.  
29  
30  
31  
32

33 As the characteristics of sediment from different locations and sources vary (e.g. arable land, road  
34 verges), methods have been developed to identify and apportion the contributions from the various  
35 potential sources of both the inorganic (e.g.<sup>12,13</sup>) and organic (e.g.<sup>14</sup>) components of fine sediment  
36 sampled from rivers. Such source apportionment studies indicate that some sources contribute  
37 disproportionately relative to their area in the catchment, particularly those that are well-connected,  
38 either arising close to the river or involving pathways over impervious surfaces, or both (e.g. direct  
39 inputs from sewage treatment works, damaged road verges)<sup>14–16</sup>. Part of the yield is derived from  
40 natural processes such as unenhanced background erosion of soils and river banks. However, the  
41 dominance of human influences on the landscape has affected fine sediment dynamics since early  
42 agriculture<sup>17–19</sup>. More recently, the total yield of fine sediment has increased from historical levels,  
43 particularly in the post-war years with the intensification of agriculture and expansion of urban areas<sup>20</sup>,  
44 such that agricultural, industrial and urban sources now contribute a substantial proportion, if not the  
45 majority of the yield of fine sediment to rivers<sup>2</sup>. As such, both the quantity and composition of fine  
46 (inorganic and organic) sediment delivered to rivers far exceeds modern background levels; [the  
47 estimated total loss of sediment in England and Wales was 1,389,818 t yr<sup>-1</sup> in excess of target modern  
48 background delivery rates<sup>21</sup>, defined as the early twentieth century up to ~1940<sup>20</sup>. \(defined as the early  
49 twentieth century up to ~1940\)<sup>20,21</sup>.](#)  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## **ABIOTIC CONTROLS ON FINE SEDIMENT TRANSPORT AND STORAGE**

### *Controls on fine sediment: Hydraulics and geomorphology*

Transport and storage of fines in streambeds plays a key role in the natural development of the sediment matrix<sup>22–24</sup>, contributing to habitat quality and quantity by influencing the hydrological and geomorphological behaviour of a river<sup>25</sup>. Fine sediments are composed of organic and inorganic components that are transported in rivers as solid load or may precipitate from the dissolved load (<0.45 µm). The majority of fines move through river systems as solid suspended load<sup>26</sup> in the form of flocculated or aggregated particles due to their cohesive nature<sup>27</sup>. The load of fine sediments in a river will be determined by the relationship between flow conditions and the structure, density and size of the sediment particles.

The magnitude and timing of events exhibiting high suspended sediment loads ~~is~~ ~~are~~ closely related to the hydrological regime. In rivers with a near-natural flow regime, sediment transport is positively correlated with flow intensity<sup>28</sup>. Human alterations, such as dams and channelisation, that modify or disrupt the flow and sediment longitudinal continuity, have multiple effects on sediment dynamics and the morphology of downstream reaches<sup>29</sup>. Moreover, sediment flushing operations adopted for the maintenance of reservoirs can lead to extreme events characterised by high suspended sediment loads and the potential for widespread deposition downstream<sup>30</sup>.

Most accumulation of fines in the streambed occurs as a result of gravitational deposition from the suspended load<sup>5,22,23,31–33</sup>. Hydraulic conditions in the water column determine whether such deposited fines are stored on the bed surface, resuspended<sup>34</sup>, or infiltrated into the bed<sup>11</sup>. Pressure gradients at the bed sediment-water interface<sup>35,36</sup> may promote suspended sediment advection into the porous matrix via downwelling and upwelling mechanisms<sup>23,33,37,38</sup>. Such pressure gradients are induced by bedforms<sup>35,39</sup>, local heterogeneity or roughness and permeability changes in the streambed<sup>40</sup>, including those created by biotic activity. Fines infiltrating into the streambed may settle within the pore spaces of the sediment matrix<sup>41</sup>, or be transported laterally and longitudinally through the streambed<sup>42</sup>. In the upper bed layers, such horizontal transport ('Brinkman load' in Figure 2) may result from turbulent mixing promoted by shear instability above the bed<sup>5,43,44</sup>. In the lower layers, subsurface (Darcian) flow driven by hydraulic gradients<sup>45</sup> may transport fines through the interstitial spaces of the sediment matrix<sup>44</sup> ('interstitial load' in Figure 2).




Figure 2 - Fine sediment transport (black arrows) and deposition (grey arrows) processes in gravel beds.

These processes occur in three key load regions (suspended and dissolved load, Brinkman load and interstitial load) within the water and sediment columns in which such processes occur (left). The model proposes distinct modes of sediment transport in these three regions identified by transitions in the velocity profile  $V(y)$  (right). Modified from<sup>44</sup>.

#### Controls on fine sediment: Physico-chemistry

The physico-chemical properties of fines are typically considered to be secondary in importance to their physical characteristics in terms of generating ecological impacts, yet there are clear feedbacks between the sediment, its chemical constituents and those of the water-column, and biotic activity. Indeed, the chemical composition of fine sediment may be more influential on invertebrate assemblage structure<sup>3,46,47</sup> and the survival of fish eggs<sup>48,49</sup> than the volume of fines deposited on the benthos<sup>50</sup>. However, it is generally acknowledged that combined multiple stressor effects are complex and poorly understood<sup>51,52</sup>. Some of the better understood feedbacks between the physico-chemical properties of fines and their dynamics are related to biological processes. For example, the scattering and prevention of solar radiation by turbid conditions caused by excess suspended sediment is known to be detrimental to benthic ecology by reducing rates of photosynthesis<sup>53,54</sup>. Similarly, high suspended sediment concentrations (SSC) drive temperature gradients with depth in the water column<sup>55</sup>, with consequences for biotic activity. Another well-studied impact on the benthic community is a reduction in hyporheic exchange flow due to the deposition of excess fine sediment<sup>11,56</sup>. Within the interstices, the breakdown of fine particulate organic matter (FPOM) increases sediment oxygen demand, exacerbating the hypoxia initiated by physical clogging<sup>49,57-60</sup>. Under such conditions, methanogenic microbial communities develop at the oxic-anoxic interface, resulting in the precipitation of ochreous masses of iron hydroxides<sup>61</sup> and further saturating the bed matrix with fine particulate matter.

Many of the direct physico-chemical impacts of fines on stream biota are ecotoxicological in nature, driving feedbacks by increasing or decreasing the biological controls exerted on fine sediment dynamics

(Figure 1). Nutrients, heavy metals and organic pollutants can adsorb to and desorb from fine particles due to their high affinity as carriers<sup>25,62,63</sup>. Their sorption potential depends upon the properties of the sediment, i.e. grain size, mineralogical composition, organic matter content and cation exchange capacity<sup>64</sup>. The physico-chemical weathering of sediments during their 'life-cycle' is driven by water column concentrations of major ions and trace chemical constituents<sup>65</sup>, in addition to biological and hydrodynamic processing and sorting. However, additional physico-chemical factors, such as dissolved oxygen, pH and temperature have also been implicated in controlling the sorption dynamics of potentially harmful substances, including heavy metals<sup>66</sup> and nutrients<sup>67</sup>.

The desorption of these substances may increase bioavailability, allowing contaminants to enter the food web along with those contaminants adsorbed to fines that are ingested by biota. For example, the exchange of various phosphorus (P) forms between the particulate (bound) and dissolved (free) pools within freshwaters is now well-documented to be influenced by pH, temperature and dissolved oxygen<sup>68</sup>. These interactions can either promote or inhibit growth depending on the stoichiometric abundance of nutrients or the toxicity of contaminants<sup>69–71</sup>. Other physico-chemical factors affecting fines may disrupt their transport and deposition mechanics. An increase in the deposition of fines due to flocculation and precipitation has been seen due to modification of pH and ionic salt concentrations in freshwaters<sup>72–74</sup>. The most widely known examples are changes in redox potential leading to the deposition of calcium salts (tufa) and ocherous masses. Temperature regulated **stream**water viscosity is also known to increase particulate deposition<sup>75</sup>. These physico-chemical feedbacks between fines, the water-column and biotic activity have complex, cascading effects on geomorphic processes.

### **BIOLOGICAL-BIOTIC CONTROLS ON FINE SEDIMENT TRANSPORT AND STORAGE**

The two-way interactions between biota and the environment have long been of interest to both physical and biological scientists. However, it was not until recent decades that they were more intensively investigated under a multitude of frameworks, including zoogeomorphology, ecogeomorphology, biogeomorphology and ecosystem engineering<sup>76–78</sup>. Below we briefly review the body of research that has emerged from application of these frameworks to biofilms, aquatic macrophytes and riparian vegetation, invertebrates and fish. In doing so, we highlight the key *effect* traits<sup>79</sup> that determine the potential of biota to influence fine sediment dynamics (Table 1).

#### *Controls on fine sediment: Biofilms*

Biofilms, referred to as 'cities of microbes'<sup>80,81</sup> comprising bacteria, fungi and algae<sup>82</sup>, are known to affect and be affected by fine sediment dynamics<sup>83–86</sup>. Although the important function biofilms fulfil in wastewater processing has long been known, it was not until more recently that research began to focus on their role in the processing and fate of fine sediment-bound contaminants in freshwater ecosystems<sup>87–91</sup>. The matrix of the biofilm microenvironment is comprised mostly of **extracellular polymeric substances (EPS)EPS**<sup>92,93</sup>, which are synthesised by the biofilm microbial community. The properties of these EPS are influential on the sediments upon which they establish, altering their physico-chemical characteristics and, therefore, their dynamics through bioflocculation<sup>94</sup>. An experimental study<sup>95</sup> found that the erodibility and resulting resuspension of settled cohesive sediments was reduced with weak biofilm EPS

growth when exposed to low shear stress (<0.1 Pa). Longer periods (up to 12 days) of undisturbed growth under low shear stresses (<0.1 Pa) allowed the establishment of a biofilm-EPS structure that could reduce erosion under increased shear stresses of up to 0.4 Pa. These results, along with other studies in agreement<sup>85,96,97</sup>, demonstrate that biofilm growth and the corresponding EPS synthesis allows for increased cohesion between fine sediment grains (Figure 3). This potentially increases the deposition of fines and reduces their resuspension<sup>98</sup>.




Figure 3 - Controls on fine sediment deposition and resuspension exerted through bioflocculation by extracellular polymeric substances (EPS) in biofilms.

Fine sediments also interact with sediment-water exchanges of nutrients associated with biofilm communities<sup>27</sup>, typically increasing community growth and EPS synthesis. During a flume study<sup>87</sup>, concentrations of soluble reactive phosphorus (SRP) were reduced through the trapping of fine particulates in biofilms. Bed sediments with larger surface areas for biofilm and filamentous algal growth resulted in a greater flux of SRP than fine sediments alone, which are unable to support filamentous growth. Furthermore, evidence that nutrient stoichiometry is influenced by fine sediment dynamics has been linked to changes in biofilm growth and species composition<sup>99,100</sup>; although it is acknowledged that physico-chemical factors also play a role<sup>101–103</sup>. The growth of biofilms has also been found to influence fine sediment transport and bed morphology<sup>85,104</sup>. Modelling following experimental observations has

1  
2  
3 demonstrated that biofilm-coated fines have a greater saltation length-height ratio, suggesting that  
4 particle deposition and settling is more likely than in uncoated fines, depending on water velocity<sup>105</sup>. More  
5 recently, a microbiological-physical model (BFLOC2) of suspended sediment aggregation and settling  
6 velocity was established<sup>106</sup>. The model shows that microbial biomass, cell motility and aggregate-attached  
7 food web interactions are significant controls on fine sediment dynamics. In summary, the interactions  
8 between biofilms and fine sediments are driven by a balance between physico-chemical and biotic  
9 influences on the processing, transport and fate of particles.  
10  
11

### 12 *Controls on fine sediment: Macrophytes and riparian vegetation*

13  
14  
15 Awareness of the intimate links between vegetation and river landforms and processes emerged during the  
16 last century<sup>107–109</sup>. However, recognition of the complex interactions between riparian vegetation and  
17 fluvial processes that result in sediment retention and landform building has only emerged strongly in  
18 the last two decades<sup>110–112</sup>. It is now recognised that particular plant types, materials and species are key  
19 to these interactions, driving meso- to macro- scale sediment retention and building landforms.  
20 Examples of the broad types of plants and plant-derived material include emergent<sup>113,114</sup> and  
21 submerged<sup>115</sup> aquatic macrophytes; dead<sup>116</sup> and living deposited trees<sup>117,118</sup>, tree fragments<sup>119</sup> and  
22 accumulations of fragments<sup>120</sup>.  
23  
24  
25

26 Interactions between plants and any fine sediment transported by rivers are particularly important  
27 because small sediment particles are easily mobilised, even at low flow velocities, and so their storage  
28 relies on them being retained in sheltered locations. The above-ground biomass of plants offers such  
29 shelter to exert control on fine sediment retention. The extent to which plants drive such retention is  
30 dependent on their ability to affect local hydraulic conditions, and hence their morphology and the  
31 stiffness and density of their stems<sup>9</sup>. The greater the resistance the plant presents to flow, the greater  
32 the retention of sediment<sup>121,122</sup>. Once a critical flow resistance (frontal area drag coefficient) has been  
33 achieved, plants will tend to retain sediment<sup>123</sup>. Aquatic plants have the ability to grow through the  
34 retained sediment, reinforcing it with their stems, roots, rhizomes and other below-ground organs and  
35 driving the further accumulation of fine sediment into erosion-resistant, aggrading landforms such as  
36 bars, benches, river banks, islands and floodplains<sup>124</sup>. In lower energy river systems, aquatic  
37 macrophytes may be more important as ecosystem engineers than riparian plants<sup>9,121,125</sup>. For example,  
38 the emergent macrophyte *Sparganium erectum* is a very effective river ecosystem engineer in British  
39 lowland rivers<sup>126</sup>. A beneficial effect of sediment retention by plants can be that, as the fine sediment  
40 landforms emerge from the ~~river bedstreambed~~, they narrow the width of river channel available for  
41 water, leading to increased flow velocities in the area occupied by water and thus increased transport  
42 and (re)mobilisation of fine sediment from any coarser bed material. The result is a more heterogeneous  
43 ~~river bedstreambed~~ supporting patches of contrasting particle size. Conversely, extreme macrophyte  
44 growth in dense stands may reduce flow velocities throughout the river channel, causing [more sediment](#)  
45 [retention and](#) homogenisation of the physical habitat<sup>127</sup>. [This process could eventually lead to infilling of](#)  
46 [the channel. Indeed, it could be argued that such circumstances often arise as a response to the artificial](#)  
47 [creation or enlargement of stream channels, flow regulation and/or excessive inputs of fine sediment.](#)  
48 [Whatever the cause, available stream power is insufficient to control macrophyte growth and flush fine](#)  
49 [sediments](#)<sup>114,128</sup>. ~~Regardless of the cause~~[In any case, the fine sediments retained by macrophytes](#)  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

frequently display a high organic content<sup>129</sup>, illustrating that this interaction between plants, fluvial processes and river morphology may have a notable role in organic matter retention. Such effects are likely to be more pronounced in nutrient-enriched waters.

As river energy increases, macrophytes are unable to resist the increasing shear stresses imposed by river flows, and their engineering role is taken up by riparian plants, particularly trees. All tree species growing along river margins interact with fluvial forms and processes through a 'large wood cycle'. Tree or branch fall introduces large pieces of wood into rivers, where they can snag and accumulate into wood jams or dams, providing retention structures for other organic material, sediments and seeds<sup>116</sup>. In unmanaged situations, these processes drive a cycle of sediment and seed retention, germination and growth, and incorporation of the aggrading vegetated landforms into islands and floodplains where they support the growth of trees and thus the delivery of more large wood to the river<sup>130</sup>. Across the temperate zone of the Northern Hemisphere, the Salicaceae (willows and poplars) are particularly important river ecosystem engineers<sup>131</sup>. Riparian species of willow and poplar not only drive a 'dead' wood cycle, as described above, but the ability of wood from these species to sprout and grow remarkably rapidly, makes them particularly effective river ecosystem engineers. Furthermore, these species can produce roots from buried stems and shoots from roots allowing them to strongly reinforce sediments to depths of several metres<sup>132–134</sup>.

Recent reviews<sup>124,135</sup> conceptualise the crucial interactions between plants and fluvial processes that influence sediment retention and reinforcement, the building of landforms such as scroll bars<sup>136</sup>, islands, river banks and floodplains that are the result of these interactions, and the ways in which these processes and forms vary through time and across space. During the storage period in these landforms, the properties of fine sediments may change under the influence of biogeochemical processes and the activities other biota that colonise the new landforms. This could include invasive species, such as the tall growing annual Himalayan balsam (*Impatiens glandulifera*) which may increase the risk of bank erosion during the winter period<sup>137,138</sup>. The controls on fine sediment transport and storage exerted by plants, therefore, may be shifting as invasive species become established.

#### *Controls on fine sediment: Invertebrates*

Lotic invertebrates are key ecosystem engineers, being ubiquitous in rivers and often present at considerable densities. For this reason, despite their typically small body sizes (e.g. <50 mm), invertebrates can have substantial cumulative impacts on the storage and transport of fines<sup>139</sup>. Crustaceans are perhaps the most widely studied geomorphic agents in rivers and can influence all aspects of fine sediment dynamics<sup>140</sup>. Crayfish can destabilise river bedstreambeds during activities such as foraging and fighting, and a number of studies have documented an increase in turbidity attributed to bioturbation by crayfish<sup>140,141</sup>. High densities of invasive The presence of crayfish can result in a 50–75% reduction in the shear stress required to entrain sand, leading to increases in sediment fluxes of as much as 32%<sup>142</sup>. Invasive signal crayfish (*Pacifastacus leniusculus*) can also enhance the delivery of fine sediment to rivers through increasing bank erosion with high densities of burrows (up to 14 burrows m<sup>-1</sup>)<sup>142,143</sup>.

1  
2  
3 Aquatic insects can increase the suspension of fine sediment by winnowing it from interstitial spaces  
4 during foraging activity<sup>144,145</sup>. The stonefly (*Dinocras cephalotes*) has the potential to erode 200-400 kg  
5 m<sup>-2</sup> yr<sup>-1</sup> of sand, with reduced prey availability increasing rates of sand erosion<sup>146</sup>. The mayfly species  
6 *Pseudiron centralis* manipulates near-bed hydraulics to disturb sand and facilitate capture of its prey<sup>147</sup>.  
7 This latter example demonstrates that some invertebrates are actively seeking to control sediment  
8 dynamics, an instance of extended phenotype engineering, whereby organisms create structures that  
9 directly influence their fitness and survival<sup>78</sup>. Invertebrate bioturbation has also been documented to  
10 influence sediment structure. Upward conveyors (Oligochaeta) and gallery-diffusers (Chironomidae) can  
11 reduce the clogging of ~~bed-sediments~~streambeds, helping to restore hyporheic exchange flow and  
12 physiochemical conditions at the surface-subsurface interface<sup>148-153</sup>. Molluscs have been reported to  
13 both disturb sediment during movement and consolidate it whilst stationary<sup>154</sup>. Flow resistance  
14 generated by the shells of bivalve molluscs can also promote localised scour<sup>155</sup>. Different modes of  
15 bioturbation (sediment reworking, biogenic structure building, burrowing depth, bioirrigation) can also  
16 determine the impact of benthic invertebrates on microbial activities and biogeochemical processes in  
17 the sediment<sup>156</sup>.

23 Hydropsychidae is one of the most abundant families of lotic insects worldwide, often accounting for as  
24 much as 80% of invertebrate biomass in some streams and, consequently, exerting substantial controls  
25 on geomorphic processes<sup>157-159</sup>. Hydropsychids construct nets to filter FPOM from the water column,  
26 increasing the force required to entrain bed sediments by 10-30%<sup>142,160</sup>. This effect increases with the  
27 density of individuals and the local richness of species exhibiting similar behaviours<sup>161</sup>. Through net-  
28 spinning, hydropsychid caddisflies increase the force required to erode armour layers and therefore  
29 reduce exposure of underlying fine sediments to entraining flows<sup>162</sup>. Many other species of caddisfly are  
30 also known for their construction of cases from mineral and organic sediment. These cases utilise a wide  
31 range of particle sizes and may be responsible for the storage of significant quantities of temporary  
32 storage of fine sediment<sup>163,164</sup>. Blackflies (Simuliidae), bivalve molluscs and other filter feeding  
33 invertebrates are also involved in driving storage of fine sediment by consolidating suspended particles  
34 from the water into faeces or pseudofaeces, thus increasing the rate of sedimentation and altering the  
35 composition of suspended solids<sup>165,166</sup>.

41 The sheer diversity of aquatic invertebrate species (85% of global freshwater animal diversity<sup>167</sup>), and  
42 the densities in which they are often present, complicate understanding of their role in the transport of  
43 fines. In particular, interactions between coexisting species probably play an important role in mediating  
44 the impacts of excess fine sediment, yet these interactions are rarely considered<sup>161</sup>. It should also be  
45 noted that the effect of biota on fine sediment dynamics will vary spatially and temporally. ~~Four main~~  
46 ~~factors have been suggested as primary controls on the impact that animals have on the environment;~~  
47 ~~behaviour, body size, density and abiotic context~~<sup>165,166</sup>. Despite management advances made in recent  
48 decades, especially in relation to organic pollution, freshwater biota face increasing challenges for  
49 survival, including habitat degradation, species invasions, pollution, climate change, and increasing  
50 urbanisation<sup>168,169</sup>. Invertebrates are one of the most affected taxonomic groups<sup>170,171</sup> and this could  
51 have consequences, not just for biodiversity, but also fine sediment dynamics in the future.

56 *Controls on fine sediment: Fish*

1  
2  
3 Fish can exert controls on fine sediment dynamics through a number of activities. Relative to other  
4 taxonomic groups, the controls exerted by fish are well-documented, most notably feeding and  
5 spawning<sup>140</sup>. We therefore provide only a brief summary here. To date, salmonid spawning has  
6 attracted the most attention. During spawning, female salmon create redds (nests) by disturbing the  
7 bed with strong undulations of their tails<sup>172</sup>. This activity leads to the suspension of fines and modifies  
8 bed stability by vertically mixing gravels and fine sediments and disturbing existing armour layers<sup>173–175</sup>.  
9 Tail slips (material excavated from redds) are particularly vulnerable to scour<sup>175,176</sup>. Spawning salmon  
10 substantially modify topography across large areas of the bed<sup>177</sup>. This effect persists until the next high  
11 flow event of sufficient magnitude<sup>178</sup>, exerting a strong, temporary control on the hydraulic processes  
12 driving fine sediment dynamics.  
13  
14  
15

16  
17 Put into perspective, the spawning activities of salmon and other lithophilous fish can have considerable  
18 geomorphic impacts at large spatial scales. The vertical mixing of material by salmon may be at a  
19 comparable magnitude to that driven by floods, resulting in lower critical shear stresses required to  
20 initiate bed movement<sup>179</sup>. In some cases, salmon can be responsible for a substantial proportion of  
21 annual sediment transport. A study in British Columbia ~~study~~ estimated that salmon account for one  
22 third to one half of total annual sediment flux in gravel bed streams<sup>178</sup>. So strong is this effect that the  
23 ~~evolution~~ evolution of freshwater fish may of multiple Pacific salmon species from a common ancestor  
24 may have consequences for landscape evolution over geological timescales profoundly influence channel  
25 erosion processes and stream profiles over geological timescales, as suggested by a coupled biological-  
26 landscape evolution model focusing on the descent of Pacific salmon species from a common  
27 ancestor<sup>180</sup>.  
28  
29  
30

31  
32 Benthivorous fish also have significant impacts on fine sediment suspension while foraging. Cyprinid  
33 species can substantially alter sediment dynamics, increasing mixing depth and suspended  
34 sedimentSSC<sup>181,182</sup>. Foraging barbel, for example, can increase total sediment yields by as much as  
35 82%<sup>183</sup>. The geomorphic impacts of benthivorous foraging increase with fish density<sup>184–187</sup>, with the body  
36 size and mass of individuals<sup>188</sup>, and with the presence of specific feeding behaviours. European barbel  
37 (*Barbus barbus*), for example, feed in both gravel and sand sized substrates, while gudgeon (*Gobio*  
38 *gobio*) feed largely in sand habitats and, therefore, their geomorphic impacts are spatially limited<sup>189</sup>. Fish  
39 can also exert controls on fine sediment dynamics indirectly through their influence on other  
40 biogeomorphic agents. For example, bed disturbance and predation by fish may reduce the controls  
41 exerted by net-spinning caddisflies, macrophytes and algae<sup>189</sup>. Moreover, fish can have considerable  
42 impacts on the biogeochemical processes in rivers. For example, spawning migrations of anadromous  
43 fish (e.g. *Oncorhynchus* spp.) may be a potential important source of marine-derived nitrogen  
44 and phosphorus to nutrient-poor habitats<sup>190,191</sup>, profoundly influencing the structure and function of  
45 further influencing aquatic and riparian biota aquatic and terrestrial ecosystems<sup>185</sup>.  
46  
47  
48  
49  
50

## 51 **CONTROLS ON ECOLOGICAL RESPONSES TO FINE SEDIMENT**

52  
53 Whilst the impacts of excess fines on biota can be pervasive<sup>7–10</sup> not all communities will be equally  
54 sensitive. In particular, the functional traits of the local biotic community, as well as those of the species  
55 comprising the regional species pool, will constrain the ecological response. Response traits determine  
56  
57  
58  
59  
60

an individual's ability to survive in different environmental conditions through characteristics that promote resistance and resilience to disturbances. Typically, such traits include information on morphology, phenology, behaviour and resource use<sup>192,193</sup>. Below we identify some of the major traits that are commonly associated with responses to fines (Table 1) and put them into context with biogeographic processes.


Table 1 - Traits involved in the effect of biota on fine sediment dynamics and the response of biota to excess fine sediment.

<b>Taxonomic group</b>	<b>Effect traits</b>	<b>Response traits</b>
<i>All biota</i>		- Dispersal capacity
<i>Biofilms</i>	<ul style="list-style-type: none"> <li>- Extracellular polymeric substance (EPS) production</li> <li>- Colony structure</li> <li>- Growth rate</li> </ul>	<ul style="list-style-type: none"> <li>- Colony structure</li> <li>- Motility</li> <li>- Size and shape</li> <li>- Cell wall structure (rigidity)</li> <li>- Photosynthetic capacity</li> </ul>
<i>Plants</i>	<ul style="list-style-type: none"> <li>- Shoot density</li> <li>- Shoot and root structure</li> <li>- Shoot flexibility</li> </ul>	<ul style="list-style-type: none"> <li>- Fecundity</li> <li>- Asexual (vegetative) reproduction</li> <li>- Dispersal mode (hydrochory, anemochory)</li> <li>- Reproductive timing</li> <li>- Growth rate</li> <li>- Shoot flexibility in high-flow periods</li> <li>- Root structure and binding capacity</li> </ul>
<i>Invertebrates</i>	<ul style="list-style-type: none"> <li>- Feeding behaviour</li> <li>- Burrowing activity</li> <li>- Construction of feeding and protective structures (e.g. nets, cases)</li> <li>- Body size</li> </ul>	<ul style="list-style-type: none"> <li>- Feeding mode and diet</li> <li>- Body size and shape</li> <li>- Respiration mode</li> <li>- Locomotion (e.g. burrowing, crawling)</li> <li>- Reproduction mode (e.g. parental care, oviviparity)</li> <li>- Reproductive timing</li> <li>- Voltinism (number of generations per year)</li> </ul>

<i>Fish</i>	<ul style="list-style-type: none"> <li>- Foraging behaviour</li> <li>- Diet</li> <li>- Spawning behaviour</li> </ul>	<ul style="list-style-type: none"> <li>- Trophic guild</li> <li>- Foraging behaviour</li> <li>- Gill morphology and physiology (e.g. increased mucus secretion, gill epithelium thickness)</li> <li>- Mechanosensory system development (e.g. vision, lateral line)</li> <li>- Reproductive timing</li> <li>- Spawning behaviour</li> </ul>
-------------	--	---

### Controls on ecological responses: Biogeography

In rivers, the spatial distribution of organisms is highly dependent on the dendritic network structure<sup>194</sup>. Thus, at a broad level, ecological responses to fine sediment depend on the location of the site under consideration, as well as the topology and connectivity of the whole network<sup>195</sup>. The responses will also depend on the spatial structure of populations present within the network; if a population can be divided into subpopulations connected via dispersal, we may call it a 'metapopulation'<sup>196</sup>. 'Metacommunity' theory extends this idea to the community level by incorporating the effects of species sorting ('environmental filtering'<sup>197</sup>) and biotic interactions<sup>198</sup>. The theory, as applied to river networks, predicts systematic variation in the forces driving community assembly with stream order (Figure 4). All else being equal, the impacts of stressors such as fine sediment are predicted to reduce alpha diversity in headwaters, where colonisation of adapted-species and phenotypes adapted to stress caused by excess fine sediment is limited. Mid-basin locations, on the other hand, are expected to exhibit high rates of turnover in species composition, as efficient species sorting leads to rapid selection of adapted taxa from the species pool. Further downstream, the effects of the stressor are predicted to be dampened by the influx of organisms from upstream, to some extent regardless of the environment. Superimposed onto this trend is a natural tendency for the importance of fine particulate matter as a basal resource to increase downstream through the river network<sup>199</sup>. Fine sediment storage dynamics in high order river-floodplain systems are also tightly linked to flood pulses that are relatively predictable compared to events in low order streams<sup>200</sup>. These spatial patterns are further complicated by the discontinuities in fluxes of matter and energy caused by riverine barriers such as dams<sup>201</sup>.



1  
2  
3 Figure 4 - Metacommunity theory predicts patterns in dominant community assembly processes in river  
4 networks (a) due to variation in spatial, environmental and demographic influences with stream order  
5 (b), with implications for ecosystem responses to stressors. Headwater communities are likely to follow  
6 the patch dynamics paradigm, whereby local species diversity is limited by dispersal and community  
7 structure becomes a function of stochastic demographic processes involving local extinction and  
8 colonisation<sup>198</sup>. The structure of lower mainstem communities is predicted to more closely resemble the  
9 regional species pool since high dispersal rates drown out the influence of the local environment  
10 through 'mass effects'<sup>198</sup>. In mid-basin locations, species sorting or 'environmental filtering'<sup>197</sup> is made  
11 possible as the influence of dispersal is at an optimal level to allow for efficient niche-based  
12 processes<sup>202</sup>.  
13  
14  
15  
16  
17  
18

19 Whilst metacommunity theory provides a useful lens with which to study ecological responses to  
20 stressors, there are other regional-scale factors involved. Among these are biogeographical factors  
21 linked to the size of the species pool<sup>203</sup>. Relative to the number of taxa found at a given site, larger  
22 species pools are likely to contain species better adapted to a given stressor. Sites that are more  
23 saturated with respect to the species pool, on the other hand, will have less potential for adaptation. In  
24 the former case, high rates of turnover would be expected, whereas in the saturated case greater  
25 reductions in alpha (local) diversity are likely to ensue. These tendencies are further dependent on the  
26 traits expressed by taxa in the local assemblage and the wider species pool. In particular, the dispersal  
27 capacities of the regional fauna will, to a large extent, influence the relationship between the species  
28 pool, the local assemblage and stressor-driven changes<sup>204</sup>. In general, a weaker dispersing taxon will be  
29 less likely to colonise a newly suitable habitat patch after some environmental change, the likelihood  
30 also being contingent on the distance of established populations of that taxon from the impacted site<sup>205</sup>.  
31 Species pools are thus best represented as species- and site- specific probabilistic functions<sup>206</sup>. We  
32 consider other important traits controlling ecological responses below.  
33  
34  
35  
36  
37

#### 38 *Controls on ecological responses: Biofilm traits*

39  
40 Biofilms comprise a mixture of heterotrophic (e.g. fungi, bacteria, protozoa) and autotrophic (e.g.  
41 cyanobacteria, algae, chemosynthetic bacteria) organisms, with many of the latter highly dependent on  
42 light. As such, the attenuation of light caused by increased sediment loads (either suspended or  
43 deposited) adversely affects the photosynthetic component of biofilms<sup>207</sup>, shifting the balance of these  
44 two traits, and thus the net primary production of affected biofilms<sup>208–211</sup>. For the sessile photosynthetic  
45 components, the absence of light caused by burial beneath depositing sediment can be catastrophic and  
46 the only option can be to produce resting stages to endure the period until erosion brings them to the  
47 surface once more. However, for motile taxa (e.g. raphid diatoms, ciliates, flagellates) shading from  
48 deposited fine sediment may not present a substantial problem, as they can move to higher light  
49 intensities at the streambed surface<sup>212,213</sup>. Hence, diatom assemblages tend to become dominated by  
50 motile taxa where rates of deposition of fine sediments are high<sup>214–216</sup>, although this trait (motility) may  
51 offer benefits under other conditions (e.g. nutrient rich conditions) such that the relationship with fine  
52 sediment is not straightforward<sup>217</sup>.  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 The fact that deposited fine sediments are relatively unstable (compared with larger particles) renders  
4 them unsuitable for the attachment of long-lived sedentary species, particularly slow-growing and  
5 chain-forming taxa, pushing assemblages towards rapidly growing, single celled taxa. Whilst the lack of  
6 stability tends to result in reduced diatom taxon richness and biomass compared with more stable  
7 sediments<sup>218,219</sup>, patch disturbance history has a large influence on the development of biofilm  
8 communities<sup>220</sup>. In turn, those taxa that exude mucilage are favoured, at least initially, as they tend to  
9 stabilise sediments (see discussion on EPS above).  
10  
11

12  
13 The risk of being dislodged<sup>221</sup> or **suffering-undergoing** physical damage<sup>222</sup> from suspended and  
14 particularly saltating sediment particles further selects against tall taxa (chain forming, filamentous,  
15 stalked and upright forms), and pushes communities towards adpressed forms and those that strongly  
16 adhere to the substrate (e.g. with mucilage pads), influencing traits associated with both growth form  
17 and attachment. More robust cell walls (e.g. thick walls, heavy silicification, costae) enable taxa to  
18 withstand physical damage. As such, species with thicker, more rigid cells walls are selected for where  
19 the suspended and saltating loads of inorganic sediment are high<sup>223</sup>. However, even these species are  
20 lost if disturbance is frequent<sup>224</sup>. Under more benign conditions, rapid growth rates may compensate for  
21 losses, with associated traits such as nutrient affinity being more apparent where deposited fine  
22 sediments are abundant<sup>217</sup>.  
23  
24  
25

#### 26 *Controls on ecological responses: Plant traits*

27  
28 River engineer plants possess important traits that allow them to establish and persist in riverine and  
29 riparian environments<sup>9,225</sup>. These traits include: (i) an ability to reproduce vegetatively, both to colonise  
30 new patches and to extend and consolidate patches once established; (ii) an ability to grow rapidly in  
31 order to maximise their chances of becoming established, and to cope with burial by deposited material;  
32 (iii) above-ground biomass with sufficient rigidity (either as individual shoots or collectively as stands)  
33 and stand density to present a resistance to flow and, thus, retain sediment, but avoids breakage under  
34 higher flows; and (iv) below-ground biomass with sufficient strength and appropriate architecture to  
35 anchor plants and resist uprooting (e.g. stolons, rhizomes).  
36  
37  
38  
39

40  
41 Reproduction and dispersal traits allow plant propagules to reach appropriate sites for germination and  
42 establishment. Both aquatic macrophyte and riparian tree species take advantage of sexual and asexual  
43 means to ensure reproductive success, with different reproductive strategies allowing successful annual  
44 recruitment at different locations. However, riparian and aquatic engineer species generally devote  
45 considerable resources to asexual reproduction, in particular through the production of adventitious  
46 roots from stem fragments, allowing them to expand their cover locally as well as colonising new  
47 areas<sup>226</sup>. Asexual reproduction by riparian tree species largely depends on the ability of uprooted trees  
48 and wood fragments to sprout both roots and shoots once deposited. Some aquatic macrophyte  
49 species, particularly those with emergent growth forms (e.g. *Sparganium erectum*, *Glyceria maxima*),  
50 produce dense networks of rhizomes or stolons that support the spread of patches, as well as dispersal  
51 if the plant is fragmented, whilst others regenerate freely from stem fragments<sup>227</sup>. The development of  
52 adventitious roots, stolons and rhizomes allows plants to cope with burial and to consolidate the  
53 deposited material (e.g. <sup>126</sup>). Many aquatic plants and riparian shrubs (e.g. Salicaceae) develop such  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 structures, which enter and reinforce aggrading sediments while maintaining reinforcement and erosion  
4 resistance in buried substrates (e.g. <sup>132</sup>). Furthermore, plants can exploit the nutrients available from  
5 deposited material by rooting into it: much of the organic material deposited in stands of *Ranunculus*  
6 *penicillatus* is remineralised and used for growth of the plant, further accentuating deposition<sup>228</sup>, a  
7 characteristic likely to be true of other species.  
8  
9

10 Once deposited, propagules of engineer plants germinate or sprout and typically grow rapidly to anchor  
11 themselves and support their survival in disturbed river environments. Rapid shoot and root growth is  
12 particularly important for species both to exploit opportunities for growth before disturbance removes  
13 them and to cope with burial. For example, field observations along the Tagliamento River, Italy, have  
14 recorded main shoot growth in *Populus nigra*, *Salix alba* and *Salix eleagnos* seedlings, cuttings and from  
15 uprooted, deposited trees of up to 3 mm day<sup>-1</sup>, 10 mm day<sup>-1</sup> and 15 mm day<sup>-1</sup>, respectively<sup>229,230</sup>.  
16 Furthermore, greenhouse experiments on cuttings of *Salix eleagnos* and *Populus nigra* have revealed  
17 vertical root penetration of 27 and 15 mm day<sup>-1</sup> respectively for sand substrates, and 20 and 10 mm day  
18 <sup>1</sup>, respectively for gravel substrates under a water table falling at 30 mm day<sup>-1</sup> <sup>231</sup>. Many aquatic plants  
19 avoid disturbance during winter and spring high flows by producing dense networks of rhizomes that  
20 persist through the winter, anchoring the plants and reinforcing penetrated sediments, whilst following  
21 an annual above-ground growth cycle whereby shoots emerge in the spring. Peak biomass is reached in  
22 mid to late summer and then senesce during autumn, leaving little (if any) above-ground biomass  
23 exposed during winter high flow events (e.g. <sup>126,232,233</sup>). During the growth period, species succession in  
24 aquatic plant stands is closely related to the ability to outpace the rate of deposition, favouring species  
25 with fast growth rates (including the ability to exploit nutrients from deposited material) and the  
26 capacity to adopt an emergent strategy<sup>234,235</sup>.  
27  
28  
29  
30  
31  
32

33 As stated above, the extent to which plants retain sediment is related to the resistance they present to  
34 the flow of water. However, this resistance coupled with the instability of deposited sediment increases  
35 the likelihood of plants being uprooted, particularly during flood conditions. The prevalence of traits  
36 associated with rapid growth, either to cope with burial, patch instability or to exploit nutrients from  
37 deposited material, together with a high dependency on vegetative growth, leads to a plant trait  
38 syndrome similar to that seen in highly competitive taxa typical of nutrient rich conditions<sup>236</sup>. Hence, it is  
39 not surprising that taxa indicative of high nutrient conditions thrive in rivers with large amounts of  
40 deposited sediment<sup>9,237</sup>.  
41  
42  
43

#### 44 *Controls on ecological responses: Invertebrate traits*

45

46 Excess fine sediment is widely acknowledged to have deleterious effects on the structure and function  
47 of invertebrate communities<sup>8,54,60,238</sup>. Substrates with a high volume of fines typically support  
48 homogenous communities that are dominated by relatively few taxa<sup>47,239–241</sup>. The composition of  
49 invertebrate taxa present in different environmental settings is strongly constrained by the functional  
50 traits they possess<sup>242</sup>, resulting in variable tolerance or sensitivity to excess fines<sup>243,244</sup>.  
51  
52

53 High sediment loads reduce the quality and availability of trophic resources which may reduce feeding  
54 efficiency, most notably for shredders<sup>244–247</sup>. Other feeding modes, including algal scrapers and filter  
55  
56  
57  
58  
59  
60

1  
2  
3 feeders, also demonstrate some sensitivity to fine sediment<sup>248–250</sup> but the effect lacks consistency,  
4 suggesting that other factors interact with these traits to determine the response. Increasing infiltration  
5 of fines fills interstitial pore spaces and reduces dissolved oxygen concentrations, limiting the ability of  
6 many taxa to persist due to their body size, shape and respiratory requirements<sup>251,252</sup>. As such,  
7 streambeds subjected to high levels of deposition and infiltration are frequently characterised by taxa  
8 with small body sizes and lower densities of interstitial dwellers<sup>60,246,253,254</sup>.

11 Locomotion traits may also be implicated in the impacts of fine sediment deposition on invertebrate  
12 communities. Burrowing taxa associated with depositional habitats are well adapted to fine sediment,  
13 whilst crawlers may be adversely affected during deposition events due to their reduced locomotive  
14 capacity<sup>251,255–257</sup>. Species with certain respiration modes can also be highly sensitive to fine sediment  
15 transported in the suspended or saltating load. Delicate gill structures, for example, may become  
16 physically abraded or clogged by fine sediment, limiting the exchange of oxygen<sup>258,259</sup>. Reproduction  
17 mode may also exert a strong control over which taxa are able to persist in substrates with high fine  
18 sediment content. Unattended eggs deposited onto the stream bed, for instance, may become  
19 smothered or abraded by fine sediment<sup>260</sup>. As a result, the prevalence of ovoviviparity, [as found in](#)  
22 [diverse taxa \(e.g. \*Sphaerium spp.\*, \*Asellus aquaticus\*, \*Cloëon spp.\*\)](#), may increase in streams subjected to  
23 high sediment loads<sup>246,261,262</sup>. Voltinism also influences a taxon's ability to recover from disturbances  
24 such as those wrought by excess fine sediment, with multivoltine taxa recovering more rapidly  
25 compared to univoltine taxa<sup>257,263</sup>. This resilience to disturbance is associated with a trait syndrome that  
26 includes small body sizes and short life cycle lengths.

30 Despite trait-based ecology gaining increasing recognition for its ability to predict tolerance and  
31 sensitivity to a range of stressors, further research is required to strengthen the mechanistic basis  
32 behind the use of invertebrate traits in fine sediment-specific biomonitoring applications<sup>244</sup>. The  
33 consideration of traits should be evaluated with caution as a number of recent studies have reported  
34 inconsistent responses of invertebrates to fine sediment deposition<sup>47,246,254,257</sup>. Traits are unlikely to act  
35 in isolation but rather as combinations of traits describing life-history strategies of varying resistance  
36 and resilience to stressors such as fine sediment<sup>47,264</sup>. Furthermore, the timing of fine sediment events  
37 relative to a taxon's life-cycle will play an important role<sup>254</sup> as will a taxon's preferred habitat and  
38 substrate composition<sup>244</sup>.

#### 42 *Controls on ecological responses: Fish traits*

43  
44  
45 A range of fish traits have been implicated in community responses to fine sediment carried in  
46 suspension, in the saltating load, surface deposited or infiltrated<sup>7</sup>. Well-documented, direct negative  
47 impacts include the clogging and abrasion of soft tissues, especially gills<sup>265,266</sup>. Fish adaptations to  
48 enhance resistance to this include increased mucus secretion and a thickening of the gill epithelium<sup>267</sup>.  
49 However, under high SSC, mucus secretions can result in increased susceptibility to clogging of the gill  
50 surface and ultimately suffocation of the fish<sup>268</sup>. Further adaptations to living in turbid conditions are  
51 associated with the mechanosensory system, including development of the lateral line system and inner  
52 ear used to sense hydrodynamic cues, specialised reflective structures in the eye, and tactile and  
53 olfactory senses<sup>269–271</sup>.

1  
2  
3 By far the most well-known impact of deposited and infiltrated fine sediment is the smothering of  
4 salmon redds<sup>49</sup>. However, the majority of freshwater fish species are not lithophilous. Phytophilous  
5 species are also likely to be indirectly impacted due to reductions in submerged vegetation under high  
6 fine sediment loadings, whereas reproduction in psammophilous taxa may be favoured if deposition is  
7 associated mainly with the sand fraction. Some species avoid the potential impacts of fine sediment on  
8 eggs by ovipositing on riparian vegetation at high flows, with eggs developing out of the water until the  
9 next high flow (e.g. *Galaxias argentus*<sup>272</sup>). The timing of life-history events relative to periods of high fine  
10 sediment loadings can play an important role in fish responses to fines. In many temperate systems, for  
11 example, fine sediment accumulation in spawning gravels will likely be highest during the summer  
12 baseflow period<sup>54</sup>. Spawning outside of this period is therefore one way to avoid the most deleterious  
13 impacts on eggs and embryos<sup>7</sup>.

14  
15  
16  
17  
18 Herbivorous fish are indirectly impacted by reductions in submerged macrophyte abundance and  
19 changes in plant species composition due to light attenuation and abrasive forces of suspended and  
20 saltating sediment, as well as the deposition of fine material on plant surfaces, which reduces plant  
21 growth and nutritional quality<sup>9,273</sup>. This generates further impacts on other species by reducing or  
22 changing the nature of available cover. Zooplankton feeders may benefit from enhanced foraging  
23 efficiency under high SSC as their prey congregate near the water surface<sup>274</sup>, although the reduced  
24 visibility associated with such conditions is likely to result in net-negative impacts on feeding<sup>275</sup>.  
25 Similarly, high turbidity events may enhance prey availability due to increased invertebrate drift rates.  
26 However, foraging efficiency for drift-feeding fish decline under even mildly turbid conditions<sup>276</sup> and, in  
27 the long-term, invertebrate drift events in response to high SSC are likely to contribute to a change in  
28 invertebrate community composition which would negatively impact invertivorous fish<sup>277</sup>.

## 32 33 Conclusion

34  
35 The presence of excess fine sediment in rivers is a major reason for failure of EU member states to  
36 achieve Good Ecological Status under the Water Framework Directive. In this context, we have  
37 presented a broad synthesis of current understanding in the belief that this can help to drive the  
38 production of new scientific knowledge and inspire management innovation. In particular, we have  
39 highlighted biogeomorphic processes that control the transport and storage of fine sediment, and the  
40 mechanisms by which this elicits ecological responses and feedbacks. The fine sediment 'problem'<sup>11</sup>  
41 represents a major challenge for applied environmental science worldwide but progress is being made  
42 towards a more complete understanding of the physical and biological processes involved. This progress  
43 should be maintained and accelerated by continuing the open dialogue between management agencies  
44 and scientists working within the broad disciplines of ecology, hydraulic engineering, fluvial  
45 geomorphology and hydrochemistry. [In Sidebar 1, we briefly describe the outcomes of a workshop  
46 intended to address this need. Based on the experiences of environmental managers present, the  
47 workshop participants identified the most urgent knowledge gaps. These ranged from the collation of  
48 existing data and case studies to the investigation of responses to fine sediment at the population and  
49 whole ecosystem levels. In this way](#)Through initiatives such as this, key management challenges,  
50 scientific research priorities, and the production of new knowledge can all inform and shape one  
51 another [\(see sidebar\)](#).

### SIDEBAR 1 - SUPPORTING GOOD ECOLOGICAL STATUS THROUGH RESEARCH

A British Ecological Society-funded workshop in July 2017 brought together early career researchers working on fine sediment internationally along with UK-based environmental managers and senior scientists in the field to discuss how to support the achievement of GES through research. The discussions indicated that fine sediment-related failures to achieve GES in England and Wales are largely due to agricultural and rural land management, followed by urbanisation and transport. Measures to address the problem for point and diffuse sources exist but there are gaps in the evidence, including information on the risk of further deterioration, impacts on protected areas, and the interaction of fine sediment with other pressures. More catchment-scale data is needed to inform management, along with greater collation of existing case studies. Evidence is required for the effectiveness of alternative measures under future climate change and socio-economic scenarios. Further development and testing of pressure-specific biomonitoring metrics will help to link ecological changes to causal factors.

Specifically, environmental managers needed answers to the following questions:

- What are sediment yields and dynamics in rivers under different land use and management scenarios?
- How do we define 'natural' sediment conditions in aquatic ecosystems?
- How much do we need to reduce the sediment input by to see an ecological benefit?
- How can we evaluate the benefits of reducing excess fine sediment in rivers?
- What are the impacts of invasive species on sediment regimes? – especially plants such as Himalayan balsam and burrowing animals such as crayfish.
- Are there thresholds of fine sediment loadings beyond which ecological impacts are more severe?
- How can the risks of fine sediment impacts on sensitive species, such as the pearl mussels (*Margaritifera margaritifera*), be assessed most effectively?
- Which management measures are most cost-effective?
- What are the risks of dispersive dredging from contaminated sediments?
- What is the role of fine sediment dynamics in the transport of microplastics?

### Acknowledgments

This paper resulted from a workshop funded by the British Ecological Society (“Early careers, fine sediment and hydroecology: Supporting Good Ecological Status through research”, July 2017). We would like to thank early career scientists who participated in the workshop but did not contribute directly to

1  
2  
3 the preparation of this paper, namely Leticia Miguel (Environment Agency), Zarah Pattison (University of  
4 Stirling), Anna Lavelle and Eleanore Heasley (King's College London), James Smith (Loughborough  
5 University), Hazel Wilson (University of Nottingham), Matt Turley (University of Brighton), Tea Basic and  
6 Catherine Gutmann Roberts (Bournemouth University), and George Bunting (University of Worcester).  
7  
8

## 9 **Figures and Tables**

## 11 **References**

- 13 1. Collins AL, Anthony SG, Hawley J, Turner T. Predicting potential change in agricultural sediment  
14 inputs to rivers across England and Wales by 2015. *Mar Freshw Res* [Internet]. 2009 [cited 2018  
15 Jul 26];60(7):626–37. Available from: <http://www.publish.csiro.au/mf/mf08033>
- 17 2. Zhang Y, Collins AL, Murdoch N, Lee D, Naden PS. Cross sector contributions to river pollution in  
18 England and Wales: Updating waterbody scale information to support policy delivery for the  
19 Water Framework Directive. *Environ Sci Policy* [Internet]. 2014 Oct [cited 2018 Jul 4];42:16–32.  
20 Available from: <http://linkinghub.elsevier.com/retrieve/pii/S1462901114000823>
- 22 3. Naden PS, Murphy JF, Old GH, Newman J, Scarlett P, Harman M, et al. Understanding the controls  
23 on deposited fine sediment in the streams of agricultural catchments. *Sci Total Environ*  
24 [Internet]. 2016 Mar 15 [cited 2018 Jul 4];547:366–81. Available from:  
25 <https://www.sciencedirect.com/science/article/pii/S004896971531233X>
- 27 4. Walling DE, Collins AL. Fine sediment transport and management. In: *River science : research and*  
28 *management for the 21st Century* [Internet]. Wiley Blackwell; 2016 [cited 2018 Jul 4]. p. 37–60.  
29 Available from: [https://www.wiley.com/en-](https://www.wiley.com/en-gb/River+Science%3A+Research+and+Management+for+the+21st+Century-p-9781119994343)  
30 [gb/River+Science%3A+Research+and+Management+for+the+21st+Century-p-9781119994343](https://www.wiley.com/en-gb/River+Science%3A+Research+and+Management+for+the+21st+Century-p-9781119994343)
- 32 5. Boano F, Revelli R, Ridolfi L. Water and solute exchange through flat streambeds induced by large  
33 turbulent eddies. *J Hydrol* [Internet]. 2011 May 25 [cited 2018 Jul 30];402(3–4):290–6. Available  
34 from: <https://www.sciencedirect.com/science/article/pii/S0022169411002034>
- 36 6. Atkinson CL, Allen DC, Davis L, Nickerson ZL. Incorporating ecogeomorphic feedbacks to better  
37 understand resiliency in streams: A review and directions forward. *Geomorphology* [Internet].  
38 2017 [cited 2018 Jul 30]; Available from:  
39 <https://www.sciencedirect.com/science/article/pii/S0169555X17300090>
- 41 7. Kemp P, Sear D, Collins A, Naden P, Jones I. The impacts of fine sediment on riverine fish. *Hydrol*  
42 *Process* [Internet]. 2011 May 30 [cited 2018 Jul 30];25(11):1800–21. Available from:  
43 <http://doi.wiley.com/10.1002/hyp.7940>
- 44 8. Jones JI, Murphy JF, Collins AL, Sear DA, Naden PS, Armitage PD. The Impact Of Fine Sediment On  
45 Macro-Invertebrates. *River Res Appl* [Internet]. 2012 Oct [cited 2018 Jul 30];28(8):1055–71.  
46 Available from: <http://doi.wiley.com/10.1002/rra.1516>
- 47 9. Jones JI, Collins AL, Naden PS, Sear DA. The relationship between fine sediment and macrophytes  
48 in rivers. *River Res Appl* [Internet]. 2012 Sep [cited 2018 Jul 30];28(7):1006–18. Available from:  
49 <http://doi.wiley.com/10.1002/rra.1486>
- 51 10. Jones JI, Duerdoth CP, Collins AL, Naden PS, Sear DA. Interactions between diatoms and fine  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 sediment. *Hydrol Process* [Internet]. 2014 Jan 30 [cited 2018 Jul 30];28(3):1226–37. Available  
4 from: <http://doi.wiley.com/10.1002/hyp.9671>  
5
- 6 11. Wharton G, Mohajeri SH, Righetti M. The pernicious problem of streambed colmatation: a multi-  
7 disciplinary reflection on the mechanisms, causes, impacts, and management challenges. *Wiley*  
8 *Interdiscip Rev Water* [Internet]. 2017 Sep [cited 2018 Jul 11];4(5):e1231. Available from:  
9 <http://doi.wiley.com/10.1002/wat2.1231>  
10
- 11 12. Collins AL, Zhang YS, Duethmann D, Walling DE, Black KS. Using a novel tracing-tracking  
12 framework to source fine-grained sediment loss to watercourses at sub-catchment scale. *Hydrol*  
13 *Process* [Internet]. 2013 Mar 15 [cited 2018 Jul 30];27(6):959–74. Available from:  
14 <http://doi.wiley.com/10.1002/hyp.9652>  
15
- 16 13. Nosrati K, Collins AL, Madankan M. Fingerprinting sub-basin spatial sediment sources using  
17 different multivariate statistical techniques and the Modified MixSIR model. *Catena* [Internet].  
18 2018 [cited 2018 Jul 30];164:32–43. Available from:  
19 <https://www.sciencedirect.com/science/article/pii/S0341816218300031>  
20  
21
- 22 14. Collins AL, Williams LJ, Zhang YS, Marius M, Dungait JAJ, Smallman DJ, et al. Catchment source  
23 contributions to the sediment-bound organic matter degrading salmonid spawning gravels in a  
24 lowland river, southern England. *Sci Total Environ* [Internet]. 2013 Jul 1 [cited 2018 Jul 30];456–  
25 457:181–95. Available from:  
26 <https://www.sciencedirect.com/science/article/pii/S0048969713003987>  
27
- 28 15. Collins AL, Walling DE, Stroud RW, Robson M, Peet LM. Assessing damaged road verges as a  
29 suspended sediment source in the Hampshire Avon catchment, southern United Kingdom. *Hydrol*  
30 *Process* [Internet]. 2010 Apr 30 [cited 2018 Jul 30];24(9):1106–22. Available from:  
31 <http://doi.wiley.com/10.1002/hyp.7573>  
32  
33
- 34 16. Collins AL, Zhang YS, Hickinbotham R, Bailey G, Darlington S, Grenfell SE, et al. Contemporary  
35 fine-grained bed sediment sources across the River Wensum Demonstration Test Catchment, UK.  
36 *Hydrol Process* [Internet]. 2013 Mar 15 [cited 2018 Jul 30];27(6):857–84. Available from:  
37 <http://doi.wiley.com/10.1002/hyp.9654>  
38
- 39 17. Edwards KJ, Whittington G. Lake sediments, erosion and landscape change during the Holocene  
40 in Britain and Ireland. *Catena* [Internet]. 2001 Jan 20 [cited 2018 Jul 30];42(2–4):143–73.  
41 Available from: <https://www.sciencedirect.com/science/article/pii/S0341816200001363>  
42
- 43 18. Chiverrell RC, Oldfield F, Appleby PG, Barlow D, Fisher E, Thompson R, et al. Evidence for changes  
44 in Holocene sediment flux in Semer Water and Raydale, North Yorkshire, UK. *Geomorphology*  
45 [Internet]. 2008 Aug 1 [cited 2018 Jul 30];100(1–2):70–82. Available from:  
46 <https://www.sciencedirect.com/science/article/pii/S0169555X08001712>  
47
- 48 19. Macklin MG, Jones AF, Lewin J. River response to rapid Holocene environmental change:  
49 evidence and explanation in British catchments. *Quat Sci Rev* [Internet]. 2010 Jun 1 [cited 2018  
50 Jul 30];29(13–14):1555–76. Available from:  
51 <https://www.sciencedirect.com/science/article/pii/S0277379109002030>  
52  
53
- 54 20. Foster IDL, Collins AL, Naden PS, Sear DA, Jones JI, Zhang Y. The potential for paleolimnology to  
55 determine historic sediment delivery to rivers. *J Paleolimnol* [Internet]. 2011 Feb 20 [cited 2018  
56  
57  
58  
59  
60

- 1  
2  
3 Jul 30];45(2):287–306. Available from: <http://link.springer.com/10.1007/s10933-011-9498-9>  
4
- 5 21. Collins AL, Zhang Y. Exceedance of modern “background” fine-grained sediment delivery to rivers  
6 due to current agricultural land use and uptake of water pollution mitigation options across  
7 England and Wales. *Environ Sci Policy* [Internet]. 2016;61:61–73. Available from:  
8 <https://www.sciencedirect.com/science/article/pii/S1462901116300740>  
9
- 10 22. Frostick LE, Lucas PM, Reid I. The infiltration of fine matrices into coarse-grained alluvial  
11 sediments and its implications for stratigraphical interpretation. *J Geol Soc London* [Internet].  
12 1984 [cited 2018 Jul 30];141(6):955–65. Available from:  
13 <http://jgs.lyellcollection.org/cgi/doi/10.1144/gsjgs.141.6.0955>  
14
- 15 23. Lisle TE. Sediment transport and resulting deposition in spawning gravels, north coastal  
16 California. *Water Resour Res* [Internet]. 1989 Jun [cited 2018 Jul 30];25(6):1303–19. Available  
17 from: <http://doi.wiley.com/10.1029/WR025i006p01303>  
18
- 19 24. Sear DA. Fine sediment infiltration into gravel spawning beds within a regulated river  
20 experiencing floods: Ecological implications for salmonids. *Regul Rivers Res Manag* [Internet].  
21 1993 Dec [cited 2018 Jul 30];8(4):373–90. Available from:  
22 <http://doi.wiley.com/10.1002/rrr.3450080407>  
23
- 24 25. Owens PN, Batalla RJ, Collins AJ, Gomez B, Hicks DM, Horowitz AJ, et al. Fine-grained sediment in  
25 river systems: environmental significance and management issues. *River Res Appl* [Internet].  
26 2005 Sep [cited 2018 Jul 30];21(7):693–717. Available from:  
27 <http://doi.wiley.com/10.1002/rra.878>  
28
- 29 26. Walling DE, Moorehead PW. The particle size characteristics of fluvial suspended sediment: an  
30 overview. In: *Hydrobiologia* [Internet]. Dordrecht: Springer Netherlands; 1989 [cited 2018 Jul 30].  
31 p. 125–49. Available from: [http://www.springerlink.com/index/10.1007/978-94-009-2376-8\\_12](http://www.springerlink.com/index/10.1007/978-94-009-2376-8_12)  
32
- 33 27. Droppo IG. Rethinking what constitutes suspended sediment. *Hydrol Process* [Internet]. 2001 Jun  
34 30 [cited 2018 Jul 30];15(9):1551–64. Available from: <http://doi.wiley.com/10.1002/hyp.228>  
35
- 36 28. Wohl E, Bledsoe BP, Jacobson RB, Poff NL, Rathburn SL, Walters DM, et al. The natural sediment  
37 regime in rivers: Broadening the foundation for ecosystem management. *Bioscience* [Internet].  
38 2015 Apr 1 [cited 2018 Jul 30];65(4):358–71. Available from:  
39 [http://academic.oup.com/bioscience/article/65/4/358/254680/The-Natural-Sediment-Regime-](http://academic.oup.com/bioscience/article/65/4/358/254680/The-Natural-Sediment-Regime-in-Rivers-Broadening)  
40 [in-Rivers-Broadening](http://academic.oup.com/bioscience/article/65/4/358/254680/The-Natural-Sediment-Regime-in-Rivers-Broadening)  
41
- 42 29. Petts GE, Gurnell AM. Dams and geomorphology: Research progress and future directions.  
43 *Geomorphology* [Internet]. 2005 Oct 1 [cited 2018 Jul 30];71(1–2):27–47. Available from:  
44 <https://www.sciencedirect.com/science/article/pii/S0169555X05000875>  
45
- 46 30. Crosa G, Castelli E, Gentili G, Espa P. Effects of suspended sediments from reservoir flushing on  
47 fish and macroinvertebrates in an alpine stream. *Aquat Sci* [Internet]. 2009 Jan 13 [cited 2018 Jul  
48 30];72(1):85–95. Available from: <http://link.springer.com/10.1007/s00027-009-0117-z>  
49
- 50 31. Einstein HA. Deposition of suspended particles in a gravel bed. *J Hydraul Div* [Internet]. 1968  
51 [cited 2018 Jul 30];1197–205. Available from:  
52 <http://cedb.asce.org/CEDBsearch/record.jsp?dockkey=0015321>  
53
- 54  
55  
56  
57  
58  
59  
60

- 1
  - 2
  - 3
  - 4
  - 5
  - 6
  - 7
  - 8
  - 9
  - 10
  - 11
  - 12
  - 13
  - 14
  - 15
  - 16
  - 17
  - 18
  - 19
  - 20
  - 21
  - 22
  - 23
  - 24
  - 25
  - 26
  - 27
  - 28
  - 29
  - 30
  - 31
  - 32
  - 33
  - 34
  - 35
  - 36
  - 37
  - 38
  - 39
  - 40
  - 41
  - 42
  - 43
  - 44
  - 45
  - 46
  - 47
  - 48
  - 49
  - 50
  - 51
  - 52
  - 53
  - 54
  - 55
  - 56
  - 57
  - 58
  - 59
  - 60
32. Beschta RL, Jackson WL. The Intrusion of Fine Sediments into a Stable Gravel Bed. *J Fish Res Board Canada* [Internet]. 1979 Feb [cited 2018 Jul 30];36(2):204–10. Available from: <http://www.nrcresearchpress.com/doi/10.1139/f79-030>
33. Schälchli U. The clogging of coarse gravel river beds by fine sediment. *Hydrobiologia* [Internet]. 1992 Jul [cited 2018 Jul 30];235–236(1):189–97. Available from: <http://link.springer.com/10.1007/BF00026211>
34. Schindler Wildhaber Y, Michel C, Epting J, Wildhaber RA, Huber E, Huggenberger P, et al. Effects of river morphology, hydraulic gradients, and sediment deposition on water exchange and oxygen dynamics in salmonid redds. *Sci Total Environ* [Internet]. 2014 Feb 1 [cited 2018 Jul 30];470–471:488–500. Available from: <https://www.sciencedirect.com/science/article/pii/S0048969713011388>
35. Elliott AH, Brooks NH. Transfer of nonsorbing solutes to a streambed with bed forms: Theory. *Water Resour Res* [Internet]. 1997 Jan [cited 2018 Jul 30];33(1):123–36. Available from: <http://doi.wiley.com/10.1029/96WR02783>
36. Mathers KL, Wood PJ. Fine sediment deposition and interstitial flow effects on macroinvertebrate community composition within riffle heads and tails. *Hydrobiologia* [Internet]. 2016 Aug 31 [cited 2018 Jul 30];776(1):147–60. Available from: <http://link.springer.com/10.1007/s10750-016-2748-0>
37. Brunke M. Colmation and depth filtration within streambeds: Retention of particles in hyporheic interstices. *Int Rev Hydrobiol* [Internet]. 1999 [cited 2018 Jul 30];84(2):99–117. Available from: <https://onlinelibrary.wiley.com/doi/abs/10.1002/iroh.199900014>
38. Savant SA, Reible DD, Thibodeaux LJ. Convective transport within stable river sediments. *Water Resour Res* [Internet]. 1987 Sep [cited 2018 Jul 30];23(9):1763–8. Available from: <http://doi.wiley.com/10.1029/WR023i009p01763>
39. Packman AI, Salehin M, Zaramella M. Hyporheic Exchange with Gravel Beds: Basic Hydrodynamic Interactions and Bedform-Induced Advective Flows. *J Hydraul Eng* [Internet]. 2004 Jul [cited 2018 Jul 30];130(7):647–56. Available from: <http://ascelibrary.org/doi/10.1061/%28ASCE%290733-9429%282004%29130%3A7%28647%29>
40. Cardenas MB, Wilson JL, Zlotnik VA. Impact of heterogeneity, bed forms, and stream curvature on subchannel hyporheic exchange. *Water Resour Res* [Internet]. 2004 Aug [cited 2018 Jul 30];40(8). Available from: <http://doi.wiley.com/10.1029/2004WR003008>
41. Stewardson MJ, Datry T, Lamouroux N, Pella H, Thommeret N, Valette L, et al. Variation in reach-scale hydraulic conductivity of streambeds. *Geomorphology* [Internet]. 2016 [cited 2018 Jul 30];259:70–80. Available from: <https://www.sciencedirect.com/science/article/pii/S0169555X16300216>
42. Casas-Mulet R, Lakhanpal G, Stewardson MJ. The relative contribution of near-bed vs. intragravel horizontal transport to fine sediment accumulation processes in river gravel beds. *Geomorphology* [Internet]. 2018 [cited 2018 Aug 6];303:299–308. Available from: <https://www.sciencedirect.com/science/article/pii/S0169555X17305147>
43. Manes C, Pokrajac D, McEwan I, Nikora V. Turbulence structure of open channel flows over

- permeable and impermeable beds: A comparative study. *Phys Fluids* [Internet]. 2009 Dec [cited 2018 Jul 30];21(12):1–12. Available from: <http://aip.scitation.org/doi/10.1063/1.3276292>
44. Casas-Mulet R, Alfredsen KT, McCluskey AH, Stewardson MJ. Key hydraulic drivers and patterns of fine sediment accumulation in gravel streambeds: A conceptual framework illustrated with a case study from the Kiewa River, Australia. *Geomorphology* [Internet]. 2017 [cited 2018 Jul 30];299:152–64. Available from: <https://www.sciencedirect.com/science/article/pii/S0169555X17303380>
45. Hassan MA, Tonina D, Beckie RD, Kinnear M. The effects of discharge and slope on hyporheic flow in step-pool morphologies. *Hydrol Process* [Internet]. 2015 Jan 30 [cited 2018 Jul 30];29(3):419–33. Available from: <http://doi.wiley.com/10.1002/hyp.10155>
46. Murphy JF, Jones JI, Pretty JL, Duerdoth CP, Hawczak A, Arnold A, et al. Development of a biotic index using stream macroinvertebrates to assess stress from deposited fine sediment. *Freshw Biol* [Internet]. 2015 Oct 1 [cited 2018 Aug 6];60(10):2019–36. Available from: <http://doi.wiley.com/10.1111/fwb.12627>
47. Murphy JF, Jones JI, Arnold A, Duerdoth CP, Pretty JL, Naden PS, et al. Can macroinvertebrate biological traits indicate fine-grained sediment conditions in streams? *River Res Appl* [Internet]. 2017 Dec 1 [cited 2018 Aug 1];33(10):1606–17. Available from: <http://doi.wiley.com/10.1002/rra.3194>
48. Sear DA, Jones JI, Collins AL, Hulin A, Burke N, Bateman S, et al. Does fine sediment source as well as quantity affect salmonid embryo mortality and development? *Sci Total Environ* [Internet]. 2016 Jan 15 [cited 2018 Aug 6];541:957–68. Available from: <https://www.sciencedirect.com/science/article/pii/S0048969715308160>
49. Sear DA, Pattison I, Collins AL, Smallman DJ, Jones JI, Naden PS. The magnitude and significance of sediment oxygen demand in gravel spawning beds for the incubation of salmonid embryos. *River Res Appl* [Internet]. 2017 Dec [cited 2018 Jul 30];33(10):1642–54. Available from: <http://doi.wiley.com/10.1002/rra.3212>
50. Von Bertrab MG, Krein A, Stendera S, Thielen F, Hering D. Is fine sediment deposition a main driver for the composition of benthic macroinvertebrate assemblages? *Ecol Indic* [Internet]. 2013 [cited 2018 Jul 30];24:589–98. Available from: <https://www.sciencedirect.com/science/article/pii/S1470160X12002920>
51. Elbrecht V, Beermann AJ, Goessler G, Neumann J, Tollrian R, Wagner R, et al. Multiple-stressor effects on stream invertebrates: A mesocosm experiment manipulating nutrients, fine sediment and flow velocity. *Freshw Biol* [Internet]. 2016 [cited 2018 Jul 30];61(4):362–75. Available from: <https://www.sciencedirect.com/science/article/pii/S0048969717320818>
52. Davis SJ, Ó hUallacháin D, Mellander PE, Kelly AM, Matthaei CD, Piggott JJ, et al. Multiple-stressor effects of sediment, phosphorus and nitrogen on stream macroinvertebrate communities. *Sci Total Environ* [Internet]. 2018 [cited 2018 Jul 30];637–638(May):577–87. Available from: <https://doi.org/10.1016/j.scitotenv.2018.05.052>
53. Kirk JTO. Effects of suspensoids (turbidity) on penetration of solar radiation in aquatic ecosystems. *Hydrobiologia* [Internet]. 1985 Jun [cited 2018 Jul 30];125(1):195–208. Available

- 1  
2  
3 from: <http://link.springer.com/10.1007/BF00045935>  
4
- 5 54. Wood PJ, Armitage PD. Biological effects of fine sediment in the lotic environment. *Environ*  
6 *Manage* [Internet]. 1997 Mar 1 [cited 2018 Jul 19];21(2):203–17. Available from:  
7 <http://link.springer.com/10.1007/s002679900019>  
8
- 9 55. Paaijmans KP, Takken W, Githeko AK, Jacobs AFG. The effect of water turbidity on the near-  
10 surface water temperature of larval habitats of the malaria mosquito *Anopheles gambiae*. *Int J*  
11 *Biometeorol* [Internet]. 2008 Nov 17 [cited 2018 Jul 30];52(8):747–53. Available from:  
12 <http://link.springer.com/10.1007/s00484-008-0167-2>  
13
- 14 56. Veličković B. Colmation as one of the processes in interaction between the groundwater and  
15 surface water. *Facta Univ - Ser Archit Civ Eng* [Internet]. 2005 [cited 2018 Jul 30];3(2):165–72.  
16 Available from: <http://scindeks.ceon.rs/article.aspx?artid=0354-46050502165V>  
17
- 18 57. Soulsby C, Youngson AF, Moir HJ, Malcolm IA. Fine sediment influence on salmonid spawning  
19 habitat in a lowland agricultural stream: A preliminary assessment. *Sci Total Environ* [Internet].  
20 2001 [cited 2018 Jul 30];265(1–3):295–307. Available from:  
21 <https://www.sciencedirect.com/science/article/pii/S0048969700006720>  
22
- 23 58. Greig SM, Sear DA, Carling PA. A review of factors influencing the availability of dissolved oxygen  
24 to incubating salmonid embryos. *Hydrol Process* [Internet]. 2007 Jan 30 [cited 2018 Jul  
25 30];21(3):323–34. Available from: <http://doi.wiley.com/10.1002/hyp.6188>  
26
- 27 59. Chapman JM, Proulx CL, Veilleux MAN, Levert C, Bliss S, André MÈ, et al. Clear as mud: A meta-  
28 analysis on the effects of sedimentation on freshwater fish and the effectiveness of sediment-  
29 control measures. *Water Res* [Internet]. 2014 [cited 2018 Jul 30];56:190–202. Available from:  
30 <https://www.sciencedirect.com/science/article/pii/S0043135414001754>  
31
- 32 60. Jones I, Grouns I, Arnold A, McCall S, Bowes M. The effects of increased flow and fine sediment  
33 on hyporheic invertebrates and nutrients in stream mesocosms. *Freshw Biol* [Internet]. 2015 Apr  
34 1 [cited 2018 Aug 2];60(4):813–26. Available from: <http://doi.wiley.com/10.1111/fwb.12536>  
35
- 36 61. Emerson D, Revsbech NP. Investigation of an iron-oxidizing microbial mat community located  
37 near Aarhus, Denmark: Field studies. *Appl Environ Microbiol* [Internet]. 1994 Nov 1 [cited 2018  
38 Jul 30];60(11):4022–31. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/16349433>  
39
- 40 62. Waters TF. *Sediment in Streams: Source, Biological Effects and Control*. American Fisheries  
41 Society monograph (USA). Bethesda: American Fisheries Society; 1995. 251 p.  
42
- 43 63. Westrich B, Förstner U. *Sediment dynamics and pollutant mobility in rivers: an interdisciplinary*  
44 *approach* [Internet]. Berlin: Springer; 2007 [cited 2018 Jul 30]. Available from:  
45 <https://books.google.co.uk/books?hl=en&lr=&id=UxbcOmW9pvQC&oi=fnd&pg=PA1&dq=Sediment+Dynamics+and+Pollutant+Mobility+in+Rivers:+An+Interdisciplinary+Approach&ots=t2X5okd4S8&sig=5Z7OOxwbjjV0qTO63RQH4G6lvKo#v=onepage&q=Sediment+Dynamics+and+Pollutant+Mobility>  
46
- 47 64. Ji Z. *Hydrodynamics and Water Quality: Modeling Rivers, Lakes, and Estuaries*. Second ed.  
48 Hoboken: John Wiley & Sons; 2017.  
49
- 50 65. Hossain HMZ, Kawahata H, Roser BP, Sampei Y, Manaka T, Otani S. *Geochemical characteristics*  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

- of modern river sediments in Myanmar and Thailand: Implications for provenance and weathering. *Chemie der Erde - Geochemistry* [Internet]. 2017 [cited 2018 Jul 30];77(3):443–58. Available from: <https://www.sciencedirect.com/science/article/pii/S0009281916302343>
66. Huang Y, Zhang D, Xu Z, Yuan S, Li Y, Wang L. Effect of overlying water pH, dissolved oxygen and temperature on heavy metal release from river sediments under laboratory conditions. *Arch Environ Prot* [Internet]. 2017 [cited 2018 Jul 30];43(2):28–36. Available from: <https://www.degruyter.com/view/j/aep.2017.43.issue-2/aep-2017-0014/aep-2017-0014.xml>
67. Liikanen, ALiikanen, A., T. Murtoniemi, H. Tanskanen, T. Väisänen and PJM 2002. E of T and OA on GG and ND in S of a EM-BL greenhouse gas and nutrient dynamics in sediment of. *B 59*:269–286., Murtoniemi T, Tanskanen H, Väisänen T, Martikainen PJ. Effects of Temperature and Oxygen Availability on Greenhouse Gas and Nutrient Dynamics in Sediment of a Eutrophic Mid-Boreal Lake greenhouse gas and nutrient dynamics in sediment of. *Biogeochemistry* [Internet]. 2002 [cited 2018 Jul 30];59(3):269–86. Available from: <https://link.springer.com/article/10.1023/A:1016015526712>
68. Wu Y, Wen Y, Zhou J, Wu Y. Phosphorus release from lake sediments: Effects of pH, temperature and dissolved oxygen. *KSCE J Civ Eng* [Internet]. 2014 Jan 27 [cited 2018 Aug 2];18(1):323–9. Available from: <http://link.springer.com/10.1007/s12205-014-0192-0>
69. Tsai LJ, Ho ST, Yu KC. Correlations of extractable heavy metals with organic matters in contaminated river sediments. *Water Sci Technol* [Internet]. 2003 [cited 2018 Jul 30];47(9):101–7. Available from: <https://iwaponline.com/wst/article-abstract/47/9/101/10564>
70. Fojut TL, Young TM. Pyrethroid sorption to Sacramento River suspended solids and bed sediments. *Environ Toxicol Chem* [Internet]. 2011 Apr [cited 2018 Jul 30];30(4):787–92. Available from: <http://doi.wiley.com/10.1002/etc.448>
71. Duong HT, Kadokami K, Pan S, Matsuura N, Nguyen TQ. Screening and analysis of 940 organic micro-pollutants in river sediments in Vietnam using an automated identification and quantification database system for GC-MS. *Chemosphere* [Internet]. 2014 [cited 2018 Jul 30];107:462–72. Available from: <http://dx.doi.org/10.1016/j.chemosphere.2014.01.064>
72. Van Leussen W. Aggregation of particles, settling velocity of mud flocs a review. In: *Physical processes in estuaries* [Internet]. Berlin, Heidelberg: Springer Berlin Heidelberg; 1988 [cited 2018 Jul 31]. p. 347–403. Available from: [http://link.springer.com/10.1007/978-3-642-73691-9\\_19](http://link.springer.com/10.1007/978-3-642-73691-9_19)
73. Bainbridge ZT, Wolanski E, Álvarez-Romero JG, Lewis SE, Brodie JE. Fine sediment and nutrient dynamics related to particle size and floc formation in a Burdekin River flood plume, Australia. *Mar Pollut Bull* [Internet]. 2012 [cited 2018 Jul 31];65(4–9):236–48. Available from: <https://www.sciencedirect.com/science/article/pii/S0025326X12000665>
74. Casas-Ruiz JP, Catalán N, Gómez-Gener L, von Schiller D, Obrador B, Kothawala DN, et al. A tale of pipes and reactors: Controls on the in-stream dynamics of dissolved organic matter in rivers. *Limnol Oceanogr* [Internet]. 2017 Nov [cited 2018 Jul 31];62(S1):S85–94. Available from: <http://doi.wiley.com/10.1002/lno.10471>
75. Lau YL. Temperature effect on settling velocity and deposition of cohesive sediments. *J Hydraul Eng* [Internet]. 1994 Jan [cited 2018 Jul 31];32(1):41–51. Available from:

- 1  
2  
3 <http://www.tandfonline.com/doi/abs/10.1080/00221689409498788>  
4
- 5 76. Butler DR. Zoogeomorphology: Animals as Geomorphic Agents [Internet]. New York: Cambridge  
6 University Press; 1995 [cited 2018 Aug 1]. Available from: <http://doi.wiley.com/10.2307/2265638>  
7
- 8 77. Jones CG, Lawton JH, Shachak M. Organisms as Ecosystem Engineers. *Oikos* [Internet].  
9 1994;69:373–86. Available from: <http://www.jstor.org/stable/3545850?origin=JSTOR-pdf>  
10
- 11 78. Wright JP, Jones CG. The Concept of Organisms as Ecosystem Engineers Ten Years On: Progress,  
12 Limitations, and Challenges. *AIBS Bull* [Internet]. 2006 Mar 1 [cited 2018 Aug 1];56(3):203–9.  
13 Available from: <https://academic.oup.com/bioscience/article/56/3/203-209/333061>  
14
- 15 79. Pakeman J. Multivariate identification of plant functional response and effect traits in an  
16 agricultural landscape. *Ecology* [Internet]. 2011 [cited 2018 Aug 1];92(6):1353–65. Available  
17 from: <https://esajournals.onlinelibrary.wiley.com/doi/abs/10.1890/10-1728.1>  
18
- 19 80. Watnick P, Kolter R. Biofilm, city of microbes. *J Bacteriol* [Internet]. 2000 May 15 [cited 2018 Jul  
20 31];182(10):2675–9. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/10781532>  
21
- 22 81. Nikolaev YA, Plakunov VK. Biofilm—“City of microbes” or an analogue of multicellular organisms?  
23 *Microbiology* [Internet]. 2007 Apr [cited 2018 Jul 31];76(2):125–38. Available from:  
24 <http://link.springer.com/10.1134/S0026261707020014>  
25
- 26 82. Lock M. Attached microbial communities in rivers. *Aquat Microbiol An Ecol Approach* [Internet].  
27 1993 [cited 2018 Aug 1];113–138. Available from: <https://ci.nii.ac.jp/naid/10009490929/>  
28
- 29 83. Madsen EL. Microorganisms and their roles in fundamental biogeochemical cycles. *Curr Opin*  
30 *Biotechnol* [Internet]. 2011 Jun 1 [cited 2018 Jul 31];22(3):456–64. Available from:  
31 <https://www.sciencedirect.com/science/article/pii/S095816691100022X>  
32
- 33 84. Aspray KL, Holden J, Ledger ME, Mainstone CP, Brown LE. Organic sediment pulses impact rivers  
34 across multiple levels of ecological organization. *Ecohydrology* [Internet]. 2017 Sep 1 [cited 2018  
35 Jul 31];10(6):e1855. Available from: <http://doi.wiley.com/10.1002/eco.1855>  
36
- 37 85. Cheng W, Fang H, Lai H, Huang L, Dey S. Effects of biofilm on turbulence characteristics and the  
38 transport of fine sediment. *J Soils Sediments* [Internet]. 2017 Nov 10 [cited 2018 Jul 31];1–15.  
39 Available from: <http://link.springer.com/10.1007/s11368-017-1859-1>  
40
- 41 86. Fang H, Chen Y, Huang L, He G. Analysis of biofilm bacterial communities under different shear  
42 stresses using size-fractionated sediment. *Sci Rep* [Internet]. 2017 Dec 2 [cited 2018 Jul  
43 31];7(1):1299. Available from: <http://www.nature.com/articles/s41598-017-01446-4>  
44
- 45 87. Gainswin BE, House WA, Leadbeater BSC, Armitage PD, Patten J. The effects of sediment size  
46 fraction and associated algal biofilms on the kinetics of phosphorus release. *Sci Total Environ*  
47 [Internet]. 2006 May 1 [cited 2018 Jul 31];360(1–3):142–57. Available from:  
48 <https://www.sciencedirect.com/science/article/pii/S0048969705005735>  
49
- 50 88. Garwood JC, Hill PS, Law BA. Biofilms and Size Sorting of Fine Sediment During Erosion in  
51 Intertidal Sands. *Estuaries and Coasts* [Internet]. 2013 Sep 17 [cited 2018 Jul 31];36(5):1024–36.  
52 Available from: <http://link.springer.com/10.1007/s12237-013-9618-z>  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 89. Lu H, Wan J, Li J, Shao H, Wu Y. Periphytic biofilm: A buffer for phosphorus precipitation and  
4 release between sediments and water. *Chemosphere* [Internet]. 2016 Feb 1 [cited 2018 Jul  
5 31];144:2058–64. Available from:  
6 <https://www.sciencedirect.com/science/article/pii/S0045653515303246>  
7
- 8 90. Horton AA, Walton A, Spurgeon DJ, Lahive E, Svendsen C. Microplastics in freshwater and  
9 terrestrial environments: Evaluating the current understanding to identify the knowledge gaps  
10 and future research priorities. *Sci Total Environ* [Internet]. 2017 May 15 [cited 2018 Jul  
11 31];586:127–41. Available from:  
12 <https://www.sciencedirect.com/science/article/pii/S0048969717302073>  
13
- 14 91. Rummel CD, Jahnke A, Gorokhova E, Kühnel D, Schmitt-Jansen M. Impacts of biofilm formation  
15 on the fate and potential effects of microplastic in the aquatic environment. *Environ Sci Technol*  
16 *Lett* [Internet]. 2017 Jul 11 [cited 2018 Jul 31];4(7):258–67. Available from:  
17 <http://pubs.acs.org/doi/10.1021/acs.estlett.7b00164>  
18
- 19 92. Wingender J, Neu TR, Flemming H-C. What are Bacterial Extracellular Polymeric Substances? In:  
20 *Microbial Extracellular Polymeric Substances* [Internet]. Berlin, Heidelberg: Springer Berlin  
21 Heidelberg; 1999 [cited 2018 Jul 31]. p. 1–19. Available from:  
22 [http://link.springer.com/10.1007/978-3-642-60147-7\\_1](http://link.springer.com/10.1007/978-3-642-60147-7_1)  
23
- 24 93. Flemming HC, Neu TR, Wozniak DJ. The EPS matrix: The “House of Biofilm Cells.” *J Bacteriol*  
25 [Internet]. 2007 Nov 15 [cited 2018 Jul 31];189(22):7945–7. Available from:  
26 <http://www.ncbi.nlm.nih.gov/pubmed/17675377>  
27
- 28 94. Lai H, Fang H, Huang L, He G, Reible D. A review on sediment bioflocculation: Dynamics,  
29 influencing factors and modeling. *Sci Total Environ* [Internet]. 2018 [cited 2018 Aug 1];642:1184–  
30 200. Available from: <https://www.sciencedirect.com/science/article/pii/S0048969718321776>  
31
- 32 95. Valentine K, Mariotti G, Fagherazzi S. Repeated erosion of cohesive sediments with biofilms. *Adv*  
33 *Geosci* [Internet]. 2014 Apr 1 [cited 2018 Jul 31];39:9–14. Available from: [https://www.adv-  
34 geosci.net/39/9/2014/](https://www.adv-geosci.net/39/9/2014/)  
35
- 36 96. Tolhurst TJ, Gust G, Paterson DM. The influence of an extracellular polymeric substance (EPS) on  
37 cohesive sediment stability. *Proc Mar Sci* [Internet]. 2002 Jan 1 [cited 2018 Jul 31];5(C):409–25.  
38 Available from: <https://www.sciencedirect.com/science/article/pii/S1568269202800304>  
39
- 40 97. Gerbersdorf SU, Wieprecht S. Biostabilization of cohesive sediments: Revisiting the role of abiotic  
41 conditions, physiology and diversity of microbes, polymeric secretion, and biofilm architecture.  
42 *Geobiology* [Internet]. 2015 Jan 1 [cited 2018 Jul 31];13(1):68–97. Available from:  
43 <http://doi.wiley.com/10.1111/gbi.12115>  
44
- 45 98. Lee BJ, Hur J, Toorman EA. Seasonal Variation in Flocculation Potential of River Water: Roles of  
46 the Organic Matter Pool. *Water (Switzerland)* [Internet]. 2017 May 8 [cited 2018 Jul 31];9(5):335.  
47 Available from: <http://www.mdpi.com/2073-4441/9/5/335>  
48
- 49 99. Passy SI, Larson CA. Succession in Stream Biofilms is an Environmentally Driven Gradient of Stress  
50 Tolerance. *Microb Ecol* [Internet]. 2011 Aug 27 [cited 2018 Jul 31];62(2):414–24. Available from:  
51 <http://link.springer.com/10.1007/s00248-011-9879-7>  
52
- 53 100. Dodds WK, Smith VH. Nitrogen, phosphorus, and eutrophication in streams. *Inl Waters* [Internet].  
54  
55  
56  
57  
58  
59  
60

- 2016 Apr 1 [cited 2018 Jul 31];6(2):155–64. Available from:  
<http://www.tandfonline.com/doi/full/10.5268/IW-6.2.909>
101. Olapade OA, Leff LG. Seasonal response of stream biofilm communities to dissolved organic matter and nutrient enrichments. *Appl Environ Microbiol* [Internet]. 2005 May 1 [cited 2018 Jul 31];71(5):2278–87. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/15870312>
102. Besemer K, Singer G, Limberger R, Chlup AK, Hochedlinger G, Hödl I, et al. Biophysical controls on community succession in stream biofilms. *Appl Environ Microbiol* [Internet]. 2007 Aug 1 [cited 2018 Jul 31];73(15):4966–74. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/17557861>
103. Prieto DM, Devesa-Rey R, Rubinos DA, Díaz-Fierros F, Barral MT. Biofilm Formation on River Sediments Under Different Light Intensities and Nutrient Inputs: A Flume Mesocosm Study. *Environ Eng Sci* [Internet]. 2016 Apr 8 [cited 2018 Jul 31];33(4):250–60. Available from: <http://online.liebertpub.com/doi/10.1089/ees.2015.0427>
104. Fang H, Cheng W, Fazeli M, Dey S. Bedforms and Flow Resistance of Cohesive Beds with and without Biofilm Coating. *J Hydraul Eng* [Internet]. 2017 Aug [cited 2018 Jul 31];143(8):06017010. Available from: <http://ascelibrary.org/doi/10.1061/%28ASCE%29HY.1943-7900.0001313>
105. Fang H, Fazeli M, Cheng W, Dey S. Transport of biofilm-coated sediment particles. *J Hydraul Res* [Internet]. 2016 Nov 23 [cited 2018 Jul 31];54(6):631–45. Available from: <https://www.tandfonline.com/doi/full/10.1080/00221686.2016.1212938>
106. Nguyen TH, Tang FHM, Maggi F. Micro food web networks on suspended sediment. *Sci Total Environ* [Internet]. 2018 Dec 1 [cited 2018 Jul 31];643:1387–99. Available from: <https://www.sciencedirect.com/science/article/pii/S0048969718323143>
107. Shelford VE. Some Lower Mississippi Valley Flood Plain Biotic Communities: Their age and Elevation. *Ecology* [Internet]. 1954 Apr [cited 2018 Aug 2];35(2):126. Available from: <http://www.jstor.org/stable/1931109?origin=crossref>
108. Hupp CR. Vegetation Pattern on Channel Features in the Passage Creek Gorge, Virginia. *Castanea* [Internet]. 1983 [cited 2018 Aug 2];48(2):62–72. Available from: <http://www.jstor.org/stable/4033067>
109. Hupp CR. Upstream variation in bottomland vegetation patterns, northwestern Virginia. *Bull Torrey Bot Club* [Internet]. 1986 [cited 2018 Aug 2];113(4):421–30. Available from: <https://www.jstor.org/stable/2996435>
110. Corenblit D, Tabacchi E, Steiger J, Gurnell AM. Reciprocal interactions and adjustments between fluvial landforms and vegetation dynamics in river corridors: A review of complementary approaches. *Earth-Science Rev* [Internet]. 2007 Sep 1 [cited 2018 Jul 31];84(1–2):56–86. Available from: <https://www.sciencedirect.com/science/article/pii/S0012825207000682>
111. Corenblit D, Steiger J, Gurnell AM, Naiman RJ. Plants intertwine fluvial landform dynamics with ecological succession and natural selection: A niche construction perspective for riparian systems. *Glob Ecol Biogeogr* [Internet]. 2009 Jul 1 [cited 2018 Jul 31];18(4):507–20. Available from: <http://doi.wiley.com/10.1111/j.1466-8238.2009.00461.x>
112. Gurnell A. Plants as river system engineers. *Earth Surf Process Landforms* [Internet]. 2014 Jan 1

- [cited 2018 Jul 31];39:4–25. Available from: <https://doi.org/10.1002/esp.3397>
113. Asaeda T, Rajapakse L, Kanoh M. Fine sediment retention as affected by annual shoot collapse: *Sparganium erectum* as an ecosystem engineer in a lowland stream. *River Res Appl* [Internet]. 2010 Nov 1 [cited 2018 Jul 31];26(9):1153–69. Available from: <http://doi.wiley.com/10.1002/rra.1322>
114. Liffen T, Gurnell AM, O'Hare MT, Pollen-Bankhead N, Simon A. Biomechanical properties of the emergent aquatic macrophyte *Sparganium erectum*: Implications for fine sediment retention in low energy rivers. *Ecol Eng* [Internet]. 2011 Nov 1 [cited 2018 Jul 31];37(11):1925–31. Available from: <https://www.sciencedirect.com/science/article/pii/S0925857411002023>
115. Cotton JA, Wharton G, Bass JAB, Heppell CM, Wotton RS. The effects of seasonal changes to in-stream vegetation cover on patterns of flow and accumulation of sediment. *Geomorphology* [Internet]. 2006 Jul 30 [cited 2018 Jul 31];77(3–4):320–34. Available from: <https://www.sciencedirect.com/science/article/pii/S0169555X06000201>
116. Abbe TB, Montgomery DR. Patterns and processes of wood debris accumulation in the Queets river basin, Washington. *Geomorphology* [Internet]. 2003 Mar 20 [cited 2018 Jul 31];51(1–3):81–107. Available from: <https://www.sciencedirect.com/science/article/pii/S0169555X02003264>
117. Nachtergaele J, Poesen J, Vandekerckhove L, Oostwoud Wijdenes D, Roxo M. Riparian vegetation and island formation along the gravel-bed Fiume Tagliamento, Italy. *Earth Surf Process Landforms* [Internet]. 2001 Jan 1 [cited 2018 Jul 31];26(1):31–62. Available from: <http://doi.wiley.com/10.1002/1096-9837%28200101%2926%3A1%3C31%3A%3AAID-ESP155%3E3.0.CO%3B2-Y>
118. Gurnell A, Tockner K, Edwards P, Petts G. Effects of deposited wood on biocomplexity of river corridors. *Front Ecol Environ* [Internet]. 2005 Sep 1 [cited 2018 Jul 31];3(7):377–82. Available from: [https://esajournals.onlinelibrary.wiley.com/doi/abs/10.1890/1540-9295\(2005\)003%5B0377:EODWOB%5D2.0.CO%3B2](https://esajournals.onlinelibrary.wiley.com/doi/abs/10.1890/1540-9295(2005)003%5B0377:EODWOB%5D2.0.CO%3B2)
119. Rood SB, Goater LA, Gill KM, Braatne JH. Sand and sandbar willow: A feedback loop amplifies environmental sensitivity at the riparian interface. *Oecologia* [Internet]. 2011 Jan 29 [cited 2018 Jul 31];165(1):31–40. Available from: <http://link.springer.com/10.1007/s00442-010-1758-2>
120. Pettit NE, Latterell JJ, Naiman RJ. Formation, distribution and ecological consequences of flood-related wood debris piles in a bedrock confined river in semi-arid South Africa. *River Res Appl* [Internet]. 2006 Dec 1 [cited 2018 Jul 31];22(10):1097–110. Available from: <http://doi.wiley.com/10.1002/rra.959>
121. Sand-Jensen K. Influence of submerged macrophytes on sediment composition and near-bed flow in lowland streams. *Freshw Biol* [Internet]. 1998 Jun 1 [cited 2018 Jul 31];39(4):663–79. Available from: <http://doi.wiley.com/10.1046/j.1365-2427.1998.00316.x>
122. Gurnell AM, Morrissey IP, Boitsidis AJ, Bark T, Clifford NJ, Petts GE, et al. Initial adjustments within a new river channel: Interactions between fluvial processes, colonizing vegetation, and bank profile development. *Environ Manage* [Internet]. 2006 Oct 23 [cited 2018 Aug 2];38(4):580–96. Available from: <http://link.springer.com/10.1007/s00267-005-0190-6>
123. Luhar M, Rominger J, Nepf H. Interaction between flow, transport and vegetation spatial

- 1  
2  
3 structure. *Environ Fluid Mech* [Internet]. 2008 Dec 29 [cited 2018 Aug 2];8(5–6):423–39.  
4 Available from: <http://link.springer.com/10.1007/s10652-008-9080-9>  
5
- 6 124. Gurnell AM, Bertoldi W, Corenblit D. Changing river channels: The roles of hydrological  
7 processes, plants and pioneer fluvial landforms in humid temperate, mixed load, gravel bed  
8 rivers. *Earth-Science Rev* [Internet]. 2012 Feb 1 [cited 2018 Jul 31];111(1–2):129–41. Available  
9 from: <https://www.sciencedirect.com/science/article/pii/S0012825211001693>  
10
- 11 125. Schoelynck J, De Groot T, Bal K, Vandenbruwaene W, Meire P, Temmerman S. Self-organised  
12 patchiness and scale-dependent bio-geomorphic feedbacks in aquatic river vegetation.  
13 *Ecography (Cop)* [Internet]. 2012 Aug 1 [cited 2018 Jul 31];35(8):760–8. Available from:  
14 <http://doi.wiley.com/10.1111/j.1600-0587.2011.07177.x>  
15
- 16 126. Liffen T, Gurnell AM, O’Hare MT, Pollen-Bankhead N, Simon A. Associations between the  
17 morphology and biomechanical properties of *Sparganium erectum*: Implications for survival and  
18 ecosystem engineering. *Aquat Bot* [Internet]. 2013 Feb 1 [cited 2018 Jul 31];105:18–24. Available  
19 from: <https://www.sciencedirect.com/science/article/pii/S0304377012001829>  
20
- 21 127. Ghisalberti M, Nepf H. Shallow flows over a permeable medium: The hydrodynamics of  
22 submerged aquatic canopies. *Transp Porous Media* [Internet]. 2009 Jun 10 [cited 2018 Jul  
23 31];78(3 SPEC. ISS.):385–402. Available from: [http://link.springer.com/10.1007/s11242-008-](http://link.springer.com/10.1007/s11242-008-9305-x)  
24 [9305-x](http://link.springer.com/10.1007/s11242-008-9305-x)  
25
- 26 128. Gurnell AM, O’Hare JM, O’Hare MT, Dunbar MJ, Scarlett PM. An exploration of associations  
27 between assemblages of aquatic plant morphotypes and channel geomorphological properties  
28 within British rivers. *Geomorphology* [Internet]. 2010 [cited 2018 Oct 22];116(1–2):135–44.  
29 Available from: <https://www.sciencedirect.com/science/article/pii/S0169555X09004504>  
30
- 31 129. Gurnell AM, O’Hare MT, O’Hare JM, Scarlett P, Liffen TMR. The geomorphological context and  
32 impact of the linear emergent macrophyte, *Sparganium erectum* L.: A statistical analysis of  
33 observations from British rivers. *Earth Surf Process Landforms* [Internet]. 2013 [cited 2018 Oct  
34 22];38(15):1869–80. Available from: <https://onlinelibrary.wiley.com/doi/abs/10.1002/esp.3473>  
35
- 36 130. Collins BD, Montgomery DR, Fetherston KL, Abbe TB. The floodplain large-wood cycle hypothesis:  
37 A mechanism for the physical and biotic structuring of temperate forested alluvial valleys in the  
38 North Pacific coastal ecoregion. *Geomorphology* [Internet]. 2012 Feb 15 [cited 2018 Jul 31];139–  
39 140:460–70. Available from:  
40 <https://www.sciencedirect.com/science/article/pii/S0169555X11006088>  
41
- 42 131. Karrenberg S, Edwards PJ, Kollmann J. The life history of Salicaceae living in the active zone of  
43 floodplains. *Freshw Biol* [Internet]. 2002 Apr 1 [cited 2018 Jul 31];47(4):733–48. Available from:  
44 <http://doi.wiley.com/10.1046/j.1365-2427.2002.00894.x>  
45
- 46 132. Holloway J V., Rillig MC, Gurnell AM. Underground riparian wood: Reconstructing the processes  
47 influencing buried stem and coarse root structures of Black Poplar (*Populus nigra* L.).  
48 *Geomorphology* [Internet]. 2017 Feb 15 [cited 2018 Jul 31];279:199–208. Available from:  
49 <https://www.sciencedirect.com/science/article/pii/S0169555X16306869>  
50
- 51 133. Pasquale N, Perona P, Francis R, Burlando P. Effects of streamflow variability on the vertical root  
52 density distribution of willow cutting experiments. *Ecol Eng* [Internet]. 2012 Mar 1 [cited 2018 Jul  
53 54  
55  
56  
57  
58  
59  
60

- 31];40:167–72. Available from:  
<https://www.sciencedirect.com/science/article/pii/S0925857411003697>
134. Tron S, Perona P, Gorla L, Schwarz M, Laio F, Ridolfi L. The signature of randomness in riparian plant root distributions. *Geophys Res Lett* [Internet]. 2015 Sep 16 [cited 2018 Jul 31];42(17):7098–106. Available from: <http://doi.wiley.com/10.1002/2015GL064857>
135. Gurnell AM, Corenblit D, García de Jalón D, González del Tánago M, Grabowski RC, O'Hare MT, et al. A Conceptual Model of Vegetation-hydrogeomorphology Interactions Within River Corridors. *River Res Appl* [Internet]. 2016 Feb 1 [cited 2018 Jul 31];32(2):142–63. Available from: <http://doi.wiley.com/10.1002/rra.2928>
136. Zen S, Gurnell AM, Zolezzi G, Surian N. Exploring the role of trees in the evolution of meander bends: The Tagliamento River, Italy. *Water Resour Res* [Internet]. 2017 Jul 1 [cited 2018 Jul 31];53(7):5943–62. Available from: <http://doi.wiley.com/10.1002/2017WR020561>
137. Greenwood P, Kuhn NJ. Does the invasive plant, *Impatiens glandulifera*, promote soil erosion along the riparian zone? An investigation on a small watercourse in northwest Switzerland. *J Soils Sediments* [Internet]. 2014 Mar 20 [cited 2018 Jul 31];14(3):637–50. Available from: <http://link.springer.com/10.1007/s11368-013-0825-9>
138. Greenwood P, Baumann P, Pulley S, Kuhn NJ. The invasive alien plant, *Impatiens glandulifera* (Himalayan Balsam), and increased soil erosion: causation or association? Case studies from a river system in Switzerland and the UK. *J Soils Sediments* [Internet]. 2018 Jun 4 [cited 2018 Jul 31];1–15. Available from: <http://link.springer.com/10.1007/s11368-018-2041-0>
139. Albertson LK, Allen DC, Trexler JC. Meta-analysis: Abundance, behavior, and hydraulic energy shape biotic effects on sediment transport in streams. *Ecology* [Internet]. 2015 May [cited 2018 Aug 2];96(5):1329–39. Available from: <http://doi.wiley.com/10.1890/13-2138.1>
140. Statzner B. Geomorphological implications of engineering bed sediments by lotic animals. *Geomorphology* [Internet]. 2012 Jul 1 [cited 2018 Aug 1];157–158:49–65. Available from: <https://www.sciencedirect.com/science/article/pii/S0169555X11003060>
141. Albertson LK, Daniels MD. Effects of invasive crayfish on fine sediment accumulation, gravel movement, and macroinvertebrate communities. *Freshw Sci* [Internet]. 2016 Jun 12 [cited 2018 Aug 1];35(2):644–53. Available from: <http://www.journals.uchicago.edu/doi/10.1086/685860>
142. Rice SP, Johnson MF, Extence C, Reeds J, Longstaff H. Diel patterns of suspended sediment flux and the zoogeomorphic agency of invasive crayfish. *Cuad Investig Geográfica* [Internet]. 2014 Mar 6 [cited 2018 Aug 1];40(1):7. Available from: <https://publicaciones.unirioja.es/ojs/index.php/cig/article/view/2508>
143. Holdich DM, James J, Jackson C, Peay S. The North American signal crayfish, with particular reference to its success as an invasive species in Great Britain. *Ethol Ecol Evol* [Internet]. 2014 Jul 3 [cited 2018 Aug 1];26(2–3):232–62. Available from: <http://www.tandfonline.com/doi/abs/10.1080/03949370.2014.903380>
144. Pringle CM, Blake GA, Covich AP, Buzby KM, Finley A. Effects of omnivorous shrimp in a montane tropical stream: sediment removal, disturbance of sessile invertebrates and enhancement of understory algal biomass. *Oecologia* [Internet]. 1993 Feb [cited 2018 Aug 2];93(1):1–11. Available

- 1  
2  
3 from: <http://link.springer.com/10.1007/BF00321183>  
4
- 5 145. Visoni SBC, Moulton TP. Effects of shrimp on periphyton and sediments in Atlantic forest  
6 streams: an exclusion experiment. *Acta Limnol Bras*. 2003;15(1):19–26.  
7
- 8 146. Statzner B, Fuchs U, Higler LWG. Sand erosion by mobile predaceous stream insects: Implications  
9 for ecology and hydrology. *Water Resour Res* [Internet]. 1996 Jul 1 [cited 2018 Aug  
10 1];32(7):2279–87. Available from: <http://doi.wiley.com/10.1029/96WR00977>  
11
- 12 147. Soluk DA, Craig DA. Digging with a vortex: Flow manipulation facilitates prey capture by a  
13 predatory stream mayfly. *Limnol Oceanogr* [Internet]. 1990 Jul 1 [cited 2018 Aug 1];35(5):1201–  
14 6. Available from: <http://doi.wiley.com/10.4319/lo.1990.35.5.1201>  
15
- 16 148. Mermillod-Blondin F, Gérino M, Degrange V. Functional Diversity among 3 Detritivorous  
17 Hyporheic Invertebrates: An Experimental Study in Microcosms. *J North Am Benthol Soc*  
18 [Internet]. 2002 Mar 21 [cited 2018 Aug 1];21(1):132–49. Available from:  
19 <http://www.journals.uchicago.edu/doi/10.2307/1468305>  
20
- 21 149. Mermillod-Blondin F, Gaudet JP, Gérino M, Desrosiers G, Creuzé des Châtelliers M. Influence of  
22 macroinvertebrates on physico-chemical and microbial processes in hyporheic sediments. *Hydrol*  
23 *Process* [Internet]. 2003 Mar 1 [cited 2018 Aug 1];17(4):779–94. Available from:  
24 <http://doi.wiley.com/10.1002/hyp.1165>  
25
- 26 150. Mermillod-Blondin F, Gaudet JP, Gerino M, Desrosiers G, Jose J, Creuzé Des Châtelliers M.  
27 Relative influence of bioturbation and predation on organic matter processing in river sediments:  
28 A microcosm experiment. *Freshw Biol* [Internet]. 2004 Jul 1 [cited 2018 Aug 1];49(7):895–912.  
29 Available from: <http://doi.wiley.com/10.1111/j.1365-2427.2004.01233.x>  
30
- 31 151. Mermillod-Blondin F, Rosenberg R. Ecosystem engineering: The impact of bioturbation on  
32 biogeochemical processes in marine and freshwater benthic habitats. *Aquat Sci* [Internet]. 2006  
33 Dec 4 [cited 2018 Aug 1];68(4):434–42. Available from: [http://link.springer.com/10.1007/s00027-](http://link.springer.com/10.1007/s00027-006-0858-x)  
34 [006-0858-x](http://link.springer.com/10.1007/s00027-006-0858-x)  
35
- 36 152. Nogaro G, Mermillod-Blondin F, François-Carcaillet F, Gaudet JP, Lafont M, Gibert J. Invertebrate  
37 bioturbation can reduce the clogging of sediment: An experimental study using infiltration  
38 sediment columns. *Freshw Biol* [Internet]. 2006 Aug 1 [cited 2018 Aug 1];51(8):1458–73.  
39 Available from: <http://doi.wiley.com/10.1111/j.1365-2427.2006.01577.x>  
40
- 41 153. Nogaro G, Mermillod-Blondin F, Valett MH, François-Carcaillet F, Gaudet JP, Lafont M, et al.  
42 Ecosystem engineering at the sediment-water interface: Bioturbation and consumer-substrate  
43 interaction. *Oecologia* [Internet]. 2009 Aug 22 [cited 2018 Aug 1];161(1):125–38. Available from:  
44 <http://link.springer.com/10.1007/s00442-009-1365-2>  
45
- 46 154. Zimmerman GF, de Szalay FA. Influence of unionid mussels (Mollusca: Unionidae) on sediment  
47 stability: an artificial stream study. *Fundam Appl Limnol / Arch für Hydrobiol* [Internet]. 2007  
48 [cited 2018 Aug 2];168(4):299–306. Available from:  
49 [http://openurl.ingenta.com/content/xref?genre=article&issn=1863-](http://openurl.ingenta.com/content/xref?genre=article&issn=1863-9135&volume=168&issue=4&spage=299)  
50 [9135&volume=168&issue=4&spage=299](http://openurl.ingenta.com/content/xref?genre=article&issn=1863-9135&volume=168&issue=4&spage=299)  
51
- 52 155. Allen JRL. Experiments on the settling, overturning and entrainment of bivalve shells and related  
53 models. *Sedimentology* [Internet]. 1984 Apr 1 [cited 2018 Aug 2];31(2):227–50. Available from:  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 <http://doi.wiley.com/10.1111/j.1365-3091.1984.tb01961.x>  
4
- 5 156. Nogaro G, Mermillod-Blondin F, Montuelle B, Boisson JC, Gibert J. Chironomid larvae stimulate  
6 biogeochemical and microbial processes in a riverbed covered with fine sediment. *Aquat Sci*  
7 [Internet]. 2008 May 5 [cited 2018 Aug 2];70(2):156–68. Available from:  
8 <http://link.springer.com/10.1007/s00027-007-7032-y>  
9
- 10 157. Statzner B, Arens MF, Champagne JY, Morel R, Herouin E. Silk-producing stream insects and  
11 gravel erosion: Significant biological effects on critical shear stress. *Water Resour Res* [Internet].  
12 1999 Nov 1 [cited 2018 Aug 1];35(11):3495–506. Available from:  
13 <http://doi.wiley.com/10.1029/1999WR900196>  
14
- 15 158. Takao A, Negishi JN, Nunokawa M, Gomi T, Nakahara O. Potential influences of a net-spinning  
16 caddisfly (Trichoptera: *Stenopsyche marmorata*) on stream substratum stability in heterogeneous  
17 field environments. *Benthol Soc* [Internet]. 2006 [cited 2018 Aug 2];25(3):545–55. Available from:  
18 [https://www.journals.uchicago.edu/doi/abs/10.1899/0887-](https://www.journals.uchicago.edu/doi/abs/10.1899/0887-3593(2006)25%5B545%3APIOANC%5D2.O.CO%3B2)  
19 [3593\(2006\)25%5B545%3APIOANC%5D2.O.CO%3B2](https://www.journals.uchicago.edu/doi/abs/10.1899/0887-3593(2006)25%5B545%3APIOANC%5D2.O.CO%3B2)  
20  
21
- 22 159. Nunokawa M, Gomi T, Negishi JN, Nakahara O. A new method to measure substrate coherent  
23 strength of *Stenopsyche marmorata*. *Landsc Ecol Eng* [Internet]. 2008 Nov 19 [cited 2018 Aug  
24 2];4(2):125–31. Available from: <http://link.springer.com/10.1007/s11355-008-0044-5>  
25
- 26 160. Cardinale BJ, Gelmann ER, Palmer MA. Net spinning caddisflies as stream ecosystem engineers:  
27 The influence of Hydropsyche on benthic substrate stability. *Funct Ecol* [Internet]. 2004 Jun 1  
28 [cited 2018 Aug 1];18(3):381–7. Available from: [http://doi.wiley.com/10.1111/j.0269-](http://doi.wiley.com/10.1111/j.0269-8463.2004.00865.x)  
29 [8463.2004.00865.x](http://doi.wiley.com/10.1111/j.0269-8463.2004.00865.x)  
30
- 31 161. Albertson LK, Cardinale BJ, Sklar LS. Non-additive increases in sediment stability are generated by  
32 macroinvertebrate species interactions in laboratory streams. Fukami T, editor. *PLoS One*  
33 [Internet]. 2014 Aug 7 [cited 2018 Aug 2];9(8):e103417. Available from:  
34 <http://dx.plos.org/10.1371/journal.pone.0103417>  
35  
36
- 37 162. Johnson MF, Reid I, Rice SP, Wood PJ. Stabilization of fine gravels by net-spinning caddisfly  
38 larvae. *Earth Surf Process Landforms* [Internet]. 2009 Mar 15 [cited 2018 Aug 1];34(3):413–23.  
39 Available from: <http://doi.wiley.com/10.1002/esp.1750>  
40
- 41 163. Statzner B, Mérigoux S, Leichtfried M. Mineral grains in caddisfly pupal cases and streambed  
42 sediments: Resource use and its limitation through conflicting resource requirements. *Limnol*  
43 *Oceanogr* [Internet]. 2005 Mar [cited 2018 Aug 2];50(2):713–21. Available from:  
44 <http://doi.wiley.com/10.4319/lo.2005.50.2.0713>  
45
- 46 164. Statzner B. Mineral grains in caddisfly pupal cases and streambed sediments- assessing resource  
47 use and its limitations across various river types. *cambridge.org* [Internet]. [cited 2018 Aug 2];  
48 Available from: [https://www.cambridge.org/core/journals/annales-de-limnologie-international-](https://www.cambridge.org/core/journals/annales-de-limnologie-international-journal-of-limnology/article/mineral-grains-in-caddisfly-pupal-cases-and-streambed-sediments-assessing-resource-use-and-its-limitation-across-various-river-types/EC6C90B23F2EA52DF)  
49 [journal-of-limnology/article/mineral-grains-in-caddisfly-pupal-cases-and-streambed-sediments-](https://www.cambridge.org/core/journals/annales-de-limnologie-international-journal-of-limnology/article/mineral-grains-in-caddisfly-pupal-cases-and-streambed-sediments-assessing-resource-use-and-its-limitation-across-various-river-types/EC6C90B23F2EA52DF)  
50 [assessing-resource-use-and-its-limitation-across-various-river-types/EC6C90B23F2EA52DF](https://www.cambridge.org/core/journals/annales-de-limnologie-international-journal-of-limnology/article/mineral-grains-in-caddisfly-pupal-cases-and-streambed-sediments-assessing-resource-use-and-its-limitation-across-various-river-types/EC6C90B23F2EA52DF)  
51  
52
- 53 165. Wharton G, Cotton JA, Wotton RS, Bass JAB, Heppell CM, Trimmer M, et al. Macrophytes and  
54 suspension-feeding invertebrates modify flows and fine sediments in the Frome and Piddle  
55 catchments, Dorset (UK). *J Hydrol* [Internet]. 2006 Oct 30 [cited 2018 Aug 1];330(1–2):171–84.  
56  
57  
58  
59  
60

- 1  
2  
3 Available from: <https://www.sciencedirect.com/science/article/pii/S0022169406001995>  
4
- 5 166. Wotton RS, Warren LL. Impacts of suspension feeders on the modification and transport of  
6 stream seston. *Fundam Appl Limnol*. 2007 Jul 1;169(3):231–6.  
7
- 8 167. Balian E V., Segers H, Martens K, L  v  que C. The Freshwater Animal Diversity Assessment: an  
9 overview of the results. In: *Freshwater Animal Diversity Assessment* [Internet]. Dordrecht:  
10 Springer Netherlands; [cited 2018 Aug 2]. p. 627–37. Available from:  
11 [http://link.springer.com/10.1007/978-1-4020-8259-7\\_61](http://link.springer.com/10.1007/978-1-4020-8259-7_61)  
12
- 13 168. Strayer DL, Dudgeon D. Freshwater biodiversity conservation: recent progress and future  
14 challenges. *J North Am Benthol Soc* [Internet]. 2010 Mar [cited 2018 Aug 2];29(1):344–58.  
15 Available from: <http://www.journals.uchicago.edu/doi/10.1899/08-171.1>  
16
- 17 169. WWF (World Wide Fund for Nature). *Living Planet Report 2016. Risk and resilience in a new era*  
18 [Internet]. 2016 [cited 2018 Aug 2]. Available from:  
19 [http://awsassets.panda.org/downloads/lpr\\_living\\_planet\\_report\\_2016.pdf](http://awsassets.panda.org/downloads/lpr_living_planet_report_2016.pdf)  
20
- 21 170. Poff NL, Olden JD, Strayer DL. Climate Change and Freshwater Fauna Extinction Risk. In: *Saving a*  
22 *Million Species* [Internet]. Washington, DC: Island Press/Center for Resource Economics; 2012  
23 [cited 2018 Aug 2]. p. 309–36. Available from: [http://www.springerlink.com/index/10.5822/978-](http://www.springerlink.com/index/10.5822/978-1-61091-182-5_17)  
24 [1-61091-182-5\\_17](http://www.springerlink.com/index/10.5822/978-1-61091-182-5_17)  
25
- 26 171. Strayer DL. Challenges for freshwater invertebrate conservation. *J North Am Benthol Soc*  
27 [Internet]. 2006 [cited 2018 Aug 2];25(2):271–87. Available from:  
28 [http://www.journals.uchicago.edu/doi/10.1899/0887-3593\(2006\)25\[271:CFFIC\]2.0.CO;2](http://www.journals.uchicago.edu/doi/10.1899/0887-3593(2006)25[271:CFFIC]2.0.CO;2)  
29
- 30 172. Burner CJ. Characteristics of spawning nests of Columbia River salmon [Internet]. Vol. 52, U.S.  
31 Fish and Wildlife Service Fishery Bulletin. 1951 [cited 2018 Aug 2]. 94–110 p. Available from:  
32 [https://www.nwfsc.noaa.gov/assets/2/9171\\_07142017\\_163818\\_Burner.1951-Fish-Bull-61-52-](https://www.nwfsc.noaa.gov/assets/2/9171_07142017_163818_Burner.1951-Fish-Bull-61-52-1.pdf)  
33 [1.pdf](https://www.nwfsc.noaa.gov/assets/2/9171_07142017_163818_Burner.1951-Fish-Bull-61-52-1.pdf)  
34
- 35 173. DeVries P. Salmonid influences on rivers: A geomorphic fish tail. *Geomorphology* [Internet]. 2012  
36 [cited 2018 Aug 2];157–158:66–74. Available from:  
37 <https://www.sciencedirect.com/science/article/pii/S0169555X11003011>  
38
- 39 174. Rice S, Stoffel M, Turowski JM, Wolf A. Disturbance regimes at the interface of geomorphology  
40 and ecology. *Earth Surf Process Landforms* [Internet]. 2012 Dec 1 [cited 2018 Aug  
41 2];37(15):1678–82. Available from: <http://doi.wiley.com/10.1002/esp.3326>  
42
- 43 175. Rennie CD, Millar RG. Spatial variability of stream bed scour and fill: A comparison of scour depth  
44 in chum salmon (*Oncorhynchus keta*) redds and adjacent bed. *Can J Fish Aquat Sci* [Internet].  
45 2000 May [cited 2018 Aug 2];57(5):928–38. Available from: [http://narwhal.cisti.nrc.ca/cgi-](http://narwhal.cisti.nrc.ca/cgi-bin/rp/servlets/ppv/RPViewDoc?_handler_=HandleInitialGet&issn=0706652X&volume=57&issue=5&startPage=928%5Cnhttp://www.nrc.ca/cgi-bin/cisti/journals/rp/rp2*desc*e?cjfas)  
46 [bin/rp/servlets/ppv/RPViewDoc?\\_handler\\_=HandleInitialGet&issn=0706652X&volume=57&issue](http://narwhal.cisti.nrc.ca/cgi-bin/rp/servlets/ppv/RPViewDoc?_handler_=HandleInitialGet&issn=0706652X&volume=57&issue=5&startPage=928%5Cnhttp://www.nrc.ca/cgi-bin/cisti/journals/rp/rp2*desc*e?cjfas)  
47 [=5&startPage=928%5Cnhttp://www.nrc.ca/cgi-bin/cisti/journals/rp/rp2\\*desc\\*e?cjfas](http://narwhal.cisti.nrc.ca/cgi-bin/rp/servlets/ppv/RPViewDoc?_handler_=HandleInitialGet&issn=0706652X&volume=57&issue=5&startPage=928%5Cnhttp://www.nrc.ca/cgi-bin/cisti/journals/rp/rp2*desc*e?cjfas)  
48
- 49 176. Buxton TH, Buffington JM, Yager EM, Hassan MA, Fremier AK. The relative stability of salmon  
50 redds and unspawned streambeds. *Water Resour Res* [Internet]. 2015 Aug 1 [cited 2018 Aug  
51 2];51(8):6074–92. Available from: <http://doi.wiley.com/10.1002/2015WR016908>  
52
- 53 177. Montgomery DR, Buffington JM, Peterson NP, Schuett-Hames D, Quinn TP. Stream-bed scour,  
54  
55  
56  
57  
58  
59  
60

- egg burial depths, and the influence of salmonid spawning on bed surface mobility and embryo survival. *Can J Fish Aquat Sci* [Internet]. 1996 May [cited 2018 Aug 2];53(5):1061–70. Available from: <http://www.nrcresearchpress.com/doi/abs/10.1139/f96-028>
178. Hassan MA, Gottesfeld AS, Montgomery DR, Tunnicliffe JF, Clarke GKC, Wynn G, et al. Salmon-driven bed load transport and bed morphology in mountain streams. *Geophys Res Lett* [Internet]. 2008 Feb 22 [cited 2018 Aug 2];35(4):L04405. Available from: <http://doi.wiley.com/10.1029/2007GL032997>
179. Gottesfeld AS, Hassan MA, Tunnicliffe JF, Poirier RW. Sediment dispersion in salmon spawning streams: The influence of floods and salmon redd construction. *J Am Water Resour Assoc* [Internet]. 2004 Aug 1 [cited 2018 Aug 2];40(4):1071–86. Available from: <http://doi.wiley.com/10.1111/j.1752-1688.2004.tb01068.x>
180. Fremier AK, Yanites BJ, Yager EM. Sex that moves mountains: The influence of spawning fish on river profiles over geologic timescales. *Geomorphology* [Internet]. 2017 [cited 2018 Aug 2]; Available from: <https://www.sciencedirect.com/science/article/pii/S0169555X17302155>
181. Wright RM. Aspects of the ecology of bream, *Abramis brama* (L.), in a gravel pit lake and the effects of reducing the population density. *J Fish Biol* [Internet]. 1990 Jan 24 [cited 2018 Aug 2];37(4):629–34. Available from: <http://doi.wiley.com/10.1111/j.1095-8649.1990.tb05896.x>
182. Huser BJ, Bajer PG, Chizinski CJ, Sorensen PW. Effects of common carp (*Cyprinus carpio*) on sediment mixing depth and mobile phosphorus mass in the active sediment layer of a shallow lake. *Hydrobiologia* [Internet]. 2016 Jan 12 [cited 2018 Aug 2];763(1):23–33. Available from: <http://link.springer.com/10.1007/s10750-015-2356-4>
183. Pledger AG, Rice SP, Millett J. Reduced bed material stability and increased bedload transport caused by foraging fish: A flume study with juvenile Barbel (*Barbus barbus*). *Earth Surf Process Landforms* [Internet]. 2014 May 15 [cited 2018 Aug 2];39(11):1500–13. Available from: <http://doi.wiley.com/10.1002/esp.3592>
184. Barton DR, Kelton N, Eedy RI. The effects of carp (*Cyprinus carpio* L.) on sediment export from a small urban impoundment. *J Aquat Ecosyst Stress Recover (Formerly J Aquat Ecosyst Heal)* [Internet]. 2000 [cited 2018 Aug 2];8(2):155–9. Available from: <https://link.springer.com/article/10.1023/A:1011423432727>
185. Breukelaar AW, Lammens EHRR, Breteler JGPK, Tátri I. Effects of benthivorous bream (*Abramis brama*) and carp (*Cyprinus carpio*) on sediment resuspension and concentrations of nutrients and chlorophyll a. *Freshw Biol* [Internet]. 1994 Aug [cited 2018 Aug 2];32(1):113–21. Available from: <http://doi.wiley.com/10.1111/j.1365-2427.1994.tb00871.x>
186. Cahn AR. The Effect of Carp on a Small Lake : the Carp as a Dominant. *Ecology* [Internet]. 1929 Jul [cited 2018 Aug 2];10(3):271–4. Available from: <http://doi.wiley.com/10.2307/1929502>
187. Parkos III JJ, Santucci, Jr. VJ, Wahl DH. Effects of adult common carp (*Cyprinus carpio*) on multiple trophic levels in shallow mesocosms. *Can J Fish Aquat Sci* [Internet]. 2003 [cited 2018 Aug 2];60(2):182–92. Available from: <http://www.nrcresearchpress.com/doi/abs/10.1139/f03-011>
188. Pledger AG, Rice SP, Millett J. Bed disturbance via foraging fish increases bedload transport during subsequent high flows and is controlled by fish size and species. *Geomorphology*

- [Internet]. 2016 [cited 2018 Aug 2];253:83–93. Available from: <https://www.sciencedirect.com/science/article/pii/S0169555X15301562>
189. Statzner B, Sagnes P, Champagne JY, Viboud S. Contribution of benthic fish to the patch dynamics of gravel and sand transport in streams. *Water Resour Res* [Internet]. 2003 Nov [cited 2018 Aug 2];39(11). Available from: <http://doi.wiley.com/10.1029/2003WR002270>
190. Moore JW, Schindler DE, Carter JL, Fox J, Griffiths J, Holtgrieve GW. Biotic control of stream fluxes: spawning salmon drive nutrient and matter export. *Ecology* [Internet]. 2007;88(5):1278–91. Available from: <https://esajournals.onlinelibrary.wiley.com/doi/abs/10.1890/06-0782>
191. Cak AD, Chaloner DT, Lamberti GA. Effects of spawning salmon on dissolved nutrients and epilithon in coupled stream-estuary systems of southeastern Alaska. *Aquat Sci* [Internet]. 2008;70(2):169–178. Available from: <https://link.springer.com/article/10.1007/s00027-008-8090-5>
192. Usseglio-Polatera P, Bournaud M, Richoux P, Tachet H. Biological and ecological traits of benthic freshwater macroinvertebrates: Relationships and definition of groups with similar traits. *Freshw Biol* [Internet]. 2000 Feb 1 [cited 2018 Aug 1];43(2):175–205. Available from: <http://doi.wiley.com/10.1046/j.1365-2427.2000.00535.x>
193. Lamouroux N, Dolédec S, Gayraud S. Biological traits of stream macroinvertebrate communities: effects of microhabitat, reach, and basin filters. *J North Am Benthol Soc* [Internet]. 2004 Sep [cited 2018 Aug 1];23(3):449–66. Available from: [http://www.journals.uchicago.edu/doi/10.1899/0887-3593\(2004\)023%3C0449:BTOSMC%3E2.0.CO;2](http://www.journals.uchicago.edu/doi/10.1899/0887-3593(2004)023%3C0449:BTOSMC%3E2.0.CO;2)
194. Altermatt F. Diversity in riverine metacommunities: A network perspective. *Aquat Ecol* [Internet]. 2013 Sep 14 [cited 2018 Jul 26];47(3):365–77. Available from: <http://link.springer.com/10.1007/s10452-013-9450-3>
195. Prunier JG, Dubut V, Loot G, Tudesque L, Blanchet S. The relative contribution of river network structure and anthropogenic stressors to spatial patterns of genetic diversity in two freshwater fishes: A multiple-stressors approach. *Freshw Biol* [Internet]. 2018 Jan 1 [cited 2018 Aug 1];63(1):6–21. Available from: <http://doi.wiley.com/10.1111/fwb.13034>
196. Hanski I. Metapopulation dynamics. *Nature* [Internet]. 1998 Nov 5 [cited 2018 Aug 1];396(6706):41–9. Available from: <http://www.nature.com/articles/23876>
197. Poff NL. Landscape Filters and Species Traits: Towards Mechanistic Understanding and Prediction in Stream Ecology. *J North Am Benthol Soc* [Internet]. 1997 Jun [cited 2018 Aug 2];16(2):391–409. Available from: <http://www.journals.uchicago.edu/doi/10.2307/1468026>
198. Leibold MA, Holyoak M, Mouquet N, Amarasekare P, Chase JM, Hoopes MF, et al. The metacommunity concept: A framework for multi-scale community ecology. *Ecol Lett* [Internet]. 2004 Jun 4 [cited 2018 Aug 1];7(7):601–13. Available from: <http://doi.wiley.com/10.1111/j.1461-0248.2004.00608.x>
199. Vannote RL, Minshall GW, Cummins KW, Sedell JR, Cushing CE. The River Continuum Concept. *Can J Fish Aquat Sci* [Internet]. 1980 Jan [cited 2018 Aug 1];37(1):130–7. Available from: <http://www.nrcresearchpress.com/doi/abs/10.1139/f80-017>

- 1  
2  
3 200. Junk W, Bayley PB, Sparks RE. The flood pulse concept in river-floodplain-systems. *Can Spec Publ Fish Aquat Sci.* 1989;106:110–27.
- 4  
5  
6 201. Ward J V., Stanford JA. The serial discontinuity concept: Extending the model to floodplain rivers. *Regul Rivers Res Manag* [Internet]. 1995 Aug [cited 2018 Aug 1];10(2–4):159–68. Available from: <http://doi.wiley.com/10.1002/rrr.3450100211>
- 7  
8  
9  
10 202. Winegardner AK, Jones BK, Ng ISY, Siqueira T, Cottenie K. The terminology of metacommunity ecology. *Trends Ecol Evol* [Internet]. 2012 May 1 [cited 2018 Aug 1];27(5):253–4. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/22325446>
- 11  
12  
13  
14 203. Gaston KJ. Global patterns in biodiversity. *Nature* [Internet]. 2000 May 11 [cited 2018 Aug 1];405(6783):220–7. Available from: <http://www.nature.com/articles/35012228>
- 15  
16  
17  
18 204. Brown LE, Khamis K, Wilkes M, Blaen P, Brittain JE, Carrivick JL, et al. Functional diversity and community assembly of river invertebrates show globally consistent responses to decreasing glacier cover. *Nat Ecol Evol* [Internet]. 2018 Feb 18 [cited 2018 Aug 2];2(2):325–33. Available from: <http://www.nature.com/articles/s41559-017-0426-x>
- 19  
20  
21  
22  
23 205. Radinger J, Essl F, Hölker F, Horký P, Slavík O, Wolter C. The future distribution of river fish: The complex interplay of climate and land use changes, species dispersal and movement barriers. *Glob Chang Biol* [Internet]. 2017 Nov 1 [cited 2018 Aug 2];23(11):4970–86. Available from: <http://doi.wiley.com/10.1111/gcb.13760>
- 24  
25  
26  
27  
28 206. Karger DN, Cord AF, Kessler M, Kreft H, Kühn I, Pompe S, et al. Delineating probabilistic species pools in ecology and biogeography. *Glob Ecol Biogeogr* [Internet]. 2016 Apr 1 [cited 2018 Aug 1];25(4):489–501. Available from: <http://doi.wiley.com/10.1111/geb.12422>
- 29  
30  
31  
32 207. Vermaat JE, De Bruyne RJ. Factors limiting the distribution of submerged waterplants in the lowland River Vecht (The Netherlands). *Freshw Biol* [Internet]. 1993 Aug 1 [cited 2018 Aug 2];30(1):147–57. Available from: <http://doi.wiley.com/10.1111/j.1365-2427.1993.tb00795.x>
- 33  
34  
35  
36  
37 208. LaPerriere JD, Bjerklie DM, Simmons RC, Van Nieuwenhuys EE, Wagener SM, Reynolds JB. Effects of gold placer mining on interior Alaskan stream ecosystems. *Manag water Resour Alaska's Dev proceedings* University Alaska, Inst Water Resour Rep IWR-105, Fairbanks, AK [Internet]. 1983 [cited 2018 Aug 2];12–34. Available from: <https://core.ac.uk/download/pdf/130253707.pdf#page=251>
- 38  
39  
40  
41  
42  
43 209. Rivier B, Segulier J. Physical and biological effects of gravel extraction in river beds. *Habitat Modif Freshw Fish Proc a Symp Eur Inl Fish Advis Comm* [Internet]. 1985 [cited 2018 Aug 2];131–45. Available from: <http://agris.fao.org/agris-search/search.do?recordID=XF2016073716>
- 44  
45  
46  
47  
48 210. Van Nieuwenhuys EE, LaPerriere JD. Effects of Placer Gold Mining on Primary Production in Subarctic Streams of Alaska. *JAWRA J Am Water Resour Assoc* [Internet]. 1986 Feb [cited 2018 Aug 2];22(1):91–9. Available from: <http://doi.wiley.com/10.1111/j.1752-1688.1986.tb01864.x>
- 49  
50  
51  
52 211. Lloyd DS, Koenings JP, Laperriere JD. Effects of Turbidity in Fresh Waters of Alaska. *North Am J Fish Manag* [Internet]. 1987 Jan [cited 2018 Aug 2];7(1):18–33. Available from: <http://www.tandfonline.com/doi/abs/10.1577/1548-8659%281987%297%3C18%3AEOTIFW%3E2.0.CO%3B2>
- 53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 212. Harper MA. Migration rhythm of the benthic diatom *pinnularia viridis* on pond silt (Note). *New Zealand J Mar Freshw Res* [Internet]. 1976 Jun 30 [cited 2018 Aug 2];10(2):381–4. Available from: <http://www.tandfonline.com/doi/abs/10.1080/00288330.1976.9515622>
- 7 213. Hay SI, Maitland TC, Paterson DM. The speed of diatom migration through natural and artificial substrata. *Diatom Res* [Internet]. 1993 Nov [cited 2018 Aug 2];8(2):371–84. Available from: <http://www.tandfonline.com/doi/abs/10.1080/0269249X.1993.9705268>
- 11 214. Yamada H, Nakamura F. Effect of fine sediment deposition and channel works on periphyton biomass in the Makomanai River, northern Japan. *River Res Appl* [Internet]. 2002 Sep [cited 2018 Aug 2];18(5):481–93. Available from: <http://doi.wiley.com/10.1002/rra.688>
- 16 215. Dickman MD, Peart MR, Wai-Shu Yim W. Benthic Diatoms as Indicators of Stream Sediment Concentration in Hong Kong. *Int Rev Hydrobiol* [Internet]. 2005 Aug [cited 2018 Aug 2];90(4):412–21. Available from: <http://doi.wiley.com/10.1002/iroh.200410806>
- 20 216. Izagirre O, Serra A, Guasch H, Elosegi A. Effects of sediment deposition on periphytic biomass, photosynthetic activity and algal community structure. *Sci Total Environ* [Internet]. 2009 Oct 15 [cited 2018 Aug 2];407(21):5694–700. Available from: <https://www.sciencedirect.com/science/article/pii/S0048969709006299>
- 25 217. Jones JI, Douthwright TA, Arnold A, Duerdoth CP, Murphy JF, Edwards FK, et al. Diatoms as indicators of fine sediment stress. *Ecohydrology* [Internet]. 2017 Jul 1 [cited 2018 Aug 2];10(5):e1832. Available from: <http://doi.wiley.com/10.1002/eco.1832>
- 29 218. Biggs BJF, Kilroy C, Lowe RL. Periphyton development in three valley segments of a New Zealand grassland river: test of a habitat matrix conceptual model within a catchment. *Arch für Hydrobiol* [Internet]. 1998 Oct 20 [cited 2018 Aug 2];143(2):147–77. Available from: [http://www.schweizerbart.de/papers/fal/detail/143/86229/Periphyton\\_development\\_in\\_three\\_valley\\_segments\\_of?af=crossref](http://www.schweizerbart.de/papers/fal/detail/143/86229/Periphyton_development_in_three_valley_segments_of?af=crossref)
- 36 219. Biggs BJF, Smith RA. Taxonomic richness of stream benthic algae: Effects of flood disturbance and nutrients. *Limnol Oceanogr* [Internet]. 2002 Jul 1 [cited 2018 Aug 2];47(4):1175–86. Available from: <http://doi.wiley.com/10.4319/lo.2002.47.4.1175>
- 40 220. Matthaei CD, Guggelberger C, Hubre H. Local disturbance history affects patchiness of benthic river algae. *Freshw Biol* [Internet]. 2003 Sep 1 [cited 2018 Aug 2];48(9):1514–26. Available from: <http://doi.wiley.com/10.1046/j.1365-2427.2003.01103.x>
- 44 221. Francoeur SN, Biggs BJF. Short-term effects of elevated velocity and sediment abrasion on benthic algal communities. *Hydrobiologia* [Internet]. 2006 May [cited 2018 Aug 2];561(1):59–69. Available from: <http://link.springer.com/10.1007/s10750-005-1604-4>
- 48 222. Delgado M, De Jonge VN, Peletier H. Effect of sand movement on the growth of benthic diatoms. *J Exp Mar Bio Ecol* [Internet]. 1991 Mar 8 [cited 2018 Aug 2];145(2):221–31. Available from: <https://www.sciencedirect.com/science/article/pii/002209819190177X>
- 52 223. Jewson DH, Lowry SF, Bowen R. Co-existence and survival of diatoms on sand grains. *Eur J Phycol* [Internet]. 2006 May [cited 2018 Aug 2];41(2):131–46. Available from: <http://www.tandfonline.com/doi/abs/10.1080/09670260600652903>
- 56  
57  
58  
59  
60

- 1  
2  
3 224. Vilbaste S, Truu J. Distribution of benthic diatoms in relation to environmental variables in  
4 lowland streams. *Hydrobiologia* [Internet]. 2003 [cited 2018 Aug 2];493(1/3):81–93. Available  
5 from: <http://link.springer.com/10.1023/A:1025401716441>  
6
- 7 225. O’Hare MT, Mountford JO, Maroto J, Gunn IDM. Plant Traits Relevant To Fluvial Geomorphology  
8 and Hydrological Interactions. *River Res Appl* [Internet]. 2016 Feb 1 [cited 2018 Aug 1];32(2):179–  
9 89. Available from: <http://doi.wiley.com/10.1002/rra.2940>  
10
- 11 226. Riis T, Sand-Jensen K. Dispersal of plant fragments in small streams. *Freshw Biol* [Internet]. 2006  
12 Feb 1 [cited 2018 Aug 1];51(2):274–86. Available from: <http://doi.wiley.com/10.1111/j.1365->  
13 2427.2005.01496.x  
14
- 15 227. Riis T, Madsen T V., Sennels RSH. Regeneration, colonisation and growth rates of allofragments in  
16 four common stream plants. *Aquat Bot* [Internet]. 2009 Feb 1 [cited 2018 Aug 1];90(2):209–12.  
17 Available from: <https://www.sciencedirect.com/science/article/pii/S0304377008001423>  
18
- 19 228. Trimmer M, Sanders IA, Heppell CM. Carbon and nitrogen cycling in a vegetated lowland chalk  
20 river impacted by sediment. *Hydrol Process* [Internet]. 2009 Jul 15 [cited 2018 Aug  
21 1];23(15):2225–38. Available from: <http://doi.wiley.com/10.1002/hyp.7276>  
22
- 23 229. Francis R a., Gurnell AM. Initial establishment of vegetative fragments within the active zone of a  
24 braided gravel-bed river (River Tagliamento, NE Italy). *Wetlands* [Internet]. 2006 [cited 2018 Aug  
25 1];26(3):641–8. Available from: <https://link.springer.com/article/10.1672/0277->  
26 5212(2006)26[641:IEOVFW]2.0.CO;2  
27
- 28 230. Moggridge HL, Gurnell AM. Controls on the sexual and asexual regeneration of Salicaceae along a  
29 highly dynamic, braided river system. *Aquat Sci* [Internet]. 2009 Oct 17 [cited 2018 Aug  
30 1];71(3):305–17. Available from: <http://link.springer.com/10.1007/s00027-009-9193-3>  
31
- 32 231. Francis RA, Gurnell AM, Petts GE, Edwards PJ. Survival and growth responses of *Populus nigra*,  
33 *Salix elaeagnos* and *Alnus incana* cuttings to varying levels of hydric stress. *For Ecol Manage*  
34 [Internet]. 2005 May 16 [cited 2018 Aug 1];210(1–3):291–301. Available from:  
35 <https://www.sciencedirect.com/science/article/pii/S0378112705001283>  
36
- 37 232. Dawson FH. The seasonal effects of aquatic plant growth on the flow of water in a stream. 5th Int  
38 Symp Aquat Weeds Eur Weed Res Soc [Internet]. 1978 [cited 2018 Aug 2];71–8. Available from:  
39 <http://agris.fao.org/agris-search/search.do?recordID=XE7980599>  
40
- 41 233. Kleeberg A, Köhler J, Sukhodolova T, Sukhodolov A. Effects of aquatic macrophytes on organic  
42 matter deposition, resuspension and phosphorus entrainment in a lowland river. *Freshw Biol*  
43 [Internet]. 2010 Feb 1 [cited 2018 Aug 2];55(2):326–45. Available from:  
44 <http://doi.wiley.com/10.1111/j.1365-2427.2009.02277.x>  
45
- 46 234. Dawson FH, Castellano E, Ladle M. The concept of species succession in relation to river  
47 vegetation and management. *Verhandlungen Int Verein für Theor und Angew Limnol* [Internet].  
48 1978 Oct [cited 2018 Aug 1];20(JANUARY 1978):1451–6. Available from:  
49 <https://www.tandfonline.com/doi/full/10.1080/03680770.1977.11896707>  
50
- 51 235. Sand-Jensen K. Macrophytes as biological engineers in the ecology of Danish streams. *Freshw*  
52 *Biol Priorities Dev Danish Res* [Internet]. 1997 [cited 2018 Aug 2];74–101. Available from:  
53 <http://www.forskningsdatabasen.dk/en/catalog/2398156347>  
54
- 55  
56  
57  
58  
59  
60

- 1  
2  
3 236. Grime JP. *Plant Strategies, Vegetation Processes, and Ecosystem Properties*. Second ed. New  
4 York: John Wiley & Sons; 2001. 417 p.  
5
- 6 237. Jones JI, Murphy JF, Anthony SG, Arnold A, Blackburn JH, Duerdoth CP, et al. Do agri-environment  
7 schemes result in improved water quality? McKenzie A, editor. *J Appl Ecol* [Internet]. 2017 Apr 1  
8 [cited 2018 Aug 2];54(2):537–46. Available from: <http://doi.wiley.com/10.1111/1365-2664.12780>  
9
- 10 238. Grown I, Murphy JF, Jones JI. The effects of altered flow and bed sediment on  
11 macroinvertebrates in stream mesocosms. *Mar Freshw Res* [Internet]. 2017 Mar 21 [cited 2018  
12 Aug 6];68(3):496–505. Available from: <http://www.publish.csiro.au/?paper=MF15160>  
13
- 14 239. Carl Richards KLB. Influence of fine sediment on macroinvertebrate colonization of surface and  
15 hyporheic stream substrates. *Gt Basin Nat* [Internet]. 1994 [cited 2018 Aug 1];54(2):106–13.  
16 Available from: <http://www.jstor.org/stable/41712819>  
17
- 18 240. Descloux S, Datry T, Marmonier P. Benthic and hyporheic invertebrate assemblages along a  
19 gradient of increasing streambed colmation by fine sediment. *Aquat Sci* [Internet]. 2013 Oct 30  
20 [cited 2018 Aug 1];75(4):493–507. Available from: [http://link.springer.com/10.1007/s00027-013-](http://link.springer.com/10.1007/s00027-013-0295-6)  
21 [0295-6](http://link.springer.com/10.1007/s00027-013-0295-6)  
22  
23
- 24 241. Larsen S, Vaughan IP, Ormerod SJ. Scale-dependent effects of fine sediments on temperate  
25 headwater invertebrates. *Freshw Biol* [Internet]. 2009 Jan 1 [cited 2018 Aug 2];54(1):203–19.  
26 Available from: <http://doi.wiley.com/10.1111/j.1365-2427.2008.02093.x>  
27
- 28 242. Townsend CR, Hildrew AG. Species traits in relation to a habitat templet for river systems. *Freshw*  
29 *Biol* [Internet]. 1994 Jun 1 [cited 2018 Aug 1];31(3):265–75. Available from:  
30 <http://doi.wiley.com/10.1111/j.1365-2427.1994.tb01740.x>  
31
- 32 243. Doretto A, Bona F, Piano E, Zanin I, Eandi AC, Fenoglio S. Trophic availability buffers the  
33 detrimental effects of clogging in an alpine stream. *Sci Total Environ* [Internet]. 2017 Aug 15  
34 [cited 2018 Aug 1];592:503–11. Available from:  
35 <https://www.sciencedirect.com/science/article/pii/S0048969717306228>  
36  
37
- 38 244. Wilkes MA, Mckenzie M, Murphy JF, Chadd RP. Assessing the Mechanistic Basis for Fine Sediment  
39 Biomonitoring: Inconsistencies among the Literature, Traits and Indices. *River Res Appl* [Internet].  
40 2017 Dec 1 [cited 2018 Aug 1];33(10):1618–29. Available from:  
41 <http://doi.wiley.com/10.1002/rra.3139>  
42
- 43 245. Couceiro SRM, Hamada N, Forsberg BR, Padovesi-Fonseca C. Effects of anthropogenic silt on  
44 aquatic macroinvertebrates and abiotic variables in streams in the Brazilian Amazon. *J Soils*  
45 *Sediments* [Internet]. 2009 Jan 10 [cited 2018 Aug 1];10(1):89–103. Available from:  
46 <http://link.springer.com/10.1007/s11368-009-0148-z>  
47
- 48 246. Descloux S, Datry T, Usseglio-Polatera P. Trait-based structure of invertebrates along a gradient  
49 of sediment colmation: Benthos versus hyporheos responses. *Sci Total Environ* [Internet]. 2014  
50 Jan 1 [cited 2018 Jul 18];466–467:265–76. Available from:  
51 <https://www.sciencedirect.com/science/article/pii/S0048969713007341>  
52  
53
- 54 247. Bona F, Doretto A, Falasco E, La Morgia V, Piano E, Ajassa R, et al. Increased Sediment Loads in  
55 Alpine Streams: An Integrated Field Study. *River Res Appl* [Internet]. 2016 Jul [cited 2018 Jul  
56 18];32(6):1316–26. Available from: <http://doi.wiley.com/10.1002/rra.2941>  
57  
58  
59  
60

- 1  
2  
3 248. Relyea CD, Minshall GW, Danehy RJ. Stream Insects As Bioindicators of Fine Sediment. *Proc*  
4 *Water Environ Fed* [Internet]. 2000 Jan 1 [cited 2018 Aug 1];2000(6):663–86. Available from:  
5 <http://www.ingentaconnect.com/content/10.2175/193864700785150123>  
6  
7 249. Rabení CF, Doisy KE, Zweig LD. Stream invertebrate community functional responses to deposited  
8 sediment. *Aquat Sci* [Internet]. 2005 Dec 11 [cited 2018 Aug 1];67(4):395–402. Available from:  
9 <http://link.springer.com/10.1007/s00027-005-0793-2>  
10  
11 250. Mathers KL, Hill MJ, Wood PJ. Benthic and hyporheic macroinvertebrate distribution within the  
12 heads and tails of riffles during baseflow conditions. *Hydrobiologia* [Internet]. 2017 Jun 19 [cited  
13 2018 Aug 1];794(1):17–30. Available from: <http://link.springer.com/10.1007/s10750-017-3092-8>  
14  
15 251. Bo T, Fenoglio S, Malacarne G, Pessino M, Sgariboldi F. Effects of clogging on stream  
16 macroinvertebrates: An experimental approach. *Limnologica* [Internet]. 2007 May 30 [cited 2018  
17 Aug 1];37(2):186–92. Available from:  
18 <https://www.sciencedirect.com/science/article/pii/S0075951107000060>  
19  
20 252. Datry T, Lamouroux N, Thivin G, Descloux S, Baudoin JM. Estimation of Sediment Hydraulic  
21 Conductivity in River Reaches and its Potential Use to Evaluate Streambed Clogging. *River Res*  
22 *Appl* [Internet]. 2015 Sep 1 [cited 2018 Aug 1];31(7):880–91. Available from:  
23 <http://doi.wiley.com/10.1002/rra.2784>  
24  
25 253. Gayraud S, Philippe M. Does subsurface interstitial space influence general features and  
26 morphological traits of the benthic macroinvertebrate community in streams? *Arch Fur Hydrobiol*  
27 [Internet]. 2001 Jul 9 [cited 2018 Aug 1];151(4):667–86. Available from:  
28 [http://www.schweizerbart.de/papers/fal/detail/151/85869/Does\\_subsurface\\_interstitial\\_space\\_](http://www.schweizerbart.de/papers/fal/detail/151/85869/Does_subsurface_interstitial_space_influence_gener?af=crossref)  
29 [influence\\_gener?af=crossref](http://www.schweizerbart.de/papers/fal/detail/151/85869/Does_subsurface_interstitial_space_influence_gener?af=crossref)  
30  
31 254. Mathers KL, Rice SP, Wood PJ. Temporal effects of enhanced fine sediment loading on  
32 macroinvertebrate community structure and functional traits. *Sci Total Environ* [Internet]. 2017  
33 Dec 1 [cited 2018 Aug 1];599–600:513–22. Available from:  
34 <https://www.sciencedirect.com/science/article/pii/S0048969717309312>  
35  
36 255. Braccia A, Voshell JR. Environmental factors accounting for benthic macroinvertebrate  
37 assemblage structure at the sample scale in streams subjected to a gradient of cattle grazing.  
38 *Hydrobiologia* [Internet]. 2006 Dec 2 [cited 2018 Aug 1];573(1):55–73. Available from:  
39 <http://link.springer.com/10.1007/s10750-006-0257-2>  
40  
41 256. Cover MR, May CL, Dietrich WE, Resh VH. Quantitative linkages among sediment supply,  
42 streambed fine sediment, and benthic macroinvertebrates in northern California streams. *J North*  
43 *Am Benthol Soc* [Internet]. 2008 Mar 19 [cited 2018 Jul 19];27(1):135–49. Available from:  
44 <http://www.journals.uchicago.edu/doi/10.1899/07-032.1>  
45  
46 257. Buendia C, Gibbins CN, Vericat D, Batalla RJ, Douglas A. Detecting the structural and functional  
47 impacts of fine sediment on stream invertebrates. *Ecol Indic* [Internet]. 2013 Feb 1 [cited 2018 Jul  
48 19];25:184–96. Available from:  
49 <https://www.sciencedirect.com/science/article/pii/S1470160X12003512>  
50  
51 258. Lemly AD. Modification of benthic insect communities in polluted streams: combined effects of  
52 sedimentation and nutrient enrichment. *Hydrobiologia* [Internet]. 1982 Mar [cited 2018 Aug  
53  
54  
55  
56  
57  
58  
59  
60


- 2];87(3):229–45. Available from: <http://link.springer.com/10.1007/BF00007232>
259. Townsend CR, Uhlmann SS, Matthaei CD. Individual and combined responses of stream ecosystems to multiple stressors. *J Appl Ecol* [Internet]. 2008 Dec 1 [cited 2018 Aug 2];45(6):1810–9. Available from: <http://doi.wiley.com/10.1111/j.1365-2664.2008.01548.x>
260. Overall NC, Johnson MF, Wood P, Mattingley L. Sensitivity of the early life stages of a mayfly to fine sediment and orthophosphate levels. *Environ Pollut* [Internet]. 2018 Jun 1 [cited 2018 Aug 2];237:792–802. Available from: <https://www.sciencedirect.com/science/article/pii/S0269749117329792>
261. Dolédec S, Phillips N, Scarsbrook MR, Riley RH, Townsend CR, Dolédec S, et al. Comparison of structural and functional approaches to determining landuse effects on grassland stream invertebrate communities. *J North Am Benthol Soc* [Internet]. 2006;25(1):44–60. Available from: [isi:000236565400005](http://www.jstor.org/stable/3066400)
262. Lange K, Townsend CR, Matthaei CD. Can biological traits of stream invertebrates help disentangle the effects of multiple stressors in an agricultural catchment? *Freshw Biol* [Internet]. 2014 Dec 1 [cited 2018 Aug 1];59(12):2431–46. Available from: <http://doi.wiley.com/10.1111/fwb.12437>
263. Larsen S, Pace G, Ormerod SJ. Experimental effects of sediment deposition on the structure and function of macroinvertebrate assemblages in temperate streams. *River Res Appl* [Internet]. 2011 Feb 1 [cited 2018 Aug 2];27(2):257–67. Available from: <http://doi.wiley.com/10.1002/rra.1361>
264. Verberk WCEP, van Noordwijk CGE, Hildrew AG. Delivering on a promise: integrating species traits to transform descriptive community ecology into a predictive science. *Freshw Sci* [Internet]. 2013 Jun 17 [cited 2018 Aug 1];32(2):531–47. Available from: <http://www.journals.uchicago.edu/doi/10.1899/12-092.1>
265. Redding J, Schreck C. Mount St. Helens ash causes sublethal stress response in steelhead trout. In *Washington Water Research Centre proceeding from the Conference, Mt. St. Helens—effects on water resources: Report 41*. Washington Water Research Center: Pullman, WA; 1982.
266. Bond NR, Downes BJ. The independent and interactive effects of fine sediment and flow on benthic invertebrate communities characteristic of small upland streams. *Freshw Biol* [Internet]. 2003 Mar [cited 2018 Jul 18];48(3):455–65. Available from: <http://doi.wiley.com/10.1046/j.1365-2427.2003.01016.x>
267. Sutherland AB, Meyer JL. Effects of increased suspended sediment on growth rate and gill condition of two southern Appalachian minnows. *Environ Biol Fishes* [Internet]. 2007 Oct 22 [cited 2018 Aug 2];80(4):389–403. Available from: <http://link.springer.com/10.1007/s10641-006-9139-8>
268. McCubbin R, Case A, Rowe D, Scruton D. Environmental effects of sediments on aquatic habitats and suggested mitigation. Appendix IV. In: *Resource Road Construction: Fish Habitat Protection Guidelines*. St John's, New Foundland: Canadian Forest Service and Department of Fisheries and Oceans; 1990. p. 65.
269. Ahlbert I. The organisation of the cone cells in the retinae of four teleosts with different feeding

- habits (*Perca fluviatilis* L., *Lucioperca lucioperca* L., *Acerina cernua* L. and *Coregonus albula* L.). *Ark Zool* [Internet]. 1969 [cited 2018 Aug 2];22(11):445–81. Available from: <https://ci.nii.ac.jp/naid/10020860423/>
270. Ali MA, Ryder RA, Anctil M. Photoreceptors and Visual Pigments as Related to Behavioral Responses and Preferred Habitats of Perches (*Perca* spp.) and Pikeperches (*Stizostedion* spp.). *J Fish Res Board Canada* [Internet]. 1977 Oct [cited 2018 Aug 2];34(10):1475–80. Available from: <http://www.nrcresearchpress.com/doi/abs/10.1139/f77-212>
271. Bergman E. Foraging Abilities and Niche Breadths of Two Percids, *Perca Fluviatilis* and *Gymnocephalus Cernua*, Under Different Environmental Conditions. *J Anim Ecol* [Internet]. 1988 Jun [cited 2018 Aug 2];57J(57):443–53. Available from: <http://www.jstor.org/stable/4916%0Ahttp://www.jstor.org/page/info/about/policies/terms.jsp>  
<http://www.jstor.org>
272. Franklin PA, Smith J, Baker CF, Bartels B, Reeve K. First observations on the timing and location of giant kokopu (*Galaxias argenteus*) spawning. *New Zeal J Mar Freshw Res* [Internet]. 2015 Jul 3 [cited 2018 Aug 2];49(3):419–26. Available from: <http://www.tandfonline.com/doi/full/10.1080/00288330.2015.1045004>
273. Power ME. The importance of sediment in the grazing ecology and size class interactions of an armored catfish, *Ancistrus spinosus*. *Environ Biol Fishes* [Internet]. 1984 Apr [cited 2018 Aug 2];10(3):173–81. Available from: <http://link.springer.com/10.1007/BF00001124>
274. Doan KH. Relation of Sauger Catch to Turbidity in Lake Erie. *Ohio J Sci* [Internet]. 1941 [cited 2018 Aug 2];41(6):449–52. Available from: [https://kb.osu.edu/dspace/bitstream/handle/1811/3200/1/V41N06\\_449.pdf](https://kb.osu.edu/dspace/bitstream/handle/1811/3200/1/V41N06_449.pdf)
275. Kulíšková P, Horký P, Slavík O, Jones JI. Factors influencing movement behaviour and home range size in ide *Leuciscus idus*. *J Fish Biol* [Internet]. 2009 Apr 1 [cited 2018 Aug 2];74(6):1269–79. Available from: <http://doi.wiley.com/10.1111/j.1095-8649.2009.02198.x>
276. Zamor RM, Grossman GD. Turbidity Affects Foraging Success of Drift-Feeding Rosyside Dace. *Trans Am Fish Soc* [Internet]. 2007 Jan 9 [cited 2018 Aug 2];136(1):167–76. Available from: <http://www.tandfonline.com/doi/abs/10.1577/T05-316.1>
277. Fairchild JF, Boyle T, English WR, Rabeni C. Effects of sediment and contaminated sediment on structural and functional components of experimental stream ecosystems. *Water Air Soil Pollut* [Internet]. 1987 [cited 2018 Aug 2];36(3–4):271–93. Available from: <http://link.springer.com/10.1007/BF00229673>

### Further Reading

None. The bibliography is extensive.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



114x46mm (300 x 300 DPI)

Peer Review




Figure 1 – Controls on the delivery, transport and storage of fine sediment and the resulting ecological responses in river networks. Natural and anthropogenic factors control the quantity and quality of sediment potentially available for delivery to the river channel. Whether or not this potential is realised, and the quantity and composition of sediment conserved during the pluvial phase, depends on delivery pathways that connect the source to the river channel. Once delivered to the river channel, particles may be transported through suspension or saltation, temporarily stored as surface-deposited material, or become infiltrated in the river bed and stored on a longer-term basis. Which compartment particles are transported or stored depends on a number of controls, including the inherent properties of the particles, hydraulics, and the activities of biota. The latter is in turn influenced by physico-chemical factors (e.g. oxygen, nutrients) that work to limit or enhance biological processes. The hydraulic and biological controls determine the compartments within which particles are transported or stored, structuring the physical habitat, modifying available resources and potentially filtering out unsuitable species from the species pool on the basis of their traits. Such environmental filtering effects may be interfered with by biogeographical processes related to dispersal. The fate of fine sediment, and the extent and nature of the impact it causes is subject to potential lags as sediment is temporarily stored within the landscape. Ecological impacts may also be delayed as time is taken for local populations to reach critical life-stages where fine sediment limits survival.

188x124mm (300 x 300 DPI)




Figure 2 - Fine sediment transport (black arrows) and deposition (grey arrows) processes in gravel beds.

These processes occur in three key load regions (suspended and dissolved load, Brinkman load and interstitial load) within the water and sediment columns in which such processes occur (left). The model proposes distinct modes of sediment transport in these three regions identified by transitions in the velocity profile  $V(y)$  (right). Modified from<sup>43</sup>.

326x175mm (300 x 300 DPI)




Figure 3 - Controls on fine sediment deposition and resuspension exerted through bioflocculation by extracellular polymeric substances (EPS) in biofilms.

218x164mm (300 x 300 DPI)




Figure 4 - Metacommunity theory predicts patterns in dominant community assembly processes in river networks (a) due to variation in spatial, environmental and demographic influences with stream order (b), with implications for ecosystem responses to stressors. Headwater communities are likely to follow the patch dynamics paradigm, whereby local species diversity is limited by dispersal and community structure becomes a function of stochastic demographic processes involving local extinction and colonisation<sup>195</sup>. The structure of lower mainstem communities is predicted to more closely resemble the regional species pool since high dispersal rates drown out the influence of local environment through 'mass effects'<sup>195</sup>. In mid-basin locations, species sorting or 'environmental filtering'<sup>194</sup> is made possible as the influence of dispersal is at an optimal level to allow for efficient niche-based processes<sup>199</sup>.

91x41mm (300 x 300 DPI)