

Design for Mobile Mental Health: Exploring the Informed Participation Approach

Abstract

Mobile applications (apps) have the potential to improve mental health services. However, there is limited evidence of efficacy or responsiveness to user needs for existing apps. A lack of design methods has contributed to this issue. Developers view mental health apps as standalone products and dismiss the complex context of use. Participatory design, particularly an informed participation approach, has potential to improve the design of mental health apps. In this study, we worked with young mobile users and mental health practitioners to examine the informed participation approach for designing apps. Using auto-ethnography and a set of design workshops, the project focused on eliciting design requirements as a factor for successful implementation. We compared resultant ideas and designs with existing apps. Many user requirements revealed were absent in existing apps, suggesting potential advantages to informed participation. The observation of the process, however, showed challenges in engagement which need to be overcome.

Design for Mobile Mental Health: Exploring the Informed Participation Approach

Introduction

The availability of smartphones makes them an attractive option for providing health services because they make digital and connectivity tools accessible to the public with relatively low cost.¹ The high penetration of smartphones internationally means that mobile health, with its capacity for so-called nomadic modes of interaction between users and mobile devices, presents opportunities for addressing existing healthcare challenges.^{2,3} Mobile solutions have the potential to revolutionize mental health services, with new mental health-focused applications (apps) added to app stores on a daily basis. Often, these apps are based on a physical health model, substituting food or exercise tracking for mood or sleep tracking. However, there is limited evidence of efficacy for existing mobile mental health (mMH) apps in real world contexts.⁴⁻⁶ Even apps with proven clinical efficacy do not achieve outcomes in real world settings, as users may not continuously engage with the app or adhere to the recommended/ planned usage scenarios.^{6,7} Apps are not stand-alone digital solutions; they have the potential to be touch points for mental health services, as part of a complex ecosystem of various stakeholders including the user, their family and friends, mental health practitioners, and relevant private and public health organizations.

Although there is a rich literature around technologies and policies that can empower mMH,^{4,7} we still do not know much about suitable mMH “design” approaches. Design methodology has the potential to significantly improve the success of mMH solutions, if one defines design as devising courses of action aimed at changing existing situations into preferred ones,⁸ aiming to

solve wicked and multidimensional problems.⁹ Considering the complexity of mental health services, and existence of multiple stakeholder groups with various requirements, using an appropriate design methodology that takes into account these complexities is vital.

In this research we aimed to assess the feasibility and effectiveness of using an informed participation approach in designing an mMH app with two user groups who are part of the complex mental health ecosystem: young people (users) and mental health practitioners (stakeholders).

Using design approaches in healthcare

Though there are many existing mMH apps, there is little regulation of this market or research into efficacy. Seeking healthcare information in a poorly-regulated digital space may mean people risk of accessing poor quality or inaccurate resources.^{10,11} Attempts for reviewing and regulating available products have failed to this point,¹² though there are some promising new approaches.¹³ Existing research on mMH apps typically views them using technological¹⁴ or clinical¹⁵ lenses, and few rigorous and evidence-based studies have focused on mMH.^{16–20} Only a small number of available mMH apps can provide evidence of efficacy,^{18,19,21} and there is evidence that some commercial apps could be potentially harmful.^{22,23} As a result, public health organizations including the UK's National Institute for Health and Care Excellence (NICE) have trialled standards or programmes for improving user experience.^{24–26} Using and improving design processes and tools rather than just regulating numerous end products may have benefit for healthcare consumers.

In healthcare, design often uses an evidence-based approach built on a core principle of making decisions based on the best available research evidence.²⁷ Evidence-based design includes a mix of methodologies and tools, and is best viewed as a problem solving approach² rather than a method for shaping physical objects and spaces. As healthcare evolves, design methods need to do the same. Existing evidence-based healthcare design approaches do not take into account some subjective, hard-to-measure, and tacit aspects such as user experience. Poor design may affect dropout rate, future treatment-seeking behaviour and even quality of life after treatment.²⁸ Previous emphasis has been on adjusting existing design methodologies for developing mMH apps, including rethinking participatory design for users with serious mental health illnesses^{29,30} and adapting user-centred design process by defining multiple players in the process.^{31,32,33}

Participatory design and informed participation

In light of the challenges of evidence-based design, regulation of apps and adapting design approaches, participatory design is a promising approach for this domain³⁴ with a strong connection to healthcare design and delivery via evidence-based practice.^{35,36} Like other established design methodologies, it can be problematic if the goal is to develop high-quality solutions and implementable results within limited time constraints.³⁷ Using an informed participation approach has helped to overcome this in some domains.^{38–41} In informed participation, participants use the information and tools provided by the designers to

incrementally obtain ownership of problems and to contribute actively to their solutions.⁴² This co-design approach encourages engagement in real-world settings.

In informed participation, the research agenda is open and transparent throughout, and the motives and objectives of data collection methods known to participants.⁴³ Traditional science and engineering processes used for designing systems,⁴⁴ and even some common user-centred design methods such as shadowing observations⁴⁵ cannot be categorised as informed participation. In contrast co-design as an interactive process, evolving with participant engagement, can be considered informed participation.⁴⁴

Informed participation allows less engaged or unrepresented communities to have a voice in design.^{46,47} For mMH research, enabling informed participation presents opportunities,⁴⁸ but also challenges. Smartphones provide users with a high level of customization and flexibility of use, allowing participants to direct the research. However, there are ethical concerns, as these devices are also effective tools for covert data collection.⁴⁹

To assess the feasibility and effectiveness of using an informed participation approach, we worked with young people (users) and mental health practitioners (stakeholders). The design project focused on the promotion of good wellbeing and prevention of mental health issues for young people, including increasing problem solving and coping skills. We hypothesised that concentrating on feasible requirements-gathering practices within the participatory design approach may help overcome the challenge of involvement in design and development of usable systems.⁵⁰ Because it was not possible to fully develop and test the app, we compared the design requirements and concepts developed with existing mMH apps to assess whether the informed participation approach revealed design requirements that have not been considered before.

Method

To ascertain if a participatory design approach was effective and feasible, we used an action research approach. Action research involves changing practice and improving knowledge about a practice or social system, in which the action researcher is also a practitioner (in this case, a designer, and an ethnographer with no design background who participated in the workshops) and the research is not separated from the action itself.^{51–53} Participatory design has deep roots in action research and can be considered as reflective practice.^{54,55} Action research is frequently used to improve and manage changes in healthcare systems and services.^{56–58} By using a reflective action research approach, we were able to examine the design process as observers engaged in the process, considering its feasibility and effectiveness.

Study Design

The study comprised three elements: an auto-ethnography and design learning process with two user groups and a review of currently available mMH apps informed by the findings from the first two elements. The study aimed to analyse the process of using an informed participation approach and not just the products of doing so, meaning that diverse data were gathered.

Two user groups participated:

- Young people/users: Seven university students aged 18-20 not currently enrolled in a design course.
- Mental health practitioners/stakeholders: Ten members of the university's counselling and mental health services team.

Users were invited to participate by an email invitation distributed via undergraduate student mailing lists. We recruited from a sample population across a university (all students aged 18-20 who were not on a design course) and took all volunteers. Young people were identified as a relevant and underrepresented group within design processes. On the recommendation of the university research ethics committee, we sampled from the general student population rather than working specifically with participants with mental health issues. This was an appropriate user group to work with, as we planned to have a well-being and prevention focus, rather than treating specific mental illnesses. Mental health practitioners agreed that coping and problem solving were core skills for improving mental wellbeing. Despite its importance, wellbeing and prevention are relatively less explored aspects of mMH.⁵⁹

To recruit stakeholders, the research team attended a meeting of the university counselling and mental health services team and explained the requirements of the study; ten practitioners volunteered to participate. Informed consent was given by all participants.

Two novel components in the study design specifically engaged with the informed participation approach:

Auto-ethnography: As an instructional tool, this method helps researchers and participants to gain profound understanding of the self and others.⁶⁰ The participant retroactively and selectively writes about experiences, combining autobiography and ethnography, and considers systemic relationships.⁶¹ It provided participants with a high level of control and self-awareness.

Design learning workshops: To design *with* participants, we integrated practical design learning components (design education delivered via lectures and practical sessions) which enabled participants to view themselves as designers. We shared our definition of design and objectives at the start of the project. As a result, participants developed design briefs, ideated, and prototyped design concepts for mMH apps.

The study comprised two separate pathways for young people as primary users of mMH apps (table 1) and practitioners as secondary users/ stakeholders (table 2). Evidence suggests that young people and mental health practitioners are not generally involved in the mMH app design process, and so we considered them to be legitimate participants in an informed participation approach. Time constraints meant that stakeholders were able to attend fewer workshops than users, though the nature of activities and objectives remained broadly the same.

Finally, we compared the pathway outcomes with existing products, looking at mMh apps in the Google Play app store using the keywords “problem solving” and “coping” in the health and fitness, lifestyle and medical categories. Eighty-seven apps were found based on this search criteria. We searched this app store because Android was the most common mobile operating system worldwide at the time of study.⁶² The mMh app content was analysed using a content-driven framework based on classifying the features in the apps themselves, then via another framework constructed from the findings from the user and stakeholder design process.

Table 1: User/young people pathway

Step	Activities
<i>Auto-ethnography (three weeks)</i>	Participants recorded their daily experiences along the themes of solving problems or coping with difficulties, including minor concerns as well as major issues. The main focus was on methods, strategies, tools, or resources used for solving problems or coping with difficulties rather than details of the problems themselves, e.g. if a participant successfully coped with an emotional problem by speaking to a friend, auto-ethnographic notes highlighted ‘sharing and communicating’ as a coping strategy, rather than the details of emotional problem itself. Participants were asked to choose their recording tool - paper and digital notes, audio and video recordings, and photos.
<i>Workshop 1: auto-ethnography review</i>	Teaching activity: reframing and redefining problems Participants shared strategies identified in the auto-ethnography.
<i>Workshop 2: design brief</i>	Teaching activity: the definition of design Using our definition of design, we explored the design process using the double diamond model (discover, define, develop, and deliver). ⁶³ The auto-ethnography summary drafted in workshop 1 was used to shape a design brief for a “problem solver” app.
<i>Workshop 3: ideation</i>	Teaching activity: ideation and creativity as the starting point of developing solutions in a design process Using brainstorming and visual thinking activities, participants generated ideas for the app, without considering their feasibility.
<i>Workshop 4: concept development</i>	Teaching activity: examples of low fidelity prototyping Looking at the initial design brief (workshop 2), participants were asked to develop more feasible concepts based on ideas in workshop 3. This involved evaluating or filtering ideas, as well as merging similar ideas collaboratively. Participants started to identify previously unapparent design requirements, which they used to inform relatively realistic design concepts.

<i>Interim activity</i>	Participants were asked to create a low-fidelity paper prototype
<i>Workshop 5: prototyping and testing</i>	Prototypes were peer tested using a simple scenario, in which a typical user approached the app for the first time to think about facing a minor problem. Tests were filmed and reviewed by the participants. Based on the evaluation, participants improved their prototypes.

Table 2: Stakeholder/ mental health practitioner pathway

<i>Step</i>	<i>Activities</i>
<i>Auto-ethnography (three weeks)</i>	Participants were asked to record daily experiences of working with students to identify solutions to problems or cope with difficulties, including minor concerns as well as major issues. A proforma was used to structure the data collection, asking for a brief outline of the presenting issue (excluding sensitive information/ identifiable detail), supportive resources, and strategies for solving problems or coping with difficulties.
<i>Interim activity</i>	The research team prepared a problem summary based on the most common strategies, solutions and problems.
<i>Workshop 1: design brief</i>	Teaching activity: the definition of design and the main steps of a design process Participants reflected on the problem summary. This reflection validated the findings and helped to understand some of contextual challenges. From this the research team constructed a design brief.
<i>Workshop 2: ideation</i>	Participants used storytelling methods as an ideation activity to develop solutions. Similar concepts were grouped to outline a set of design requirements.

Data Analysis

A three-step qualitative analysis was used to gather and reduce (code) the data, visualise the data, and draw conclusions (figure 1).⁶⁴ This was based on a data-driven constant comparison approach,⁶⁵ which was closely grounded in observations of the workshops and was also responsive to the action research approach. Data mapping and visualisation helped us to compare results from different data sources and relate outcomes to the research aims.

[Figure 1 about here]

Results

In summary, data collected were auto-ethnographic data and participants' reflections on these data, ideas in textual and visual formats, low-fidelity prototypes, observation notes mapping the

procedure and reflections on each workshop, video recordings of prototype tests and comparison between prototypes, and a review of available mMH apps (table 3). The fast-paced nature of developing technologies means that descriptions of current technologies quickly become obsolete; for this reason, we have provided an outline of the features included rather than the specific apps.

Table 3: Features of reviewed Android apps

Feature	Apps which have this feature	
	Number of apps	Approximate percentage (of total 87 reviewed apps)
Glossary of mental health information	50	57%
Informative multimedia contents including guided meditations and brain waves music, hypnosis	30	34%
(Self) mood tracking	22	25%
Learning skills by practice	21	24%
Using visual metaphors	13	14%
Building habits using reminders and practices	11	12%
Mental health self-assessment	11	12%
Social networking, collaborating, and sharing	10	11%
Gamification	7	8%
Mental health assessment using voice recognition technology	2	2%
Using artificial intelligence for advice and coaching	2	2%
Online counselling	1	1%
Cognitive Behavioral Therapy	1	1%

From the data collected, we were able to extract a number of *core concepts* associated with problem solving and coping, which were then used to identify *high-level* and *detailed design requirements* for the users (table 4) and stakeholders (table 5). High-level requirements are broad themes identified by the participants. Detailed requirements expand on these high-level themes to provide greater clarity about how these high-level requirements will be achieved. For the users we worked with, we were able to extend this process to identify *concept features* that an mMH app might have, and start to perform *concept evaluation* with prototypes (figure 2). Key themes emerging across the data from both participant groups were then compared and similarities and differences highlighted. We then compared the themes in user and stakeholder data with existing mMH apps (table 6 and table 7). From this process, we were able to address our aim of investigating whether the informed participation approach was effective in identifying design requirements.

[Figure 2 about here]

By examining the core concepts found in user and practitioner auto-ethnographies, we found that core concepts were described differently, but addressed similar issues. For example, stakeholders noted the importance of asking for help and connecting with others. Users focused

on problem solving as a shared practice. Detailed design requirements showed greater overlap, and some detailed requirements from stakeholders can be linked to concept features found in the users' planning of apps. Most features in the final design concepts were directly linked to design requirements outlined earlier in the process. In mapping existing app features, user and stakeholder design requirements, and user concept features, we found that not all requirements and features were present in existing apps.

Data collection also allowed us to address the question of whether an informed participation process was feasible. The data collection process showed that the challenges of implementing the approach include the length of time required for the process. For both users and stakeholders, although there was enthusiasm for the project, limited time was available to learn techniques, generate ideas and design requirements and produce prototypes. Users, who were unfamiliar with the design approach, also struggled with the openness and ambiguity within the process. This had a negative impact on participation, with one participant (an engineering student) withdrawing from the study because they felt that the process was too vague.

Table 4: User/ young people pathway summary of data and themes

<i>Step</i>	Key themes
<i>Workshop 1: Auto-ethnography review</i>	<p>Data gathered: Participants insights about auto- ethnographic experience</p> <p>Core concepts associated with problem solving and coping</p> <ul style="list-style-type: none"> ● Analytical thinking can help when coping with emotional problems e.g. when we overestimate the likelihood of failure ● Holistic thinking can also help, as we may miss the big picture because of focusing on details ● Self-awareness is key to solving problems ● Solving problems can be a shared practice ● We often know about the solutions, but the problem is choosing the best solution ● Many problems can be solved by long term plans and habits rather than instant solutions
<i>Workshop 2: design brief</i>	<p>Data gathered: Design brief</p> <p>High-level design requirements</p> <p>Ability to:</p> <ul style="list-style-type: none"> ● evaluate the situation and the users' emotional status ● assist user to use both analytical and holistic thinking ● assist in making decisions ● monitor and facilitate the progress ● connect with other who can help

<p><i>Workshop 3: ideation</i></p>	<p>Data gathered: Ideas (text, visuals)</p> <p>Detailed design requirements Ability to:</p> <ul style="list-style-type: none"> ● build habits ● plan ahead: identify milestones and manage time ● enhance the problem solving and coping capabilities by improving physical health ● learn from solving problems and recall solutions in future ● break down complex problems into simpler ones ● identify the right time for making decisions e.g. when user in a stable emotional mode and has necessary information ● evaluate users' emotional state ● prioritize problems and objectives, and help users to focus on solving the most impactful parts of a problem ● help users to decide quickly when necessary ● find the "optimum" solution ● assist users to have an "out of the box" view on problems ● link users to those who successfully managed to solve problems: building a network and a directory of experiences and solutions ● make the process of "asking for help" easier ● visualize information ● communicate and interact in a "natural way" [NB: This was a phrase used by students, who felt that communication should be naturalistic and colloquial not feel forced or artificial.]
<p><i>Workshop 4: concept development</i></p>	<p>Data gathered: Selected ideas represented by users' sketches for the app user interface (two ideas represented)</p> <p>Concept features</p> <ul style="list-style-type: none"> ● evaluate users' emotional state ● find the "optimum" solution ● plan ahead: identify milestones and manage time ● track progress towards solution ● learn from solving problems and recall solutions in future ● visualize information (using metaphors for emotions, problems, solutions etc.) ● communicate and interact in a "natural" way ● gamify the problem solving process ● help user to be achieve a stable emotional state (e.g. by meditation) ● identify the right time for making decisions (e.g. when user in a stable emotional mode and has necessary information) ● break down complex problems into simpler ones ● prioritize problems and objectives, and help users to focus on solving the most impactful parts of a problem ● build habits ● link users to those who successfully managed to solve problems: building a network and a directory of experiences and solutions
<p><i>Workshop 5: prototyping and testing</i></p>	<p>Data gathered: Paper prototypes; Test videos (two final ideas)</p> <p>Concept evaluation</p> <ul style="list-style-type: none"> ● Visual metaphors can be culturally or context specific ● Too many features were defined in workshops 3 and 4, considering all of them in one product might make it too complex ● Although visually different, both ideas had similar user scenarios, starting with evaluating users emotions, and ending with recording progress and rewarding them for their achievements

Table 5: Stakeholder/ mental health practitioner pathway summary of data and themes

Step	Key themes
<i>Auto-ethnography (three weeks)</i>	<p>Data gathered: 131 records of student encounters, thematically analysed</p> <p>Presenting mental health problems were diverse and often overlapping, but could be broadly categorised as emotional problems (e.g. anxiety, depression, panic), educational issues (e.g. exam anxiety, transition to university, perfectionism) and problems related to events and people, including relationship breakdown, bereavement and dealing with past traumatic events.</p> <p>A number of core concepts associated with problem solving and coping were suggested:</p> <ul style="list-style-type: none"> • Seeking help from others (e.g. academic department) provided practical support in the face of difficulties • Connecting with others (e.g. friends/family) could be a solution • Making positive behavioural changes including exercise and sleep • Learning more about the problem could help to identify potential solutions • Using grounding and thought change techniques helped to cope with emotional problems
<i>Workshop 1: design brief</i>	<p>Data gathered: Stakeholder reflections on auto-ethnography and contextual challenges</p> <p>High-level design requirements</p> <p>An app needs the ability to:</p> <ul style="list-style-type: none"> • facilitate time for stakeholders’ own self-care, learning, reflection and analysis • enable efficient administrative processes • evaluate user readiness and suitability for face-to-face intervention • monitor, motivate and facilitate user progress • connect with other local services who can help
<i>Workshop 2: ideation</i>	<p>Data gathered: Stakeholder creation of design requirements</p> <p>Detailed design requirements</p> <p>Ability to help users to:</p> <ul style="list-style-type: none"> • build habits and set targets • track progress towards a solution, including self-review • plan ahead: identify milestones and manage time, including reminders • break down complex problems into simpler ones (take a stepwise approach) • link users to those who successfully managed to solve problems: building a network and a directory of experiences and solutions • learn from solving problems and recall solutions in future • visualize information (using metaphors for emotions, problems, solutions, habits, etc.) • access grounding and thought change techniques • access personalised recommended information-based resources • normalise problems appropriately • understand the counselling/ mental health support process • access longer-term support to prevent relapse • engage with appropriate university processes (e.g. mitigating circumstances)

Table 6: Comparison between requirements in users/ young people pathway and existing apps

Source	Requirement	Apps having this feature	
		Number of apps	Approx. % of reviewed apps
<i>Core concepts</i>	Facilitating analytical thinking	0	0%
	Facilitating holistic thinking	0	0%
	Bringing self awareness	0	0%
	Developing long term plans	0	0%
	Building positive habits	11	12%
	Comparing solutions	0	0%
	Collective problem solving (problem solving as a shared practice)	10	11%
<i>High-level design requirements</i>	Evaluating the solution	0	0%
	Combining analytical and holistic thinking	0	0%
	Facilitating decision making	0	0%
	Monitoring progress	0	0%
<i>Detailed design requirements</i>	Building habits	11	12%
	Planning	0	0%
	Improving physical health	0	0%
	Evaluating the users' emotional state	0	0%
	Identifying the right time for making decisions	0	0%
	Breaking down the problems	0	0%
	Facilitating decision making	0	0%
	Giving an out-of-the-box view	0	0%
	Prioritizing problems	0	0%
	Finding the optimum solution	0	0%
	Learning and recalling solutions	0	0%
	Building a network and a directory of experiences and solutions	10	11%
	Asking for help when necessary	0	0%
	Visualizing information	13	14%
Interacting naturally	2	2%	
<i>Concept features</i>	Help users to achieve a stable emotional state	0	0%
	Gamifying the problem solving process	7	8%
	Simplifying the problem solving process (avoiding complexity)	0	0%
	Using culturally- specific metaphors	13	14%

Table 7: Comparison between requirements in stakeholder/ mental health practitioner pathway and existing apps

Source	Requirement	Apps having this feature	
		Number of apps	Approx. % of reviewed apps)
<i>Core concepts</i>	Grounding and thought change techniques	30	34%
	Learning more about the problem	0	0
	Making positive behavioral changes	1	1%
	Connecting with others	10	11%
	Asking for help when necessary	0	0%
<i>High-level design</i>	Monitoring and motivating patients' progress	25*	22%
	Handling administrative processes	0	0%

<i>requirements</i>	Facilitating stakeholders' own time management	0	0%
	Evaluating users' readiness for intervention	0	0%
	Connecting with other local services	0	0%
<i>Detailed design requirements</i>	Building habits	11	12%
	Planning	0	0%
	Tracking progress	25*	22%
	Self review	25*	22%
	Reminding tasks and milestones	11**	11%
	Learning and recalling solutions	0	0%
	Building a network and a directory of experiences and solutions	10	11%
	Facilitating access to longer-term support to prevent relapse	0	0%
	Clarifying the counselling /mental health support process	30†	34%
	Visualising information	13	14%
	Normalising problems	0	0%
	Grounding and thought change techniques	30†	34%
<i>*mood tracking apps; **habit builder apps; †using informative contents</i>			

In summary, using an informed participation approach identified design requirements and concepts not currently commonplace in existing mMh apps, which predominantly provided information and/or multimedia content like guided meditations. Users focused much more on expanding their options for problem-solving by link building and they wanted help to identify the right time for making decisions. Users wanted an app to help them to identify milestones, manage time and track progress towards a solution, and to be iterative in learning, recalling solutions in the future. All of this had to be achieved with the mMh app communicating and interacting in a “natural” (note: this was a phrase coined by users, who felt that communication should be naturalistic and colloquial not feel forced or artificial) way.

Mental health practitioners had similar requirements, but also wanted an app to have a focus on helping young people they worked with to normalise problems appropriately, to understand the counselling/ mental health support process and to provide access to grounding and thought change techniques and personalised recommended information-based resources. They considered when an app might be useful, for example in accessing longer-term support to prevent relapse, and to help users engage with appropriate processes (e.g. for a university context, exceptional circumstances committees) to support them.

Discussion

Our experience of implementing an informed participation design process demonstrates that it is a promising approach which is effective at outlining user requirements that could not be identified in currently available apps. We particularly focused on design requirements and product features, as we recognised that successful requirements gathering can significantly impact successful implementation.⁶⁶ However, the length and ambiguity of the process may still impact user engagement, leading to questions about its feasibility. This mirrors previous studies which have highlighted that engagement in processes is challenging.⁶⁷ Analysis showed that 11 of the 13 features identified in the user pathway final concepts were identified in the earlier

steps. Focusing on the auto-ethnography and requirements gathering steps may be a solution for shortening the process, and consequently improving user engagement.

Users and stakeholders identified different requirements and features, but expressed similar core concepts. These core concepts emphasise a need to learn different thinking and grounding patterns, looking at problem solving as a shared practice, and empowering positive habits and behaviour changes. When online reviews of mMH apps have been analysed, similar calls for discrete social networks within apps have been noted.⁶⁸ Both participant groups required a combination of features in their ideal solution, helping them through the whole problem-solving process.

Existing apps are often designed to contain one or two features, without supporting a sustainable and holistic process, which is a weakness of current provision. Apps on the market were mainly information giving (57%) or informative multimedia content (34%). Participants in our research identified much more diverse requirements involving collective problem solving and thinking about how to think and behave differently. When we compared existing apps to the requirements gathered, one significant finding was that our users and stakeholders wanted one app that covered all their requirements, whereas apps tended to concentrate on one or two specific features. Apps that took a more holistic approach were not available.

Paper prototyping showed that sustaining the complexity of these features within one mMH app may be difficult. Instead, creating a suite of apps with a hub may be an option.⁶⁹ The requirements identified here were able to address scenarios of use in a detailed, everyday and comprehensive way. For users, these requirements included the ability to identify and evaluate emotional states to consider when it was an appropriate time to make a decision to solve a problem, and tools to break down complex problems into simple ones. For the stakeholders, it was important that the tool could be appropriately personalised, connecting both to local services and appropriate information resources, and could help the user to understand their problems and the process for solving them in context. Many stakeholder requirements initially focused on addressing service processes, something which is missing in the existing apps. However, their overall design requirements showed that encouraging independence, self-review and engagement would meet many of these needs.

As we were not able to develop and test beyond the prototyping stage in this small pilot study, future work could consider how open innovation methods,⁷⁰ such as innovation jams,⁷¹ may have a role to play in further development. Innovation jams encourage the rapid generation of many ideas by a large cohort, and thus speed up the process of getting from initial idea to finished product. This may assist with engagement.

Strengths and limitations

This was a small-scale study, focused on one particular under-represented group (young people) and it is difficult to generalize the results for designing mMH apps. We did include two major stakeholder groups (young people and mental health service providers) and reflecting on the differences between these two groups brings valuable insight into the use of the methods.

As the study was conducted in students without mental health problems, this is also a limitation, and using informed participation approach would require further testing in this population. Future research could examine the informed participation approach at scale, e.g. at an organizational level, or with other groups. A strength of the study is that we collected diverse and rich qualitative data and supplemented it with a quantitative review. However, one of the limitations of participatory design in general is a lack of quantitative evaluation methods.

Conclusion

Informed participation can close the gap between ideas, final concepts and prototypes by enabling users to generate feasible, relevant, and detailed ideas which can communicate anticipated design requirements in further concepts and prototypes. Informed participation had a positive impact on implementability of designs, but this exploratory study also demonstrates the necessity of further rigorous work to developing and evaluate market-ready mMh apps.

References

1. Luxton DD, McCann RA, Bush NE, et al. mHealth for mental health: Integrating smartphone technology in behavioral healthcare. *Prof Psychol Res Pr* 2011; 42: 505.
2. Mastors P. *Design to Survive: 9 Ways an IKEA Approach Can Fix Health Care and Save Lives*. Morgan James Publishing, 2013.
3. Silberman MJ, Clark L. M-health: the union of technology and healthcare regulations. *J Med Pract Manage* 2012; 28: 118–120.
4. Firth J, Torous J, Nicholas J, et al. Can smartphone mental health interventions reduce symptoms of anxiety? A meta-analysis of randomized controlled trials. *J Affect Disord* 2017; 218: 15–22.
5. Firth J, Torous J, Nicholas J, et al. The efficacy of smartphone-based mental health interventions for depressive symptoms: a meta-analysis of randomized controlled trials. *World Psychiatry* 2017; 16: 287–298.
6. Chan S, Torous J, Hinton L, et al. Towards a Framework for Evaluating Mobile Mental Health Apps. *Telemed J E Health* 2015; 21: 1038–1041.
7. Price M, Yuen EK, Goetter EM, et al. mHealth: a mechanism to deliver more accessible, more effective mental health care. *Clin Psychol Psychother* 2014; 21: 427–436.
8. Simon HA. *The Sciences of the Artificial*. MIT Press, 1996.
9. Adam RJ. *Education for Wicked Problems and the Reconciliation of Opposites: A Theory of Bi-relational Development*. Routledge, 2016.
10. Xiao N, Sharman R, Rao HR, et al. Factors influencing online health information search: An empirical analysis of a national cancer-related survey. *Decis Support Syst* 2014; 57: 417–427.

11. Fergus TA, Dolan SL. Problematic internet use and internet searches for medical information: the role of health anxiety. *Cyberpsychol Behav Soc Netw* 2014; 17: 761–765.
12. Hickey E, McMillan B, Mitchell C. Practitioners should embrace, not ignore, health apps. *BMJ* 2015; 350: h2336.
13. Healthier Lancashire and South Cumbria ORCHA (the Organisation for the Review of Care and Health Applications). An App a day. *Lancashire ORCHA*, <https://lancashire.orcha.co.uk/> (2017, accessed 6 July 2018).
14. Proudfoot J. The future is in our hands: the role of mobile phones in the prevention and management of mental disorders. *Aust N Z J Psychiatry* 2013; 47: 111–113.
15. Harrison V, Proudfoot J, Wee PP, et al. Mobile mental health: review of the emerging field and proof of concept study. *J Ment Health* 2011; 20: 509–524.
16. Hind J, Sibbald SL. Smartphone Applications for Mental Health—A Rapid Review. *Western Undergraduate Research Journal Health and Natural Sciences* 2015; 1–9.
17. Bakker D, Kazantzis N, Rickwood D, et al. Mental health smartphone apps: review and evidence-based recommendations for future developments. *JMIR mental health*; 3, <https://www.ncbi.nlm.nih.gov/pmc/articles/pmc4795320/> (2016).
18. Leigh S, Flatt S. App-based psychological interventions: friend or foe?: Table 1. *Evid Based Ment Health* 2015; 18: 97–99.
19. Donker T, Petrie K, Proudfoot J, et al. Smartphones for smarter delivery of mental health programs: a systematic review. *J Med Internet Res* 2013; 15: e247.
20. Heron KE, Smyth JM. Ecological momentary interventions: Incorporating mobile technology into psychosocial and health behaviour treatments. *Br J Health Psychol* 2010; 15: 1–39.
21. Olff M. Mobile mental health: a challenging research agenda. *Eur J Psychotraumatol* 2015; 6: 27882.
22. Nicholas J, Larsen ME, Proudfoot J, et al. Mobile Apps for Bipolar Disorder: A Systematic Review of Features and Content Quality. *J Med Internet Res* 2015; 17: e198.
23. Gajecki M, Berman AH, Sinadinovic K, et al. Mobile phone brief intervention applications for risky alcohol use among university students: a randomized controlled study. *Addict Sci Clin Pract* 2014; 9: 11.
24. Bruyneel L, Van Houdt S, Coeckelberghs E, et al. Patient experiences with care across various types of mental health care: Questionnaire development, measurement invariance, and patients' reports. *Int J Methods Psychiatr Res*; 27. Epub ahead of print March 2018. DOI: 10.1002/mpr.1592.
25. Bate P, Robert G. Toward More User-Centric OD. *J Appl Behav Sci* 2007; 43: 41–66.
26. Government WWA, Wales. Health and Social Services. *Care Programme Approach/ Unified Assessment Process: Interim Approach*. 2010.
27. Kirk Hamilton D, Watkins DH. *Evidence-Based Design for Multiple Building Types*. John

Wiley & Sons, 2009.

28. Brownell T, Schrank B, Jakaite Z, et al. Mental health service user experience of positive psychotherapy. *J Clin Psychol* 2015; 71: 85–92.
29. Pelletier J-F, Rowe M, François N, et al. No personalization without participation: on the active contribution of psychiatric patients to the development of a mobile application for mental health. *BMC Med Inform Decis Mak* 2013; 13: 78.
30. Mulvenna MD, Nugent CD. *Supporting People with Dementia Using Pervasive Health Technologies*. Springer Science & Business Media, 2010.
31. Marcu G, Bardram J, Gabrielli S. A Framework for Overcoming Challenges in Designing Persuasive Monitoring and Feedback Systems for Mental Illness. In: *Proceedings of the 5th International ICST Conference on Pervasive Computing Technologies for Healthcare*. Epub ahead of print 2011. DOI: 10.4108/icst.pervasivehealth.2011.246097.
32. Frost M, Marcu G, Hansen R, et al. The MONARCA Self-assessment System: Persuasive Personal Monitoring for Bipolar Patients. In: *Proceedings of the 5th International ICST Conference on Pervasive Computing Technologies for Healthcare*. Epub ahead of print 2011. DOI: 10.4108/icst.pervasivehealth.2011.246050.
33. European Federation for Medical Informatics. International Congress, Andersen SK. *EHealth Beyond the Horizon - Get IT There: Proceedings of Mie2008*. 2008.
34. Doherty G, Coyle D, Matthews M. Design and evaluation guidelines for mental health technologies. *Interact Comput* 2010; 22: 243–252.
35. Luck R. Learning to talk to users in participatory design situations. *Design Studies* 2007; 28: 217–242.
36. Bate P, Robert G. Experience-based design: from redesigning the system around the patient to co-designing services with the patient. *Qual Saf Health Care* 2006; 15: 307–310.
37. Robertson T, Simonsen J. Challenges and Opportunities in Contemporary Participatory Design. *Design Issues* 2012; 28: 3–9.
38. Grundy S. Principles and possibilities¹. *Action research in practice* 2002; 37.
39. Jarvenpaa SL, Ives B. Executive Involvement and Participation in the Management of Information Technology. *Miss Q* 1991; 15: 205–227.
40. Julie Meyer U of M-A, Authors. Participation in the Planning and Design of Public Open Space, https://scholarworks.umass.edu/larp_ms_projects/32/ (2011, accessed 6 February 2018).
41. Fischer G. Meta-design: Beyond user-centered and participatory design. In: *Proceedings of HCI International*. l3d.cs.colorado.edu, 2003, pp. 88–92.
42. Fischer G, Ostwald J. Seeding, evolutionary growth, and reseeding: Enriching participatory design with informed participation. In: *Proceedings of the participatory design conference (PDC'02)*. Malmö University, Sweden, 2002, pp. 135–143.

43. Tucker WD. Beyond Traditional Ethics when Developing Assistive Technology for and with Deaf People in Developing Regions. In: Hersh M (ed) *Ethical Engineering for International Development and Environmental Sustainability*. London: Springer London, 2015, pp. 293–323.
44. Hersh MA, Tucker WD. ETHICS AND MONO-DISCIPLINARITY: POSITIVISM, INFORMED CONSENT AND INFORMED PARTICIPATION. *IFAC Proceedings Volumes 2005*; 38: 52–59.
45. Marti P, Bannon LJ. Exploring User-Centred Design in Practice: Some Caveats. *Knowledge, Technology & Policy* 2009; 22: 7–15.
46. Chatterjee S, Kieselbach B, Naik S, et al. Customising informed consent procedures for people with schizophrenia in India. *Soc Psychiatry Psychiatr Epidemiol* 2015; 50: 1527–1536.
47. Woudstra AJ, Dekker E, Essink-Bot M-L, et al. Knowledge, attitudes and beliefs regarding colorectal cancer screening among ethnic minority groups in the Netherlands--a qualitative study. *Health Expect* 2016; 19: 1312–1323.
48. Lasorsa I, D Antrassi P, Ajčević M, et al. Personalized support for chronic conditions. A novel approach for enhancing self-management and improving lifestyle. *Appl Clin Inform* 2016; 7: 633–645.
49. Albrecht U-V, Fangerau H. Do Ethics Need to be Adapted to mHealth? In: *ICIMTH*. books.google.com, 2015, pp. 219–222.
50. Kujala S, Kauppinen M. Identifying and Selecting Users for User-centered Design. In: *Proceedings of the Third Nordic Conference on Human-computer Interaction*. New York, NY, USA: ACM, 2004, pp. 297–303.
51. Craig DV. *Action Research Essentials*. John Wiley & Sons, 2009.
52. Sohng SSL. *Community-Based Participatory Research*. 2013.
53. Bowling A. *Research Methods in Health: Investigating Health and Health Services*. McGraw-Hill Education (UK), 2014.
54. Thoresen K. Principles in practice, two cases of situated participatory design. *Participatory Design, Perspectives on Systems Design*, <https://www.nr.no/publarchive?query=1562> (1992).
55. Bannon LJ, Ehn P. Design matters in participatory design. *Routledge handbook of participatory design* 2012; 37–63.
56. Koshy E, Koshy V, Waterman H. *Action Research in Healthcare*. SAGE, 2010.
57. Williamson GR, Bellman L, Webster J. *Action Research in Nursing and Healthcare*. SAGE, 2011.
58. Parkin P. *Managing Change in Healthcare: Using Action Research*. SAGE, 2009.
59. Aryana B, Brewster L, Nocera JA. Design for mobile mental health: an exploratory review.

Health Technol . Epub ahead of print 27 October 2018. DOI: 10.1007/s12553-018-0271-1.

60. Chang H. *Autoethnography as Method*. Routledge, 2016.
61. Ellis C, Adams TE, Bochner AP. Forum for Qualitative Research. *Autoethnography: An overview*; 12.
62. Mobile Operating System Market Share Worldwide | StatCounter Global Stats. *StatCounter Global Stats*, <http://gs.statcounter.com/os-market-share/mobile> (accessed 22 May 2018).
63. Hands D. *Design Management: The Essential Handbook*. Kogan Page Publishers, 2017.
64. Miles MB, Michael Huberman A. *Qualitative Data Analysis: An Expanded Sourcebook*. SAGE, 1994.
65. Charmaz K. *Constructing Grounded Theory*. SAGE, 2014.
66. Hunt EC, Sproat SB, Kitzmiller RR. *The Nursing Informatics Implementation Guide*. Springer Science & Business Media, 2013.
67. Anguera JA, Jordan JT, Castaneda D, et al. Conducting a fully mobile and randomised clinical trial for depression: access, engagement and expense. *BMJ Innov* 2016; 2: 14–21.
68. Owen JE, Jaworski BK, Kuhn E, et al. mHealth in the Wild: Using Novel Data to Examine the Reach, Use, and Impact of PTSD Coach. *JMIR Ment Health* 2015; 2: e7.
69. Lattie EG, Schueller SM, Sargent E, et al. Uptake and Usage of IntelliCare: A Publicly Available Suite of Mental Health and Well-Being Apps. *Internet Interv* 2016; 4: 152–158.
70. Chesbrough H, Vanhaverbeke W, West J. *Open Innovation: Researching a New Paradigm*. OUP Oxford, 2008.
71. The Emerging Impact of Open Innovation. *Reimagining Business Education* 2016; 41–49.