

SEARCHES FOR CONTINUOUS GRAVITATIONAL WAVES FROM FIFTEEN SUPERNOVA REMNANTS AND FOMALHAUT B WITH ADVANCED LIGO

- B. P. ABBOTT,¹ R. ABBOTT,¹ T. D. ABBOTT,² S. ABRAHAM,³ F. ACERNESE,^{4,5} K. ACKLEY,⁶ C. ADAMS,⁷
R. X. ADHIKARI,¹ V. B. ADYA,^{8,9} C. AFFELDT,^{8,9} M. AGATHOS,¹⁰ K. AGATSUMA,¹¹ N. AGGARWAL,¹² O. D. AGUIAR,¹³
L. AIELLO,^{14,15} A. AIN,³ P. AJITH,¹⁶ G. ALLEN,¹⁷ A. ALLOCCHA,^{18,19} M. A. ALOY,²⁰ P. A. ALTIN,²¹ A. AMATO,²²
A. ANANYEVA,¹ S. B. ANDERSON,¹ W. G. ANDERSON,²³ S. V. ANGELOVA,²⁴ S. ANTIER,²⁵ S. APPERT,¹ K. ARAI,¹
M. C. ARAYA,¹ J. S. AREEDA,²⁶ M. ARÈNE,²⁷ N. ARNAUD,^{25,28} K. G. ARUN,²⁹ S. ASCENZI,^{30,31} G. ASHTON,⁶
S. M. ASTON,⁷ P. ASTONE,³² F. AUBIN,³³ P. AUFMUTH,⁹ K. AULTONEAL,³⁴ C. AUSTIN,² V. AVENDANO,³⁵
A. AVILA-ALVAREZ,²⁶ S. BABAK,^{36,27} P. BACON,²⁷ F. BADARACCO,^{14,15} M. K. M. BADER,³⁷ S. BAE,³⁸ P. T. BAKER,³⁹
F. BALDACCINI,^{40,41} G. BALLARDIN,²⁸ S. W. BALLMER,⁴² S. BANAGIRI,⁴³ J. C. BARAYOGA,¹ S. E. BARCLAY,⁴⁴
B. C. BARISH,¹ D. BARKER,⁴⁵ K. BARKETT,⁴⁶ S. BARNUM,¹² F. BARONE,^{4,5} B. BARR,⁴⁴ L. BARSOTTI,¹² M. BARSUGLIA,²⁷
D. BARTA,⁴⁷ J. BARTLETT,⁴⁵ I. BARTOS,⁴⁸ R. BASSIRI,⁴⁹ A. BASTI,^{18,19} M. BAWAJ,^{50,41} J. C. BAYLEY,⁴⁴ M. BAZZAN,^{51,52}
B. BÉCSY,⁵³ M. BEJGER,^{27,54} I. BELAHcene,²⁵ A. S. BELL,⁴⁴ D. BENIWAL,⁵⁵ B. K. BERGER,⁴⁹ G. BERGMANN,^{8,9}
S. BERNUZZI,^{56,57} J. J. BERO,⁵⁸ C. P. L. BERRY,⁵⁹ D. BERSANETTI,⁶⁰ A. BERTOLINI,³⁷ J. BETZWIESER,⁷ R. BHANDARE,⁶¹
J. BIDLER,²⁶ I. A. BILENKO,⁶² S. A. BILGILI,³⁹ G. BILLINGSLEY,¹ J. BIRCH,⁷ R. BIRNEY,²⁴ O. BIRNHOLTZ,⁵⁸ S. BISCANS,^{1,12}
S. BISCOVEANU,⁶ A. BISHI,⁹ M. BITOSSI,^{28,19} M. A. BIZOUARD,²⁵ J. K. BLACKBURN,¹ C. D. BLAIR,⁷ D. G. BLAIR,⁶³
R. M. BLAIR,⁴⁵ S. BLOEMEN,⁶⁴ N. BODE,^{8,9} M. BOER,⁶⁵ Y. BOETZEL,⁶⁶ G. BOGAERT,⁶⁵ F. BONDU,⁶⁷ E. BONILLA,⁴⁹
R. BONNAND,³³ P. BOOKER,^{8,9} B. A. BOOM,³⁷ C. D. BOOTH,⁶⁸ R. BORK,¹ V. BOSCHI,²⁸ S. BOSE,^{69,3} K. BOSSIE,⁷
V. BOSSILKOV,⁶³ J. BOSVELD,⁶³ Y. BOUFFANAI,²⁷ A. BOZZI,²⁸ C. BRADASCHIA,¹⁹ P. R. BRADY,²³ A. BRAMLEY,⁷
M. BRANCHESI,^{14,15} J. E. BRAU,⁷⁰ T. BRIANT,⁷¹ J. H. BRIGGS,⁴⁴ F. BRIGHENTI,^{72,73} A. BRILLET,⁶⁵ M. BRINKMANN,^{8,9}
V. BRISSON,^{25,*} P. BROCKILL,²³ A. F. BROOKS,¹ D. D. BROWN,⁵⁵ S. BRUNETT,¹ A. BUIKEMA,¹² T. BULIK,⁷⁴
H. J. BULTEN,^{75,37} A. BUONANNO,^{36,76} D. BUSKULIC,³³ C. BUY,²⁷ R. L. BYER,⁴⁹ M. CABERO,^{8,9} L. CADONATI,⁷⁷
G. CAGNOLI,^{22,78} C. CAHILLANE,¹ J. CALDERÓN BUSTILLO,⁶ T. A. CALLISTER,¹ E. CALLONI,^{79,5} J. B. CAMP,⁸⁰
W. A. CAMPBELL,⁶ K. C. CANNON,⁸¹ H. CAO,⁵⁵ J. CAO,⁸² E. CAPOCASA,²⁷ F. CARBOGNANI,²⁸ S. CARIDE,⁸³
M. F. CARNEY,⁵⁹ G. CARULLO,¹⁸ J. CASANUEVA DIAZ,¹⁹ C. CASENTINI,^{30,31} S. CAUDILL,³⁷ M. CAVAGLIÀ,⁸⁴ F. CAVALIER,²⁵
R. CAVALIERI,²⁸ G. CELLA,¹⁹ P. CERDÁ-DURÁN,²⁰ G. CERRETANI,^{18,19} E. CESARINI,^{85,31} O. CHAIBI,⁶⁵ K. CHAKRAVARTI,³
S. J. CHAMBERLIN,⁸⁶ M. CHAN,⁴⁴ S. CHAO,⁸⁷ P. CHARLTON,⁸⁸ E. A. CHASE,⁵⁹ E. CHASSANDE-MOTTIN,²⁷
D. CHATTERJEE,²³ M. CHATURVEDI,⁶¹ B. D. CHEESEBORO,³⁹ H. Y. CHEN,⁸⁹ X. CHEN,⁶³ Y. CHEN,⁴⁶ H.-P. CHENG,⁴⁸
C. K. CHEONG,⁹⁰ H. Y. CHIA,⁴⁸ A. CHINCARINI,⁶⁰ A. CHIUMMO,²⁸ G. CHO,⁹¹ H. S. CHO,⁹² M. CHO,⁷⁶
N. CHRISTENSEN,^{65,93} Q. CHU,⁶³ S. CHUA,⁷¹ K. W. CHUNG,⁹⁰ S. CHUNG,⁶³ G. CIANI,^{51,52} A. A. CIOBANU,⁵⁵ R. CIOLFI,^{94,95}
F. CIPRIANO,⁶⁵ A. CIRONE,^{96,60} F. CLARA,⁴⁵ J. A. CLARK,⁷⁷ P. CLEARWATER,⁹⁷ F. CLEVA,⁶⁵ C. COCCHERI,⁸⁴
E. COCCIA,^{14,15} P.-F. COHADON,⁷¹ D. COHEN,²⁵ R. COLGAN,⁹⁸ M. COLLEONI,⁹⁹ C. G. COLLETTE,¹⁰⁰ C. COLLINS,¹¹
L. R. COMINSKY,¹⁰¹ M. CONSTANCIO JR.,¹³ L. CONTI,⁵² S. J. COOPER,¹¹ P. CORBAN,⁷ T. R. CORBITT,²
I. CORDERO-CARRIÓN,¹⁰² K. R. CORLEY,⁹⁸ N. CORNISH,⁵³ A. CORSI,⁸³ S. CORTESE,²⁸ C. A. COSTA,¹³ R. COTESTA,³⁶
M. W. COUGHLIN,¹ S. B. COUGHLIN,^{68,59} J.-P. COULON,⁶⁵ S. T. COUNTRYMAN,⁹⁸ P. COUVARES,¹ P. B. COVAS,⁹⁹
E. E. COWAN,⁷⁷ D. M. COWARD,⁶³ M. J. COWART,⁷ D. C. COYNE,¹ R. COYNE,¹⁰³ J. D. E. CREIGHTON,²³
T. D. CREIGHTON,¹⁰⁴ J. CRIPE,² M. CROQUETTE,⁷¹ S. G. CROWDER,¹⁰⁵ T. J. CULLEN,² A. CUMMING,⁴⁴ L. CUNNINGHAM,⁴⁴
E. CUOCO,²⁸ T. DAL CANTON,⁸⁰ G. DÁLYA,¹⁰⁶ S. L. DANILISHIN,^{8,9} S. D'ANTONIO,³¹ K. DANZMANN,^{9,8} A. DASGUPTA,¹⁰⁷
C. F. DA SILVA COSTA,⁴⁸ L. E. H. DATRIER,⁴⁴ V. DATTILO,²⁸ I. DAVE,⁶¹ M. DAVIER,²⁵ D. DAVIS,⁴² E. J. DAW,¹⁰⁸
D. DEBRA,⁴⁹ M. DEENADAYALAN,³ J. DEGALLAIX,²² M. DE LAURENTIS,^{79,5} S. DELÉGLISE,⁷¹ W. DEL POZZO,^{18,19}
L. M. DEMARCHI,⁵⁹ N. DEMOS,¹² T. DENT,^{8,9} M. DENYS,⁷⁴ R. DE PIETRI,^{109,57} J. DERBY,²⁶ R. DE ROSA,^{79,5}
C. DE ROSSI,^{22,28} R. DESALVO,¹¹⁰ O. DE VARONA,^{8,9} S. DHURANDHAR,³ M. C. DÍAZ,¹⁰⁴ T. DIETRICH,³⁷ L. DI FIORE,⁵
M. DI GIOVANNI,^{111,95} T. DI GIROLAMO,^{79,5} A. DI LIETO,^{18,19} B. DING,¹⁰⁰ S. DI PACE,^{112,32} I. DI PALMA,^{112,32}
F. DI RENZO,^{18,19} A. DMITRIEV,¹¹ Z. DOCTOR,⁸⁹ F. DONOVAN,¹² K. L. DOOLEY,^{68,84} S. DORAVARI,^{8,9} I. DORRINGTON,⁶⁸
T. P. DOWNES,²³ M. DRAGO,^{14,15} J. C. DRIGGERS,⁴⁵ Z. DU,⁸² J.-G. DUCOIN,²⁵ P. DUPEJ,⁴⁴ S. E. DWYER,⁴⁵
P. J. EASTER,⁶ T. B. EDO,¹⁰⁸ M. C. EDWARDS,⁹³ A. EFFLER,⁷ P. EHRENS,¹ J. EICHHOLZ,¹ S. S. EIKENBERRY,⁴⁸
M. EISENMANN,³³ R. A. EISENSTEIN,¹² R. C. ESSICK,⁸⁹ H. ESTELLES,⁹⁹ D. ESTEVEZ,³³ Z. B. ETIENNE,³⁹ T. ETZEL,¹
M. EVANS,¹² T. M. EVANS,⁷ V. FAFONE,^{30,31,14} H. FAIR,⁴² S. FAIRHURST,⁶⁸ X. FAN,⁸² S. FARINON,⁶⁰ B. FARR,⁷⁰
W. M. FARR,¹¹ E. J. FAUCHON-JONES,⁶⁸ M. FAVATA,³⁵ M. FAYS,¹⁰⁸ M. FAZIO,¹¹³ C. FEE,¹¹⁴ J. FEICHT,¹ M. M. FEJER,⁴⁹
F. FENG,²⁷ A. FERNANDEZ-GALIANA,¹² I. FERRANTE,^{18,19} E. C. FERREIRA,¹³ T. A. FERREIRA,¹³ F. FERRINI,²⁸
F. FIDECARO,^{18,19} I. FIORI,²⁸ D. FIORUCCI,²⁷ M. FISHBACH,⁸⁹ R. P. FISHER,^{42,115} J. M. FISHNER,¹² M. FITZ-AXEN,⁴³
R. FLAMINIO,^{33,116} M. FLETCHER,⁴⁴ E. FLYNN,²⁶ H. FONG,¹¹⁷ J. A. FONT,^{20,118} P. W. F. FORSYTH,²¹ J.-D. FOURNIER,⁶⁵
S. FRASCA,^{112,32} F. FRASCONI,¹⁹ Z. FREI,¹⁰⁶ A. FREISE,¹¹ R. FREY,⁷⁰ V. FREY,²⁵ P. FRITSCHEL,¹² V. V. FROLOV,⁷

- P. FULDA,⁴⁸ M. FYFFE,⁷ H. A. GABBARD,⁴⁴ B. U. GADRE,³ S. M. GAEBEL,¹¹ J. R. GAIR,¹¹⁹ L. GAMMAITONI,⁴⁰ M. R. GANIJA,⁵⁵ S. G. GAONKAR,³ A. GARCIA,²⁶ C. GARCÍA-QUIRÓS,⁹⁹ F. GARUFI,^{79,5} B. GATELEY,⁴⁵ S. GAUDIO,³⁴ G. GAUR,¹²⁰ V. GAYATHRI,¹²¹ G. GEMME,⁶⁰ E. GENIN,²⁸ A. GENNAI,¹⁹ D. GEORGE,¹⁷ J. GEORGE,⁶¹ L. GERGELY,¹²² V. GERMAIN,³³ S. GHONGE,⁷⁷ ABHIRUP GHOSH,¹⁶ ARCHISMAN GHOSH,³⁷ S. GHOSH,²³ B. GIACOMAZZO,^{111,95} J. A. GIAIME,^{2,7} K. D. GIARDINA,⁷ A. GIAZOTTO,^{19,†} K. GILL,³⁴ G. GIORDANO,^{4,5} L. GLOVER,¹¹⁰ P. GODWIN,⁸⁶ E. GOETZ,⁴⁵ R. GOETZ,⁴⁸ B. GONCHAROV,⁶ G. GONZÁLEZ,² J. M. GONZALEZ CASTRO,^{18,19} A. GOPAKUMAR,¹²³ M. L. GORODETSKY,⁶² S. E. GOSSAN,¹ M. GOSSELIN,²⁸ R. GOUATY,³³ A. GRADO,^{124,5} C. GRAEF,⁴⁴ M. GRANATA,²² A. GRANT,⁴⁴ S. GRAS,¹² P. GRASSIA,¹ C. GRAY,⁴⁵ R. GRAY,⁴⁴ G. GRECO,^{72,73} A. C. GREEN,^{11,48} R. GREEN,⁶⁸ E. M. GRETARSSON,³⁴ P. GROOT,⁶⁴ H. GROTE,⁶⁸ S. GRUNEWALD,³⁶ P. GRUNING,²⁵ G. M. GUIDI,^{72,73} H. K. GULATI,¹⁰⁷ Y. GUO,³⁷ A. GUPTA,⁸⁶ M. K. GUPTA,¹⁰⁷ E. K. GUSTAFSON,¹ R. GUSTAFSON,¹²⁵ L. HAEGEL,⁹⁹ O. HALIM,^{15,14} B. R. HALL,⁶⁹ E. D. HALL,¹² E. Z. HAMILTON,⁶⁸ G. HAMMOND,⁴⁴ M. HANEY,⁶⁶ M. M. HANKE,^{8,9} J. HANKS,⁴⁵ C. HANNA,⁸⁶ O. A. HANNUKSELA,⁹⁰ J. HANSON,⁷ T. HARDWICK,² K. HARIS,¹⁶ J. HARMS,^{14,15} G. M. HARRY,¹²⁶ I. W. HARRY,³⁶ C.-J. HASTER,¹¹⁷ K. HAUGHIAN,⁴⁴ F. J. HAYES,⁴⁴ J. HEALY,⁵⁸ A. HEIDMANN,⁷¹ M. C. HEINTZE,⁷ H. HEITMANN,⁶⁵ P. HELLO,²⁵ G. HEMMING,²⁸ M. HENDRY,⁴⁴ I. S. HENG,⁴⁴ J. HENNIG,^{8,9} A. W. HEPTONSTALL,¹ F. J. HERNANDEZ,⁶ M. HEURS,^{8,9} S. HILD,⁴⁴ T. HINDERER,^{127,37,128} D. HOAK,²⁸ S. HOCHHEIM,^{8,9} D. HOFMAN,²² A. M. HOLGADO,¹⁷ N. A. HOLLAND,²¹ K. HOLT,⁷ D. E. HOLZ,⁸⁹ P. HOPKINS,⁶⁸ C. HORST,²³ J. HOUGH,⁴⁴ E. J. HOWELL,⁶³ C. G. HOY,⁶⁸ A. HREIBI,⁶⁵ E. A. HUERTA,¹⁷ D. HUET,²⁵ B. HUGHEY,³⁴ M. HULKO,¹ S. HUSA,⁹⁹ S. H. HUTTNER,⁴⁴ T. HUYNH-DINH,⁷ B. IDZKOWSKI,⁷⁴ A. IESS,^{30,31} C. INGRAM,⁵⁵ R. INTA,⁸³ G. INTINI,^{112,32} B. IRWIN,¹¹⁴ H. N. ISA,⁴⁴ J.-M. ISAC,⁷¹ M. ISI,¹ B. R. IYER,¹⁶ K. IZUMI,⁴⁵ T. JACQMIN,⁷¹ S. J. JADHAV,¹²⁹ K. JANI,⁷⁷ N. N. JANTHALUR,¹²⁹ P. JARANOWSKI,¹³⁰ A. C. JENKINS,¹³¹ J. JIANG,⁴⁸ D. S. JOHNSON,¹⁷ A. W. JONES,¹¹ D. I. JONES,¹³² R. JONES,⁴⁴ R. J. G. JONKER,³⁷ L. JU,⁶³ J. JUNKER,^{8,9} C. V. KALAGHATGI,⁶⁸ V. KALOGERA,⁵⁹ B. KAMAI,¹ S. KANDHASAMY,⁸⁴ G. KANG,³⁸ J. B. KANNER,¹ S. J. KAPADIA,²³ S. KARKI,⁷⁰ K. S. KARVINEN,^{8,9} R. KASHYAP,¹⁶ M. KASPRZACK,¹ S. KATSANEVAS,²⁸ E. KATSAVOUNIDIS,¹² W. KATZMAN,⁷ S. KAUFER,⁹ K. KAWABE,⁴⁵ N. V. KEERTHANA,³ F. KÉFÉLIAN,⁶⁵ D. KEITEL,⁴⁴ R. KENNEDY,¹⁰⁸ J. S. KEY,¹³³ F. Y. KHALILI,⁶² H. KHAN,²⁶ I. KHAN,^{14,31} S. KHAN,^{8,9} Z. KHAN,¹⁰⁷ E. A. KHAZANOV,¹³⁴ M. KHURSHEED,⁶¹ N. KJBUNCHOO,²¹ CHUNGLEE KIM,¹³⁵ J. C. KIM,¹³⁶ K. KIM,⁹⁰ W. KIM,⁵⁵ W. S. KIM,¹³⁷ Y.-M. KIM,¹³⁸ C. KIMBALL,⁵⁹ E. J. KING,⁵⁵ P. J. KING,⁴⁵ M. KINLEY-HANLON,¹²⁶ R. KIRCHHOFF,^{8,9} J. S. KISSEL,⁴⁵ L. KLEYBOLTE,¹³⁹ J. H. KLIKA,²³ S. KLIMENTKO,⁴⁸ T. D. KNOWLES,³⁹ P. KOCH,^{8,9} S. M. KOEHLENBECK,^{8,9} G. KOEKOEK,^{37,140} S. KOLEY,³⁷ V. KONDRASTOV,¹ A. KONTOS,¹² N. KOPER,^{8,9} M. KOROBKO,¹³⁹ W. Z. KORTH,¹ I. KOWALSKA,⁷⁴ D. B. KOZAK,¹ V. KRINGEL,^{8,9} N. KRISHNENDU,²⁹ A. KRÓLAK,^{141,142} G. KUEHN,^{8,9} A. KUMAR,¹²⁹ P. KUMAR,¹⁴³ R. KUMAR,¹⁰⁷ S. KUMAR,¹⁶ L. KUO,⁸⁷ A. KUTYNIA,¹⁴¹ S. KWANG,²³ B. D. LACKEY,³⁶ K. H. LAI,⁹⁰ T. L. LAM,⁹⁰ M. LANDRY,⁴⁵ B. B. LANE,¹² R. N. LANG,¹⁴⁴ J. LANGE,⁵⁸ B. LANTZ,⁴⁹ R. K. LANZA,¹² A. LARTAUX-VOLLARD,²⁵ P. D. LASKY,⁶ M. LAXEN,⁷ A. LAZZARINI,¹ C. LAZZARO,⁵² P. LEACI,^{112,32} S. LEAVEY,^{8,9} Y. K. LECOEUCHE,⁴⁵ C. H. LEE,⁹² H. K. LEE,¹⁴⁵ H. M. LEE,¹⁴⁶ H. W. LEE,¹³⁶ J. LEE,⁹¹ K. LEE,⁴⁴ J. LEHMANN,^{8,9} A. LENON,³⁹ N. LEROY,²⁵ N. LETENDRE,³³ Y. LEVIN,^{6,98} J. LI,⁸² K. J. L. LI,⁹⁰ T. G. F. LI,⁹⁰ X. LI,⁴⁶ F. LIN,⁶ F. LINDE,³⁷ S. D. LINKER,¹¹⁰ T. B. LITTENBERG,¹⁴⁷ J. LIU,⁶³ X. LIU,²³ R. K. L. LO,^{90,1} N. A. LOCKERBIE,²⁴ L. T. LONDON,⁶⁸ A. LONGO,^{148,149} M. LORENZINI,^{14,15} V. LORIETTE,¹⁵⁰ M. LORMAND,⁷ G. LOSURDO,¹⁹ J. D. LOUGH,^{8,9} C. O. LOUSTO,⁵⁸ G. LOVELACE,²⁶ M. E. LOWER,¹⁵¹ H. LÜCK,^{9,8} D. LUMACA,^{30,31} A. P. LUNDGREN,¹⁵² R. LYNCH,¹² Y. MA,⁴⁶ R. MACAS,⁶⁸ S. MACFOY,²⁴ M. MACINNIS,¹² D. M. MACLEOD,⁶⁸ A. MACQUET,⁶⁵ F. MAGAÑA-SANDOVAL,⁴² L. MAGAÑA ZERTUCHE,⁸⁴ R. M. MAGEE,⁸⁶ E. MAJORANA,³² I. MAKSIMOVIC,¹⁵⁰ A. MALIK,⁶¹ N. MAN,⁶⁵ V. MANDIC,⁴³ V. MANGANO,⁴⁴ G. L. MANSELL,^{45,12} M. MANSKE,^{23,21} M. MANTOVANI,²⁸ F. MARCHESONI,^{50,41} F. MARION,³³ S. MÁRKA,⁹⁸ Z. MÁRKA,⁹⁸ C. MARKAKIS,^{10,17} A. S. MARKOSYAN,⁴⁹ A. MARKOWITZ,¹ E. MAROS,¹ A. MARQUINA,¹⁰² S. MARSAT,³⁶ F. MARTELLI,^{72,73} I. W. MARTIN,⁴⁴ R. M. MARTIN,³⁵ D. V. MARTYNOV,¹¹ K. MASON,¹² E. MASSERA,¹⁰⁸ A. MASSEROT,³³ T. J. MASSINGER,¹ M. MASSO-REID,⁴⁴ S. MASTROGIOVANNI,^{112,32} A. MATAS,^{43,36} F. MATICHARD,^{1,12} L. MATONE,⁹⁸ N. MAVALVALA,¹² N. MAZUMDER,⁶⁹ J. J. McCANN,⁶³ R. McCarthy,⁴⁵ D. E. MCCLELLAND,²¹ S. MCCORMICK,⁷ L. McCULLER,¹² S. C. McGuire,¹⁵³ J. McIVER,¹ D. J. McMANUS,²¹ T. MCRAE,²¹ S. T. McWILLIAMS,³⁹ D. MEACHER,⁸⁶ G. D. MEADORS,⁶ M. MEHMET,^{8,9} A. K. MEHTA,¹⁶ J. MEIDAM,³⁷ A. MELATOS,⁹⁷ G. MENDELL,⁴⁵ R. A. MERCER,²³ L. MERENI,²² E. L. MERILH,⁴⁵ M. MERZOUGUI,⁶⁵ S. MESHKOV,¹ C. MESSENGER,⁴⁴ C. MESSICK,⁸⁶ R. METZDORFF,⁷¹ P. M. MEYERS,⁹⁷ H. MIAO,¹¹ C. MICHEL,²² H. MIDDLETON,⁹⁷ E. E. MIKHAILOV,¹⁵⁴ L. MILANO,^{79,5} A. L. MILLER,⁴⁸ A. MILLER,^{112,32} M. MILLHOUSE,⁵³ J. C. MILLS,⁶⁸ M. C. MILOVICH-GOFF,¹¹⁰ O. MINAZZOLI,^{65,155} Y. MINENKOV,³¹ A. MISHKIN,⁴⁸ C. MISHRA,¹⁵⁶ T. MISTRY,¹⁰⁸ S. MITRA,³ V. P. MITROFANOV,⁶² G. MITSELMAKER,⁴⁸ R. MITTLEMAN,¹² G. MO,⁹³ D. MOFFA,¹¹⁴ K. MOGUSHI,⁸⁴ S. R. P. MOHAPATRA,¹² M. MONTANI,^{72,73} C. J. MOORE,¹⁰ D. MORARU,⁴⁵ G. MORENO,⁴⁵ S. MORISAKI,⁸¹ B. MOURS,³³ C. M. MOW-LOWRY,¹¹ ARUNAVA MUKHERJEE,^{8,9} D. MUKHERJEE,²³ S. MUKHERJEE,¹⁰⁴ N. MUKUND,³ A. MULLAVY,⁷ J. MUNCH,⁵⁵ E. A. MUÑIZ,⁴² M. MURATORE,³⁴ P. G. MURRAY,⁴⁴ A. NAGAR,^{85,157,158} I. NARDECCHIA,^{30,31} L. NATICCHIONI,^{112,32} R. K. NAYAK,¹⁵⁹ J. NEILSON,¹¹⁰ G. NELEMANS,^{64,37} T. J. N. NELSON,⁷ M. NERY,^{8,9} A. NEUNZERT,¹²⁵ K. Y. NG,¹² S. NG,⁵⁵ P. NGUYEN,⁷⁰ D. NICHOLS,^{127,37} S. NISSANKE,^{127,37} F. NOCERA,²⁸ C. NORTH,⁶⁸ L. K. NUTTALL,¹⁵² M. OBERGAULINGER,²⁰ J. OBERLING,⁴⁵ B. D. O'BRIEN,⁴⁸ G. D. O'DEA,¹¹⁰ G. H. OGIN,¹⁶⁰ J. J. OH,¹³⁷ S. H. OH,¹³⁷ F. OHME,^{8,9} H. OHTA,⁸¹ M. A. OKADA,¹³ M. OLIVER,⁹⁹ P. OPPERMANN,^{8,9} RICHARD J. ORAM,⁷ B. O'REILLY,⁷ R. G. ORMISTON,⁴³ L. F. ORTEGA,⁴⁸ R. O'SHAUGHNESSY,⁵⁸ S. OSSOKINE,³⁶ D. J. OTTAWAY,⁵⁵ H. OVERMIER,⁷ B. J. OWEN,⁸³ A. E. PACE,⁸⁶ G. PAGANO,^{18,19} M. A. PAGE,⁶³ A. PAI,¹²¹ S. A. PAI,⁶¹ J. R. PALAMOS,⁷⁰ O. PALASHOV,¹³⁴

- C. PALOMBA,³² A. PAL-SINGH,¹³⁹ HUANG-WEI PAN,⁸⁷ B. PANG,⁴⁶ P. T. H. PANG,⁹⁰ C. PANKOW,⁵⁹ F. PANNARALE,^{112,32}
 B. C. PANT,⁶¹ F. PAOLETTI,¹⁹ A. PAOLI,²⁸ A. PARIDA,³ W. PARKER,^{7,153} D. PASCUCCI,⁴⁴ A. PASQUALETTI,²⁸
 R. PASSAQUIETI,^{18,19} D. PASSUELLO,¹⁹ M. PATIL,¹⁴² B. PATRICELLI,^{18,19} B. L. PEARLSTONE,⁴⁴ C. PEDERSEN,⁶⁸
 M. PEDRAZA,¹ R. PEDURAND,^{22,161} A. PELE,⁷ S. PENN,¹⁶² C. J. PEREZ,⁴⁵ A. PERRECA,^{111,95} H. P. PFEIFFER,^{36,117}
 M. PHELPS,^{8,9} K. S. PHUKON,³ O. J. PICCINNI,^{112,32} M. PICHOT,⁶⁵ F. PIERGIOVANNI,^{72,73} G. PILLANT,²⁸ L. PINARD,²²
 M. PIRELLO,⁴⁵ M. PITKIN,⁴⁴ R. POGGIANI,^{18,19} D. Y. T. PONG,⁹⁰ S. PONRATHNAM,³ P. POPOLIZIO,²⁸ E. K. PORTER,²⁷
 J. POWELL,¹⁵¹ A. K. PRAJAPATI,¹⁰⁷ J. PRASAD,³ K. PRASAI,⁴⁹ R. PRASANNA,¹²⁹ G. PRATTEN,⁹⁹ T. PRESTEGARD,²³
 S. PRIVITERA,³⁶ G. A. PRODI,^{111,95} L. G. PROKHOROV,⁶² O. PUNCKEN,^{8,9} M. PUNTURO,⁴¹ P. PUPPO,³² M. PÜRRER,³⁶
 H. QI,²³ V. QUETSCHKE,¹⁰⁴ P. J. QUINONEZ,³⁴ E. A. QUINTERO,¹ R. QUITZOW-JAMES,⁷⁰ F. J. RAAB,⁴⁵ H. RADKINS,⁴⁵
 N. RADULESCU,⁶⁵ P. RAFFAI,¹⁰⁶ S. RAJA,⁶¹ C. RAJAN,⁶¹ B. RAJBHANDARI,⁸³ M. RAKHMANOV,¹⁰⁴ K. E. RAMIREZ,¹⁰⁴
 A. RAMOS-BUADES,⁹⁹ JAVED RANA,³ K. RAO,⁵⁹ P. RAPAGNANI,^{112,32} V. RAYMOND,⁶⁸ M. RAZZANO,^{18,19} J. READ,²⁶
 T. REGIMBAU,³³ L. REI,⁶⁰ S. REID,²⁴ D. H. REITZE,^{1,48} W. REN,¹⁷ F. RICCI,^{112,32} C. J. RICHARDSON,³⁴
 J. W. RICHARDSON,¹ P. M. RICKER,¹⁷ K. RILES,¹²⁵ M. RIZZO,⁵⁹ N. A. ROBERTSON,^{1,44} R. ROBIE,⁴⁴ F. ROBINET,²⁵
 A. ROCCHI,³¹ L. ROLLAND,³³ J. G. ROLLINS,¹ V. J. ROMA,⁷⁰ M. ROMANELLI,⁶⁷ R. ROMANO,^{4,5} C. L. ROMEL,⁴⁵
 J. H. ROMIE,⁷ K. ROSE,¹¹⁴ D. ROSIŃSKA,^{163,54} S. G. ROSOFSKY,¹⁷ M. P. ROSS,¹⁶⁴ S. ROWAN,⁴⁴ A. RÜDIGER,^{8,9,‡}
 P. RUGGI,²⁸ G. RUTINS,¹⁶⁵ K. RYAN,⁴⁵ S. SACHDEV,¹ T. SADECKI,⁴⁵ M. SAKELLARIADOU,¹³¹ L. SALCONI,²⁸ M. SALEEM,²⁹
 A. SAMAJDAR,³⁷ L. SAMMUT,⁶ E. J. SANCHEZ,¹ L. E. SANCHEZ,¹ N. SANCHIS-GUAL,²⁰ V. SANDBERG,⁴⁵ J. R. SANDERS,⁴²
 K. A. SANTIAGO,³⁵ N. SARIN,⁶ B. SASSOLAS,²² P. R. SAULSON,⁴² O. SAUTER,¹²⁵ R. L. SAVAGE,⁴⁵ P. SCHALE,⁷⁰
 M. SCHEEL,⁴⁶ J. SCHEUER,⁵⁹ P. SCHMIDT,⁶⁴ R. SCHNABEL,¹³⁹ R. M. S. SCHOFIELD,⁷⁰ A. SCHÖNBECK,¹³⁹ E. SCHREIBER,^{8,9}
 B. W. SCHULTE,^{8,9} B. F. SCHUTZ,⁶⁸ S. G. SCHWALBE,³⁴ J. SCOTT,⁴⁴ S. M. SCOTT,²¹ E. SEIDEL,¹⁷ D. SELLERS,⁷
 A. S. SENGUPTA,¹⁶⁶ N. SENNETT,³⁶ D. SENTENAC,²⁸ V. SEQUINO,^{30,31,14} A. SERGEEV,¹³⁴ Y. SETYAWATI,^{8,9}
 D. A. SHADDOCK,²¹ T. SHAFFER,⁴⁵ M. S. SHAHRIAR,⁵⁹ M. B. SHANER,¹¹⁰ L. SHAO,³⁶ P. SHARMA,⁶¹ P. SHAWHAN,⁷⁶
 H. SHEN,¹⁷ R. SHINK,¹⁶⁷ D. H. SHOEMAKER,¹² D. M. SHOEMAKER,⁷⁷ S. SHYAMSUNDAR,⁶¹ K. SIELLEZ,⁷⁷ M. SIENIAWSKA,⁵⁴
 D. SIGG,⁴⁵ A. D. SILVA,¹³ L. P. SINGER,⁸⁰ N. SINGH,⁷⁴ A. SINGHAL,^{14,32} A. M. SINTES,⁹⁹ S. SITMUKHAMBECHOV,¹⁰⁴
 V. SKLIRIS,⁶⁸ B. J. J. SLAGMOLEN,²¹ T. J. SLAVEN-BLAIR,⁶³ J. R. SMITH,²⁶ R. J. E. SMITH,⁶ S. SOMALA,¹⁶⁸ E. J. SON,¹³⁷
 B. SORAZU,⁴⁴ F. SORRENTINO,⁶⁰ T. SOURADEEP,³ E. SOWELL,⁸³ A. P. SPENCER,⁴⁴ A. K. SRIVASTAVA,¹⁰⁷ V. SRIVASTAVA,⁴²
 K. STAATS,⁵⁹ C. STACHIE,⁶⁵ M. STANDKE,^{8,9} D. A. STEER,²⁷ M. STEINKE,^{8,9} J. STEINLECHNER,^{139,44} S. STEINLECHNER,¹³⁹
 D. STEINMEYER,^{8,9} S. P. STEVENSON,¹⁵¹ D. STOCKS,⁴⁹ R. STONE,¹⁰⁴ D. J. STOPS,¹¹ K. A. STRAIN,⁴⁴ G. STRATTA,^{72,73}
 S. E. STRING,⁶² A. STRUNK,⁴⁵ R. STURANI,¹⁶⁹ A. L. STUVER,¹⁷⁰ V. SUDHIR,¹² T. Z. SUMMERSCALES,¹⁷¹ L. SUN,¹
 S. SUNIL,¹⁰⁷ J. SURESH,³ P. J. SUTTON,⁶⁸ B. L. SWINKELS,³⁷ M. J. SZCZEPANČZYK,³⁴ M. TACCA,³⁷ S. C. TAIT,⁴⁴
 C. TALBOT,⁶ D. TALUKDER,⁷⁰ D. B. TANNER,⁴⁸ M. TÁPAI,¹²² A. TARACCHINI,³⁶ J. D. TASSON,⁹³ R. TAYLOR,¹ F. THIES,^{8,9}
 M. THOMAS,⁷ P. THOMAS,⁴⁵ S. R. THONDAPU,⁶¹ K. A. THORNE,⁷ E. THRANE,⁶ SHUBHANSHU TIWARI,^{111,95}
 SRISHTI TIWARI,¹²³ V. TIWARI,⁶⁸ K. TOLAND,⁴⁴ M. TONELLI,^{18,19} Z. TORNASI,⁴⁴ A. TORRES-FORNÉ,¹⁷² C. I. TORRIE,¹
 D. TÖYRÄ,¹¹ F. TRAVASSO,^{28,41} G. TRAYLOR,⁷ M. C. TRINGALI,⁷⁴ A. TROVATO,²⁷ L. TROZZO,^{173,19} R. TRUDEAU,¹
 K. W. TSANG,³⁷ M. TSE,¹² R. TSO,⁴⁶ L. TSUKADA,⁸¹ D. TSUNA,⁸¹ D. TUYENBAYEV,¹⁰⁴ K. UENO,⁸¹ D. UGOLINI,¹⁷⁴
 C. S. UNNIKRISHNAN,¹²³ A. L. URBAN,² S. A. USMAN,⁶⁸ H. VAHLBRUCH,⁹ G. VAJENTE,¹ G. VALDES,² N. VAN BAKEL,³⁷
 M. VAN BEUZEKOM,³⁷ J. F. J. VAN DEN BRAND,^{75,37} C. VAN DEN BROECK,^{37,175} D. C. VANDER-HYDE,⁴²
 L. VAN DER SCHAAF,³⁷ J. V. VAN HEIJNINGEN,⁶³ A. A. VAN VEGGEL,⁴⁴ M. VARDARO,^{51,52} V. VARMA,⁴⁶ S. VASS,¹
 M. VASÚTH,⁴⁷ A. VECCHIO,¹¹ G. VEDOVATO,⁵² J. VEITCH,⁴⁴ P. J. VEITCH,⁵⁵ K. VENKATESWARA,¹⁶⁴ G. VENUGOPALAN,¹
 D. VERKINDT,³³ F. VETRANO,^{72,73} A. VICERÉ,^{72,73} A. D. VIETS,²³ D. J. VINE,¹⁶⁵ J.-Y. VINET,⁶⁵ S. VITALE,¹² T. VO,⁴²
 H. VOCCA,^{40,41} C. VORVICK,⁴⁵ S. P. VYATCHANIN,⁶² A. R. WADE,¹ L. E. WADE,¹¹⁴ M. WADE,¹¹⁴ R. WALET,³⁷
 M. WALKER,²⁶ L. WALLACE,¹ S. WALSH,²³ G. WANG,^{14,19} H. WANG,¹¹ J. Z. WANG,¹²⁵ W. H. WANG,¹⁰⁴ Y. F. WANG,⁹⁰
 R. L. WARD,²¹ Z. A. WARDEN,³⁴ J. WARNER,⁴⁵ M. WAS,³³ J. WATCHI,¹⁰⁰ B. WEAVER,⁴⁵ L.-W. WEI,^{8,9} M. WEINERT,^{8,9}
 A. J. WEINSTEIN,¹ R. WEISS,¹² F. WELLMANN,^{8,9} L. WEN,⁶³ E. K. WESSEL,¹⁷ P. WESSELS,^{8,9} J. W. WESTHOUSE,³⁴
 K. WETTE,²¹ J. T. WHELAN,⁵⁸ B. F. WHITING,⁴⁸ C. WHITTLE,¹² D. M. WILKEN,^{8,9} D. WILLIAMS,⁴⁴
 A. R. WILLIAMSON,^{127,37} J. L. WILLIS,¹ B. WILLKE,^{8,9} M. H. WIMMER,^{8,9} W. WINKLER,^{8,9} C. C. WIPF,¹ H. WITTEL,^{8,9}
 G. WOAN,⁴⁴ J. WOEHLER,^{8,9} J. K. WOFFORD,⁵⁸ J. WORDEN,⁴⁵ J. L. WRIGHT,⁴⁴ D. S. WU,^{8,9} D. M. WYSOCKI,⁵⁸ L. XIAO,¹
 H. YAMAMOTO,¹ C. C. YANCEY,⁷⁶ L. YANG,¹¹³ M. J. YAP,²¹ M. YAZBACK,⁴⁸ D. W. YEELES,⁶⁸ HANG YU,¹² HAOCUN YU,¹²
 S. H. R. YUEN,⁹⁰ M. YVERT,³³ A. K. ZADROŻNY,^{104,141} M. ZANOLIN,³⁴ T. ZELENOVA,²⁸ J.-P. ZENDRI,⁵² M. ZEVIN,⁵⁹
 J. ZHANG,⁶³ L. ZHANG,¹ T. ZHANG,⁴⁴ C. ZHAO,⁶³ M. ZHOU,⁵⁹ Z. ZHOU,⁵⁹ X. J. ZHU,⁶ M. E. ZUCKER,^{1,12} AND J. ZWEIZIG¹

THE LIGO SCIENTIFIC COLLABORATION AND THE VIRGO COLLABORATION

¹LIGO, California Institute of Technology, Pasadena, CA 91125, USA²Louisiana State University, Baton Rouge, LA 70803, USA³Inter-University Centre for Astronomy and Astrophysics, Pune 411007, India⁴Università di Salerno, Fisciano, I-84084 Salerno, Italy

- ⁵INFN, Sezione di Napoli, Complesso Universitario di Monte S.Angelo, I-80126 Napoli, Italy
- ⁶OzGrav, School of Physics & Astronomy, Monash University, Clayton 3800, Victoria, Australia
- ⁷LIGO Livingston Observatory, Livingston, LA 70754, USA
- ⁸Max Planck Institute for Gravitational Physics (Albert Einstein Institute), D-30167 Hannover, Germany
- ⁹Leibniz Universität Hannover, D-30167 Hannover, Germany
- ¹⁰University of Cambridge, Cambridge CB2 1TN, United Kingdom
- ¹¹University of Birmingham, Birmingham B15 2TT, United Kingdom
- ¹²LIGO, Massachusetts Institute of Technology, Cambridge, MA 02139, USA
- ¹³Instituto Nacional de Pesquisas Espaciais, 12227-010 São José dos Campos, São Paulo, Brazil
- ¹⁴Gran Sasso Science Institute (GSSI), I-67100 L'Aquila, Italy
- ¹⁵INFN, Laboratori Nazionali del Gran Sasso, I-67100 Assergi, Italy
- ¹⁶International Centre for Theoretical Sciences, Tata Institute of Fundamental Research, Bengaluru 560089, India
- ¹⁷NCSA, University of Illinois at Urbana-Champaign, Urbana, IL 61801, USA
- ¹⁸Università di Pisa, I-56127 Pisa, Italy
- ¹⁹INFN, Sezione di Pisa, I-56127 Pisa, Italy
- ²⁰Departamento de Astronomía y Astrofísica, Universitat de València, E-46100 Burjassot, València, Spain
- ²¹OzGrav, Australian National University, Canberra, Australian Capital Territory 0200, Australia
- ²²Laboratoire des Matériaux Avancés (LMA), CNRS/IN2P3, F-69622 Villeurbanne, France
- ²³University of Wisconsin-Milwaukee, Milwaukee, WI 53201, USA
- ²⁴SUPA, University of Strathclyde, Glasgow G1 1XQ, United Kingdom
- ²⁵LAL, Univ. Paris-Sud, CNRS/IN2P3, Université Paris-Saclay, F-91898 Orsay, France
- ²⁶California State University Fullerton, Fullerton, CA 92831, USA
- ²⁷APC, AstroParticule et Cosmologie, Université Paris Diderot, CNRS/IN2P3, CEA/Irfu, Observatoire de Paris, Sorbonne Paris Cité, F-75205 Paris Cedex 13, France
- ²⁸European Gravitational Observatory (EGO), I-56021 Cascina, Pisa, Italy
- ²⁹Chennai Mathematical Institute, Chennai 603103, India
- ³⁰Università di Roma Tor Vergata, I-00133 Roma, Italy
- ³¹INFN, Sezione di Roma Tor Vergata, I-00133 Roma, Italy
- ³²INFN, Sezione di Roma, I-00185 Roma, Italy
- ³³Laboratoire d'Annecy de Physique des Particules (LAPP), Univ. Grenoble Alpes, Université Savoie Mont Blanc, CNRS/IN2P3, F-74941 Annecy, France
- ³⁴Embry-Riddle Aeronautical University, Prescott, AZ 86301, USA
- ³⁵Montclair State University, Montclair, NJ 07043, USA
- ³⁶Max Planck Institute for Gravitational Physics (Albert Einstein Institute), D-14476 Potsdam-Golm, Germany
- ³⁷Nikhef, Science Park 105, 1098 XG Amsterdam, The Netherlands
- ³⁸Korea Institute of Science and Technology Information, Daejeon 34141, South Korea
- ³⁹West Virginia University, Morgantown, WV 26506, USA
- ⁴⁰Università di Perugia, I-06123 Perugia, Italy
- ⁴¹INFN, Sezione di Perugia, I-06123 Perugia, Italy
- ⁴²Syracuse University, Syracuse, NY 13244, USA
- ⁴³University of Minnesota, Minneapolis, MN 55455, USA
- ⁴⁴SUPA, University of Glasgow, Glasgow G12 8QQ, United Kingdom
- ⁴⁵LIGO Hanford Observatory, Richland, WA 99352, USA
- ⁴⁶Caltech CaRT, Pasadena, CA 91125, USA
- ⁴⁷Wigner RCP, RMKI, H-1121 Budapest, Konkoly Thege Miklós út 29-33, Hungary
- ⁴⁸University of Florida, Gainesville, FL 32611, USA
- ⁴⁹Stanford University, Stanford, CA 94305, USA
- ⁵⁰Università di Camerino, Dipartimento di Fisica, I-62032 Camerino, Italy
- ⁵¹Università di Padova, Dipartimento di Fisica e Astronomia, I-35131 Padova, Italy
- ⁵²INFN, Sezione di Padova, I-35131 Padova, Italy
- ⁵³Montana State University, Bozeman, MT 59717, USA
- ⁵⁴Nicolaus Copernicus Astronomical Center, Polish Academy of Sciences, 00-716, Warsaw, Poland
- ⁵⁵OzGrav, University of Adelaide, Adelaide, South Australia 5005, Australia
- ⁵⁶Theoretisch-Physikalisches Institut, Friedrich-Schiller-Universität Jena, D-07743 Jena, Germany

- ⁵⁷INFN, Sezione di Milano Bicocca, Gruppo Collegato di Parma, I-43124 Parma, Italy
⁵⁸Rochester Institute of Technology, Rochester, NY 14623, USA
⁵⁹Center for Interdisciplinary Exploration & Research in Astrophysics (CIERA), Northwestern University, Evanston, IL 60208, USA
⁶⁰INFN, Sezione di Genova, I-16146 Genova, Italy
⁶¹RRCAT, Indore, Madhya Pradesh 452013, India
⁶²Faculty of Physics, Lomonosov Moscow State University, Moscow 119991, Russia
⁶³OzGrav, University of Western Australia, Crawley, Western Australia 6009, Australia
⁶⁴Department of Astrophysics/IMAPP, Radboud University Nijmegen, P.O. Box 9010, 6500 GL Nijmegen, The Netherlands
⁶⁵Artemis, Université Côte d'Azur, Observatoire Côte d'Azur, CNRS, CS 34229, F-06304 Nice Cedex 4, France
⁶⁶Physik-Institut, University of Zurich, Winterthurerstrasse 190, 8057 Zurich, Switzerland
⁶⁷Univ Rennes, CNRS, Institut FOTON - UMR6082, F-3500 Rennes, France
⁶⁸Cardiff University, Cardiff CF24 3AA, United Kingdom
⁶⁹Washington State University, Pullman, WA 99164, USA
⁷⁰University of Oregon, Eugene, OR 97403, USA
⁷¹Laboratoire Kastler Brossel, Sorbonne Université, CNRS, ENS-Université PSL, Collège de France, F-75005 Paris, France
⁷²Università degli Studi di Urbino 'Carlo Bo,' I-61029 Urbino, Italy
⁷³INFN, Sezione di Firenze, I-50019 Sesto Fiorentino, Firenze, Italy
⁷⁴Astronomical Observatory Warsaw University, 00-478 Warsaw, Poland
⁷⁵VU University Amsterdam, 1081 HV Amsterdam, The Netherlands
⁷⁶University of Maryland, College Park, MD 20742, USA
⁷⁷School of Physics, Georgia Institute of Technology, Atlanta, GA 30332, USA
⁷⁸Université Claude Bernard Lyon 1, F-69622 Villeurbanne, France
⁷⁹Università di Napoli 'Federico II,' Complesso Universitario di Monte S.Angelo, I-80126 Napoli, Italy
⁸⁰NASA Goddard Space Flight Center, Greenbelt, MD 20771, USA
⁸¹RESCEU, University of Tokyo, Tokyo, 113-0033, Japan.
⁸²Tsinghua University, Beijing 100084, China
⁸³Texas Tech University, Lubbock, TX 79409, USA
⁸⁴The University of Mississippi, University, MS 38677, USA
⁸⁵Museo Storico della Fisica e Centro Studi e Ricerche "Enrico Fermi", I-00184 Roma, Italy
⁸⁶The Pennsylvania State University, University Park, PA 16802, USA
⁸⁷National Tsing Hua University, Hsinchu City, 30013 Taiwan, Republic of China
⁸⁸Charles Sturt University, Wagga Wagga, New South Wales 2678, Australia
⁸⁹University of Chicago, Chicago, IL 60637, USA
⁹⁰The Chinese University of Hong Kong, Shatin, NT, Hong Kong
⁹¹Seoul National University, Seoul 08826, South Korea
⁹²Pusan National University, Busan 46241, South Korea
⁹³Carleton College, Northfield, MN 55057, USA
⁹⁴INAF, Osservatorio Astronomico di Padova, I-35122 Padova, Italy
⁹⁵INFN, Trento Institute for Fundamental Physics and Applications, I-38123 Povo, Trento, Italy
⁹⁶Dipartimento di Fisica, Università degli Studi di Genova, I-16146 Genova, Italy
⁹⁷OzGrav, University of Melbourne, Parkville, Victoria 3010, Australia
⁹⁸Columbia University, New York, NY 10027, USA
⁹⁹Universitat de les Illes Balears, IAC3—IEEC, E-07122 Palma de Mallorca, Spain
¹⁰⁰Université Libre de Bruxelles, Brussels 1050, Belgium
¹⁰¹Sonoma State University, Rohnert Park, CA 94928, USA
¹⁰²Departamento de Matemáticas, Universitat de València, E-46100 Burjassot, València, Spain
¹⁰³University of Rhode Island, Kingston, RI 02881, USA
¹⁰⁴The University of Texas Rio Grande Valley, Brownsville, TX 78520, USA
¹⁰⁵Bellevue College, Bellevue, WA 98007, USA
¹⁰⁶MTA-ELTE Astrophysics Research Group, Institute of Physics, Eötvös University, Budapest 1117, Hungary
¹⁰⁷Institute for Plasma Research, Bhat, Gandhinagar 382428, India
¹⁰⁸The University of Sheffield, Sheffield S10 2TN, United Kingdom
¹⁰⁹Dipartimento di Scienze Matematiche, Fisiche e Informatiche, Università di Parma, I-43124 Parma, Italy
¹¹⁰California State University, Los Angeles, 5151 State University Dr, Los Angeles, CA 90032, USA

- ¹¹¹Università di Trento, Dipartimento di Fisica, I-38123 Povo, Trento, Italy
- ¹¹²Università di Roma 'La Sapienza,' I-00185 Roma, Italy
- ¹¹³Colorado State University, Fort Collins, CO 80523, USA
- ¹¹⁴Kenyon College, Gambier, OH 43022, USA
- ¹¹⁵Christopher Newport University, Newport News, VA 23606, USA
- ¹¹⁶National Astronomical Observatory of Japan, 2-21-1 Osawa, Mitaka, Tokyo 181-8588, Japan
- ¹¹⁷Canadian Institute for Theoretical Astrophysics, University of Toronto, Toronto, Ontario M5S 3H8, Canada
- ¹¹⁸Observatori Astronòmic, Universitat de València, E-46980 Paterna, València, Spain
- ¹¹⁹School of Mathematics, University of Edinburgh, Edinburgh EH9 3FD, United Kingdom
- ¹²⁰Institute Of Advanced Research, Gandhinagar 382426, India
- ¹²¹Indian Institute of Technology Bombay, Powai, Mumbai 400 076, India
- ¹²²University of Szeged, Dóm tér 9, Szeged 6720, Hungary
- ¹²³Tata Institute of Fundamental Research, Mumbai 400005, India
- ¹²⁴INAF, Osservatorio Astronomico di Capodimonte, I-80131, Napoli, Italy
- ¹²⁵University of Michigan, Ann Arbor, MI 48109, USA
- ¹²⁶American University, Washington, D.C. 20016, USA
- ¹²⁷GRAPPA, Anton Pannekoek Institute for Astronomy and Institute of High-Energy Physics, University of Amsterdam, Science Park 904, 1098 XH Amsterdam, The Netherlands
- ¹²⁸Delta Institute for Theoretical Physics, Science Park 904, 1090 GL Amsterdam, The Netherlands
- ¹²⁹Directorate of Construction, Services & Estate Management, Mumbai 400094 India
- ¹³⁰University of Białystok, 15-424 Białystok, Poland
- ¹³¹King's College London, University of London, London WC2R 2LS, United Kingdom
- ¹³²University of Southampton, Southampton SO17 1BJ, United Kingdom
- ¹³³University of Washington Bothell, Bothell, WA 98011, USA
- ¹³⁴Institute of Applied Physics, Nizhny Novgorod, 603950, Russia
- ¹³⁵Ewha Womans University, Seoul 03760, South Korea
- ¹³⁶Inje University Gimhae, South Gyeongsang 50834, South Korea
- ¹³⁷National Institute for Mathematical Sciences, Daejeon 34047, South Korea
- ¹³⁸Ulsan National Institute of Science and Technology, Ulsan 44919, South Korea
- ¹³⁹Universität Hamburg, D-22761 Hamburg, Germany
- ¹⁴⁰Maastricht University, P.O. Box 616, 6200 MD Maastricht, The Netherlands
- ¹⁴¹NCBJ, 05-400 Świerk-Otwock, Poland
- ¹⁴²Institute of Mathematics, Polish Academy of Sciences, 00656 Warsaw, Poland
- ¹⁴³Cornell University, Ithaca, NY 14850, USA
- ¹⁴⁴Hillsdale College, Hillsdale, MI 49242, USA
- ¹⁴⁵Hanyang University, Seoul 04763, South Korea
- ¹⁴⁶Korea Astronomy and Space Science Institute, Daejeon 34055, South Korea
- ¹⁴⁷NASA Marshall Space Flight Center, Huntsville, AL 35811, USA
- ¹⁴⁸Dipartimento di Matematica e Fisica, Università degli Studi Roma Tre, I-00146 Roma, Italy
- ¹⁴⁹INFN, Sezione di Roma Tre, I-00146 Roma, Italy
- ¹⁵⁰ESPCI, CNRS, F-75005 Paris, France
- ¹⁵¹OzGrav, Swinburne University of Technology, Hawthorn VIC 3122, Australia
- ¹⁵²University of Portsmouth, Portsmouth, PO1 3FX, United Kingdom
- ¹⁵³Southern University and A&M College, Baton Rouge, LA 70813, USA
- ¹⁵⁴College of William and Mary, Williamsburg, VA 23187, USA
- ¹⁵⁵Centre Scientifique de Monaco, 8 quai Antoine Ier, MC-98000, Monaco
- ¹⁵⁶Indian Institute of Technology Madras, Chennai 600036, India
- ¹⁵⁷INFN Sezione di Torino, Via P. Giuria 1, I-10125 Torino, Italy
- ¹⁵⁸Institut des Hautes Etudes Scientifiques, F-91440 Bures-sur-Yvette, France
- ¹⁵⁹IISER-Kolkata, Mohanpur, West Bengal 741252, India
- ¹⁶⁰Whitman College, 345 Boyer Avenue, Walla Walla, WA 99362 USA
- ¹⁶¹Université de Lyon, F-69361 Lyon, France
- ¹⁶²Hobart and William Smith Colleges, Geneva, NY 14456, USA
- ¹⁶³Janusz Gil Institute of Astronomy, University of Zielona Góra, 65-265 Zielona Góra, Poland

¹⁶⁴University of Washington, Seattle, WA 98195, USA

¹⁶⁵SUPA, University of the West of Scotland, Paisley PA1 2BE, United Kingdom

¹⁶⁶Indian Institute of Technology, Gandhinagar Ahmedabad Gujarat 382424, India

¹⁶⁷Université de Montréal/Polytechnique, Montreal, Quebec H3T 1J4, Canada

¹⁶⁸Indian Institute of Technology Hyderabad, Sangareddy, Khandi, Telangana 502285, India

¹⁶⁹International Institute of Physics, Universidade Federal do Rio Grande do Norte, Natal RN 59078-970, Brazil

¹⁷⁰Villanova University, 800 Lancaster Ave, Villanova, PA 19085, USA

¹⁷¹Andrews University, Berrien Springs, MI 49104, USA

¹⁷²Max Planck Institute for Gravitationalphysik (Albert Einstein Institute), D-14476 Potsdam-Golm, Germany

¹⁷³Università di Siena, I-53100 Siena, Italy

¹⁷⁴Trinity University, San Antonio, TX 78212, USA

¹⁷⁵Van Swinderen Institute for Particle Physics and Gravity, University of Groningen, Nijenborgh 4, 9747 AG Groningen, The Netherlands

ABSTRACT

We describe directed searches for continuous gravitational waves from sixteen well localized candidate neutron stars assuming none of the stars has a binary companion. The searches were directed toward fifteen supernova remnants and Fomalhaut b, an extrasolar planet candidate which has been suggested to be a nearby old neutron star. Each search covered a broad band of frequencies and first and second time derivatives. After coherently integrating spans of data from the first Advanced LIGO observing run of 3.5–53.7 days per search, applying data-based vetoes and discounting known instrumental artifacts, we found no astrophysical signals. We set upper limits on intrinsic gravitational wave strain as strict as 1×10^{-25} , on fiducial neutron star ellipticity as strict as 2×10^{-9} , and on fiducial r -mode amplitude as strict as 3×10^{-8} .

Keywords: gravitational waves — stars: neutron — supernova remnants

* Deceased, February 2018.

† Deceased, November 2017.

‡ Deceased, July 2018.

1. INTRODUCTION

With the detections of several binary black hole mergers (Abbott et al. 2016c,b, 2017e,g,f) and one binary neutron star merger (Abbott et al. 2017h) seen also in electromagnetic waves (Abbott et al. 2017i), Advanced LIGO and Virgo have spectacularly inaugurated the field of gravitational wave (GW) astronomy. While the binary neutron star merger has had far-reaching implications for our knowledge of neutron star matter (De et al. 2018; Abbott et al. 2018), a continuous GW signal could teach us even more—not just about bulk properties but internal magnetic fields, the extent and strength of crystalline phases, and potentially other microphysics of extreme matter (Owen 2009; Glampedakis & Gualtieri 2017).

Young isolated neutron stars are promising sources of continuous GWs. The spin-downs of young pulsars are rapid enough to include significant continuous GW emission, as shown by the latest GW search for known pulsars (Abbott et al. 2017d,c). Theoretical arguments suggest that *r*-modes (oscillations dominated by the Coriolis force) might remain unstable and detectable in neutron stars up to a few thousand years old (Owen 2010, and references therein). Most young supernova remnants (SNRs) do not contain known pulsars (Green 2014). On the other hand, many of these SNRs contain small pulsar wind nebulae (PWNe), central compact objects (CCOs), or other well localized non-pulsing candidate neutron stars. Also, some of these SNRs are young enough that a neutron star could not have been kicked far, and thus the star can be considered well localized even if it is not seen at all. GW searches directed at single sky positions can significantly improve on the sensitivities of all-sky surveys, even while needing to cover a wide band of possible GW frequencies and first and second time derivatives due to lack of pulsations from the object (Wette et al. 2008). This makes non-pulsing isolated neutron stars attractive targets for continuous GW searches if they are well localized.

Directed GW searches for isolated neutron stars have been published targeting SNRs (Abadie et al. 2010; Abadie et al. 2011; Aasi et al. 2015; Sun et al. 2016; Zhu et al. 2016; Abbott et al. 2017b) and promising locations including the galactic center (Abadie et al. 2011; Aasi et al. 2013; Abbott et al. 2017b) and the core of a nearby globular cluster—where multi-body interactions might effectively rejuvenate some neutron stars’ continuous GW emission (Abbott et al. 2017j). The only such search of data from advanced interferometers so far (Abbott et al. 2017b) employed methods from stochastic background searches which, while quick to implement, are not as sensitive as continuous wave search methods.

Here we present the first directed continuous wave searches for isolated non-pulsing neutron stars in data from the first Advanced LIGO observing run (O1). We used an extension of the coherent data analysis pipeline used in Abadie et al. (2010) and Aasi et al. (2015), to which this paper is a sequel. The improved noise curve (with respect to initial LIGO and Virgo) means that we can search more targets with sensitivity beating the indirect upper limit on GW emission due to energy conservation (Wette et al. 2008) based on the age of the neutron star (similar to the spin-down limit for known pulsars). We include not only more supernova remnants, but also the exoplanet candidate Fomalhaut b, which has been proposed to be an old nearby neutron star (Neuhäuser et al. 2015)—close enough that it is an attractive target in spite of being much older than the others. We do not include SN 1987A because it is so young that the possible spin-down parameter space is too large to cover with a coherent wide band search and a reasonable computational cost.

2. SEARCHES

2.1. Methods

These searches were based on the multi-interferometer \mathcal{F} -statistic (Jaranowski et al. 1998; Cutler & Schutz 2005). The \mathcal{F} -statistic accounts for the modulation of the signal due to the daily rotation of the detectors by adding the outputs of sinusoidal matched filters in quadrature. For these searches the frequency evolution of each filter, in the reference frame of the solar system barycenter, was given by

$$f(t) = f + \dot{f}(t - t_0) + \frac{1}{2}\ddot{f}(t - t_0)^2, \quad (1)$$

where t_0 is the beginning of the observation and the frequency derivatives are evaluated at that time and in a slight abuse of notation we use a simple f for $f(t_0)$. Hence these filters are designed to detect neutron stars without binary companions whose spin-down is not too fast (requiring third or higher frequency derivatives) or too irregular (having significant timing noise or glitches) during the observation. In stationary Gaussian noise, $2\mathcal{F}$ is drawn from a χ^2 distribution with four degrees of freedom, which for loud signals makes the amplitude signal-to-noise ratio roughly $\sqrt{\mathcal{F}/2}$. If a signal is present, the χ^2 is noncentral.

We used data from LIGO O1, but none from Virgo because that interferometer was down for upgrades during O1. At the frequencies to which LIGO was most sensitive (about 100-300 Hz), the strain noise amplitude was about 3–4 times lower than in the sixth LIGO science run (S6) (Abbott et al. 2016a). However there were

many more spectral lines due to instrumental artifacts than in S6, which complicated the analysis. We used the calibration described in Abbott et al. (2017d), which is an update of the first O1 calibration described in Abbott et al. (2017a). Hence, as in Abbott et al. (2017d), our upper limits on strain are uncertain by at least 14%. Like many other continuous GW searches, ours used data in the form of short Fourier transforms (SFTs) of duration 1800 s, high pass filtered and Tukey windowed to reduce artifacts, recording only frequencies up to 2 kHz.

While each search targeted a specific direction (right ascension and declination), each had to cover a broad band of frequencies and first and second derivatives. That is, a bank of signal templates was required, constructed to cover the parameter space (f, \dot{f}, \ddot{f}) with sufficient density (Whitbeck 2006; Wette et al. 2008). We chose coverage such that the maximum loss of power signal-to-noise ratio due to mismatch between the signal and the nearest template (Owen 1996; Brady et al. 1998) was no worse than 20%, a common choice in continuous GW analyses. Given the parameter choices described below, this resulted in 10^{12} – 10^{13} templates for most searches, with the Cas A search getting more than 10^{14} since it was allocated ten times the computing cycles of each other search.

All searches ran on the Atlas computing cluster at the Max Planck Institute for Gravitational Physics (Albert Einstein Institute) in Hanover, Germany using the same tag (S6SNRSearch) of the LALSuite software package¹ as in Aasi et al. (2015) although the controlling scripts were upgraded. Most searches used roughly 10^5 core hours (split into roughly 3×10^4 batch jobs) and Cas A used more than 10^6 (split into roughly 3×10^5 jobs). The splitting into jobs was used in the vetoes and other post-processing described in Sec. 2.4. Post-processing for each search used at most of order ten percent of the core hours dedicated to the search. Several terabytes of search results were written to disk.

2.2. Target List

Our choice of targets required that a search of fixed computational cost be sensitive enough to detect the strongest continuous GW signal consistent with broad conservation of energy considerations. As introduced by Wette et al. (2008), the strongest possible signal based on the age a and distance D of the source,

$$h_0^{\text{age}} = 1.26 \times 10^{-24} \left(\frac{3.30 \text{ kpc}}{D} \right) \left(\frac{300 \text{ yr}}{a} \right)^{1/2}, \quad (2)$$

¹ Available at <https://github.com/lscsoft/lalsuite-archive>

is analogous to the spin-down limit for known pulsars and indicates the strongest possible intrinsic strain produced by an object whose unknown spin-down is entirely due to GW emission and has been since birth. The intrinsic strain h_0 (Jaranowski et al. 1998) characterizes the GW metric perturbation without reference to any particular orientation or polarization, and hence is typically a factor 2–3 greater than the strain response measured by the interferometers. The indirect limit h_0^{age} is slightly different for r -mode emission (Owen 2010) than for the mass quadrupole source tacitly assumed above and in most of the literature, but we neglect this small difference. Due to uncertainties in the neutron star mass and equation of state, h_0^{age} is uncertain by of order 50%, which we also neglect.

To choose directions to search, we started from the Green catalog of supernova remnants (Green 2014). We picked x-ray point sources (CCOs or candidate CCOs), small PWNe, and in some cases relatively young SNRs where any neutron star could not yet have moved far. We selected only targets with age and distance estimates so that we could evaluate h_0^{age} . In some cases there is a wide range of estimates in the literature, leading to significant differences in h_0^{age} . In most cases we used the most optimistic estimates, yielding the highest h_0^{age} but also the most difficult search over the widest band of frequency and spin-down parameters. In addition to this wide search using the optimistic age and distance, we did a deep search using the most pessimistic age and distance in cases where the strain sensitivity would improve over the wide search by a factor of roughly $\sqrt{2}$.

The resulting targets and chosen parameters are shown in Table 1. We now briefly summarize each target and the provenance of the parameters used for it.

G1.9+0.3—Currently the youngest known SNR in the galaxy (Reynolds et al. 2008). Several arguments favor it being a Type Ia (Reynolds et al. 2008), which would leave no neutron star behind, but this is not definite and the remnant’s youth makes it an interesting target on the chance that it is not Type Ia. We used the position of the center of the remnant from the discovery paper (Reich et al. 1984). At maximum kick velocity any neutron star could have moved only a few arcseconds, which is not an issue for our searches. The age and distance shown are from the “rediscovery” paper (Reynolds et al. 2008), though the latter is a nominal galactic center distance.

G15.9+0.2—The CCO was discovered in *Chandra* data by Reynolds et al. (2006). We used the lower limit on age and the galactic center distance estimate from the same paper, though both quantities may be significantly greater (Klochkov et al. 2016).

Table 1. Targeted objects and astronomical parameters used in each search

SNR (G name)	parameter space	Other name	RA+dec (J2000)	<i>D</i> (kpc)	<i>a</i> (kyr)
1.9+0.3	—	—	174846.9–271016	8.5	0.1
15.9+0.2	—	—	181852.1–150214	8.5	0.54
18.9–1.1	—	—	182913.1–125113	2	4.4
39.2–0.3	3C 396	—	190404.7+052712	6.2	3
65.7+1.2	DA 495	—	195217.0+292553	1.5	20
93.3+6.9	DA 530	—	205214.0+551722	1.7	5
111.7–2.1	Cas A	—	232327.9+584842	3.3	0.3
189.1+3.0	wide	IC 443	061705.3+222127	1.5	3
189.1+3.0	deep	IC 443	061705.3+222127	1.5	20
266.2–1.2	wide	Vela Jr.	085201.4–461753	0.2	0.69
266.2–1.2	deep	Vela Jr.	085201.4–461753	0.9	5.1
291.0–0.1	—	MSH 11–62	111148.6–603926	3.5	1.2
330.2+1.0	—	—	160103.1–513354	5	1
347.3–0.5	—	—	171328.3–394953	0.9	1.6
350.1–0.3	—	—	172054.5–372652	4.5	0.6
353.6–0.7	—	—	173203.3–344518	3.2	27
354.4+0.0	wide	—	173127.5–333412	5	0.1
354.4+0.0	deep	—	173127.5–333412	8	0.5
—	wide	Fomalhaut b	225739.1–293720	0.011	316
—	deep	Fomalhaut b	225739.1–293720	0.02	3000

Values of distance *D* and age *a* are generally at the optimistic (nearby and young) end of ranges given in the literature. For some objects the range of parameters is wide enough to justify a wide search for optimistic parameter values (first entry for that object in the table) and a deep search over more pessimistic parameter values (second entry). See text for details and references.

G18.9–1.1—The position is that of the *Chandra* point source discovered by Tüllmann et al. (2010). Age and distance estimates are from the previous *ROSAT* and *ASCA* observations of Harrus et al. (2004).

G39.2–0.3—Also known as 3C 396. The PWN and embedded point source were found by Olbert et al. (2003) in *Chandra* data, the point source being localized to within 2'' in spite of the PWN. Su et al. (2011) estimate the age and distance, the latter based on the tangent point of the spiral arm.

G65.7+1.2—Also known as DA 495. Arzoumanian et al. (2008) found the *Chandra* point source in the PWN. The quoted distance (Kothes et al. 2004) and minimum age (Kothes et al. 2008) are derived slightly inconsistently due to assumed distances to the galactic center. We did not attempt to resolve the inconsistency, though we did choose the distance from the former paper since it uses the more commonly accepted galactic cen-

ter distance. The latter paper (and others) also argue that the distance could be several times higher.

G93.3+6.9—Also known as DA 530. The position and age are from Jiang et al. (2007) and the distance estimate is from Foster & Routledge (2003). Jiang et al. (2007) find no *Chandra* point source, but the X-ray intensity of the faint candidate PWN falls off on a scale of 6'', which qualifies as a point source for our purposes.

G111.7–2.1—Also known as Cas A. The position of the CCO is from the *Chandra* “first light” observation (Tananbaum 1999), the distance is from Reed et al. (1995), and the age is from Fesen et al. (2006).

G189.1+3.0—Also known as IC 443. The position is that of the *Chandra* point source found by Olbert et al. (2001) embedded in the PWN. This object is often studied, with a wide range of distance and age estimates in the literature. We used Petre et al. (1988) for an optimistic age estimate. Our pessimistic age estimate is not

Table 2. Derived parameters used in each search

SNR (G name)	parameter space	f_{\min} (Hz)	f_{\max} (Hz)	T_{span} (s)	T_{span} (days)	Start of span (UTC, 2015)	H1 SFTs	L1 SFTs	Duty factor	h_0^{age} ($\times 10^{-25}$)
1.9+0.3		38	1332	336 307	3.9	Nov 30 03:53:08	156	141	0.79	8.4
15.9+0.2		72	538	887 744	10.3	Nov 25 13:39:16	369	304	0.68	3.6
18.9−1.1		45	987	1 133 255	13.1	Nov 21 00:00:40	462	346	0.64	5.4
39.2−0.3		98	295	1 965 780	22.8	Nov 28 00:47:19	641	647	0.59	2.1
65.7+1.2		53	794	1 932 067	22.4	Dec 14 04:52:40	774	555	0.62	3.4
93.3+6.9		41	1215	1 051 764	12.2	Nov 25 12:39:16	385	354	0.63	5.9
111.7−2.1		31	1998	775 855	9.0	Nov 26 20:58:03	317	294	0.71	12.6
189.1+3.0	wide	37	1547	803 419	9.3	Nov 26 12:43:17	331	296	0.70	8.7
189.1+3.0	deep	50	805	1 933 867	22.4	Dec 14 04:52:40	775	555	0.62	3.4
266.2−1.2	wide	19	1998	462 616	5.4	Nov 28 02:17:19	191	213	0.79	136
266.2−1.2	deep	32	1998	799 819	9.3	Nov 26 12:43:17	329	294	0.70	11.2
291.0−0.1		42	987	788 409	9.1	Nov 26 18:28:03	322	295	0.70	5.9
330.2+1.0		53	731	851 744	9.9	Nov 25 23:39:16	349	302	0.69	4.5
347.3−0.5		27	1998	578 325	6.7	Nov 28 05:17:19	237	253	0.76	19.9
350.1−0.3		42	1038	637 577	7.4	Nov 28 02:17:19	257	271	0.75	6.5
353.6−0.7		132	275	3 762 662	43.5	Nov 21 02:30:40	1339	1078	0.58	1.4
354.4+0.0	wide	36	1677	301 250	3.5	Nov 28 02:17:19	125	152	0.83	14.4
354.4+0.0	deep	62	635	790 209	9.1	Nov 26 17:58:03	323	295	0.70	4.0
Fomalhaut b	wide	19	1998	2 492 267	28.8	Sep 18 20:08:24	955	799	0.63	116
Fomalhaut b	deep	22	1998	4 639 371	53.7	Nov 19 23:13:10	1626	1295	0.57	20.7

The span reported is the final one, including the possible extension to the end of an SFT in progress at the end of the originally requested span. The duty factor reported is total SFT time divided by T_{span} divided by the number of interferometers (two). As in the previous table, for objects with two entries the first is a wide search (optimistic parameter estimates) and the second is a deep search (pessimistic parameter estimates). In some cases the frequency ranges for wide and deep searches are nearly identical, but the ranges of spin-down parameters (described in the text) are not.

quite the most extreme in the literature, but rather a best fit for a pessimistic scenario from relatively recent modeling (Swartz et al. 2015). We did not use the most optimistic distance quoted, but rather the assumed association with the I Gem cluster from Fesen & Kirshner (1980) which is common in the literature.

G266.2−1.2—Also known as Vela Jr. The position is that of the CCO found by Pavlov et al. (2001). We used Iyudin et al. (1998) for the most optimistic age and distance estimates. The pessimistic age estimate is from Allen et al. (2015), which was published too recently for the previous paper in this series (Aasi et al. 2015). Allen et al. (2015) also discuss the possible association of several surrounding objects with the nearer concentration of the Vela Molecular Ridge, at a spread of distances providing our pessimistic distance estimate

(Liseau et al. 1992) and rendering the more pessimistic ones unlikely.

G291.0−0.1—Also known as MSH 11−62. The position and age are from the *Chandra* point source discovery paper (Slane et al. 2012). The distance is from Moffett et al. (2001). The age and distance are derived in slightly inconsistent ways, but rather than attempt to repeat the calculations we used the numbers quoted in the literature.

G330.2+1.0—The CCO was discovered by Park et al. (2006) in *Chandra* data with sub-arcsecond position accuracy. We used a distance estimate from radio observations (McClure-Griffiths et al. 2001) and an age estimate from the x-ray spectrum (Park et al. 2009).

G347.3−0.5—Mignani et al. (2008) obtained the sub-arcsecond position from archival *Chandra* data, although the CCO had been identified in *ASCA* data




Figure 1. Direct observational 95% confidence upper limits on intrinsic strain as a function of frequency in 1 Hz bands for four searches. The horizontal line indicates the indirect limit from energy conservation. Scattered points on a higher line indicate 1 Hz bands where no upper limit was set due to data quality issues. All figures trace a slightly distorted version of the noise curve, with G39.2–0.3 appearing flat because it covers only the bottom of the curve.

earlier (Slane et al. 1999). We used the distance from Cassam-Chenai et al. (2004) and the age from the proposed identification with a possible SN 393 (Wang et al. 1997). Although this identification may be problematic given the inferred properties of such a supernova, other age estimates are comparable (Fesen et al. 2012).

G350.1–0.3—Position and distance estimates are from the discovery paper of the CCO candidate by Gaensler et al. (2008). The age is from *Chandra* observations Lovchinsky et al. (2011).

G353.6–0.7—Halpern & Gotthelf (2010) identified the most likely of several candidate CCOs. The age estimate (Tian et al. 2008) makes this CCO candidate the only one that is almost certainly too old for *r*-modes, although we still set upper limits on *r*-mode amplitude. The distance estimate is also from Tian et al. (2008).

We used the first-observation position contained in the name of the candidate CCO rather than the slightly better *Chandra* position reported by Halpern & Gotthelf (2010); the roughly 1'' difference is not significant for GW integration times used in this paper.

G354.4+0.0—All parameters are from the discovery paper (Roy & Pal 2013). No associated point source has been detected yet, but if the remnant's age is correct any young neutron star should be within roughly 20'' of the center (whose location we used for the GW search). Such a position error is not significant for the integration times used here.

Fomalhaut b—Considered an extrasolar planet candidate since its discovery in visible light (Kalas et al. 2008), this has been proposed based on a lack of infrared detection to be a serendipitous discovery of a




Figure 2. Same as the previous figure for four more searches.

nearby neutron star (Neuhäuser et al. 2015). Parameters are taken from Neuhäuser et al. (2015), with the maximum distance an attempt to balance the uncertainties in the scenarios discussed there. After this search was run, Poppenhaeger et al. (2017) searched for and did not find the object with *Chandra*. If the object is a neutron star, this somewhat reduced the possible distance and significantly increased the minimum age.

2.3. Parameter Space

After sky position, the key parameters for each search were the GW frequency band (f_{\min}, f_{\max}) and time span of integration T_{span} . As in Aasi et al. (2015), these parameters were determined in an iterative process intended to produce a search more sensitive than h_0^{age} over as wide a frequency band as possible for a fixed computational cost. Due to Doppler shifts and several features of the analysis, we capped the maximum frequency at 1998 Hz rather than the 2 kHz in the SFTs. The cost,

approximated as proportional to $a^{-1.1} f_{\max}^{2.2} T_{\text{span}}^4$, was kept comparable to Abadie et al. (2010) for most targets, but Cas A was allocated ten times as many computational cycles due to its status as youngest known neutron star in the galaxy. Due to some inaccuracy in the power-law fit used for computational cost as a function of the key parameters, the computational cost and sensitivity varied by up to 20–30% from these goals. For a given frequency f , as in Abadie et al. (2010) and Aasi et al. (2015), we searched

$$-\frac{f}{a} \leq \dot{f} \leq -\frac{1}{6} \frac{f}{a}; \quad (3)$$

and for a given \ddot{f} we searched

$$2 \frac{\dot{f}^2}{f} \leq \ddot{f} \leq 7 \frac{\dot{f}^2}{f}. \quad (4)$$

These ranges and the computational cost fixed f_{\min} , f_{\max} , and T_{span} for each search.




Figure 3. Same as the previous figure for four more searches.

We then chose the start time of each search by the same method as Abadie et al. (2010) and Aasi et al. (2015), minimizing the harmonic mean of the strain noise power spectral density during the span over the frequency band (f_{\min}, f_{\max}). Neglecting the small effect of the declination of the target, this corresponds to maximizing the search sensitivity for a fixed T_{span} —which is roughly a fixed computational cost. Hence the algorithm chose spans when both interferometers had good noise performance and little down time, usually later in O1. The resulting search parameters are described in Table 2.

We applied the same consistency checks as in previous searches: For each search we checked using the parameter space metric (Whitbeck 2006; Wette et al. 2008) that neglect of the third frequency derivative in Eq. (1) did not significantly reduce $2\mathcal{F}$, even in the worst case (G1.9+0.3). We also checked that the position uncer-

tainties of the targets also did not significantly reduce $2\mathcal{F}$. A simple approximation (Whitbeck 2006) suggests that the sky resolution of these searches is an arcminute or two at 2 kHz and a 10 day integration, and it scales inversely with f_{\max} and T_{span} . We spot checked this with injection studies and found it to be accurate. Given the integration times in Table 2, even the worst position uncertainty (20'' for G354.4+0.0) is well within bounds for a single directed search. Finally, we checked that the standard 1800 s SFT duration did not diminish sensitivity to signals with \dot{f} high enough that the frequency could move to another SFT frequency bin over the duration of the SFT. This effect was negligible except for SNR G1.9+0.3, where it could reduce the sensitivity (raise the detectable h_0) by of order 10% at frequencies above 1 kHz.

2.4. Post-processing




Figure 4. Same as the previous figure for four more searches.

Each search recorded a list of candidates with high values of $2\mathcal{F}$, which was then pared using two automated vetoes designed for instrumental artifacts, as used in Abadie et al. (2010) and Aasi et al. (2015). The “Fscan veto” used a normalized spectrogram formed from the SFTs to detect and veto spectral lines and nonstationary noise. Its implementation and parameters were the same as in Aasi et al. (2015) except that we fixed a bug in the old code whereby the Doppler shift due to the Earth’s orbital motion was not applied. (This bug allowed more noise lines to pass the automated vetoes and require manual scrutiny, but had a negligible effect on the false dismissal rate.) The “interferometer consistency veto” ruled out candidates for which a single-interferometer $2\mathcal{F}$ exceeded the two-interferometer $2\mathcal{F}$ for the same event, indicating a disturbance present in only one interferometer. It also vetoed entire search jobs if the number of candidates vetoed was high enough.

This veto was also applied in the same way as in Aasi et al. (2015), except that the threshold for vetoing an entire search job was 5% of the templates in that job. Unlike in previous papers in this series, we also vetoed a list of known instrumental spectral lines compiled from studies of the interferometers (Covas et al. 2018).

After these steps, including fixing the Doppler bug, the searches still had almost two thousand jobs containing non-vetoed outliers above the 95% confidence level for Gaussian noise. All of these jobs were examined by hand. As in Aasi et al. (2015), two plots were made and inspected for each job. (See Figure 1 of that paper for illustrative examples.) In case of a real or injected signal the first plot, of $2\mathcal{F}$ vs. frequency for all loud candidates in the job, would show a δ -function like spike even for very loud signals, as verified by studying hardware injections. The candidates generally showed broad bands of high noise, occupying a fraction of order unity




Figure 5. Same as the previous figure for four more searches.

of the search job frequency band except for a handful which occupied a few percent of the search band. These few candidates, which were still of order one hundred times broader than a real signal would be, were verified to be hardware injected test signals detectable in the wrong sky location due to their huge amplitudes. The second plot for each search job containing candidates was a semilog histogram of loud candidates, which on inspection typically showed the tail of a $\chi^2(4)$ distribution with the wrong amplitude, indicative of a broad band disturbance in the noise spectrum. See [Aasi et al. \(2015\)](#) for examples and further details.

No candidates survived inspection of these plots, and therefore we conclude that no astrophysical signal was detected.

3. UPPER LIMITS

Our method of setting upper limits was almost the same as in previous papers ([Abadie et al. 2010; Aasi et al. 2015](#)).

In each 1 Hz band searched, we estimated the value of h_0 that would be detected 95% of the time by our search (assuming random variation of other signal parameters such as inclination of the star’s rotation axis to the line of sight) at a louder value than the loudest $2\mathcal{F}$ actually recorded by the search in that band. We made an initial estimate from a semianalytic integration of the expected $2\mathcal{F}$ distribution. Then we injected simulated signals with different values of h_0 near this value to refine the location of the 95% confidence (5% false dismissal) threshold. We reduced the number of injections per band to 1000 (from 6000 in previous papers) due to the computational cost of setting upper limits on wider bands.

For each search we pared the list of upper limits on h_0 versus frequency. We dropped bands where the injections indicated the false dismissal rate was more than




Figure 6. Upper limits on fiducial ellipticity (left panel) and r -mode amplitude (right panel) for a representative sample of searches.

5% and we dropped ± 1 Hz bands around harmonics of the 60 Hz power mains up to 300 Hz.

The resulting upper limits on h_0 , in 1 Hz frequency bands, are plotted in Figs. 1–5. Each curve has roughly the same shape as the amplitude spectral density of the strain noise. The line of dots near the top of each plot corresponds to bands where no upper limit was set. Some features such as the “violin modes” of the interferometer test mass suspension (roughly 500 Hz and harmonics) are evident. The horizontal line in each plot is h_0^{age} , the strain the search was intended to beat. In some cases the estimate of sensitivity made before performing the search was wrong by of order 10%, so the upper limits (lower dots) do not always lie below the line.

Upper limits on h_0 can be converted to upper limits on fiducial neutron-star ellipticity $\epsilon = |I_{xx} - I_{yy}|/I_{zz}$ (where I_{ab} is the moment of inertia) using (e.g. Wette et al. 2008)

$$\epsilon = 9.5 \times 10^{-5} \left(\frac{h_0}{1.2 \times 10^{-24}} \right) \left(\frac{D}{1 \text{ kpc}} \right) \left(\frac{100 \text{ Hz}}{f} \right)^2. \quad (5)$$

This number assumes $I_{zz} = 10^{45} \text{ g cm}^2$. Uncertainties in the mass, radius, and neutron star equation of state make the conversion from h_0 to ϵ uncertain by a factor of two or more. This fiducial ellipticity can be converted to the true shape of the star (Johnson-McDaniel 2013) or other quantities (Owen 2010). We plot upper limits on ϵ for a selection of searches in the left hand panel of Fig. 6. We do not plot the indirect limits on ϵ and α derived from h_0^{age} since they are close to the direct upper limits on the scale of the plot. We do not plot the remaining searches because their upper limits are close

to those of the searches plotted. The great differences between curves are mainly due to the distances to the sources; hence Fomalhaut b has the best upper limits—of order 10^{-9} at high frequencies.

Upper limits on h_0 can be converted to the common r -mode amplitude parameter α (Lindblom et al. 1998) via (Owen 2010)

$$\alpha = 0.28 \left(\frac{h_0}{10^{-24}} \right) \left(\frac{100 \text{ Hz}}{f} \right)^3 \left(\frac{D}{1 \text{ kpc}} \right). \quad (6)$$

This number assumes a fiducial set of stellar parameters described in Owen (2010) and is uncertain by a factor of up to about three depending in the neutron star mass and equation of state. We plot upper limits on α for a selection of searches in the right hand panel of Fig. 6. Again, the differences between curves are mainly due to the source distances. The best upper limits, apart from Fomalhaut b which is almost certainly too old for active r -modes, are of order 10^{-6} at high frequencies for Vela Jr.

4. DISCUSSION

These are the first directed searches of Advanced LIGO data using continuous wave analysis methods. These searches have improved on previous directed searches by covering wider parameter ranges and more targets, and by setting better upper limits on targets searched previously. Our upper limits on h_0 approach 2×10^{-25} for many targets and approach 1×10^{-25} for one—about a factor of 3 improvement on Aasi et al. (2015), due mainly to the improvement in the detectors. And our upper limits beat the indirect limit h_0^{age} over bands of 1–2 kHz for more targets than were ever published before. [Searches for some of these targets in less

sensitive S6 data for the purpose of testing code were described in an unpublished thesis (Idrisy 2015).] As with previous data runs, we improved on the sensitivity of all-sky wide-band searches (Abbott et al. 2018) but did not match the sensitivity of searches for known pulsars with full timing solutions (Abbott et al. 2017d). As before, the directed searches described here also have the caveats that there might be no neutron star present in some cases, any neutron star might be spinning too slowly to be detected, and a neutron star spinning at a detectable frequency might glitch—the latter phenomenon in a CCO is now an observation (Gotthelf & Halpern 2018) rather than a surmise, and would somewhat reduce the sensitivity of these searches (Ashton et al. 2017).

Most of our upper limits on ϵ and α are competitive with the largest numbers predicted by theory. The maximum ϵ for “mountains” supported by elastic stresses of normal neutron star matter is probably 10^{-5} – 10^{-6} (Horowitz & Kadau 2009; Johnson-McDaniel & Owen 2013; Baiko & Chugunov 2018), and for many of our searches upper limits are in this region over hundreds of Hz. The maximum α (nonlinear saturation amplitude) for r -modes is probably of order 10^{-3} (Bondarescu et al. 2009), and for many of our searches upper limits beat this over hundreds of Hz. Mountains supported by an internal magnetic field can produce ϵ of order $10^{-4}(B/10^{15} \text{ G})^2$ where B is the poloidal part of the field (e.g. Ciolfi & Rezzolla 2013). Since, unlike elastic mountains, magnetic mountains are likely to be within about an order of magnitude of this limit for a given internal field, depending on its configuration, our upper limits on ϵ translate into rough limits on internal magnetic field—if a neutron star is present and spinning rapidly enough to emit GWs in band.

More data from Advanced LIGO and Advanced Virgo is now available, with more live time and lower noise amplitude than before. This makes more targets feasible for directed searches at greater sensitivity, increasing the chances of a detection of continuous GWs. Such searches will be done in the near future.

The authors gratefully acknowledge the support of the United States National Science Foundation (NSF) for the construction and operation of the LIGO Laboratory and Advanced LIGO as well as the Science and Technology Facilities Council (STFC) of the United Kingdom, the Max-Planck-Society (MPS), and the State of Niedersachsen/Germany for support of the construction of Advanced LIGO and construction and operation of the GEO600 detector. Additional support for Advanced LIGO was provided by the Australian Research Council. The authors gratefully acknowledge the Italian Istituto Nazionale di Fisica Nucleare (INFN), the French Centre National de la Recherche Scientifique (CNRS) and the Foundation for Fundamental Research on Matter supported by the Netherlands Organisation for Scientific Research, for the construction and operation of the Virgo detector and the creation and support of the EGO consortium. The authors also gratefully acknowledge research support from these agencies as well as by the Council of Scientific and Industrial Research of India, the Department of Science and Technology, India, the Science & Engineering Research Board (SERB), India, the Ministry of Human Resource Development, India, the Spanish Agencia Estatal de Investigación, the Vicepresidència i Conselleria d'Innovació, Recerca i Turisme and the Conselleria d'Educació i Universitat del Govern de les Illes Balears, the Conselleria d'Educació, Investigació, Cultura i Esport de la Generalitat Valenciana, the National Science Centre of Poland, the Swiss National Science Foundation (SNSF), the Russian Foundation for Basic Research, the Russian Science Foundation, the European Commission, the European Regional Development Funds (ERDF), the Royal Society, the Scottish Funding Council, the Scottish Universities Physics Alliance, the Hungarian Scientific Research Fund (OTKA), the Lyon Institute of Origins (LIO), the Paris Île-de-France Region, the National Research, Development and Innovation Office Hungary (NKFIH), the National Research Foundation of Korea, Industry Canada and the Province of Ontario through the Ministry of Economic Development and Innovation, the Natural Science and Engineering Research Council Canada, the Canadian Institute for Advanced Research, the Brazilian Ministry of Science, Technology, Innovations, and Communications, the International Center for Theoretical Physics South American Institute for Fundamental Research (ICTP-SAIFR), the Research Grants Council of Hong Kong, the National Natural Science Foundation of China (NSFC), the Leverhulme Trust, the Research Corporation, the Ministry of Science and Technology (MOST), Taiwan and the Kavli Foundation. The authors gratefully acknowledge the support of the NSF,

STFC, MPS, INFN, CNRS and the State of Niedersachsen/Germany for provision of computational resources.

This paper has been assigned document number LIGO-P1800333.

REFERENCES

- Aasi, J., Abadie, J., Abbott, B. P., et al. 2013, *PhRvD*, 88, 102002
- Aasi, J., Abbott, B. P., Abbott, R., et al. 2015, *ApJ*, 813, 39
- Abadie, J., et al. 2010, *ApJ*, 722, 1504
- Abadie, J., Abbott, B. P., Abbott, R., et al. 2011, *Physical Review Letters*, 107, 271102
- Abbott, B. P., Abbott, R., Abbott, T. D., et al. 2016a, *Physical Review Letters*, 116, 131103
- . 2016b, *Physical Review Letters*, 116, 241103
- . 2016c, *Physical Review Letters*, 116, 061102
- . 2017a, *PhRvD*, 95, 062003
- . 2017b, *Physical Review Letters*, 118, 121102
- . 2017c, *ApJ*, 851, 71
- . 2017d, *ApJ*, 839, 12
- . 2017e, *Physical Review Letters*, 118, 221101
- . 2017f, *ApJL*, 851, L35
- . 2017g, *Physical Review Letters*, 119, 141101
- . 2017h, *Physical Review Letters*, 119, 161101
- . 2017i, *ApJL*, 848, L12
- . 2017j, *PhRvD*, 95, 082005
- . 2018, *PhRvD*, 97, 102003
- Abbott, B. P., et al. 2018, *Phys. Rev. Lett.*, 121, 161101
- Allen, G., Chow, K., DeLaney, T., et al. 2015, *Astrophys. J.*, 798, 82
- Arzoumanian, Z., Safi-Harb, S., Landecker, T. L., Kothes, R., & Camilo, F. 2008, *ApJ*, 687, 505
- Ashton, G., Prix, R., & Jones, D. I. 2017, *PhRvD*, 96, 063004
- Baiko, D. A., & Chugunov, A. I. 2018, *MNRAS*, 480, 5511
- Bondarescu, R., Teukolsky, S. A., & Wasserman, I. 2009, *PhRvD*, 79, 104003
- Brady, P. R., Creighton, T., Cutler, C., & Schutz, B. F. 1998, *PhRvD*, 57, 2101
- Cassam-Chenaï, G., Decourchelle, A., Ballet, J., et al. 2004, *A&A*, 427, 199
- Ciolfi, R., & Rezzolla, L. 2013, *MNRAS*, 435, L43
- Covas, P. B., Effler, A., Goetz, E., et al. 2018, *PhRvD*, 97, 082002
- Cutler, C., & Schutz, B. F. 2005, *PhRvD*, 72, 063006
- De, S., Finstad, D., Lattimer, J. M., et al. 2018, *Physical Review Letters*, 121, 091102
- Fesen, R. A., & Kirshner, R. P. 1980, *ApJ*, 242, 1023
- Fesen, R. A., Kremer, R., Patnaude, D., & Milisavljevic, D. 2012, *AJ*, 143, 27
- Fesen, R. A., et al. 2006, *ApJ*, 645, 283
- Foster, T., & Routledge, D. 2003, *ApJ*, 598, 1005
- Gaensler, B. M., Tanna, A., Slane, P. O., et al. 2008, *ApJL*, 680, L37
- Glampedakis, K., & Gualtieri, L. 2017, ArXiv e-prints, arXiv:1709.07049
- Gotthelf, E. V., & Halpern, J. P. 2018, *ApJ*, 866, 154
- Green, D. A. 2014, *Bulletin of the Astronomical Society of India*, 42, 47, 2017 update at <http://www.mrao.cam.ac.uk/surveys/snrs/>
- Halpern, J. P., & Gotthelf, E. V. 2010, *ApJ*, 710, 941
- Harrus, I. M., Slane, P. O., Hughes, J. P., & Plucinsky, P. P. 2004, *ApJ*, 603, 152
- Horowitz, C. J., & Kadau, K. 2009, *Physical Review Letters*, 102, 191102
- Idrisy, A. 2015, PhD thesis, The Pennsylvania State University
- Iyudin, A. F., Schönfelder, V., Bennett, K., et al. 1998, *Nature*, 396, 142
- Jaranowski, P., Królak, A., & Schutz, B. F. 1998, *PhRvD*, 58, 063001
- Jiang, B., Chen, Y., & Wang, Q. D. 2007, *ApJ*, 670, 1142
- Johnson-McDaniel, N. K. 2013, *PhRvD*, 88, 044016
- Johnson-McDaniel, N. K., & Owen, B. J. 2013, *PhRvD*, 88, 044004
- Kalas, P., Graham, J. R., Chiang, E., et al. 2008, *Science*, 322, 1345
- Klochkov, D., Suleimanov, V., Sasaki, M., & Santangelo, A. 2016, *A&A*, 592, L12
- Kothes, R., Landecker, T. L., Reich, W., Safi-Harb, S., & Arzoumanian, Z. 2008, *ApJ*, 687, 516
- Kothes, R., Landecker, T. L., & Wolleben, M. 2004, *ApJ*, 607, 855
- Lindblom, L., Owen, B. J., & Morsink, S. M. 1998, *Physical Review Letters*, 80, 4843
- Liseau, R., Lorenzetti, D., Nisini, B., Spinoglio, L., & Moneti, A. 1992, *A&A*, 265, 577
- Lovchinsky, I., Slane, P., Gaensler, B. M., et al. 2011, *ApJ*, 731, 70
- McClure-Griffiths, N. M., Green, A. J., Dickey, J. M., et al. 2001, *ApJ*, 551, 394
- Mignani, R. P., Zaggia, S., de Luca, A., et al. 2008, *A&A*, 484, 457

- Moffett, D., Gaensler, B., & Green, A. 2001, in AIP Conf. Proc., Vol. 565, Young Supernova Remnants: Eleventh Astrophysics Conference, ed. S. S. Holt & U. Hwang (Melville, NY: AIP), 333–336
- Neuhäuser, R., Hohle, M. M., Ginski, C., et al. 2015, MNRAS, 448, 376
- Olbert, C. M., Clearfield, C. R., Williams, N. E., Keohane, J. W., & Frail, D. A. 2001, ApJL, 554, L205
- Olbert, C. M., Keohane, J. W., Arnaud, K. A., et al. 2003, ApJL, 592, L45
- Owen, B. J. 1996, PhRvD, 53, 6749
- . 2009, arXiv:0903.2603
- . 2010, PhRvD, 82, 104002
- Park, S., Kargaltsev, O., Pavlov, G. G., et al. 2009, ApJ, 695, 431
- Park, S., Mori, K., Kargaltsev, O., et al. 2006, ApJL, 653, L37
- Pavlov, G. G., Sanwal, D., Kızıltan, B., & Garmire, G. P. 2001, ApJL, 559, L131
- Petre, R., Szymkowiak, A. E., Seward, F. D., & Willingale, R. 1988, ApJ, 335, 215
- Poppenhaeger, K., Auchettl, K., & Wolk, S. J. 2017, MNRAS, 468, 4018
- Reed, J. E., Hester, J. J., Fabian, A. C., & Winkler, P. F. 1995, ApJ, 440, 706
- Reich, W., Fuerst, E., Haslam, C. G. T., Steffen, P., & Reif, K. 1984, A&AS, 58, 197
- Reynolds, S. P., Borkowski, K. J., Green, D. A., et al. 2008, ApJL, 680, L41
- Reynolds, S. P., Borkowski, K. J., Hwang, U., et al. 2006, ApJL, 652, L45
- Roy, S., & Pal, S. 2013, ApJ, 774, 150
- Slane, P., Gaensler, B. M., Dame, T. M., et al. 1999, ApJ, 525, 357
- Slane, P., Hughes, J. P., Temim, T., et al. 2012, ApJ, 749, 131
- Su, Y., Chen, Y., Yang, J., et al. 2011, ApJ, 727, 43
- Sun, L., Melatos, A., Lasky, P. D., Chung, C. T. Y., & Darman, N. S. 2016, PhRvD, 94, 082004
- Swartz, D. A., Pavlov, G. G., Clarke, T., et al. 2015, ApJ, 808, 84
- Tananbaum, H. 1999, IAUC, 7246, 1
- Tian, W. W., Leahy, D. A., Havercorn, M., & Jiang, B. 2008, ApJL, 679, L85
- Tüllmann, R., Plucinsky, P. P., Gaetz, T. J., et al. 2010, ApJ, 720, 848
- Wang, Z. R., Qu, Q., & Chen, Y. 1997, A&A, 318, L59
- Wette, K., Owen, B. J., Allen, B., et al. 2008, Classical and Quantum Gravity, 25, 235011
- Whitbeck, D. M. 2006, PhD thesis, The Pennsylvania State University
- Zhu, S. J., Papa, M. A., Eggenstein, H.-B., et al. 2016, PhRvD, 94, 082008