

1 **The effect of organic acids on the behaviour and biodegradation of <sup>14</sup>C-phenanthrene in**  
2 **contaminated soil**

3

4 Gabriela M. Vázquez-Cuevas <sup>a,c</sup>, Alfonso J. Lag-Brotons <sup>a</sup>, Jose J. Ortega-Calvo <sup>b</sup>, Carly J.  
5 Stevens <sup>a</sup> and Kirk T. Semple <sup>a\*</sup>

6 <sup>a</sup>Lancaster Environment Centre, Lancaster University, Lancaster, LA1 4YQ, United Kingdom

7 <sup>b</sup>Instituto de Recursos Naturales y Agrobiología, C.S.I.C., Apartado 1052, 41080 Seville,  
8 Spain

9 <sup>c</sup>Área Académica de Biología, Instituto de Ciencias Básicas e Ingeniería. Universidad  
10 Autónoma del Estado de Hidalgo, 42010, Hidalgo, México

11

12

13

14

15

16

17 \*Corresponding author: Phone no. +44 (0)1524 510554; email: k.semple@lancaster.ac.uk

18

19 **Abstract**

20 The interaction between root exudates and soil microbes has been hypothesised as the  
21 primary mechanism for the biodegradation of organic pollutants in the rhizosphere. However,  
22 the mechanisms governing this loss process are not completely understood. This study aimed  
23 to investigate the effect of two important compounds within root exudates (citric and malic  
24 acid) on <sup>14</sup>C-phenanthrene desorption and bioaccessibility in soil. Overall results showed that  
25 the presence of both citric and malic acid (> 100 mmol l<sup>-1</sup>) enhanced the desorption of <sup>14</sup>C-  
26 phenanthrene; this appeared to be concentration dependant. Increases in extractability were  
27 not reflected in a higher bioaccessibility. Despite enhancing the desorption of <sup>14</sup>C-  
28 phenanthrene in soil, there is no direct evidence indicating that citric or malic acid have the  
29 ability to promote the biodegradation of <sup>14</sup>C-phenanthrene from soil. Results from this study  
30 provide a novel understanding of the role that substrates, typically found within the  
31 rhizosphere due to root exudation, play in the bioaccessibility and biodegradation of  
32 hydrocarbons in contaminated soil.

33

34 **Keywords:** Phenanthrene, organic acids, root exudates, desorption, bioavailability, ageing

35

## 36 **1 Introduction**

37 The rhizosphere is defined as the soil in closest proximity to plant roots and has been  
38 hypothesised to enhance the biodegradation of organic contaminants such as aliphatic and  
39 aromatic hydrocarbons through different mechanisms (Anderson *et al.*, 1993; Pilon-Smits,  
40 2005). These include the promotion of (1) larger microbial populations (Anderson *et al.*,  
41 1993) and shifts on their community composition (Joner *et al.*, 2002), (2) source of  
42 biologically important substrates including nutrients and readily available sources of carbon  
43 (Reilley *et al.*, 1996; Dakora & Phillips, 2002; Martin *et al.*, 2014; Sivaram *et al.*, 2019), and  
44 (3) increasing the bioavailability of the contaminants due to root exudates, decay and  
45 turnover (Siciliano & Germida, 1998; Martin *et al.*, 2014; Sivaram *et al.*, 2019). The amount  
46 and type of substances released by roots is highly dependent on a series of factors including  
47 plant species and age, as well as particular soil and environmental conditions (Jones, 1998;  
48 Shukla *et al.*, 2011; Agnello *et al.*, 2014; Martin *et al.*, 2014). However, a number of low  
49 molecular weight compounds, such as amino acids, sugars and organic acids, have been  
50 identified as common constituents of root exudates (Jones *et al.*, 2003; van Hees *et al.*, 2005).  
51 Organic acids, including citric, malic and oxalic are reported to be the most abundant (Jones  
52 & Brassington, 1998; Ling *et al.*, 2015); therefore, it is reasonable to consider the role that  
53 these acids might play in influencing the extractability and the bioavailability of different  
54 contaminants and how this might influence their biodegradation.

55 The use of root exudates for the dissipation of organic contaminants in soil has been reported  
56 (Miya & Firestone, 2001; Yoshitomi & Shann, 2001; Joner *et al.*, 2002). These investigations  
57 have used simulated rhizosphere conditions by the introduction of artificial or natural root  
58 exudates in order to approach the subject in a more controlled manner (Miya & Firestone,  
59 2001; Joner *et al.*, 2002). From these studies, research has been developed to consider the

60 effect of these substances on the shifts of the microbial populations and/or communities  
61 (Joner *et al.*, 2002; Shukla *et al.*, 2011), overall dissipation of contaminants (Joner *et al.*,  
62 2002), and their effect on soil physical and chemical properties (Shukla *et al.*, 2011). Authors  
63 such as Sun *et al.* (2013), Martin *et al.* (2014) and Gao *et al.* (2015) have pointed out that  
64 although efforts have been directed towards investigating the effect of root exudates on the  
65 biodegradation of hydrocarbons in contaminated soil, information regarding the role of single  
66 compounds from this solution is scarce. Within these few studies, it has been reported that  
67 organic acids commonly found in root exudates can promote the desorption of phenanthrene  
68 from soil (Gao *et al.*, 2010b, 2015b; Ling *et al.*, 2015).

69 It has been observed that changes in the extractability of polycyclic aromatic hydrocarbons  
70 (PAHs) might act as a predictor of the microbial degradability of different species of PAHs.  
71 Specifically, the rates of desorption of some PAHs have successfully been used as predictors  
72 of their biodegradation (Cornelissen *et al.*, 1998a). As fractions of hydrocarbons are  
73 transferred from soil to solution through the desorption process; these can also become more  
74 bioaccessible and susceptible to be metabolised by the soil microbial community (Semple *et*  
75 *al.*, 2003, 2007, 2013). Therefore, the possibility of enhancing the desorption of PAHs by  
76 using organic acids to promote or enhance biodegradation in soil has been identified as a  
77 promising strategy, but remains poorly explored (Martin *et al.*, 2014). In addition, the extent  
78 into which these organic acids affect the biodegradation of the desorbed hydrocarbon has not  
79 been considered. Therefore, the aim of this study was to investigate the effect of two low  
80 molecular weight organic acids (LOAs) commonly found within root exudates in the  
81 extractability and bioaccessibility of <sup>14</sup>C-phenanthrene contaminated soil. Phenanthrene was  
82 selected as a model PAH given its widespread distribution, biodegradability and persistent  
83 properties in soil. For this, mineralisation, hydroxypropyl-β-cyclodextrin (HPCD)  
84 extractability and desorption kinetics of <sup>14</sup>C-phenanthrene were assessed in the presence of

85 organic acids at a range of concentrations. Results from this experiment provide a novel  
86 perspective of the effect of organic acids on the fate of  $^{14}\text{C}$ -phenanthrene soil by investigating  
87 (1) the desorbing capacity of citric and malic acid and (2) the extent by which these can  
88 promote a higher bioavailability and mineralisation.

## 89 **2 Methodology**

### 90 **2.1 Soil preparation and spiking**

91 An uncontaminated clay loam soil (top 20 cm, 2.7 % organic matter) was collected from  
92 Myerscough Agricultural College, Preston, U.K. Partially air-dried soil (24 h) was passed  
93 through a 2 mm sieve and stored in the dark at  $4 \pm 1$  °C until needed. Main soil physical and  
94 chemical characteristics have been described by Towell *et al.* (2011). Sieved soil was  
95 rehydrated (50% water holding capacity (whc)) and spiked following the procedure proposed  
96 by Doick *et al.* (2003). In short, this approach consists on the application of the standards to a  
97 fraction of the total amount of soil (inoculum) followed by gradual mixing and incorporation  
98 of the remaining soil with a stainless steel spoon. Standards used for spiking contained  $^{12/14}\text{C}$   
99 phenanthrene dissolved in acetone to deliver a final concentration of  $100 \text{ mg kg}^{-1}$  (dw)  
100 phenanthrene with an associated  $^{14}\text{C}$ -activity of  $83 \text{ Bq g}^{-1}$  (dw). Spiked soil was placed in  
101 sealed sterilized amber jars and incubated in the dark at  $21 \pm 1$  °C in a controlled environment  
102 room for up to 15 weeks. Determination of the total  $^{14}\text{C}$ -phenanthrene associated activity in  
103 the soil was assessed at every time point by sample oxidation following the methodology  
104 described by Rhodes *et al.* (2012).

105

## 106 **2.2 Influence of organic acids on the mineralisation of <sup>14</sup>C-phenanthrene**

107 *Mycobacterium gilvum* has been previously shown to degrade a range of hydrocarbons,  
108 including PAHs such as naphthalene, fluorine, phenanthrene, anthracene, fluoranthene and  
109 pyrene and as sole or primary source of carbon (Xiong *et al.*, 2017; Posada-Baquero *et al.*,  
110 2019). In addition, this strain has also been isolated from chronically contaminated  
111 environments such as soil from a former coking plant (Xiong *et al.*, 2017), indicating its  
112 ability to adapt to these type of conditions, utilising hydrocarbons as their primary source of  
113 carbon under nutrient limiting conditions, which are typically observed on highly  
114 contaminated sites. To assess this, the mineralisation of <sup>14</sup>C-phenanthrene was measured  
115 using the methodology developed and tested by Reid *et al.* (2001) and Semple *et al.* (2006).  
116 Soil (10 g dw) aged over 14 and 50 days was placed into 250 ml modified Schott bottles  
117 fitted with a 1 M NaOH <sup>14</sup>CO<sub>2</sub> trap ( $n = 3$ ). These ageing times were selected based on that  
118 reported by Kelsey *et al.* (1997), where <sup>14</sup>C-phenanthrene mineralisation was significantly  
119 reduced within the first few weeks and then remained stable for 50 days. To assess the effect  
120 of organic acids towards the mineralisation process, citric, malic, oxalic and succinic acids  
121 were selected as representative LOAs often observed in the rhizosphere (van Hees *et al.*,  
122 2005). Solutions containing individual LOAs within its naturally appearing range in the  
123 rhizosphere soil solution (0.1 and 0.5 mmol l<sup>-1</sup>) (van Hees *et al.*, 2005) were used. These were  
124 incorporated into the minimal basal salts (MBS) medium and used for the mineralisation  
125 assay. For the mineralisation assays, soil was mixed with the MBS containing the organic  
126 acids (25 ml) and 5 ml of a bacterial inoculum of *M. gilvum* (10<sup>5</sup> cells ml<sup>-1</sup>) to achieve a final  
127 3:1 liquid:soil ratio. Bottles were then placed onto an orbital shaker at 100 rpm in a controlled  
128 environment room at 21 ± 1 °C in the dark. <sup>14</sup>CO<sub>2</sub> evolution was assessed by periodically (up  
129 to every 24 h) replacing the trap, mixing with 5 ml liquid scintillation cocktail and assessed  
130 by liquid scintillation counting (LSC) (10 min - Canberra Packard Tri-Carb 2300, U.K.).

131

## 132 **2.3 Influence of organic acids on the extractability of <sup>14</sup>C-phenanthrene**

### 133 *2.3.1 Preliminary tests*

134 A series of preliminary tests were carried in order to optimize the general experimental  
135 parameters and design of the extraction assays. Solutions of deionised water containing citric,  
136 malic, oxalic and succinic acids solutions of individual organic acids were prepared at 0.1 and  
137 0.5 mmol l<sup>-1</sup>. Desorption kinetics of <sup>14</sup>C-phenanthrene with these solutions (*n* = 3) were  
138 assessed from spiked soil following the methodology described below.

139 The temporal effect of organic acids on the bioaccessibility of <sup>14</sup>C-phenanthrene was also  
140 assessed (*n* = 3). Soil was saturated with malic acid solution (100 % whc) at two  
141 concentrations (0.5 and 500 mmol l<sup>-1</sup>) and incubated in a controlled environment room 21 ± 1  
142 °C for 1, 3, 6, 8 or 24 h. Each experimental unit was also fitted with a 1 M NaOH <sup>14</sup>CO<sub>2</sub> trap  
143 for the assessment of any possible dissipation of <sup>14</sup>C-phenanthrene by microbial respiration  
144 during the incubation time. Soil was extracted with 50 mM HPCD solutions after each  
145 incubation time following the methodology described below. <sup>14</sup>CO<sub>2</sub> traps were assessed by  
146 adding 5 ml of liquid scintillation cocktail and assessed by LSC as previously described.

147 Citric, malic, oxalic and succinic acids did not impact significantly on the desorption of <sup>14</sup>C-  
148 phenanthrene from the soil (Table SI-1) when compared against the control (*p* > 0.05).

149 Despite of this, soil extracted with citric and malic acid at 0.5 mmol l<sup>-1</sup> were the only two  
150 treatments presenting higher rapidly desorbing fractions (*F<sub>rap</sub>*), with (44.16 %) and (49.76 %)   
151 respectively, than the control (40.94 %). Based on this, these two organic acids were selected  
152 for further investigation at a wider range of concentrations (0.5, 100, 250, 500 and 1000  
153 mmol l<sup>-1</sup>) in order to assess the full potential of these compounds to impact the desorption of

154 <sup>14</sup>C-phenanthrene in soil. Selected concentrations ranged from naturally appearing LOAs  
155 concentrations (van Hees *et al.*, 2005) up to maximum tested concentrations within  
156 experiments with similar aims (Gao *et al.*, 2015a; Ling *et al.*, 2015).

157

### 158 2.3.2 HPCD extraction of <sup>14</sup>C-phenanthrene from soil

159 Changes in the bioaccessibility of <sup>14</sup>C-phenanthrene were measured by HPCD extractions  
160 from soil aged over 1 and 15 weeks. At each time point, 1.25 g soil (dw) were placed into  
161 Teflon centrifuge tubes ( $n = 5$ ); soil was saturated (100 % whc) with citric and malic acid  
162 solution (0.5, 100, 250, 500 and 1000 mmol l<sup>-1</sup>). Sealed tubes were incubated in a controlled  
163 environment room ( $21 \pm 1$  °C) for 8 h. Then, 25 ml of 50 mM HPCD solution was added.  
164 Tubes were placed onto an orbital shaker (100 rpm) for 22 h at  $21 \pm 1$  °C. Afterwards, samples  
165 were centrifuged (3000  $\times$  g for 1 h) and 5 ml of the supernatant was placed in a glass  
166 scintillation vial and mixed with 15 ml liquid scintillation cocktail. Samples were assessed  
167 through LSC as described previously. Remaining <sup>14</sup>C-associated activity in the soil was  
168 assessed by sample oxidation (Rhodes *et al.*, 2012).

169

### 170 2.3.3 Desorption of <sup>14</sup>C-phenanthrene by organic acids

171 Tests were performed following a randomized design ( $n = 5$ ) and blind sampling. Desorption  
172 kinetics were assessed after 1 and 15 weeks soil-contaminant contact time; at each time point,  
173 4 g soil (dw) was placed into Teflon centrifuge tubes and mixed with 25 ml of organic acid  
174 solution at a given concentration (0.5, 100, 250, 500 and 1000 mmol l<sup>-1</sup>). Tubes were placed  
175 onto an orbital shaker (100 rpm) in a controlled environment room at  $21 \pm 1$  °C. Soil samples  
176 were sequentially extracted after 1, 4, 6, 12, 24, 45, 90, 180 and 360 h by centrifuging at 3000



177  $x$  g for 1 h. Aliquots (5 ml) were mixed with 15 ml liquid scintillation cocktail in a glass  
178 scintillation vial and assessed by LSC. Residual activity in the soil after the last extraction  
179 was assessed by sample oxidation as described by Rhodes *et al.* (2012).

180 Desorption of  $^{14}\text{C}$ -phenanthrene was examined by two- (Equation 1) and three-compartment  
181 (Equation 2) first-order kinetics (Cornelissen *et al.*, 1998b; Rhodes *et al.*, 2010):

182 *Equation 1:*

$$183 \quad S_t / S_0 = [F_{rap} \cdot \exp(-k_{rap} \cdot t)] + [F_{slow} \cdot \exp(-k_{slow} \cdot t)]$$

184 *Equation 2:*

$$185 \quad S_t / S_0 = [F_{rap} \cdot \exp(-k_{rap} \cdot t)] + [F_{slow} \cdot \exp(-k_{slow} \cdot t)] + [F_{very\ slow} \cdot \exp(-k_{very\ slow} \cdot t)]$$

186 where  $S_t$  represents the amount of  $^{14}\text{C}$ -phenanthrene sorbed to the soil at the desorption time  $t$   
187 (h) and  $S_0$  is the initial total amount of  $^{14}\text{C}$ -phenanthrene at the beginning of the assay (time  
188 0).  $F_{rap}$ ,  $F_{slow}$  and  $F_{very\ slow}$  (%) are the rapid, slow and very slow desorbing fractions and  $k_{rap}$ ,  
189  $k_{slow}$  and  $k_{very\ slow}$  ( $\text{h}^{-1}$ ) are the rate constants for the rapid, slow and very slow desorption,  
190 respectively. The model assumes that  $k_{very\ slow} \leq k_{slow} \leq k_{rap}$  (Rhodes *et al.*, 2010; Clegg *et al.*,  
191 2014), and that the addition of the desorbing fractions equals 100 % (Clegg *et al.*, 2014). The  
192 values of  $F_{rap}$ ,  $F_{slow}$ ,  $F_{very\ slow}$ ,  $k_{rap}$ ,  $k_{slow}$  and  $k_{very\ slow}$  were obtained by exponential curve  
193 fitting using Excel Solver add-in, using a non-linear least squares method.

194

## 195 **2.4 Statistical analysis**

196 Statistical analyses were carried using the SPSS 21 (95 % confidence interval). Normality of  
197 the data was verified by Shapiro-Wilk tests, transformations were applied in cases where a

198 normal distribution was not observed. Analyses of the differences across time were carried by  
199 Student's t-test and Wilcoxon test for normal and not normally distributed data respectively.  
200 Differences between the treatments at each time point were analysed using One-Way  
201 ANOVA (Tukey) or Kruskal-Wallis test for normal and not-normal distributed data  
202 respectively. Graphical representations of the results were done with the software Sigma Plot  
203 2000.

## 204 **3 Results**

### 205 **3.1 Short-term impact of organic acids on the mineralisation of <sup>14</sup>C-phenanthrene in soil**

206 The impact of citric, malic, oxalic and succinic acids within a naturally occurring range of  
207 concentrations was tested on the mineralisation of <sup>14</sup>C-phenanthrene. Organic acids were only  
208 observed to produce significant differences on the mineralisation of <sup>14</sup>C-phenanthrene after  
209 14 days soil-PAH contact time, while remaining unaffected after 50 days of soil-PAH contact  
210 time. The data showed that after a short soil-PAH contact time (14 d), the presence of citric  
211 acid (0.1 mmol l<sup>-1</sup>) resulted in a significantly faster rate of mineralisation (29.52 % d<sup>-1</sup>) than  
212 the control (20.99 % d<sup>-1</sup>) ( $F = 2.795$ ,  $p = 0.016$ ) (Table 1). At this same time point, although  
213 not significant ( $p = 0.077$ ), the lag phase of the control soil was longer (18.28 h) than in soil  
214 incubated with organic acids (4.11 – 5.42 h).

215

### 216 **3.2 Preliminary tests for the selection of organic acids and soil-organic acid contact time** 217 **for the assessment of <sup>14</sup>C-phenanthrene bioaccessibility in soil**

218 As significant differences were not observed within the different organic acids used in the  
219 preliminary assay looking at their impact in <sup>14</sup>C-phenanthrene desorption kinetics; malic acid

220 was selected as a representative organic acid for the optimisation of the methodology for the  
221 assessment of bioaccessibility. Results from the test looking at the temporal impact of malic  
222 acid in HPCD-extractable <sup>14</sup>C-phenanthrene fraction showed that soil-organic acid contact  
223 time did not have a significant effect on the bioaccessible fraction of this hydrocarbon ( $p >$   
224 0.05). However, data showed that the largest extractable proportion of <sup>14</sup>C-phenanthrene was  
225 obtained after 8 h of soil-organic acid incubation (control soil, 8.15 %), compared to the  
226 lowest value presented after 48 hours (control soil, 1.10 %). Therefore, 8 h soil-organic acid  
227 incubation was considered to be the most suitable contact time and consequently selected for  
228 further investigation. Furthermore, mineralisation from the HPCD extractable experimental  
229 units within the incubation time was observed to be negligible.

230

### 231 **3.3 Bioaccessibility of <sup>14</sup>C-phenanthrene in soil**

232 Changes on the bioaccessibility of <sup>14</sup>C-phenanthrene in soil was assessed by HPCD  
233 extractions and were observed to be significantly different over time (Table 2,  $t = 66.682$ ,  $p <$   
234 0.001). After one week of soil-PAH contact time; saturation of soil with organic acids (100 %  
235 whc, 8 h) did not have significant effects on the bioaccessibility of <sup>14</sup>C-phenanthrene ( $F =$   
236 1.981,  $p = 0.059$ ). In the case of soil incubated for 15 weeks, the addition of 500 mmol l<sup>-1</sup>  
237 citric acid significantly enhanced the bioaccessible fraction of <sup>14</sup>C-phenanthrene (14.92 %)  
238 compared to the control (6.72 %) ( $F = 4.513$ ,  $p = 0.003$ ).

239

### 240 **3.4 Desorption of <sup>14</sup>C-phenanthrene with organic acids**

241 The amount of <sup>14</sup>C-phenanthrene that was desorbed from soil was significantly affected by  
242 the presence of organic acids ( $p < 0.001$ ) (Table 3). Citric acid at the highest concentration

243 (1000 mmol l<sup>-1</sup>) consistently produced a significantly higher desorption than any other  
244 treatment after 1 week (39.27 %) and 15 weeks (47.86 %) soil-PAH contact time ( $p < 0.001$ ).  
245 Furthermore, this was the only treatment capable of enhancing the desorption of <sup>14</sup>C-  
246 phenanthrene after 1 week soil-PAH contact time. The presence of citric acid (0.5 - 250  
247 mmol l<sup>-1</sup>) and malic acid (0.5 - 500 mmol l<sup>-1</sup>) significantly reduced the total desorbable  
248 fraction of <sup>14</sup>C-phenanthrene soil after one-week soil-PAH contact time ( $p < 0.001$ ). In  
249 contrast, only the lowest concentrations (0.5 mmol l<sup>-1</sup>) of citric acid (13.04 %) and malic acid  
250 (12.51 %) produced significantly lower levels of desorption of <sup>14</sup>C-phenanthrene than the  
251 control (18.96 %) after 15 weeks soil-PAH contact time. This was in contrast to the  
252 desorption behaviour observed at concentrations above 100 mmol l<sup>-1</sup> citric acid and 250  
253 mmol l<sup>-1</sup> malic acid, where desorbed <sup>14</sup>C-phenanthrene was significantly higher ( $p < 0.001$ ).

254

### 255 3.4.1 Impact of organic acids on <sup>14</sup>C-phenanthrene desorption kinetics

256 Desorbing fractions ( $F_{rap}$  and  $F_{slow}$ ;  $F_{rap}$ ,  $F_{slow}$  and  $F_{very\ slow}$ ) and rate constants ( $k_{rap}$  and  $k_{slow}$ ;  
257  $k_{rap}$ ,  $k_{slow}$  and  $k_{very\ slow}$ ) from the two- and three-compartment model fitting, respectively, are  
258 presented on Tables 4 and 5. Squared deviations data showed a better fit by the three-  
259 compartment one (Table SI-3,  $p < 0.001$ ); therefore, further analysis was focused on the  
260 values estimated by this desorption model. Desorbing fractions (%) and rate constants (h<sup>-1</sup>)  
261 obtained by the three-compartment model (Figures SI 4-9) showed significant differences for  
262 all cases ( $p < 0.001$ ). After one-week soil-PAH contact time, significantly higher fractions of  
263 <sup>14</sup>C-phenanthrene were rapidly desorbed by 1000 mmol l<sup>-1</sup> citric (19.22 %) and malic acid  
264 (20.20%) than in the control soil (12.08 %). In contrast, lower concentrations of malic acid  
265 (100 and 250 mmol l<sup>-1</sup>) and citric acid (100 mmol l<sup>-1</sup>) significantly reduced the rapidly  
266 desorbing fractions. Rapidly desorbing rate constants were not affected by the majority of the

267 treatments with the exception of the effect produced by citric acid at 100 mmol l<sup>-1</sup>. Slowly  
268 desorbing fractions were significantly reduced by all treatments apart from citric acid (1000  
269 mmol l<sup>-1</sup>), which was found to be similar to the control. Furthermore, rate constants of this  
270 fraction ( $k_{slow}$ ) were significantly enhanced in most of the treatments (except 0.5 and 1000  
271 mmol l<sup>-1</sup> citric acid), with a longest slowly desorbing phase produced in the present of malic  
272 acid (0.139 – 0.146 h<sup>-1</sup>) when compared against the control (0.013 h<sup>-1</sup>). Very slowly  
273 desorbing fractions accounted for the largest phase in all of the treatments. Moreover, organic  
274 acids significantly increased this fraction in all treatments (except 1000 mmol l<sup>-1</sup> citric acid),  
275 ranging from 77.1 to 88.09 % against the 72.12 % when dH<sub>2</sub>O was used as extractant. Very  
276 slowly desorbing rate constants were also significantly higher in the presence of citric ( $\geq$  500  
277 mmol l<sup>-1</sup>) and malic acid at all tested concentrations.

278 After 15 weeks incubation, high concentrations of citric (500 – 1000 mmol l<sup>-1</sup>) and malic  
279 (500 mmol l<sup>-1</sup>) acid were found to significantly enhance the rapidly desorbing fraction of <sup>14</sup>C-  
280 phenanthrene, representing up to 25.12 % compared to the control (13.11 %). Moreover, low  
281 concentrations of both organic acids (0.5 mmol l<sup>-1</sup>) had the opposite effect, significantly  
282 reducing the fraction of <sup>14</sup>C-phenanthrene desorbed to 3.38 and 5.50 % respectively.

283 Similarly, rapidly desorbing rate constants were also significantly larger when soil was  
284 extracted with citric (0.5 – 1000 mmol l<sup>-1</sup>) and malic acid (1000 mmol l<sup>-1</sup>). Slowly desorbing  
285 fractions ( $F_{slow}$ ) were significantly reduced by all tested concentrations of citric acid and 0.5,  
286 100 and 1000 mmol l<sup>-1</sup> malic acid, while the corresponding desorption rate constants  
287 displayed the opposite behaviour. Fractions desorbed in the very slow phase were  
288 significantly increased by all treatments (except 500 mmol l<sup>-1</sup> malic acid) going from 3.23 %  
289 in the control up to 92.50% when soil was treated with 0.5 mmol l<sup>-1</sup> malic acid. Very slowly  
290 desorbing rate constants were similar to the control with the exception of 100 mmol l<sup>-1</sup> malic  
291 acid where significantly higher values were observed ( $p < 0.001$ ).

## 292 4 Discussion

### 293 4.1 Effect of organic acids in the bioaccessibility of <sup>14</sup>C-phenanthrene in soil

294 The bioaccessibility of PAHs can be quantified using different biological and chemical  
295 approaches. For the purposes of this study, the mineralisation and extraction of <sup>14</sup>C-  
296 phenanthrene by *M. gilvum* and HPCD were used, respectively. As a whole, these two  
297 methods are considered acceptable methodologies to assess not only the fraction of the  
298 hydrocarbon that is freely available to microorganisms, but also encompasses the fraction of  
299 the contaminant that may become bioavailable, and therefore removed from the soil (Semple  
300 *et al.*, 2004). The general absence of effects by organic acids on the mineralisation of <sup>14</sup>C-  
301 phenanthrene reported in this study has also been observed by Cébron *et al.* (2011) and  
302 Louvel *et al.* (2011), both of whom worked with root exudates containing mixtures of organic  
303 acids. Despite this trend, both authors were able to observe an initial acceleration of the  
304 mineralisation process (Cébron *et al.*, 2011; Louvel *et al.*, 2011), as was the case of citric acid  
305 (100 mmol l<sup>-1</sup>) in this present study. Cébron *et al.* (2011) further discussed that this general  
306 absence of effects might be the consequence of enhanced sorption of phenanthrene to SOM  
307 and other soil inorganic fractions such as mineral clays caused by the organic acids, and that  
308 ultimately reflected in a reduction in the availability of phenanthrene for microbial  
309 degradation. Given the chemical characteristics of citric acid, other authors suggest that this  
310 LOA is capable of forming more stable complexes with different compounds in soil (An *et*  
311 *al.*, 2011; Ling *et al.*, 2015).

312 Similar trends were also observed when the bioaccessibility of <sup>14</sup>C-phenanthrene was  
313 measured through its HPCD extractability. Bioaccessibility was only significantly higher in  
314 one of the treatments (500 mmol l<sup>-1</sup> citric and malic acid after 15 weeks soil-PAH contact  
315 time). These findings contrast with that reported by other authors where PAH availability can

316 be significantly promoted by different LOAs assessed through *n*-butanol extractions (Ling *et*  
317 *al.*, 2009; Sun *et al.*, 2012, 2013; Kong *et al.*, 2013; Gao *et al.*, 2015b). Disagreement  
318 between these two trends is suggested to be due to differences in the methodologies used for  
319 this purpose, where *n*-butanol extracted PAH is not only the bioaccessible fraction of the  
320 hydrocarbon, but also a portion of the non-bioaccessible PAH residues, as pointed by Ling *et*  
321 *al.* (2009). Although *n*-butanol has been proposed to act as a predictor for the bioavailability  
322 of PAHs in soil (Kelsey *et al.*, 1997; Liste & Alexander, 2002), this extractant has also been  
323 observed to exhibit greater extraction efficiencies when compared against HPCD  
324 extractability (Swindell & Reid, 2006). This has become important when comparing these  
325 two methods to the mineralisation of phenanthrene in soil (Reid *et al.*, 2000; Rhodes *et al.*,  
326 2010), where close linear 1:1 relationships have been observed for the case of HPCD  
327 extracted PAH. Furthermore, *n*-butanol has also been demonstrated to act as a more  
328 exhaustive extractant than HPCD (Reid *et al.*, 2000; Swindell & Reid, 2006), even extracting  
329 similar quantities of PAHs than DCM, which is often used to determine total concentrations of  
330 contaminants in soil (Reid *et al.*, 2000).

331 The impact that organic acids from the rhizosphere might have towards the biodegradation of  
332 PAHs remains a poorly explored area; however, this is one of the presumed mechanisms by  
333 which plant-enhanced bioremediation is thought to occur (Pilon-Smits, 2005). Changes in the  
334 physico-chemical conditions in soil, such as pH, may play an important role in the microbial  
335 degradation of PAHs (Kästner *et al.*, 1998). Specifically, the acidic ranges of pH that the  
336 presence of high concentrations of organic acids will produce, associated to the SOM-bound  
337 PAHs that have been discussed, may be responsible to limit the microbial activity. This may  
338 explain why, despite the fact that larger amounts of <sup>14</sup>C-phenanthrene can be extracted with  
339 high concentrations of organic acids, the PAHs are not being biodegraded by soil bacteria and  
340 metabolised to CO<sub>2</sub>.

## 341 4.2 Impact of LOAs on the desorption of <sup>14</sup>C-phenanthrene in soil

342 The total desorbable fraction of <sup>14</sup>C-phenanthrene did not decrease as a function of time over  
343 the course of the incubation when soil was extracted with organic acids ( $\geq 100 \text{ mmol l}^{-1}$ ).  
344 Despite the general acknowledgement of the negative correlation between the extractability  
345 of organic contaminants and contact time (Hatzinger & Alexander, 1995; Semple *et al.*,  
346 2003), this behaviour was only observed in the control and the lowest tested concentration of  
347 organic acids after 15 weeks of soil-PAH contact time. This trend suggests that high amounts  
348 of organic acids could potentially restrict the reduction of bioaccessibility of <sup>14</sup>C-  
349 phenanthrene, therefore limiting the ageing process. Although not observed before, this  
350 behaviour could be the reflection of a dual effect of organic acids on phenanthrene sorption  
351 reported by Ouvrard *et al.* (2006) who described the impact of LOAs as a combined process  
352 characterised by an initial short term enhanced sorption of phenanthrene by SOM, followed  
353 by an increased mass transfer of the hydrocarbon due to the destabilisation of this soil  
354 fraction. Similarly, data from the present study showed a general reduction of the extractable  
355 <sup>14</sup>C-phenanthrene after a short period of ageing while organic acids were consistently  
356 observed to promote a larger desorption after 15 weeks of soil-PAH contact time when  
357 compared against the control. This increase in the extractability of PAHs by LOAs has also  
358 been reported by other authors (Ling *et al.*, 2009, 2015; Gao *et al.*, 2010a; b, 2015b; Kong *et*  
359 *al.*, 2013), but rarely considered the impact of soil-PAH ageing included in the present study.  
360 Although not common, the reduction of <sup>14</sup>C-phenanthrene desorption in the presence of  
361 organic acids observed after a short soil-PAH contact time in the present study has been  
362 reported before (Ouvrard *et al.*, 2006; Zhu *et al.*, 2009; Gao *et al.*, 2015b). This initial  
363 behaviour has been associated with the capacity of small amounts of oxalate, citrate and  
364 malate to promote the sorption of anions to the soil (Jones & Brassington, 1998; Jones *et al.*,  
365 2003). In a similar way, phenanthrene has been hypothesised to be also sorbed through the



366 development of new sorption sites by these sorbed organic acids (Ouvrard *et al.*, 2006; Gao *et*  
367 *al.*, 2015a).

368 LOAs have been acknowledged to significantly influence the physical, chemical and  
369 biological properties of soil (Jones & Darrah, 1994; Jones, 1998). As such, the main  
370 mechanism behind the enhancement of the desorption of PAHs in soil impacted by organic  
371 acids has been proposed to be the solubilisation of soil organic matter (SOM) with a  
372 subsequent release SOM-associated hydrocarbons (Ouvrard *et al.*, 2006; Agnello *et al.*,  
373 2014). This explanation is supported by findings from different authors who have reported  
374 consistently higher amounts of dissolved organic matter and certain minerals when artificial  
375 root exudates (Gao *et al.*, 2010a) and single LOAs (Ling *et al.*, 2009, 2015; Gao *et al.*, 2010b,  
376 2015a; Sun *et al.*, 2012; Kong *et al.*, 2013) were used to extract PAHs from contaminated  
377 soil. In a similar way, previously immobilised aromatic compounds have been observed to be  
378 released from soil to pore water after the introduction of organic acids solution (White *et al.*,  
379 2003; Gao *et al.*, 2015b; Keiluweit *et al.*, 2015).

#### 380 4.2.1 Desorption kinetics

381 <sup>14</sup>C-Phenanthrene desorption kinetics in the presence of the organic acids displayed a 3-  
382 compartment desorption behaviour. Although desorption of organic contaminants has been  
383 widely observed to behave with an initial rapid desorption followed by a slower phase  
384 (Cornelissen *et al.*, 1998b), this rapid/slow/very slow release from the soil has also been  
385 observed (Rhodes *et al.*, 2010). Typically, studies investigating desorption kinetics are  
386 performed using extractants that are known (or suspected) to correlate with the biodegradable  
387 fraction of the contaminant in question (Cornelissen *et al.*, 2001; Rhodes *et al.*, 2010).  
388 However, this current research study was focused on the assessment of the desorbing  
389 potential that organic acids might be able to provide. Bearing this in mind, the proportion of

390 <sup>14</sup>C-phenanthrene desorbed at each of these phases should be considered as a measure of the  
391 behaviour of this PAH under the influence of organic acids rather than an indication of its  
392 bioaccessibility.

393 The different fractions described by the desorption kinetics can be interpreted as the  
394 biodegradable ( $F_{rap}$ ), and less accessible  $F_{slow}$  and/or  $F_{very\ slow}$  fractions of the organic  
395 contaminant (Cornelissen *et al.*, 1998b; Rhodes *et al.*, 2010). Results from this investigation  
396 showed that the majority of the treatments had a tendency to enhance the very slowly  
397 desorbing fraction ( $F_{very\ slow}$ ). These results could be interpreted as that the presence of organic  
398 acids might be able to mobilise a significant proportion of the readily bioaccessible fraction  
399 of <sup>14</sup>C-phenanthrene ( $F_{rap}$ ) towards a less accessible form ( $F_{slow}$  and  $F_{very\ slow}$ ), therefore  
400 limiting the biological degradation of the contaminant or the rate at which this process takes  
401 place (Pignatello & Xing, 1996; Clegg *et al.*, 2014). Moreover, similar behaviour has been  
402 observed to occur during the mineralisation of organic acids, where these compounds have  
403 been observed to induce shifts of <sup>14</sup>CO<sub>2</sub> production from a rapid to a slower phase (Oburger  
404 *et al.*, 2009).

## 405 **5 Conclusions**

406 Organic acids found within the rhizosphere play an important role on the behaviour of  
407 phenanthrene in soil. It was found that the total extractable fraction of <sup>14</sup>C-phenanthrene can  
408 be significantly enhanced by citric and malic acid. This effect is most likely to be observed at  
409 a longer soil-PAH contact time, where organic acids showed to restrict the ageing effect.  
410 Despite these enhancing effects, desorption kinetics indicated that the desorbed phenanthrene  
411 was readily available given the behaviour as slow and very slow desorbing fractions. These  
412 trends were confirmed when accessibility and mineralisation of <sup>14</sup>C-phenanthrene were  
413 assessed. In this case, despite the enhancement of the total hydrocarbon extractable fractions

414 in the presence of citric and malic acid; there is no clear evidence suggesting that this  
415 condition can promote the microbial utilisation of <sup>14</sup>C-phenanthrene. This study contributes to  
416 the understanding of the role of root exudation within the rhizosphere towards the  
417 bioaccessibility and biodegradation of hydrocarbons in contaminated soil. It is important to  
418 note that organic acids may be able to remobilise contaminants, which were considered to be  
419 non-bioaccessible. This may be important from a risk assessment perspective; however, the  
420 concentrations of remobilised PAHs may be low and not represent a risk to environmental or  
421 human health (Umeh et al., 2018).

## 422 **Acknowledgements**

423 GVC would like to thank the Mexican National Council for Science and Technology  
424 (CONACyT) for their financial support through a scholarship (No. 313807) to pursue  
425 postgraduate studies at the Lancaster Environment Centre, Lancaster University.

426

427 **References**

- 428 Agnello, A.C., Huguenot, D., Van Hullebusch, E.D. & Esposito, G. 2014. Enhanced  
429 phytoremediation: A review of low molecular weight organic acids and surfactants used  
430 as amendments. *Critical Reviews in Environmental Science and Technology*, **44**, 2531–  
431 2576.
- 432 An, C.J., Huang, G.H., Wei, J. & Yu, H. 2011. Effect of short-chain organic acids on the  
433 enhanced desorption of phenanthrene by rhamnolipid biosurfactant in soil-water  
434 environment. *Water Research*, **45**, 5501–5510, (At:  
435 <http://dx.doi.org/10.1016/j.watres.2011.08.011> ).
- 436 Anderson, T.A., Guthrie, E.A. & Walton, B.T. 1993. Bioremediation in the Rhizosphere:  
437 Plant roots and associated microbes clean contaminated soil. *Environmental Science &*  
438 *Technology*, **27**, 2630–2636.
- 439 Cébron, A., Louvel, B., Faure, P., France-Lanord, C., Chen, Y., Murrell, J.C. & Leyval, C.  
440 2011. Root exudates modify bacterial diversity of phenanthrene degraders in PAH-  
441 polluted soil but not phenanthrene degradation rates. *Environmental Microbiology*, **13**,  
442 722–736.
- 443 Clegg, H., Riding, M.J., Oliver, R., Jones, K.C. & Semple, K.T. 2014. The impact of soil  
444 organic matter and soil sterilisation on the bioaccessibility of <sup>14</sup>C-azoxystrobin  
445 determined by desorption kinetics. *Journal of Hazardous Materials*, **278**, 336–342.
- 446 Cornelissen, G., Van Noort, P.C.M. & Covers, H.A.J. 1998a. Mechanism of slow desorption  
447 of organic compounds from sediments: A study using model sorbents. *Environmental*  
448 *Science & Technology*, **32**, 3124–3131.

- 449 Cornelissen, G., Rigterink, H., Ferdinandy, M.M.A. & Van Noort, P.C.M. 1998b. Rapidly  
450 desorbing fractions of PAHs in contaminated sediments as a predictor of the extent of  
451 bioremediation. *Environmental Science & Technology*, **32**, 966–970.
- 452 Cornelissen, G., Rigterink, H., ten Hulscher, D.E.M., Vrind, B.A. & van Noort, P.C.M. 2001.  
453 A simple tenax® extraction method to determine the availability of sediment-sorbed  
454 organic compounds. *Environmental Toxicology and Chemistry*, **20**, 706–711.
- 455 Dakora, F.D. & Phillips, D.A. 2002. Root exudates as mediators of mineral acquisition in  
456 low-nutrient environments. *Plant and Soil*, **245**, 35–47.
- 457 Doick, K.J., Lee, P.H. & Semple, K.T. 2003. Assessment of spiking procedures for the  
458 introduction of a phenanthrene-LNAPL mixture into field-wet soil. *Environmental*  
459 *Pollution*, **126**, 399–406.
- 460 Gao, Y., Ren, L., Ling, W., Gong, S., Sun, B. & Zhang, Y. 2010a. Desorption of  
461 phenanthrene and pyrene in soils by root exudates. *Bioresource Technology*, **101**, 1159–  
462 1165.
- 463 Gao, Y., Ren, L., Ling, W., Kang, F., Zhu, X. & Sun, B. 2010b. Effects of low-molecular-  
464 weight organic acids on sorption–desorption of phenanthrene in soils. *Soil Science*  
465 *Society of America Journal*, **74**, 51.
- 466 Gao, Y., Wang, N., Li, H., Hu, X. & Goikavi, C. 2015a. Low-molecular-weight organic acids  
467 influence the sorption of phenanthrene by different soil particle size fractions. *Journal of*  
468 *Environment Quality*, **44**, 219.
- 469 Gao, Y., Yuan, X., Lin, X., Sun, B. & Zhao, Z. 2015b. Low-molecular-weight organic acids  
470 enhance the release of bound PAH residues in soils. *Soil and Tillage Research*, **145**,

471 103–110.

472 Hatzinger, P.B. & Alexander, M. 1995. Effect of ageing of chemicals in soil on their  
473 biodegradability and extractability. *Environmental Science & Technology*, **29**, 537–545.

474 van Hees, P.A.W., Jones, D.L., Finlay, R., Godbold, D.L. & Lundström, U.S. 2005. The  
475 carbon we do not see—the impact of low molecular weight compounds on carbon  
476 dynamics and respiration in forest soils: a review. *Soil Biology and Biochemistry*, **37**, 1–  
477 13.

478 Joner, E.J., Corgié, S.C. & Leyval, C. 2002. Nutritional constraints to degradation of  
479 polycyclic aromatic hydrocarbons in a simulated rhizosphere. *Soil Biology and*  
480 *Biochemistry*, **34**, 859–864.

481 Jones, D. 1998. Organic acids in the rhizosphere - a critical review. *Plant and Soil*, **205**, 25–  
482 44.

483 Jones, D.L. & Brassington, D.S. 1998. Sorption of organic acids in acid soils and its  
484 implications in the rhizosphere. *European Journal of Soil Science*, **49**, 447–455.

485 Jones, D.L. & Darrah, P.R. 1994. Role of root derived organic acids in the mobilization of  
486 nutrients from the rhizosphere. *Plant and Soil*, **166**, 247–257.

487 Jones, D.L.L., Dennis, P.G.G., Owen, a. G.G. & van Hees, P. a. W. a W. 2003. Organic acid  
488 behavior in soils - misconceptions and knowledge gaps. *Plant and Soil*, **248**, 31–41.

489 Kästner, M., Breuer-Jammali, M. & Mahro, B. 1998. Impact of inoculation protocols,  
490 salinity, and pH on the degradation of polycyclic aromatic hydrocarbons (PAHs) and  
491 survival of PAH-degrading bacteria introduced into soil. *Applied and Environmental*  
492 *Microbiology*, 359–362.

- 493 Keiluweit, M., Bougoure, J.J., Nico, P.S., Pett-Ridge, J., Weber, P.K. & Kleber, M. 2015.  
494 Mineral protection of soil carbon counteracted by root exudates. *Nature Climate*  
495 *Change*, **5**, 588–595.
- 496 Kelsey, J.W., Kottler, B.D. & Alexander, M. 1997. Selective chemical extractants to predict  
497 bioavailability of soil-aged organic chemicals. *Environmental Science & Technology*, **31**,  
498 214–217.
- 499 Kong, H., Sun, R., Gao, Y. & Sun, B. 2013. Elution of polycyclic aromatic hydrocarbons in  
500 soil columns using low-molecular-weight organic acids. *Soil Science Society of America*  
501 *Journal*, **77**, 72–82.
- 502 Ling, W., Ren, L., Gao, Y., Zhu, X. & Sun, B. 2009. Impact of low-molecular-weight organic  
503 acids on the availability of phenanthrene and pyrene in soil. *Soil Biology and*  
504 *Biochemistry*, **41**, 2187–2195.
- 505 Ling, W., Sun, R., Gao, X., Xu, R. & Li, H. 2015. Low-molecular-weight organic acids  
506 enhance desorption of polycyclic aromatic hydrocarbons from soil. *European Journal of*  
507 *Soil Science*, **66**, 339–347.
- 508 Liste, H.H. & Alexander, M. 2002. Butanol extraction to predict bioavailability of PAHs in  
509 soil. *Chemosphere*, **46**, 1011–1017.
- 510 Louvel, B., Cébron, A. & Leyval, C. 2011. Root exudates affect phenanthrene  
511 biodegradation, bacterial community and functional gene expression in sand  
512 microcosms. *International Biodeterioration and Biodegradation*, **65**, 947–953.
- 513 Martin, B.C., George, S.J., Price, C.A., Ryan, M.H. & Tibbett, M. 2014. The role of root  
514 exuded low molecular weight organic anions in facilitating petroleum hydrocarbon

515 degradation: Current knowledge and future directions. *Science of the Total Environment*,  
516 **472**, 642–653.

517 Miya, R.K. & Firestone, M.K. 2001. Enhanced phenanthrene biodegradation in soil by  
518 slender oat root exudates and root debris. *Journal of Environment Quality*, **30**, 1911–  
519 1918.

520 Oburger, E., Kirk, G.J.D., Wenzel, W.W., Puschenreiter, M. & Jones, D.L. 2009. Interactive  
521 effects of organic acids in the rhizosphere. *Soil Biology and Biochemistry*, **41**, 449–457.

522 Ouvrard, S., Lapole, D. & Morel, J.L. 2006. Root exudates impact on phenanthrene  
523 availability. *Water, Air, and Soil Pollution: Focus*, **6**, 343–352.

524 Pignatello, J.J. & Xing, B. 1996. Mechanisms of slow sorption of organic chemicals to  
525 natural particles. *Environmental Science & Technology*, **30**, 1–11.

526 Pilon-Smits, E. 2005. Phytoremediation. *Annual Review of Plant Biology*, **56**, 15–39.

527 Posada-Baquero, R., Grifoll, M. & Ortega-Calvo, J.J. 2019. Rhamnolipid-enhanced  
528 solubilization and biodegradation of PAHs in soils after conventional bioremediation.  
529 *Science of the Total Environment*, **668**, 790–796.

530 Reid, B.J., MacLeod, C.J.A., Lee, P.H., Morriss, A.W.J., Stokes, J.D. & Semple, K.T. 2001.  
531 A simple <sup>14</sup>C-respirometric method for assessing microbial catabolic potential and  
532 contaminant bioavailability. *FEMS Microbiology Letters*, **196**, 141–146.

533 Reid, B.J., Stokes, J.D., Jones, K.C. & Semple, K.T. 2000. Nonexhaustive cyclodextrin-based  
534 extraction technique for the evaluation of PAH bioavailability. *Environmental Science &*  
535 *Technology*, **34**, 3174–3179.



- 536 Reilley, K.A., Banks, M.K. & Schwab, A.P. 1996. Organic chemicals in the environment:  
537 Dissipation of polycyclic aromatic hydrocarbons in the rhizosphere. *Journal of*  
538 *Environmental Quality*, **25**, 212–219.
- 539 Rhodes, A.H., McAllister, L.E. & Semple, K.T. 2010. Linking desorption kinetics to  
540 phenanthrene biodegradation in soil. *Environmental Pollution*, **158**, 1348–1353.
- 541 Rhodes, A.H., Riding, M.J., McAllister, L.E., Lee, K. & Semple, K.T. 2012. Influence of  
542 activated charcoal on desorption kinetics and biodegradation of phenanthrene in soil.  
543 *Environmental Science & Technology*, **46**, 12445–12451.
- 544 Semple, K.T., Dew, N.M., Doick, K.J. & Rhodes, A.H. 2006. Can microbial mineralization  
545 be used to estimate microbial availability of organic contaminants in soil?  
546 *Environmental Pollution*, **140**, 164–172.
- 547 Semple, K.T., Doick, K.J., Jones, K.C., Burauel, P., Craven, A. & Harms, H. 2004. Defining  
548 bioavailability and bioaccessibility of contaminated soil and sediment is complicated.  
549 *Environmental Science & Technology*, **38**, 228A-231A.
- 550 Semple, K.T., Doick, K.J., Wick, L.Y. & Harms, H. 2007. Microbial interactions with  
551 organic contaminants in soil: Definitions, processes and measurement. *Environmental*  
552 *Pollution*, **150**, 166–176.
- 553 Semple, K.T., Morriss, a. W.J. & Paton, G.I. 2003. Bioavailability of hydrophobic organic  
554 contaminants in soils: fundamental concepts and techniques for analysis. *European*  
555 *Journal of Soil Science*, **54**, 809–818.
- 556 Semple, K.T., Riding, M.J., McAllister, L.E., Sopena-Vazquez, F. & Bending, G.D. 2013.  
557 Impact of black carbon on the bioaccessibility of organic contaminants in soil. *Journal*

558 *of Hazardous Materials*, **261**, 808–816.

559 Shukla, K.P., Sharma, S., Singh, N.K., Singh, V. & Tiwari, K. 2011. Nature and role of root  
560 exudates : Efficacy in bioremediation. *African Journal of Biotechnology*, **10**, 9717–  
561 9724.

562 Siciliano, S.D. & Germida, J.J. 1998. Mechanisms of phytoremediation: biochemical and  
563 ecological interactions between plants and bacteria. *Environmental Reviews*, **6**, 65–79.

564 Sivaram, A.K., Logeshwaran, P., Lockington, R., Naidu, R. & Megharaj, M. 2019. Low  
565 molecular weight organic acids enhance the high molecular weight polycyclic aromatic  
566 hydrocarbons degradation by bacteria. *Chemosphere*, **222**, 132–140.

567 Sun, B., Gao, Y., Liu, J. & Sun, Y. 2012. The impact of different root exudate components on  
568 phenanthrene availability in soil. *Soil Science Society of America Journal Abstract - Soil  
569 Chemistry*, **76**, 2041–2050.

570 Sun, B., Ling, W. & Wang, Y. 2013. Can root exudate components influence the availability  
571 of pyrene in soil? *Journal of Soils and Sediments*, **13**, 1161–1169.

572 Swindell, A.L. & Reid, B.J. 2006. Comparison of selected non-exhaustive extraction  
573 techniques to assess PAH availability in dissimilar soils. *Chemosphere*, **62**, 1126–1134.

574 Towell, M.G., Browne, L.A., Paton, G.I. & Semple, K.T. 2011. Impact of carbon  
575 nanomaterials on the behaviour of <sup>14</sup>C-phenanthrene and <sup>14</sup>C-benzo-[a] pyrene in soil.  
576 *Environmental Pollution*, **159**, 706–715.

577 Umeh, U.C., Duan, L., Naidu, R. and Semple, K.T., 2018. Remobilisation of Non-Extractable  
578 Benzo[a]pyrene Residues in Contrasting Australian Soils: Effects of Aging, Spiked  
579 Concentration, and Soil Properties. *Environmental Science and Technology* **52**, 12295-

580 12305.

581 White, J.C., Mattina, M.I., Lee, W.Y., Eitzer, B.D. & Iannucci-Berger, W. 2003. Role of  
582 organic acids in enhancing the desorption and uptake of weathered p,p'-DDE by  
583 *Cucurbita pepo*. *Environmental Pollution*, **124**, 71–80.

584 Xiong, B., Zhang, Y., Hou, Y., Arp, H.P.H., Reid, B.J. & Cai, C. 2017. Enhanced  
585 biodegradation of PAHs in historically contaminated soil by *M. gilvum* inoculated  
586 biochar. *Chemosphere*, **182**, 316–324.

587 Yoshitomi, K. & Shann, J. 2001. Corn (*Zea mays* L.) root exudates and their impact on 14C-  
588 pyrene mineralization. *Soil Biology and Biochemistry*, **33**, 1769–1776.

589 Zhu, Y., Zhang, S., Huang, H. & Wen, B. 2009. Effects of maize root exudates and organic  
590 acids on the desorption of phenanthrene from soils. *Journal of Environmental Sciences*,  
591 **21**, 920–926.

592

Table 1. Mineralisation kinetics of  $^{14}\text{C}$ -phenanthrene from soil affected by organic acids after 14 and 50 days ageing. Values of the lag phases (h), maximum rates ( $\% \text{ d}^{-1}$ ) and total extents (%) represent the mean  $\pm$  standard error of the mean ( $n = 3$ ). Different letters indicate significant differences between the treatments at each time point assessed by post hoc Tukey tests.

14 days ageing				
Treatment	Concentration (mmol l <sup>-1</sup> )	Lag phase*	Fastest rate	Total extent
Control	0	18.288 $\pm$ 0.358	<sup>a</sup> 20.991 $\pm$ 0.564	<sup>ab</sup> 57.483 $\pm$ 0.617
Citric acid	0.1	4.104 $\pm$ 0.417	<sup>b</sup> 29.520 $\pm$ 2.316	<sup>ab</sup> 65.325 $\pm$ 3.590
	0.5	4.728 $\pm$ 0.382	<sup>ab</sup> 25.519 $\pm$ 1.439	<sup>ab</sup> 63.863 $\pm$ 2.559
Malic acid	0.1	4.440 $\pm$ 0.347	<sup>ab</sup> 27.114 $\pm$ 1.268	<sup>ab</sup> 63.741 $\pm$ 1.430
	0.5	4.320 $\pm$ 0.364	<sup>ab</sup> 27.902 $\pm$ 1.050	<sup>b</sup> 68.370 $\pm$ 0.101
Oxalic acid	0.1	5.424 $\pm$ 0.564	<sup>ab</sup> 22.562 $\pm$ 2.125	<sup>a</sup> 54.085 $\pm$ 3.386
	0.5	4.632 $\pm$ 0.397	<sup>ab</sup> 26.093 $\pm$ 1.576	<sup>ab</sup> 58.915 $\pm$ 1.694
Succinic acid	0.1	4.656 $\pm$ 0.260	<sup>ab</sup> 25.806 $\pm$ 0.664	<sup>ab</sup> 63.543 $\pm$ 2.409
	0.5	4.800 $\pm$ 0.445	<sup>ab</sup> 25.270 $\pm$ 1.938	<sup>ab</sup> 60.683 $\pm$ 3.183
50 days ageing				
Treatment	Concentration (mmol l <sup>-1</sup> )	Lag phase	Fastest rate	Total extent
Control	0	<sup>a</sup> 84.350 $\pm$ 2.400	<sup>a</sup> 1.613 $\pm$ 0.257	<sup>a</sup> 11.581 $\pm$ 1.321
Citric acid	0.1	<sup>a</sup> 81.128 $\pm$ 1.945	<sup>a</sup> 1.607 $\pm$ 0.128	<sup>a</sup> 11.419 $\pm$ 0.560
	0.5	<sup>a</sup> 48.488 $\pm$ 1.400	<sup>a</sup> 1.651 $\pm$ 0.020	<sup>a</sup> 11.688 $\pm$ 0.332
Malic acid	0.1	<sup>a</sup> 69.532 $\pm$ 2.020	<sup>a</sup> 1.592 $\pm$ 0.108	<sup>a</sup> 12.535 $\pm$ 0.474
	0.5	<sup>a</sup> 85.519 $\pm$ 2.319	<sup>a</sup> 1.345 $\pm$ 0.089	<sup>a</sup> 10.590 $\pm$ 0.667
Oxalic acid	0.1	<sup>a</sup> 69.922 $\pm$ 0.894	<sup>a</sup> 1.553 $\pm$ 0.026	<sup>a</sup> 12.387 $\pm$ 0.410
	0.5	<sup>a</sup> 72.237 $\pm$ 1.494	<sup>a</sup> 1.670 $\pm$ 0.128	<sup>a</sup> 12.959 $\pm$ 0.935
Succinic acid	0.1	<sup>a</sup> 78.770 $\pm$ 1.819	<sup>a</sup> 1.661 $\pm$ 0.140	<sup>a</sup> 11.093 $\pm$ 0.591
	0.5	<sup>a</sup> 56.177 $\pm$ 2.119	<sup>a</sup> 1.959 $\pm$ 0.154	<sup>a</sup> 13.641 $\pm$ 0.159

\*Not normally distributed data analysed by Kruskal-Wallis non-parametric test ( $p = 0.077$ )

Table 2. HPCD extractable fraction of <sup>14</sup>C-phenanthrene from soil after 1 and 15 weeks ageing following saturation with organic acids solution (100 % whc, 8 h). Values represent the mean ± standard error of the mean (*n* = 5). Different letters indicate significant differences between the treatments at each time point (Tukey)

Treatment	Concentration (mmol l <sup>-1</sup> )	Bioaccessible <sup>14</sup> C-phenanthrene (%)	
		1 week*	15 weeks
Control	0	<sup>a</sup> 79.841 ± 0.717	<sup>ab</sup> 6.721 ± 0.970
Citric acid	0.5	<sup>a</sup> 78.964 ± 2.070	<sup>a</sup> 5.033 ± 0.955
	100	<sup>a</sup> 78.769 ± 1.346	<sup>ab</sup> 6.492 ± 1.646
	250	<sup>a</sup> 79.239 ± 1.243	<sup>abc</sup> 9.804 ± 1.033
	500	<sup>a</sup> 79.852 ± 0.633	<sup>c</sup> 14.929 ± 0.582
	1000	<sup>a</sup> 76.745 ± 0.164	<sup>bc</sup> 11.223 ± 1.524
Malic acid	0.5	<sup>a</sup> 81.138 ± 0.611	<sup>ab</sup> 7.992 ± 0.690
	100	<sup>a</sup> 80.517 ± 0.366	<sup>abc</sup> 8.938 ± 1.509
	250	<sup>a</sup> 78.540 ± 0.793	<sup>ab</sup> 8.764 ± 2.035
	500	<sup>a</sup> 77.937 ± 0.579	<sup>abc</sup> 10.726 ± 0.871
	1000	<sup>a</sup> 76.737 ± 1.003	<sup>ab</sup> 8.594 ± 1.284

Table 3. Total <sup>14</sup>C-phenanthrene desorbed from soil after 1 and 15 weeks ageing. Values represent the mean ± standard error of the mean (*n* = 5). Different letters indicate significant differences between the treatments at each time point (Tukey).

Treatment	Concentration (mmol l <sup>-1</sup> )	Desorbed <sup>14</sup> C-pheanthrene (%)	
		1 week	15 weeks
Control	0	<sup>de</sup> 27.980 ± 1.636	<sup>b</sup> 18.958 ± 0.931
Citric acid	0.5	<sup>ab</sup> 20.755 ± 0.432	<sup>a</sup> 13.038 ± 1.010
	100	<sup>a</sup> 17.221 ± 0.211	<sup>cd</sup> 26.462 ± 0.431
	250	<sup>ab</sup> 19.709 ± 0.081	<sup>de</sup> 31.264 ± 1.851
	500	<sup>cd</sup> 26.579 ± 0.795	<sup>f</sup> 40.006 ± 0.655
	1000	<sup>f</sup> 39.274 ± 1.921	<sup>g</sup> 47.856 ± 1.060
Malic acid	0.5	<sup>ab</sup> 20.068 ± 0.376	<sup>a</sup> 12.507 ± 0.176
	100	<sup>a</sup> 16.552 ± 0.119	<sup>bc</sup> 21.986 ± 1.363
	250	<sup>ab</sup> 18.820 ± 0.256	<sup>cd</sup> 25.923 ± 0.983
	500	<sup>bc</sup> 22.558 ± 0.266	<sup>e</sup> 32.955 ± 1.134
	1000	<sup>e</sup> 31.720 ± 1.249	<sup>ef</sup> 36.184 ± 2.054

Table 4. Desorbing fractions ( $F_{rap}$  and  $F_{slow}$ ) and constant rates ( $k_{rap}$  and  $k_{slow}$ ) calculated by a two-compartment model. Values represent the mean  $\pm$  standard error of the mean ( $n = 5$ ). Different letters indicate significant differences between the treatments assessed by post hoc Tukey tests

Treatment	Concentration (mmol l <sup>-1</sup> )	$F_{rap}$ (%)	$k_{rap}$ (h <sup>-1</sup> )	$F_{slow}$ (%)	$k_{slow}$ (h <sup>-1</sup> )
1 week ageing					
Control	0	<sup>cd</sup> 16.571 $\pm$ 0.442	<sup>a</sup> 0.128 $\pm$ 0.004	<sup>cd</sup> 83.428 $\pm$ 0.442	<sup>c</sup> 0.001 <0.001
Citric acid	0.5	<sup>abc</sup> 14.393 $\pm$ 0.381	<sup>ab</sup> 0.142 $\pm$ 0.004	<sup>def</sup> 85.606 $\pm$ 0.381	<sup>ab</sup> <0.001 <0.001
	100	<sup>a</sup> 12.313 $\pm$ 0.160	<sup>b</sup> 0.145 $\pm$ 0.001	<sup>f</sup> 87.687 $\pm$ 0.160	<sup>a</sup> <0.001 <0.001
	250	<sup>abc</sup> 14.423 $\pm$ 0.108	<sup>ab</sup> 0.134 $\pm$ 0.001	<sup>def</sup> 85.576 $\pm$ 0.108	<sup>a</sup> <0.001 <0.001
	500	<sup>bcd</sup> 19.090 $\pm$ 0.706	<sup>ab</sup> 0.142 $\pm$ 0.006	<sup>c</sup> 80.909 $\pm$ 0.706	<sup>b</sup> 0.001 <0.001
	1000	<sup>e</sup> 28.381 $\pm$ 1.587	<sup>ab</sup> 0.140 $\pm$ 0.005	<sup>a</sup> 71.618 $\pm$ 1.587	<sup>d</sup> 0.001 <0.001
Malic acid	0.5	<sup>abc</sup> 15.123 $\pm$ 0.104	<sup>b</sup> 0.139 $\pm$ 0.002	<sup>def</sup> 84.876 $\pm$ 0.104	<sup>a</sup> <0.001 <0.001
	100	<sup>a</sup> 11.901 $\pm$ 0.143	<sup>ab</sup> 0.142 $\pm$ 0.002	<sup>f</sup> 88.098 $\pm$ 0.143	<sup>a</sup> <0.001 <0.001
	250	<sup>ab</sup> 13.447 $\pm$ 0.119	<sup>ab</sup> 0.146 $\pm$ 0.002	<sup>ef</sup> 86.553 $\pm$ 0.119	<sup>a</sup> <0.001 <0.001
	500	<sup>d</sup> 16.122 $\pm$ 0.272	<sup>ab</sup> 0.141 $\pm$ 0.003	<sup>cde</sup> 83.877 $\pm$ 0.272	<sup>ab</sup> <0.001 <0.001
	1000	<sup>f</sup> 22.899 $\pm$ 0.936	<sup>ab</sup> 0.140 $\pm$ 0.003	<sup>b</sup> 77.100 $\pm$ 0.936	<sup>c</sup> 0.001 <0.001
15 weeks ageing					
Control	0	<sup>b</sup> 14.683 $\pm$ 3.591	<sup>ab</sup> 0.195 $\pm$ 0.009	<sup>f</sup> 85.317 $\pm$ 3.591	<sup>a</sup> 0.001 <0.001
Citric acid	0.5	<sup>a</sup> 6.789 $\pm$ 0.247	<sup>ab</sup> 0.170 $\pm$ 0.012	<sup>g</sup> 93.211 $\pm$ 0.247	<sup>a</sup> <0.001 <0.001
	100	<sup>bc</sup> 14.566 $\pm$ 0.436	<sup>bc</sup> 0.217 $\pm$ 0.020	<sup>ef</sup> 85.434 $\pm$ 0.436	<sup>b</sup> 0.001 <0.001
	250	<sup>b</sup> 19.534 $\pm$ 0.493	<sup>b</sup> 0.186 $\pm$ 0.014	<sup>bc</sup> 80.466 $\pm$ 0.493	<sup>bc</sup> 0.001 <0.001
	500	<sup>f</sup> 25.150 $\pm$ 0.876	<sup>bc</sup> 0.213 $\pm$ 0.009	<sup>c</sup> 74.850 $\pm$ 0.876	<sup>cd</sup> 0.001 <0.001
	1000	<sup>g</sup> 30.907 $\pm$ 0.377	<sup>c</sup> 0.262 $\pm$ 0.006	<sup>a</sup> 69.093 $\pm$ 0.377	<sup>d</sup> 0.002 <0.001
Malic acid	0.5	<sup>a</sup> 7.533 $\pm$ 0.333	<sup>bc</sup> 0.206 $\pm$ 0.023	<sup>g</sup> 92.467 $\pm$ 0.333	<sup>a</sup> <0.001 <0.001
	100	<sup>bcd</sup> 15.906 $\pm$ 0.770	<sup>a</sup> 0.108 $\pm$ 0.021	<sup>def</sup> 84.094 $\pm$ 0.770	<sup>a</sup> <0.001 <0.001
	250	<sup>cde</sup> 17.146 $\pm$ 0.621	<sup>ab</sup> 0.156 $\pm$ 0.021	<sup>cde</sup> 82.854 $\pm$ 0.621	<sup>ab</sup> 0.001 <0.001
	500	<sup>e</sup> 20.412 $\pm$ 0.598	<sup>bc</sup> 0.209 $\pm$ 0.007	<sup>a</sup> 79.588 $\pm$ 0.598	<sup>bc</sup> 0.001 <0.001
	1000	<sup>f</sup> 24.585 $\pm$ 1.845	<sup>ab</sup> 0.156 $\pm$ 0.009	<sup>b</sup> 75.415 $\pm$ 1.845	<sup>bc</sup> 0.001 <0.001

Table 5. Desorbing fractions ( $F_{rap}$ ,  $F_{slow}$  and  $F_{very\ slow}$ ) and constant rates ( $k_{rap}$ ,  $k_{slow}$  and  $k_{very\ slow}$ ) calculated by a three-compartment model. Values represent the mean  $\pm$  standard error of the mean ( $n = 5$ ). Different letters indicate significant differences between the treatments assessed by post hoc Tukey tests.

Treatment	Concentration (mmol l <sup>-1</sup> )	$F_{rap}$ (%)	$k_{rap}$ (h <sup>-1</sup> )	$F_{slow}$ (%)	$k_{slow}$ (h <sup>-1</sup> )	$F_{very\ slow}$ (%)	$k_{very\ slow}$ (h <sup>-1</sup> )
1 week ageing							
Control	0	<sup>de</sup> 12.078 $\pm$ 0.418	<sup>ab</sup> 0.210 $\pm$ 0.008	<sup>e</sup> 15.801 $\pm$ 0.850	<sup>a</sup> 0.013 $\pm$ 0.001	<sup>b</sup> 72.122 $\pm$ 1.251	< <sup>a</sup> 0.001 <0.001
Citric acid	0.5	<sup>bcd</sup> 10.164 $\pm$ 0.564	<sup>b</sup> 0.226 $\pm$ 0.014	<sup>cd</sup> 9.162 $\pm$ 0.224	<sup>ab</sup> 0.021 $\pm$ 0.004	<sup>d</sup> 80.674 $\pm$ 0.708	< <sup>ab</sup> 0.001 <0.001
	100	<sup>a</sup> 7.099 $\pm$ 0.457	<sup>c</sup> 0.319 $\pm$ 0.037	<sup>b</sup> 7.136 $\pm$ 0.418	<sup>c</sup> 0.042 $\pm$ 0.003	<sup>efg</sup> 85.765 $\pm$ 0.224	< <sup>abc</sup> 0.001 <0.001
	250	<sup>b</sup> 8.984 $\pm$ 0.423	<sup>b</sup> 0.242 $\pm$ 0.013	<sup>bc</sup> 7.992 $\pm$ 0.219	<sup>bc</sup> 0.035 $\pm$ 0.003	<sup>de</sup> 83.025 $\pm$ 0.254	< <sup>abc</sup> 0.001 <0.001
	500	<sup>cde</sup> 11.822 $\pm$ 0.656	<sup>bc</sup> 0.271 $\pm$ 0.029	<sup>d</sup> 10.589 $\pm$ 0.536	<sup>c</sup> 0.036 $\pm$ 0.004	<sup>c</sup> 77.589 $\pm$ 0.563	< <sup>cde</sup> 0.001 <0.001
	1000	<sup>f</sup> 19.220 $\pm$ 0.344	<sup>b</sup> 0.234 $\pm$ 0.007	<sup>e</sup> 16.563 $\pm$ 0.614	<sup>abc</sup> 0.027 $\pm$ 0.005	<sup>a</sup> 64.217 $\pm$ 0.657	< <sup>cde</sup> 0.001 <0.001
Malic acid	0.5	<sup>cde</sup> 11.379 $\pm$ 0.524	<sup>a</sup> 0.139 $\pm$ 0.002	<sup>a</sup> 3.686 $\pm$ 0.473	<sup>d</sup> 0.139 $\pm$ 0.002	<sup>ef</sup> 84.935 $\pm$ 0.086	< <sup>de</sup> 0.001 <0.001
	100	<sup>bc</sup> 9.614 $\pm$ 0.136	<sup>a</sup> 0.142 $\pm$ 0.002	<sup>a</sup> 2.288 $\pm$ 0.008	<sup>d</sup> 0.142 $\pm$ 0.002	<sup>g</sup> 88.098 $\pm$ 0.143	< <sup>cde</sup> 0.001 <0.001
	250	<sup>bcd</sup> 11.074 $\pm$ 0.115	<sup>a</sup> 0.146 $\pm$ 0.002	<sup>a</sup> 2.373 $\pm$ 0.005	<sup>d</sup> 0.146 $\pm$ 0.002	<sup>fg</sup> 86.553 $\pm$ 0.119	< <sup>de</sup> 0.001 <0.001
	500	<sup>e</sup> 13.632 $\pm$ 0.260	<sup>a</sup> 0.141 $\pm$ 0.003	<sup>a</sup> 2.491 $\pm$ 0.012	<sup>d</sup> 0.141 $\pm$ 0.003	<sup>ef</sup> 83.877 $\pm$ 0.272	< <sup>e</sup> 0.001 <0.001
	1000	<sup>f</sup> 20.196 $\pm$ 0.914	<sup>a</sup> 0.140 $\pm$ 0.003	<sup>a</sup> 2.705 $\pm$ 0.022	<sup>d</sup> 0.140 $\pm$ 0.003	<sup>c</sup> 77.100 $\pm$ 0.936	< <sup>f</sup> 0.001 <0.001
15 weeks ageing							
Control	0	<sup>bcd</sup> 13.115 $\pm$ 0.712	<sup>b</sup> 0.172 $\pm$ 0.010	<sup>d</sup> 83.646 $\pm$ 3.300	< <sup>a</sup> 0.001 <0.001	<sup>a</sup> 3.239 $\pm$ 3.234	< <sup>a</sup> 0.001 <0.001
Citric acid	0.5	<sup>a</sup> 6.389 $\pm$ 0.180	<sup>bc</sup> 0.180 $\pm$ 0.019	<sup>c</sup> 14.050 $\pm$ 4.178	<sup>bc</sup> 0.005 $\pm$ 0.001	<sup>b</sup> 79.561 $\pm$ 4.036	< <sup>ab</sup> 0.001 <0.001
	100	<sup>bc</sup> 12.466 $\pm$ 0.512	<sup>cd</sup> 0.298 $\pm$ 0.023	<sup>e</sup> 16.386 $\pm$ 0.491	<sup>cd</sup> 0.010 $\pm$ 0.001	<sup>b</sup> 71.149 $\pm$ 0.609	< <sup>b</sup> 0.001 <0.001
	250	<sup>bc</sup> 11.753 $\pm$ 1.008	<sup>ef</sup> 0.511 $\pm$ 0.053	<sup>c</sup> 18.569 $\pm$ 1.781	<sup>de</sup> 0.027 $\pm$ 0.007	<sup>b</sup> 69.678 $\pm$ 2.345	< <sup>ab</sup> 0.001 <0.001
	500	<sup>ef</sup> 17.595 $\pm$ 1.768	<sup>de</sup> 0.521 $\pm$ 0.126	<sup>e</sup> 19.782 $\pm$ 1.110	<sup>d</sup> 0.020 $\pm$ 0.007	<sup>b</sup> 62.623 $\pm$ 2.536	< <sup>ab</sup> 0.001 <0.001
	1000	<sup>g</sup> 25.120 $\pm$ 1.153	<sup>de</sup> 0.398 $\pm$ 0.046	<sup>e</sup> 15.909 $\pm$ 2.951	<sup>de</sup> 0.026 $\pm$ 0.005	<sup>b</sup> 58.971 $\pm$ 2.787	<sup>ab</sup> 0.001 <0.001
Malic acid	0.5	<sup>a</sup> 5.504 $\pm$ 0.526	<sup>bc</sup> 0.190 $\pm$ 0.021	<sup>b</sup> 1.994 $\pm$ 0.305	<sup>f</sup> 0.190 $\pm$ 0.021	<sup>b</sup> 92.502 $\pm$ 0.324	< <sup>a</sup> 0.001 <0.001
	100	<sup>cde</sup> 14.762 $\pm$ 0.759	<sup>a</sup> 0.090 $\pm$ 0.012	<sup>a</sup> 1.057 $\pm$ 0.006	<sup>f</sup> 0.090 $\pm$ 0.012	<sup>b</sup> 84.181 $\pm$ 0.765	< <sup>a</sup> 0.001 <0.001
	250	<sup>def</sup> 16.916 $\pm$ 0.492	<sup>b</sup> 0.150 $\pm$ 0.010	<sup>d</sup> 58.727 $\pm$ 1.719	<sup>a</sup> 0.001 $\pm$ 0.000	<sup>b</sup> 24.357 $\pm$ 1.776	< <sup>a</sup> 0.001 <0.001
	500	<sup>f</sup> 20.465 $\pm$ 0.659	<sup>bc</sup> 0.210 $\pm$ 0.007	<sup>d</sup> 58.849 $\pm$ 1.998	<sup>ab</sup> 0.001 <0.001	<sup>ab</sup> 20.686 $\pm$ 1.732	<sup>a</sup> 0.001 <0.001
	1000	<sup>ab</sup> 9.066 $\pm$ 0.771	<sup>f</sup> 0.830 $\pm$ 0.032	<sup>e</sup> 18.118 $\pm$ 1.007	<sup>ef</sup> 0.061 $\pm$ 0.002	<sup>ef</sup> 72.816 $\pm$ 1.353	<sup>a</sup> 0.001 <0.001





## **Supplementary information**

### **The effect of organic acids on the fate of $^{14}\text{C}$ -phenanthrene in contaminated soil**

Gabriela M. Vázquez-Cuevas <sup>a,c</sup>, Alfonso J. Lag-Brotons <sup>a</sup>, Jose J. Ortega-Calvo <sup>b</sup>,  
Carly J. Stevens <sup>a</sup> and Kirk T. Semple <sup>a\*</sup>

<sup>a</sup>Lancaster Environment Centre, Lancaster University, Lancaster, LA1 4YQ, United Kingdom

<sup>b</sup>Instituto de Recursos Naturales y Agrobiología, C.S.I.C., Apartado 1052, 41080 Seville, Spain

<sup>c</sup>Área Académica de Biología, Instituto de Ciencias Básicas e Ingeniería. Universidad Autónoma del Estado de Hidalgo, 42010, Mexico

\*Corresponding author: Phone no. +44 (0)1524 510554; email:  
k.semple@lancaster.ac.uk

## Supplementary information

Table SI 1. Desorption kinetics of <sup>14</sup>C-phenanthrene from mildly aged soil (50 d). Values represent the mean ± standard error of the mean (*n* = 3). Different letters indicate significant differences between the treatments assessed by post hoc Tukey tests

Two-compartment fitting											
Treatment	Concentration (mmol l <sup>-1</sup> )	<i>F<sub>rap</sub></i> (%)	<i>k<sub>rap</sub></i> (h <sup>-1</sup> )	<i>F<sub>slow</sub></i> (%)	<i>k<sub>slow</sub></i> (h <sup>-1</sup> )						
Control	0	<sup>ab</sup> 40.939 ± 3.861	<sup>a</sup> 0.090 ± 0.019	<sup>ab</sup> 59.061 ± 3.861	<sup>a</sup> 0.001 < 0.001						
Citric acid	0.1	<sup>a</sup> 25.905 ± 6.193	<sup>a</sup> 0.133 ± 0.020	<sup>b</sup> 74.095 ± 6.193	<sup>a</sup> 0.001 < 0.001						
	0.5	<sup>ab</sup> 44.164 ± 7.875	<sup>a</sup> 0.119 ± 0.010	<sup>ab</sup> 55.836 ± 7.875	<sup>a</sup> 0.001 < 0.001						
Malic acid	0.1	<sup>a</sup> 25.356 ± 3.052	<sup>a</sup> 0.179 ± 0.007	<sup>b</sup> 74.644 ± 3.052	<sup>a</sup> 0.001 < 0.001						
	0.5	<sup>b</sup> 49.763 ± 3.847	<sup>a</sup> 0.110 ± 0.015	<sup>a</sup> 50.237 ± 3.847	<sup>a</sup> 0.001 < 0.001						
Oxalic acid	0.1	<sup>ab</sup> 37.096 ± 5.009	<sup>a</sup> 0.107 ± 0.017	<sup>ab</sup> 62.904 ± 5.009	<sup>a</sup> 0.001 < 0.001						
	0.5	<sup>ab</sup> 34.857 ± 1.603	<sup>a</sup> 0.151 ± 0.027	<sup>ab</sup> 65.143 ± 1.603	<sup>a</sup> 0.001 ± 0.001						
Succinic acid	0.1	<sup>ab</sup> 30.451 ± 2.869	<sup>a</sup> 0.115 ± 0.036	<sup>ab</sup> 69.549 ± 2.869	<sup>a</sup> 0.000 < 0.001						
	0.5	<sup>ab</sup> 28.999 ± 1.983	<sup>a</sup> 0.123 ± 0.007	<sup>ab</sup> 71.001 ± 1.983	<sup>a</sup> 0.001 < 0.001						
Three-compartment fitting											
Treatment	Concentration (mmol l <sup>-1</sup> )	<i>F<sub>rap</sub></i> (%)	<i>k<sub>rap</sub></i> (h <sup>-1</sup> )	<i>F<sub>slow</sub></i> (%)	<i>k<sub>slow</sub></i> (h <sup>-1</sup> )	<i>F<sub>very slow</sub></i> (%)	<i>k<sub>very slow</sub></i> (h <sup>-1</sup> )				
Control	0	<sup>ab</sup> 36.112 ± 5.067	<sup>ab</sup> 0.138 ± 0.023	<sup>a</sup> 56.511 ± 2.855	<sup>ab</sup> 0.003 < 0.001	<sup>a</sup> 7.377 ± 2.458	<sup>a</sup> 0.003 < 0.001				
Citric acid	0.1	<sup>a</sup> 22.920 ± 6.498	<sup>ab</sup> 0.168 ± 0.035	<sup>a</sup> 51.399 ± 18.98	<sup>ab</sup> 0.003 ± 0.001	<sup>a</sup> 25.682 ± 13.80	<sup>a</sup> 0.002 ± 0.001				
	0.5	<sup>b</sup> 56.929 ± 9.031	<sup>a</sup> 0.130 ± 0.083	<sup>a</sup> 30.697 ± 10.45	<sup>ab</sup> 0.007 ± 0.005	<sup>a</sup> 12.374 ± 12.37	<sup>a</sup> 0.001 ± 0.001				
Malic acid	0.1	<sup>a</sup> 23.902 ± 2.529	<sup>ab</sup> 0.289 ± 0.045	<sup>a</sup> 61.384 ± 9.804	<sup>ab</sup> 0.002 ± 0.001	<sup>a</sup> 14.714 ± 8.010	<sup>a</sup> 0.001 ± 0.001				
	0.5	<sup>ab</sup> 43.770 ± 2.514	<sup>ab</sup> 0.193 ± 0.035	<sup>a</sup> 46.991 ± 8.865	<sup>ab</sup> 0.005 ± 0.003	<sup>a</sup> 9.240 ± 6.792	<sup>a</sup> 0.002 ± 0.001				

Oxalic acid	0.1	<sup>ab</sup> 31.645 ± 7.967	<sup>ab</sup> 0.321 ± 0.179	<sup>a</sup> 44.376 ± 9.240	<sup>ab</sup> 0.007 ± 0.005	<sup>a</sup> 23.979 ± 15.78	<sup>a</sup> 0.001 < 0.001
	0.5	<sup>ab</sup> 29.134 ± 1.809	<sup>ab</sup> 0.375 ± 0.082	<sup>a</sup> 38.480 ± 9.898	<sup>ab</sup> 0.007 ± 0.003	<sup>a</sup> 32.386 ± 10.93	<sup>a</sup> 0.001 ± 0.001
Succinic acid	0.1	<sup>ab</sup> 29.880 ± 2.748	<sup>a</sup> 0.147 ± 0.058	<sup>a</sup> 64.678 ± 5.535	<sup>a</sup> 0.001 < 0.001	<sup>a</sup> 5.443 ± 4.019	<sup>a</sup> 0.001 ± 0.001
	0.5	<sup>a</sup> 19.530 ± 0.664	<sup>b</sup> 0.577 ± 0.106	<sup>a</sup> 24.270 ± 5.003	<sup>b</sup> 0.016 ± 0.006	<sup>a</sup> 56.200 ± 5.500	<sup>a</sup> < 0.001 < 0.001

Table SI 2. Proportion of  $^{14}\text{C}$ -phenanthrene (1) extracted with 50 mM HPCD solution, (2) mineralisation rate ( $\%, \text{h}^{-1}$ ) within the assessed contact time. Values represent the mean  $\pm$  standard error of the mean ( $n = 3$ )

Contact time (h)	Treatment	Extracted (%) <sup>1</sup>	Mineralised (%) <sup>2</sup>
1	Control	7.269 $\pm$ 0.993	1.454 $\pm$ 0.199
	0.5 mmol l <sup>-1</sup>	5.777 $\pm$ 0.767	1.155 $\pm$ 0.153
	500 mmol l <sup>-1</sup>	7.854 $\pm$ 0.257	1.571 $\pm$ 0.051
3	Control	6.062 $\pm$ 0.943	0.505 $\pm$ 0.236
	0.5 mmol l <sup>-1</sup>	6.101 $\pm$ 0.358	0.508 $\pm$ 0.089
	500 mmol l <sup>-1</sup>	7.268 $\pm$ 1.315	0.605 $\pm$ 0.329
6	Control	5.843 $\pm$ 1.035	0.243 $\pm$ 0.259
	0.5 mmol l <sup>-1</sup>	6.264 $\pm$ 0.954	0.261 $\pm$ 0.238
	500 mmol l <sup>-1</sup>	5.528 $\pm$ 0.818	0.230 $\pm$ 0.205
8	Control	8.146 $\pm$ 1.876	0.254 $\pm$ 0.469
	0.5 mmol l <sup>-1</sup>	5.978 $\pm$ 1.104	0.186 $\pm$ 0.276
	500 mmol l <sup>-1</sup>	8.495 $\pm$ 1.182	0.265 $\pm$ 0.296
24	Control	5.250 $\pm$ 3.213	0.054 $\pm$ 0.803
	0.5 mmol l <sup>-1</sup>	3.805 $\pm$ 1.050	0.039 $\pm$ 0.263
	500 mmol l <sup>-1</sup>	6.282 $\pm$ 1.327	0.065 $\pm$ 0.332
48	Control	1.102 $\pm$ 0.108	0.091 $\pm$ 0.432
	0.5 mmol l <sup>-1</sup>	0.929 $\pm$ 0.169	0.077 $\pm$ 0.676
	500 mmol l <sup>-1</sup>	1.257 $\pm$ 0.157	0.104 $\pm$ 0.628

Table SI 3. Sums of squared deviations of desorbed  $^{14}\text{C}$ -phenanthrene fitted to a two- and three-compartment model. Values represent the mean  $\pm$  standard error of the mean ( $n = 5$ )

Treatment	Concentration (mmol l <sup>-1</sup> )	Sum of squared difference 2 compartment fitting	Sum of squared difference 3-compartment fitting
1 week ageing			
Control	0	9.05E-04 $\pm$ 1.0E-04	2.24E-04 $\pm$ 2.8E-05
Citric acid	0.5	4.79E-04 $\pm$ 3.6E-05	2.53E-04 $\pm$ 7.8E-05
	100	3.97E-04 $\pm$ 2.6E-05	1.39E-04 $\pm$ 1.3E-05
	250	4.19E-04 $\pm$ 2.4E-05	1.28E-04 $\pm$ 1.1E-05
	500	8.97E-04 $\pm$ 1.0E-04	2.73E-04 $\pm$ 5.1E-05
	1000	2.40E-03 $\pm$ 2.8E-04	7.09E-04 $\pm$ 9.4E-05
Malic acid	0.5	5.44E-04 $\pm$ 4.0E-05	5.44E-04 $\pm$ 4.0E-05
	100	3.13E-04 $\pm$ 1.3E-05	3.13E-04 $\pm$ 1.3E-05
	250	4.98E-04 $\pm$ 2.0E-05	4.98E-04 $\pm$ 2.0E-05
	500	6.71E-04 $\pm$ 1.4E-05	6.71E-04 $\pm$ 1.4E-05
	1000	1.49E-03 $\pm$ 1.7E-04	1.49E-03 $\pm$ 1.7E-04
15 weeks ageing			
Control	0	4.50E-04 $\pm$ 6.0E-05	4.60E-04 $\pm$ 6.0E-05
Citric acid	0.5	9.59E-05 $\pm$ 4.6E-05	6.00E-05 $\pm$ 4.0E-05
	100	1.17E-03 $\pm$ 2.0E-04	2.20E-04 $\pm$ 9.3E-05
	250	2.50E-03 $\pm$ 3.9E-04	1.48E-04 $\pm$ 5.0E-05
	500	3.59E-03 $\pm$ 4.5E-04	9.20E-04 $\pm$ 1.9E-04
	1000	2.99E-03 $\pm$ 7.8E-04	3.70E-04 $\pm$ 4.8E-05
Malic acid	0.5	3.92E-04 $\pm$ 6.4E-05	4.20E-04 $\pm$ 6.6E-05
	100	4.63E-04 $\pm$ 1.3E-04	4.63E-04 $\pm$ 1.3E-04
	250	1.98E-03 $\pm$ 1.2E-04	1.28E-03 $\pm$ 4.1E-04
	500	1.75E-03 $\pm$ 1.0E-04	1.58E-03 $\pm$ 2.3E-04
	1000	3.70E-03 $\pm$ 4.7E-04	2.80E-04 $\pm$ 6.8E-05




Figure SI 4. Rapid desorbing fractions ( $F_{rap}$ ) of <sup>14</sup>C-phenanthrene from soil aged for 1 (A) and 15 (B) weeks extracted with citric (●) and malic (○) acid. Values were obtained using a three-compartment model fitting. Error bars represent the standard error of the mean ( $n = 5$ ).




Figure SI 5. Rate constants of the rapid desorbing fractions ( $k_{rap}$ ) of <sup>14</sup>C-phenanthrene from soil aged for 1 (A) and 15 (B) weeks extracted with citric (●) and malic (○) acid. Values were obtained using a three-compartment model fitting. Error bars represent the standard error of the mean ( $n = 5$ ).




Figure SI 6. Slow desorbing fractions ( $F_{slow}$ ) of  $^{14}\text{C}$ -phenanthrene from soil aged for 1 (A) and 15 (B) weeks extracted with citric (●) and malic (○) acid. Values were obtained using a three-compartment model fitting. Error bars represent the standard error of the mean ( $n = 5$ ).




Figure SI 7. Rate constants of the slow desorbing fractions ( $k_{slow}$ ) of  $^{14}\text{C}$ -phenanthrene from soil aged for 1 (A) and 15 (B) weeks extracted with citric (●) and malic (○) acid. Values were obtained using a three-compartment model fitting. Error bars represent the standard error of the mean ( $n = 5$ ).






Figure SI 8. Very slow desorbing fractions ( $F_{very\ slow}$ ) of  $^{14}\text{C}$ -phenanthrene from soil aged for 1 (A) and 15 (B) weeks extracted with citric (●) and malic (○) acid. Values were obtained using a three-compartment model fitting. Error bars represent the standard error of the mean ( $n = 5$ ).




Figure SI 9. Rate constants of the very slow desorbing fractions ( $k_{very\ slow}$ ) of  $^{14}\text{C}$ -phenanthrene from soil aged for 1 (A) and 15 (B) weeks extracted with citric (●) and malic (○) acid. Values were obtained using a three-compartment model fitting. Error bars represent the standard error of the mean ( $n = 5$ ).