


Organic & Biomolecular Chemistry

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: M. J. Young, G. Akien and N. H. Evans, *Org. Biomol. Chem.*, 2020, DOI: 10.1039/D0OB01190H.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

ARTICLE

An amide hydrogen bond templated [1]rotaxane displaying a peptide motif – demonstrating an expedient route to synthetic mimics of lasso peptides†

Matthew J. Young,‡^a Geoffrey R. Akien^a and Nicholas H. Evans*^aReceived 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

The rapid synthesis of an amide hydrogen bond templated [1]rotaxane is reported – demonstrating a potential pathway to synthetic analogues of lasso peptides. The structures of the [1]rotaxane and its unthreaded isomer have been characterized by NMR spectroscopy and modelled using DFT calculations.

Introduction

Lasso peptides are a fascinating class of structurally unusual peptides consisting of a polypeptide chain that threads its “tail” through a macrolactam “ring” (Figure 1).¹ Due to their entangled structure, lasso peptides have been found to evade degradation by enzymes, and examples have been shown to possess a number of useful bioactive characteristics, including antimicrobial activity,² receptor antagonism³ and enzyme inhibition.⁴ The total synthesis of lasso peptides is in its infancy, with the first chemical synthesis of a lasso peptide having only very recently been reported.⁵ However, it has been established that the conformationally constrained amino acids displayed in the “loop region” (see Figure 1) are important in determining the biological activity of a lasso peptide.⁶ Synthetic [1]rotaxanes⁷ are structurally equivalent to lasso peptides, but examples that display peptide sequences are rare.^{8,9}




Figure 1: Schematic representation of structure of a lasso peptide and [1]rotaxane.

Recently we have rapidly prepared [2]rotaxanes^{10–12} by use of hydrogen bond templation¹³ and CuAAC click chemistry to stopper a simple amide half-axle component threaded through an isophthalamide macrocycle.¹⁴ Here we are reporting an

adaptation of this method to rapidly prepare a [1]rotaxane displaying a Gly-Gly-Gly-Gly peptide sequence within the key loop region, by appending and then cyclizing a peptide chain on a novel “scaffold” [2]rotaxane. Successful preparation of the target [1]rotaxane was confirmed by NMR spectroscopy and mass spectrometry. Structural comparison with the non-interlocked isomer of the [1]rotaxane has also been undertaken by NMR spectroscopy and DFT calculations.

When compared to prior work from the laboratories of Coutrot⁸ and Bode⁹, the synthetic route reported here has the potential to generate better mimics of naturally occurring lasso peptides. In the previous cases, the entangled structure arises from the non-amide templation of a protonated ammonium passing through a 24-crown-8 macrocycle. Here, amide motifs (used to template formation of the interlocked structure through the carbonyl O atom of the axle to the isophthalamide N-Hs of the macrocycle) are to be found in both the axle and ring components – not just the peptide sequence in the loop region.

Results and discussion


The synthesis of [1]rotaxane **6** is presented in Scheme 1. All novel compounds were characterized by NMR spectroscopy (¹H, ¹³C and where applicable ¹⁹F), IR spectroscopy and high resolution mass spectrometry. Scaffold [2]rotaxane **4** was prepared using azide **1**, alkyne **2** and amino macrocycle **3**. Azide **1** was synthesized according to our previously reported procedures in two chromatography-free steps.¹⁰ Novel alkyne **2** was prepared in four chromatography-free steps (see ESI†, page S3), with inclusion of a glycine unit as part of **2** to guarantee sufficient bulk for the rotaxane stopper. Novel amino macrocycle **3** was synthesized in four steps from commercially available starting materials (see ESI†, pages S6 & S9).§

^a Department of Chemistry, Lancaster University, Lancaster, LA1 4YB. UK.

† Electronic Supplementary Information (ESI) available: Further experimental procedures; copies of spectral characterization data; crystallographic data for macrocycle **3** (CCDC 1995143); details of computational modelling. See DOI: 10.1039/x0xx00000x.

‡ Current address: Institute of Organic Chemistry, Albert-Ludwigs-University of Freiburg, Albertstrasse 21, 79104 Freiburg, Germany.





Scheme 1: Synthesis of tetra-glycine [1]rotaxane **6** via "scaffold" [2]rotaxane **4**, and structures of axle **7** and unthreaded "[1]rotaxane" isomer **8**.

[2]Rotaxane **4** was synthesized using our recently reported hydrogen bond templated strategy.¹⁰⁻¹² Macrocyclic **3** was dissolved in CH_2Cl_2 in the presence of 1.1 equivalents of azide **1** and 1.1 equivalents of alkyne **2**. Catalytic $[\text{Cu}(\text{CH}_3\text{CN})_4]\text{BF}_4$ and TBTA and 1.2 equivalents of DIPEA were added and the reaction stirred overnight at room temperature under an inert atmosphere. After aqueous work-up and careful silica chromatography, [2]rotaxane **4** was isolated in 27% yield.¹⁵

The ^1H NMR spectra of [2]rotaxane **4**, along with that of non-interlocked macrocycle **3** and axle **7** (for synthesis of axle **7** see

ESI⁺, page S30) are shown in Figure 2. The upfield shift and splitting of aromatic macrocycle protons **8** and **9** and the upfield shifts of axle protons **17**, **20**, **21**, **22**, **23** and **25** in [2]rotaxane **4** compared to macrocycle **3** and axle **7** are consistent with the threading of the axle component through the macrocycle. The downfield shift of internal isophthalamide proton **4** is indicative of hydrogen bonding to the carbonyl oxygen of the axle amide. Evidence to support the co-conformation of [2]rotaxane **4** depicted in Scheme 1 is provided by the appearance of through-space correlations in the ^1H ROESY NMR spectrum between



protons 8 and 9 of the macrocycle and axle proton environments 17, 20 and 21 (see ESI[†], page S16). For [2]rotaxane **4**, the resonances of macrocyclic protons 6 and 11 are split due to the directionality of the threaded axle that causes the protons attached to the same carbon to become inequivalent. In addition, the molecular ion peak [M + Na]⁺ for [2]rotaxane **4** was observed in the positive-ion electrospray mass spectrum at $m/z = 1209$ Da (see ESI[†], page S17).




Figure 2: ¹H NMR spectra of (a) macrocycle **3**, (b) [2]rotaxane **4** and (c) axle **7** (1:1 CDCl₃/CD₃OD, 400 MHz, 298 K). For atom labels see Scheme 1.

With [2]rotaxane **4** in-hand, preparation of the target [1]rotaxane was executed. Boc-Gly-Gly-Gly was appended to the macrocyclic amine group of [2]rotaxane **4** by use of EDC coupling in 80% yield (using three equivalents of the protected tripeptide). Treatment of Boc-peptide appended [2]rotaxane **5** with TFA in CH₂Cl₂ simultaneously removed both the Boc and *tert*-butyl ester. Cyclization was carried out using HATU¹⁶ at a dilute concentration of 2 mmol/L of deprotected **5** in CH₂Cl₂ to minimize the effect of competing oligomerization reactions, with [1]rotaxane **6** being isolated in 34% yield (over two steps from **5**) after aqueous work-up and silica gel column chromatography. Confirmation of the successful isolation of target [1]rotaxane **6** was ascertained by NMR spectroscopy and mass spectrometry – a molecular ion peak [M + Na]⁺ for **6** being observed in the positive-ion electrospray mass spectrum at $m/z = 1306$ Da (see ESI[†], page S24).

To study the structural effects of [1]rotaxane **6** being interlocked, the unthreaded analogue **8** was also prepared (for

details of synthesis see ESI[†], page S33). Inspection of the ¹H NMR spectra (recorded in D₆-DMSO due to the very limited solubility of **8**, Figure 3) reveals the anticipated shielding of protons 8, 9, 17, 20, 21, 22 and 25 (but interestingly not 23) and deshielding of proton 4 for [1]rotaxane **6** compared to unthreaded **8**. In addition, the axle amide proton *c* is notably upfield in **6** compared to **8**, consistent with being present within the shielding macrocyclic cavity of [1]rotaxane **6**. There is also the expected marked splitting of macrocyclic resonances 6 and 11 for [1]rotaxane **6** arising from the directionality of the threaded axle. Other notable differences in chemical shift between the two molecules are for amide proton *a* (significantly greater than for any of the glycine amide protons *d-g*) and aromatic axle proton 27, tentatively attributed to differences in hydrogen bonding involving these protons (either intramolecularly and/or with the strongly hydrogen bond accepting DMSO solvent).




Figure 3: ¹H NMR spectra of (a) [1]rotaxane **6** and (b) unthreaded isomer **8** (D₆-DMSO, 400 MHz, 298 K). For atom labels see Scheme 1.

NMR experiments at elevated temperatures provided no evidence of dethreading of **6** or threading of **8**. However, they were used to study variation in the chemical shift of NH resonances as a function of temperature, as has been used in the structural characterization of proteins in aqueous solution (see ESI[†], pages S45-S47).¹⁷ In aqueous solution, temperature coefficients $\Delta\delta/\Delta T > -4.5$ ppb/K are attributed to intramolecular hydrogen bonds, while those that are more negative (< -4.5 ppb/K) are deemed to be freely interacting with solvent. The limited literature data for rigid peptides in D₆-DMSO is consistent with all coefficients being slightly less negative on average.¹⁸




View Article Online
DOI: 10.1039/D0OB01190H

Figure 4: Minimum energy DFT calculated structures of (a) [1]rotaxane **6** and (b) unthreaded isomer **8** (C-H hydrogen atoms omitted for clarity).

For unthreaded **8**, all amide temperature coefficients are in the range -3.3 to -4.5 ppb/K, which is consistent with relatively unrestricted solvent accessibility. For [1]rotaxane **6**, the bulk of the NH environments have similar coefficients (in the range of -3.0 to -4.3 ppb/K) but there are two key exceptions. Axle amide *c* has a coefficient of -1.1 ppb/K implying it is inaccessible to solvent and macrocycle protons *b* have an unusually negative coefficient of -6.7 ppb/K. We attribute these deviations to the various non-covalent interactions arising from the axle component threading through the macrocycle cavity.

Further structural comparison between **6** and **8** was possible from ^1H diffusion NMR experiments (see ESI[†], pages S48 & S49).¹⁹ For [1]rotaxane **6** the diffusion coefficient $D = 1.28 \pm 0.03 \times 10^{-10} \text{ m}^2 \text{ s}^{-1}$, whereas for unthreaded analogue **8** $D = 1.10 \pm 0.03 \times 10^{-10} \text{ m}^2 \text{ s}^{-1}$. In other words, the threaded [1]rotaxane diffuses faster which is consistent with a more compact 3D structure in solution. The observed D for both **6** and **8** is consistent with monomeric forms in solution,²⁰ and is also in agreement with a comparison of the DFT-simulated (B3LYP/6-31G*) structures of [1]rotaxane **6** and its unthreaded isomer **8** (Figure 4).²¹

Conclusions

An expedient route to a peptide containing amide hydrogen bond templated [1]rotaxane has been demonstrated by the synthesis and structural characterization of a tetra-glycine containing [1]rotaxane. Looking forward, substitution of Boc-Gly-Gly-Gly by other aliphatic side-chained peptide sequences should be straightforward, and in principle with modifications to protecting group strategy more exotic amino acid side-chains (or non-peptide motifs) could be incorporated. Hence delicate bioactive functionality could be displayed in the loop region, with protection against enzymatic degradation⁹ and/or conformational control being provided by the entangled [1]rotaxane structure. Work on some of these avenues of research, as well as developing alternative synthetic routes to novel [1]rotaxanes, are ongoing in our laboratories.

Experimental

General information

Commercially available solvents and chemicals were used without further purification unless stated. Dry solvents, NEt_3 and DIPEA were purchased dry and stored under an inert atmosphere. $\text{Cu}(\text{CH}_3\text{CN})_4\text{BF}_4$ was stored in desiccator over P_4O_{10} . Deionised water was used in all cases.

Silica gel with a 60\AA particle size was used as the stationary phase for column chromatography. Analytical TLC was used to monitor the progress of column chromatography and analytical TLC plates were examined under short wavelength ($\lambda = 254 \text{ nm}$) UV light. Preparatory TLC was carried out on silica gel possessing a fluorescent indicator to allow for examination with short wavelength UV light.


IR spectra of neat samples were recorded on an Agilent Technologies Cary 630 FTIR spectrometer. NMR spectra of samples dissolved in deuterated solvent were recorded on a Bruker AVANCE III 400 spectrometer at 298 K. 1D NMR spectra were referenced using residual solvent peaks, and assigned using standard ^1H - ^1H COSY, ^1H - ^{13}C HSQC and ^1H - ^{13}C HMBC NMR experiments, with data being reported according to the atom labels as defined in the structures below. Electrospray mass spectra of dissolved samples, diluted in CH_3CN or CH_3OH , were recorded on a Shimadzu LCMS IT ToF instrument. Melting points were recorded on a Gallenkamp capillary melting point apparatus and are uncorrected.

Experimental procedures

[2]Rotaxane 4. Amino macrocycle **3** (107 mg, 0.219 mmol) and azide **1** (82 mg, 0.24 mmol) were dissolved in dry CH_2Cl_2 (10 mL) under an Ar (g) atmosphere. Then alkyne stopper **2** (86 mg, 0.24 mmol) in dry CH_2Cl_2 (1 mL), $\text{Cu}(\text{CH}_3\text{CN})_4\text{BF}_4$ (7.6 mg, 0.024 mmol), TBTA (13 mg, 0.024 mmol) and DIPEA (34 mg, 46 μL , 0.26 mmol) were added. The reaction was stirred at r.t. for 15 h under an Ar (g) atmosphere. Then, the reaction was diluted to 20 mL, washed with 0.02 M EDTA in 1M NH_3 (aq) solution ($2 \times 20 \text{ mL}$) and brine ($1 \times 20 \text{ mL}$). The organic layer was dried (MgSO_4) and solvent removed *in vacuo*. The crude material was purified by silica gel column chromatography



(CH₂Cl₂/CH₃OH 99:1 to 96:4) to yield pure product **4** as a colourless film (52 mg). In addition, axle **7** (102 mg) was also isolated. Further pure product **4** was isolated after prep TLC of mixed fractions (17 mg). Total yield of [2]rotaxane **4**: 69 mg, 27%.



R_f: 0.20 (CH₂Cl₂/CH₃OH 98:2).

v_{max}/cm⁻¹ (neat): 3350 (N–H), 2920 (C–H), 1740 (C=O ester), 1650 (C=O amide), 1600, 1520, 1450, 1360, 1280, 1230, 1130, 1030.

δ_H (400 MHz; CDCl₃/CD₃OD 1:1): 8.18 (2H, s, C¹⁷H), 8.05 (1H, s, C¹⁴H), 7.75–7.76 (2H, m, C²³H & C²⁹H), 7.67 (1H, t, ⁴J = 1.5 Hz, C⁴H), 7.64 (1H, s, C²⁷H), 7.43 (1H, t, ³J = 5.4 Hz, C²⁰NH), 7.36 (2H, d, ⁴J = 1.5 Hz, C²H), 7.32 (1H, s, C³²H), 6.88 (4H, d, ³J = 7.9 Hz, C⁸H), 6.71 (4H, d, ³J = 7.9 Hz, C⁹H), 5.02 (2H, s, C²⁵H), 4.49 (2H, d, ²J = 14 Hz, C⁶H - diastereotopic) 4.25–4.30 (4H, m, C⁶H - diastereotopic & C¹¹H - diastereotopic) 4.05–4.08 (4H, m, C¹¹H - diastereotopic & C³⁴H), 3.99 (2H, t, ³J = 7.0 Hz, C²²H), 3.54–3.72 (8H, m, C¹³H & C¹²H), 2.67 (2H, br s, C²⁰H), 1.66–1.73 (2H, m, C²¹H), 1.47 (9H, s, C³⁷H).


δ_C (100 MHz; CDCl₃/CD₃OD 1:1): 170.0 (C³⁵), 168.7 (C⁵), 167.9 (C³³), 165.0 (C¹⁹), 159.4 (C²⁶), 149.1 (C¹), 143.1 (C²⁴), 138.1 (C⁷), 137.1 (C¹⁰), 137.1 (sic, C²⁸), 136.7 (C¹⁸), 136.0 (C³), 132.8 (q, ²J = 32 Hz, C³⁰), 132.3 (q, ²J = 34 Hz, C¹⁵), 129.6 (C⁹), 129.1 (C¹⁷), 129.0 (C⁸), 125.7 (C²³), 125.3 (br, C¹⁴), 124.0 (q, ¹J = 271 Hz, C¹⁶), 118.0 (C²), 117.6 (br, C²⁹), 117.3 (C²⁷) 115.9 (br, C³²), 114.2 (C⁴), 82.9 (C³⁶), 74.3 (C¹¹), 71.6 (C¹³), 70.4 (C¹²), 62.4 (C²⁵), 48.7 (C²²), 44.8 (C⁶), 43.1 (C³⁴), 38.1 (C²⁰), 29.6 (C²¹), 28.4 (C³⁷). Quartet for C³¹ lost in baseline.

δ_F (377 MHz; CDCl₃/CD₃OD 1:1): –63.1 (C¹⁶F₃), –63.4 (C³¹F₃).

m/z (ES): 1209.4047 ([M + Na]⁺ C₅₇H₅₉F₉N₈NaO₁₀ requires 1209.4103).

Boc-Peptide [2]Rotaxane 5. [2]Rotaxane **4** (69 mg, 0.058 mmol) was dissolved in dry CH₂Cl₂ (2 mL), and cooled to 0 °C under an Ar (g) atmosphere. Then dry NEt₃ (8.8 mg, 12 μL, 0.087 mmol), triglycine *tert*-butyl ester (17 mg, 0.058 mmol), HOBt.hydrate (10 mg, 0.076 mmol) and EDCl.HCl (15 mg, 0.076 mmol) were added, maintaining the temperature at 0 °C. The reaction was stirred at r.t. for 6 h, at which time analytical TLC confirmed presence of **4**. The reaction flask was cooled to 0 °C and further dry NEt₃ (8.8 mg, 12 μL, 0.087 mmol), triglycine *tert*-butyl ester (17 mg, 0.058 mmol), HOBt.hydrate (10 mg, 0.076 mmol) and EDCl.HCl (15 mg, 0.076 mmol) were added. The reaction was stirred at r.t. for a further 16 h, at which time analytical TLC

confirmed presence of **4**. The reaction flask was cooled to 0 °C and further dry NEt₃ (8.8 mg, 12 μL, 0.087 mmol), triglycine *tert*-butyl ester (17 mg, 0.058 mmol), HOBt.hydrate (10 mg, 0.076 mmol) and EDCl.HCl (15 mg, 0.076 mmol) were added. The reaction was stirred at r.t. for a further 24 h, then diluted with CH₂Cl₂ (20 mL), and then washed with H₂O (1 × 20 mL), 10% citric acid (aq) (2 × 20 mL), saturated NaHCO₃ (aq) (2 × 20 mL) and brine (1 × 20 mL). The organic layer was dried (MgSO₄), and the solvent removed *in vacuo*. The crude material was purified by silica gel chromatography (CH₂Cl₂/CH₃OH 98:2 to 92:8) to yield pure product **5** as a white film (68 mg, 80%).



R_f: 0.22 (CH₂Cl₂/CH₃OH 96:4).

v_{max}/cm⁻¹ (neat): 3320 (N–H), 3070 (C–H), 2920 (C–H), 1650 (C=O), 1600, 1520, 1450, 1370, 1280, 1130, 1030.

δ_H (400 MHz; CDCl₃): 9.45 (1H, s, C⁹NH), 8.19 (4H, s, C¹¹H & C²⁶H), 8.07–8.10 (2H, br s, C⁵HNH or C⁷HNH & C⁴²HNH), 8.04 (1H, s, C¹³H), 8.00 (1H, s, C³²H), 7.90 (1H, br s, C⁵HNH or C⁷HNH), 7.84 (2H, br s, C¹⁴NH), 7.81 (1H, s, C³²H), 7.70 (1H, s, C³⁸H), 7.58 (1H, s, C³⁶H), 7.24 (2H, s, C²⁸NH & C⁴¹H), 6.94 (4H, d, ³J = 7.9 Hz, C¹⁷H), 6.74 (4H, d, ³J = 7.9 Hz, C¹⁸H), 6.28 (1H, br s, C³HNH), 4.99 (2H, s, C³⁴H), 4.60–4.65 (2H, m, C¹⁵H - diastereotopic), 4.29 (2H, d, ²J = 10 Hz, C²⁰H - diastereotopic), 4.20–4.25 (2H, m, C¹⁵H - diastereotopic), 4.15 (2H, d, ³J = 5.5 Hz, C⁴³H), 4.09 (2H, d, ²J = 10 Hz, C²⁰H - diastereotopic), 4.01 (6H, m, C⁶H, C⁸H & C³¹H), 3.87 (2H, d, ³J = 5.5 Hz, C⁴H), 3.56–3.75 (8H, m, C²¹H & C²²H), 2.65–2.69 (2H, app q, C²⁹H), 1.70–1.73 (2H, m, C³⁰H), 1.51 (9H, s, C⁴⁶H), 1.35 (9H, s, C¹H).

δ_C (100 MHz; CDCl₃): 171.9 (C⁵), 170.7 (C⁷ or C⁹), 169.9 (C⁴⁴), 168.0 (C⁷ or C⁹), 166.8 (C¹⁴), 166.2 (C⁴²), 164.0 (C²⁸), 158.5 (C³⁵), 156.7 (C³), 142.1 (C³³), 138.8 (*), 137.3 (C¹⁶), 136.2 (C¹⁹), 136.0 (*), 135.8 (*), 135.1 (*), 131.4 (q, ²J = 32 Hz, C²⁴ and/or C³⁹), 128.8 (C¹⁸), 128.3 (C¹⁷ & C²⁶), 124.8 (C³²), 124.5 (C²³), 123.1 (q, ¹J = 271 Hz, C²⁵ and/or C⁴⁰), 121.6 (C¹¹), 120.3 (br, C¹³), 117.0 (br, C³⁸), 116.4 (C³⁶), 115.0 (br, C⁴¹), 82.7 (C⁴⁵), 80.1 (C²), 73.6 (C²⁰), 70.8 (C²²), 69.5 (C²¹), 62.0 (C³⁴), 47.7 (C³¹), 44.5 (C¹⁵), 44.3 (C⁴), 43.7 (C⁶ or C⁸), 43.4 (C⁶ or C⁸), 42.8 (C⁴³), 37.3 (C²⁹), 28.7 (C³⁰), 28.2 (C¹), 28.1 (C⁴⁶). * = C¹⁰ or C¹² or C²⁷ or C³⁷.


δ_F (377 MHz; CDCl₃): –62.5 (C²⁵F₃), –62.7 (C⁴⁰F₃).

m/z (ES): 1480.5282 ([M + Na]⁺ C₆₈H₇₆F₉N₁₁NaO₁₅ requires 1480.5271).

[1]Rotaxane 6. [2]Rotaxane **5** (20 mg, 0.014 mmol) was dissolved in CH₂Cl₂ (0.5 mL) and cooled to 0 °C. TFA (0.25 mL)



was added dropwise and the reaction mixture was stirred at r.t. for 2 h. Volatiles were removed *in vacuo* and deprotection confirmed by ^1H NMR. The residue was re-dissolved in dry CH_2Cl_2 (7 mL), and cooled to 0°C under an Ar (g) atmosphere. Then dry DIPEA (18 mg, 24 μL , 0.14 mmol) and HATU (16 mg, 0.042 mmol) were added, maintaining the temperature at 0°C . The reaction was stirred at r.t. for 90 h, then diluted with CH_2Cl_2 to 20 mL, then washed with 1M HCl (aq) (2×20 mL), sat. NaHCO_3 (aq) (2×20 mL) and brine (1×20 mL). The organic layer was dried (MgSO_4), and the solvent removed *in vacuo*. The crude material was purified by silica gel chromatography ($\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$ 96:4 to 90:10) to yield pure product **6** as a white film (6.0 mg, 34%).



R_f : 0.31 ($\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$ 94:6).

$\nu_{\text{max}}/\text{cm}^{-1}$ (neat): 3300 (N–H), 3080 (C–H), 2920 (C–H), 2870 (C–H), 1650 (C=O), 1600, 1530, 1450, 1360, 1330, 1280, 1170, 1130, 1030.

δ_{H} (400 MHz; $\text{CDCl}_3/\text{CD}_3\text{OD}$ 1:1): 8.23 (2H, d, $^4J = 1.5$ Hz, C^2H), 8.14 (1H, s, C^{23}H), 8.11 (2H, br s, C^{17}H), 8.02 (2H, app t, C^4H & C^{14}H), 7.65 (1H, s, C^{29}H), 7.38 (1H, s, C^{27}H), 7.15 (1H, s, C^{32}H), 6.91 (4H, d, $^3J = 8.0$ Hz, C^8H), 6.76 (4H, d, $^3J = 8.0$ Hz, C^9H), 4.83 (2H, s, C^{25}H), 4.60–4.62 (2H, obscured doublet, C^6H - diastereotopic), 4.30 (2H, d, $^2J = 10$ Hz, C^{11}H - diastereotopic), 4.19 (2H, s, C^{34}H), 4.12–4.17 (4H, m, C^{22}H & C^6H - diastereotopic), 4.09 (2H, d, $^2J = 10$ Hz, C^{11}H - diastereotopic), 4.03 (2H, s, C^{38}H or C^{40}H), 3.99 (2H, s, C^{36}H), 3.92 (2H, s, C^{38}H or C^{40}H), 3.71–3.76 (2H, m, C^{13}H - diastereotopic), 3.55–3.66 (6H, C^{12}H & C^{13}H - diastereotopic), 2.61 (2H, t, $^3J = 7.4$ Hz, C^{20}H), 1.76–1.83 (2H, app quin, C^{21}H).

δ_{C} (100 MHz; $\text{CDCl}_3/\text{CD}_3\text{OD}$ 1:1): 172.3 (C^{35}), 172.1 ($\text{C}^{37}/\text{C}^{39}/\text{C}^{41}$), 171.6 ($\text{C}^{37}/\text{C}^{39}/\text{C}^{41}$), 169.4 ($\text{C}^{37}/\text{C}^{39}/\text{C}^{41}$), 168.3 (C^5 & C^{33}), 164.7 (C^{19}), 159.2 (C^{26}), 142.9 (C^{24}), 139.3 (C^1 or C^3), 138.1 (C^7), 137.1 (C^{10}), 136.8 (C^{18} or C^{28}), 136.5 (C^{18} or C^{28}), 136.1 (C^1 or C^3), 132.7 (q, $^2J = 33$ Hz, C^{30}), 132.2 (q, $^2J = 34$ Hz, C^{15}), 129.4 (C^9), 128.9 (C^8 & C^{17}), 126.5 (C^{23}), 125.3 (br, C^{14}), 122.7 (C^2), 121.5 (C^4), 117.9 (br, C^{29}), 117.0 (C^{27}), 115.2 (br, C^{32}), 74.2 (C^{11}), 71.6 (C^{13}), 70.3 (C^{12}), 62.4 (C^{25}), 48.6 (C^{22}), 45.0 (C^6), 44.3 (C^{34}), 44.3 (sic, C^{38} or C^{40}), 43.8 (C^{38} or C^{40}), 43.6 (C^{36}), 37.8 (C^{20}), 29.5 (C^{21}). Quartets for C^{16} & C^{31} not fully resolved in 120–130 ppm region.

δ_{F} (377 MHz; $\text{CDCl}_3/\text{CD}_3\text{OD}$ 1:1): -63.1 (C^{16}F_3), -63.5 (C^{31}F_3).

m/z (ES): 1306.3971 ($[\text{M} + \text{Na}]^+$ $\text{C}_{59}\text{H}_{58}\text{F}_9\text{N}_{11}\text{NaO}_{12}$ requires 1306.4015).

Conflicts of interest

There are no conflicts to declare.

View Article Online

DOI: 10.1039/D0OB01190H

Acknowledgements

N.H.E. expresses his appreciation for funding from Lancaster University (Faculty of Science & Technology Research Grant and Joy Welch Educational Charitable Trust Grant). We also wish to thank the following colleagues at Lancaster University: Dr Mike Coogan & Dr Nathan Halcovitch for the collecting and processing of X-ray crystallographic data, Dr David Rochester for the recording of mass spectra and Dr Michael Peach for computational advice.

Underlying data for this paper are provided in the experimental section and ESI[†], along with electronic copies of NMR spectra (including fid files) and computational output files being available from: DOI: 10.17635/lancaster/researchdata/372.

Notes and references

§ Single crystals of macrocycle **3** suitable for X-ray structure determination were obtained by slow evaporation of a chloroform/methanol solution. The solved structure (CCDC 1995143) and further details are to be found in ESI[†], page S50.

- For reviews about lasso peptides see: (a) M. O. Maksimov, S. J. Pan and A. J. Link, *Nat. Prod. Rep.*, 2012, **29**, 996–1006; (b) J. D. Hegemann, H. Zimmermann, X. Xie and M. A. Marahiel, *Acc. Chem. Res.*, 2015, **48**, 1909–1919; (c) H. Martín-Gómez and J. Tulla-Puche, *Org. Biomol. Chem.*, 2018, **16**, 5065–5080.
- (a) R. A. Salomón and R. N. Fariás, *J. Bacteriol.*, 1992, **174**, 7428–7435; (b) M. Iwatsuki, H. Tomoda, R. Uchida, H. Gouda, S. Hirono and S. Omura, *J. Am. Chem. Soc.*, 2006, **128**, 7486–7491; (c) T. A. Knappe, U. Linne, S. Zirah, S. Rebuffat, X. Xie and M. A. Marahiel, *J. Am. Chem. Soc.*, 2008, **130**, 11446–11454; (d) W. L. Cheung-Lee, M. E. Parry, A. J. Cartagena, S. A. Darst and A. J. Link, *J. Bio. Chem.*, 2019, **294**, 6822–6830.
- (a) R. Katahira, K. Shibata, M. Yamasaki, Y. Matsuda and M. Yoshida, *Bioorg. Med. Chem.*, 1995, **3**, 1273–1280; (b) T. A. Knappe, U. Linne, X. Xie and M. A. Marahiel, *FEBS Lett.*, 2010, **584**, 785–789.
- (a) G. Helyncck, C. Dubertret, J. F. Mayaux and J. Leboul, *J. Antibiot.*, 1993, **46**, 1756–1757; (b) N. R. Braddman, F. J. Piscotta, J. Hauver, E. A. Campbell, A. J. Link and S. A. Darst, *Proc. Nat. Acad. Sci. U.S.A.*, 2019, **116**, 1273–1278.
- M. Chen, S. Wang and X. Yu, *Chem. Commun.*, 2019, **55**, 3323–3326.
- T. A. Knappe, F. Manzenrieder, C. Mas-Moruno, U. Linne, F. Sasse, H. Kessler, X. Xie and M. A. Marahiel, *Angew. Chem. Int. Ed.*, 2011, **50**, 8714–8717.
- Selected examples of synthetic [1]rotaxanes: (a) K. Hiratani, M. Kaneyama, Y. Nagawa, E. Koyama and M. Kanosato, *J. Am. Chem. Soc.*, 2004, **126**, 13568–13569; (b) Z. Xue and M. F. Mayer, *J. Am. Chem. Soc.*, 2010, **132**, 3274–3276; (c) H. Li, H. Zhang, Q. Zhang, Q.-W. Zhang and D.-H. Qu, *Org. Lett.*, 2012, **14**, 5900–5903; (d) H. Li, J.-N. Zhang, W. Zhou, H. Zhang, Q. Zhang, D.-H. Qu and H. Tian, *Org. Lett.*, 2013, **15**, 3070–3073; (e) B. Xia and M. Xue, *Chem. Commun.*, 2014, **50**, 1021–1023; (f) P. Waelès, C. Clavel, K. Fournel-Marotte and F. Coutrot, *Chem. Sci.*, 2015, **6**, 4828–4836; (g) Y. Wang, J. Sun, Z. Liu, M. S. Nassar, Y. Y. Botros and J. F. Stoddart, *Chem. Sci.*, 2017, **8**, 2562–2568; (h) A. Saura-Sanmartin, A. Martinez-Cuevas, A. Pastor, D. Bautista and J. Berna, *Org. Biomol. Chem.*, 2018, **16**,




- 6980-6987; (i) J.-J. Yu, L.-Y. Zhao, Z.-T. Shi, Q. Zhang, G. London, W.-J. Liang, C. Gao, M.-M. Li, X.-M. Cao, H. Tian, B. L. Feringa and D.-H. Qu, *J. Org. Chem.*, 2019, **84**, 5790-5802.
- 8 C. Clavel, K. Fournel-Marotte and F. Coutrot, *Molecules*, 2013, **18**, 11553-11575.
- 9 F. Saito and J. W. Bode, *Chem. Sci.*, 2017, **8**, 2878-2884.
- 10 B. E. Fletcher, M. J. G. Peach and N. H. Evans, *Org. Biomol. Chem.*, 2017, **15**, 2797-2803.
- 11 N. H. Evans and G. R. Akien, *Supramol. Chem.*, 2018, **30**, 758-764.
- 12 C. E. Gell, T. A. McArdle-Ismaguilov and N. H. Evans, *Chem. Commun.*, 2019, **55**, 1576-1579.
- 13 For a recent review on the hydrogen bond templated synthesis of rotaxanes (and catenanes) see: N. H. Evans, *Eur. J. Org. Chem.*, 2019, 3320-3343.
- 14 Selected examples of other, non-CuAAC click, stoppering of simple amide half-axes threaded through mono-isophthalamide macrocycles to generate [2]rotaxanes: (a) D. A. Leigh and A. R. Thomson, *Org. Lett.*, 2006, **8**, 5377-5379; (b) A. Vidonne and D. Philp, *Tetrahedron*, 2008, **64**, 8464-8475; (c) T. Kosikova, N. I. Hassan, D. B. Cordes, A. M. Z. Slawin and D. Philp, *J. Am. Chem. Soc.*, 2015, **137**, 16074-16083; (d) A. Vidonne, T. Kosikova and D. Philp, *Chem. Sci.*, 2016, **7**, 2592-2603.
- 15 The modest isolated yield of [2]rotaxane **4** is attributed to oxidation of the macrocyclic amino functionality, in particular during chromatographic purification.
- 16 HATU - (1-[Bis(dimethylaminomethylene)-1H-1,2,3-triazolo[4,5-b]pyridinium oxide hexafluorophosphate) - is widely used for the macrocyclization of peptides. For reviews of this topic see: (a) C. J. White and A. K. Yudin, *Nature Chem.*, 2011, **3**, 509-524; (b) Y. H. Lau, P. de Andrade, Y. Wu and D. R. Spring, *Chem. Soc. Rev.*, 2015, **44**, 91-102; (c) D. G. Rivera, G. M. Ojeda-Carralero, L. Reguera, E. V. Van der Eycken, *Chem. Soc. Rev.*, 2020, **49**, 2039-2059.
- 17 (a) N. J. Baxter and M. P. Williamson, *J. Biomol. NMR*, 1997, **9**, 359-369; (b) T. Cierpicki and J. Otlewski, *J. Biomol. NMR*, 2001, **21**, 249-261.
- 18 M. Llinas and M. P. Klein, *J. Am. Chem. Soc.*, 1975, **97**, 4731-4737.
- 19 For reviews on the use of diffusion NMR experiments in supramolecular chemistry see: (a) S. V. Kharlamov and S. K. Latypov, *Russ. Chem. Rev.*, 2010, **79**, 635-653; (b) L. Avram and Y. Cohen, *Chem. Soc. Rev.*, 2015, **44**, 586-602.
- 20 R. Evans, Z. Deng, A. K. Rogerson, A. S. McLachlan, J. J. Richards, M. Nilsson and G. A. Morris, *Angew. Chem. Int. Ed.*, 2013, **52**, 3199-3202.
- 21 The DFT structure of [1]rotaxane **6** reveals that all amides in the loop (*a* & *d-g*) can be in a *trans* conformation. Depiction of amide *a* as *cis* in *Scheme 1* arises from attempting to achieve a 2D depiction of [1]rotaxane **6** without unusual bond lengths or angles.

View Article Online
DOI: 10.1039/D0OB01190H



Graphical Abstract

5



The rapid synthesis and characterization of an amide hydrogen bond templated [1]rotaxane – demonstrating a potential pathway to synthetic analogues of lasso peptides.

