

1
2
3
4
5
6
RESEARCH ARTICLE

The effect of *Staphylococcus aureus* carriage in late pregnancy on antibody levels to staphylococcal toxins in cord blood and breast milk

7 Linda M. Harrison¹, James A. Morris¹, Robert M. Lauder² & David R. Telford¹
 8
 9
 10

11
12
13
14
15
16
17
18
19
20
1 Department of Pathology, University Hospitals of Morecambe Bay NHS Trust, Royal Lancaster Infirmary, Lancaster LA1 4RP, UK; and ²School of Health and
Medicine, Division of Biomedical and Life Sciences, Lancaster University, Lancaster, UK
 21
 22
 23
 24
 25
 26
 27
 28
 29
 30
 31
 32
 33
 34
 35
 36
 37
 38
 39
 40
 41
 42
 43
 44
 45
 46
 47
 48
 49
 50
 51
 52
 53
 54
 55
 56
 57
 58
 59
 60
 61
 62
 63
 64
 65
 66
 67
 68
 69
 70
 71
 72
 73
 74
 75
 76
 77
 78
 79
 80
 81
 82
 83
 84
 85
 86
 87
 88
 89
 90
 91
 92
 93
 94
 95
 96
 97
 98
 99
 100
 101
 102
 103
 104
 105
 106
 107
 108
 109
 110
 111
 112
 113
 114
 115
 116
 117
 118
 119
 120
 121
 122
 123
 124
 125
 126
 127
 128
 129
 130
 131
 132
 133
 134
 135
 136
 137
 138
 139
 140
 141
 142
 143
 144
 145
 146
 147
 148
 149
 150
 151
 152
 153
 154
 155
 156
 157
 158
 159
 160
 161
 162
 163
 164
 165
 166
 167
 168
 169
 170
 171
 172
 173
 174
 175
 176
 177
 178
 179
 180
 181
 182
 183
 184
 185
 186
 187
 188
 189
 190
 191
 192
 193
 194
 195
 196
 197
 198
 199
 200
 201
 202
 203
 204
 205
 206
 207
 208
 209
 210
 211
 212
 213
 214
 215
 216
 217
 218
 219
 220
 221
 222
 223
 224
 225
 226
 227
 228
 229
 230
 231
 232
 233
 234
 235
 236
 237
 238
 239
 240
 241
 242
 243
 244
 245
 246
 247
 248
 249
 250
 251
 252
 253
 254
 255
 256
 257
 258
 259
 260
 261
 262
 263
 264
 265
 266
 267
 268
 269
 270
 271
 272
 273
 274
 275
 276
 277
 278
 279
 280
 281
 282
 283
 284
 285
 286
 287
 288
 289
 290
 291
 292
 293
 294
 295
 296
 297
 298
 299
 300
 301
 302
 303
 304
 305
 306
 307
 308
 309
 310
 311
 312
 313
 314
 315
 316
 317
 318
 319
 320
 321
 322
 323
 324
 325
 326
 327
 328
 329
 330
 331
 332
 333
 334
 335
 336
 337
 338
 339
 340
 341
 342
 343
 344
 345
 346
 347
 348
 349
 350
 351
 352
 353
 354
 355
 356
 357
 358
 359
 360
 361
 362
 363
 364
 365
 366
 367
 368
 369
 370
 371
 372
 373
 374
 375
 376
 377
 378
 379
 380
 381
 382
 383
 384
 385
 386
 387
 388
 389
 390
 391
 392
 393
 394
 395
 396
 397
 398
 399
 400
 401
 402
 403
 404
 405
 406
 407
 408
 409
 410
 411
 412
 413
 414
 415
 416
 417
 418
 419
 420
 421
 422
 423
 424
 425
 426
 427
 428
 429
 430
 431
 432
 433
 434
 435
 436
 437
 438
 439
 440
 441
 442
 443
 444
 445
 446
 447
 448
 449
 450
 451
 452
 453
 454
 455
 456
 457
 458
 459
 460
 461
 462
 463
 464
 465
 466
 467
 468
 469
 470
 471
 472
 473
 474
 475
 476
 477
 478
 479
 480
 481
 482
 483
 484
 485
 486
 487
 488
 489
 490
 491
 492
 493
 494
 495
 496
 497
 498
 499
 500
 501
 502
 503
 504
 505
 506
 507
 508
 509
 510
 511
 512
 513
 514
 515
 516
 517
 518
 519
 520
 521
 522
 523
 524
 525
 526
 527
 528
 529
 530
 531
 532
 533
 534
 535
 536
 537
 538
 539
 540
 541
 542
 543
 544
 545
 546
 547
 548
 549
 550
 551
 552
 553
 554
 555
 556
 557
 558
 559
 560
 561
 562
 563
 564
 565
 566
 567
 568
 569
 570
 571
 572
 573
 574
 575
 576
 577
 578
 579
 580
 581
 582
 583
 584
 585
 586
 587
 588
 589
 590
 591
 592
 593
 594
 595
 596
 597
 598
 599
 600
 601
 602
 603
 604
 605
 606
 607
 608
 609
 610
 611
 612
 613
 614
 615
 616
 617
 618
 619
 620
 621
 622
 623
 624
 625
 626
 627
 628
 629
 630
 631
 632
 633
 634
 635
 636
 637
 638
 639
 640
 641
 642
 643
 644
 645
 646
 647
 648
 649
 650
 651
 652
 653
 654
 655
 656
 657
 658
 659
 660
 661
 662
 663
 664
 665
 666
 667
 668
 669
 670
 671
 672
 673
 674
 675
 676
 677
 678
 679
 680
 681
 682
 683
 684
 685
 686
 687
 688
 689
 690
 691
 692
 693
 694
 695
 696
 697
 698
 699
 700
 701
 702
 703
 704
 705
 706
 707
 708
 709
 710
 711
 712
 713
 714
 715
 716
 717
 718
 719
 720
 721
 722
 723
 724
 725
 726
 727
 728
 729
 730
 731
 732
 733
 734
 735
 736
 737
 738
 739
 740
 741
 742
 743
 744
 745
 746
 747
 748
 749
 750
 751
 752
 753
 754
 755
 756
 757
 758
 759
 7510
 7511
 7512
 7513
 7514
 7515
 7516
 7517
 7518
 7519
 7520
 7521
 7522
 7523
 7524
 7525
 7526
 7527
 7528
 7529
 7530
 7531
 7532
 7533
 7534
 7535
 7536
 7537
 7538
 7539
 7540
 7541
 7542
 7543
 7544
 7545
 7546
 7547
 7548
 7549
 7550
 7551
 7552
 7553
 7554
 7555
 7556
 7557
 7558
 7559
 7560
 7561
 7562
 7563
 7564
 7565
 7566
 7567
 7568
 7569
 7570
 7571
 7572
 7573
 7574
 7575
 7576
 7577
 7578
 7579
 7580
 7581
 7582
 7583
 7584
 7585
 7586
 7587
 7588
 7589
 7590
 7591
 7592
 7593
 7594
 7595
 7596
 7597
 7598
 7599
 75100
 75101
 75102
 75103
 75104
 75105
 75106
 75107
 75108
 75109
 75110
 75111
 75112
 75113
 75114
 75115
 75116
 75117
 75118
 75119
 75120
 75121
 75122
 75123
 75124
 75125
 75126
 75127
 75128
 75129
 75130
 75131
 75132
 75133
 75134
 75135
 75136
 75137
 75138
 75139
 75140
 75141
 75142
 75143
 75144
 75145
 75146
 75147
 75148
 75149
 75150
 75151
 75152
 75153
 75154
 75155
 75156
 75157
 75158
 75159
 75160
 75161
 75162
 75163
 75164
 75165
 75166
 75167
 75168
 75169
 75170
 75171
 75172
 75173
 75174
 75175
 75176
 75177
 75178
 75179
 75180
 75181
 75182
 75183
 75184
 75185
 75186
 75187
 75188
 75189
 75190
 75191
 75192
 75193
 75194
 75195
 75196
 75197
 75198
 75199
 75200
 75201
 75202
 75203
 75204
 75205
 75206
 75207
 75208
 75209
 75210
 75211
 75212
 75213
 75214
 75215
 75216
 75217
 75218
 75219
 75220
 75221
 75222
 75223
 75224
 75225
 75226
 75227
 75228
 75229
 75230
 75231
 75232
 75233
 75234
 75235
 75236
 75237
 75238
 75239
 75240
 75241
 75242
 75243
 75244
 75245
 75246
 75247
 75248
 75249
 75250
 75251
 75252
 75253
 75254
 75255
 75256
 75257
 75258
 75259
 75260
 75261
 75262
 75263
 75264
 75265
 75266
 75267
 75268
 75269
 75270
 75271
 75272
 75273
 75274
 75275
 75276
 75277
 75278
 75279
 75280
 75281
 75282
 75283
 75284
 75285
 75286
 75287
 75288
 75289
 75290
 75291
 75292
 75293
 75294
 75295
 75296
 75297
 75298
 75299
 75300
 75301
 75302
 75303
 75304
 75305
 75306
 75307
 75308
 75309
 75310
 75311
 75312
 75313
 75314
 75315
 75316
 75317
 75318
 75319
 75320
 75321
 75322
 75323
 75324
 75325
 75326
 75327
 75328
 75329
 75330
 75331
 75332
 75333
 75334
 75335
 75336
 75337
 75338
 75339
 75340
 75341
 75342
 75343
 75344
 75345
 75346
 75347
 75348
 75349
 75350
 75351
 75352
 75353
 75354
 75355
 75356
 75357
 75358
 75359
 75360
 75361
 75362
 75363
 75364
 75365
 75366
 75367
 75368
 75369
 75370
 75371
 75372
 75373
 75374
 75375
 75376
 75377
 75378
 75379
 75380
 75381
 75382
 75383
 75384
 75385
 75386
 75387
 75388
 75389
 75390
 75391
 75392
 75393
 75394
 75395
 75396
 75397
 75398
 75399
 75400
 75401
 75402
 75403
 75404
 75405
 75406
 75407
 75408
 75409
 75410
 75411
 75412
 75413
 75414
 75415
 75416
 75417
 75418
 75419
 75420
 75421
 75422
 75423
 75424
 75425
 75426
 75427
 75428
 75429
 75430
 75431
 75432
 75433
 75434
 75435
 75436
 75437
 75438
 75439
 75440
 75441
 75442
 75443
 75444
 75445
 75446
 75447
 75448
 75449
 75450
 75451
 75452
 75453
 75454
 75455
 75456
 75457
 75458
 75459
 75460
 75461
 75462
 75463
 75464
 75465
 75466
 75467
 75468
 75469
 75470
 75471
 75472
 75473
 75474
 75475
 75476
 75477
 75478
 75479
 75480
 75481
 75482
 75483
 75484
 75485
 75486
 75487
 75488
 75489
 75490
 75491
 75492
 75493
 75494
 75495
 75496
 75497
 75498
 75499
 75500
 75501
 75502
 75503
 75504
 75505
 75506
 75507
 75508
 75509
 75510
 75511
 75512
 75513
 75514
 75515
 75516
 75517
 75518
 75519
 75520
 75521
 75522
 75523
 75524
 75525
 75526
 75527
 75528
 75529
 75530
 75531
 75532
 75533
 75534
 75535
 75536
 75537
 75538
 75539
 75540
 75541
 75542
 75543
 75544
 75545
 75546
 75547
 75548
 75549
 75550
 75551
 75552
 75553
 75554
 75555
 75556
 75557
 75558
 75559
 75560
 75561
 75562
 75563
 75564
 75565
 75566
 75567
 75568
 75569
 75570
 75571
 75572
 75573
 75574
 75575
 75576
 75577
 75578
 75579
 75580
 75581
 75582
 75583
 75584
 75585
 75586
 75587
 75588
 75589
 75590
 75591
 75592
 75593
 75594
 75595
 75596
 75597
 75598
 75599
 75600
 75601
 75602
 75603
 75604
 75605
 75606
 75607
 75608
 75609
 75610
 75611
 75612
 75613
 75614
 75615
 75616
 75617
 75618
 75619
 75620
 75621
 75622
 75623
 75624
 75625
 75626
 75627
 75628
 75629
 75630
 75631
 75632
 75633
 75634
 75635
 75636
 75637
 75638
 75639
 75640
 75641
 75642
 75643
 75644
 75645
 75646
 75647
 75648
 75649
 75650
 75651
 75652
 75653
 75654
 75655
 75656
 75657
 75658
 75659
 75660
 75661
 75662
 75663
 75664
 75665
 75666
 75667
 75668
 75669
 75670
 75671
 75672
 75673
 75674
 75675
 75676
 75677
 75678
 75679
 75680
 75681

of *S. aureus* in the later stages of pregnancy influenced the levels of antibody specific to the *S. aureus* toxins (SEB, SEC and TSST-1) in cord blood and breast milk and to explore the relationship between the levels of antibody in antenatal blood and cord blood.

Materials and methods

Population studied

Pregnant women were recruited to the study at antenatal classes. The study had ethical approval from the Preston, Chorley and South Ribble Local Research Ethics Committee and informed consent was obtained from the women.

Sample collection and preparation

Cord blood samples were collected by the midwives. The samples were separated and the serum was stored at -70°C . Blood samples taken during the second trimester, as part of the normal antenatal procedure, were processed and stored in the same manner as the cord blood samples. Women were visited at home on two occasions during the last 6 weeks of pregnancy. Before each visit, participants collected a stool sample in a sterile container and stored it in a refrigerator. A nasopharyngeal swab was taken at each visit.

Isolation and identification of *S. aureus*

Staphylococcus aureus was isolated and identified from nasopharyngeal swabs using standard clinical laboratory methods. Stool samples were diluted serially in 10-fold steps in peptone water and cultured aerobically on *Staphylococcus* medium no 110 (Oxoid) at 37°C for 2 days. Colonies of different morphology were enumerated and subcultured on blood agar. Standard methods were used to identify *S. aureus* isolates. All *S. aureus* isolates were stored at -70°C for further tests.

Detection of IgG to staphylococcal pyrogenic toxins in antenatal and cord blood

Serum samples were examined for IgG antibody to SEB, SEC and TSST-1 by an enzyme-linked immunosorbent assay (ELISA). A modification of the protocol of Al Madani *et al.* (1999) was used. Test and control wells were coated with 100 μL of SEB, SEC or TSST-1 at 1 $\mu\text{g mL}^{-1}$ in carbonate buffer (Sigma). Standard wells were coated with 100 μL of serial dilutions of human IgG (Sigma); concentrations ranged from 5 to 10 000 ng mL^{-1} . Plates were incubated overnight at 4°C . The wells were then emptied and washed four times with washing buffer [0.05% (v/v) Tween-20 in phosphate-buffered saline (PBST)]. The plates were blocked for 1 h at 37°C with a buffer containing 5% (w/v) skimmed-milk powder in PBST. Plates were washed four times. Serum

samples were diluted in buffer [2.5% (w/v) skimmed-milk powder in PBST] and 100 μL was loaded into each test well. The same buffer (100 μL) was added to standard and control wells. Plates were then incubated for 1 h at 37°C . After four washes, 100 μL of anti-human IgG, conjugated to horseradish peroxidase (HRP) (Sigma) diluted in buffer [2.5% (w/v) skimmed-milk powder in PBST], was added to the test and standard wells. HRP-labelled sheep anti-SEB (100 μL), HRP-labelled sheep anti-SEC (100 μL) or HRP-labelled sheep anti-TSST-1, diluted 1 in 300 in buffer [2.5% (w/v) skimmed-milk powder in PBST], was added to the appropriate control wells. The plates were incubated for 1 h at 37°C . After three washes, 100 μL of the substrate 3,3',5,5'-tetramethylbenzidine (TMB) (Sigma) was added to each well. The substrate contained 1 mg TMB in 10 mL of 0.05 M phosphate-citrate buffer (pH 5.0) and was activated immediately before use by adding 2 μL of 30% (v/v) H_2O_2 . The reaction was stopped after 15 min by the addition of 100 μL 2 M H_2SO_4 and the absorbance was read at 450 nm with a plate reader (Organon Teknica).

Detection of IgA to staphylococcal pyrogenic toxins in breast milk

A modification of the protocol of Gordon *et al.* (1999) was used. Test and control wells of microtitre plates were coated with 100 μL of SEB, SEC or TSST-1 at 1 $\mu\text{g mL}^{-1}$ in carbonate buffer. Standard wells were coated with 100 μL of serial dilutions of a standard human IgA derived from colostrum (Sigma); concentrations ranged from 5 to 10 000 ng mL^{-1} . Plates were incubated overnight at 4°C and then washed four times. The plates were blocked for 1 h at 37°C . After three washes, 100 μL samples of breast milk diluted in blocking buffer were loaded into the appropriate test wells. Blocking buffer (100 μL) was added to standard and control wells. Plates were incubated for 1 h at 37°C . After four washes, 100 μL of HRP-labelled goat anti-human IgA (Calbiochem) diluted in blocking buffer was added to the test and standard wells and 100 μL of HRP-labelled sheep anti-SEB, HRP-labelled sheep anti-SEC or HRP-labelled sheep anti-TSST-1, diluted 1 in 300 in blocking buffer, was added to the appropriate control wells. The plates were incubated at 37°C for 1 h. The experiment then continued as described previously.

Detection of toxin production by *S. aureus* isolates

Staphylococcus aureus isolates were cultured using the cellophane over agar technique (Newbould *et al.*, 1989) and the resulting supernatant samples were analysed by ELISA. Test and control wells were coated with 100 μL of anti-SEB, anti-SEC or anti-TSST-1 (Toxin Technology), diluted 1:1000 in carbonate buffer (Sigma). Plates were incubated overnight

at 4 °C and then emptied and washed four times with washing buffer. The plates were blocked with blocking buffer containing 2.5% (v/v) fish gelatin in PBST. Plates were washed four times. Supernatant samples and controls were diluted in PBS and 100 µL was loaded into each test well.

Q2 Controls included purified SEB, purified SEC, purified TSST-1 (Toxin Technology, Sarasota, FL) and supernatant prepared from *S. aureus* isolates previously tested for the production of SEB, SEC and TSST-1 (Colindale Public Health Laboratory). Plates were then incubated for 1 h at 37 °C. After four washes, 100 µL of the corresponding biotinylated antitoxin (Toxin Technology) diluted 1 in 2000 in blocking buffer was added to the test and control wells. The plates were incubated for 1 h at 37 °C. After four washes, 100 µL of extravidin peroxidase (Sigma) diluted 1 : 2000 was added to each well and plates were incubated for 30 min at 37 °C. The experiment then continued as described previously.

Statistical analyses

Results were analysed using SPSS software (version 13.0). Nonparametric tests were used for data that were not normally distributed. These data were not normalized by log transformation because many nil values were detected. Spearman's correlation coefficient was used to assess the association between levels of antibody in antenatal and cord blood samples. Antitoxin levels were compared between antenatal and cord blood samples using the Wilcoxon matched samples test. χ^2 Test, Mann–Whitney *U*-test and Kruskal–Wallis test were used to assess antitoxin levels in relation to other variables. Comparison of data is presented in Figs 1, 3 and 4 as box and whisker plots. The box represents the 25th–75th percentiles, and the median is represented by the line within the box. The whiskers

represent the 5th–95th percentiles and the circles and stars depict outliers for each group.

Results

Population characteristics

Ninety-six pregnant women were enrolled into the study. The mean gestation period was 281 days (range 257–296, SD ± 9.5). The average maternal age at birth was 31 years (range 18–42, SD ± 4.4). There was no significant association between length of gestation and maternal age.

Detection of IgG to staphylococcal pyrogenic toxins in blood samples

Ninety-two antenatal blood samples were available to test for the presence of IgG to SEB, SEC and TSST-1. The median level of antibody to SEB was 3799 ng mL⁻¹ (mean: 6204 ng mL⁻¹; range: 0–35 588 ng mL⁻¹; SD ± 6846). Antibody levels to SEC ranged from 0 to 21 499 ng mL⁻¹ (median: 5045 ng mL⁻¹; mean: 6248 ng mL⁻¹; SD ± 5858) and those for TSST-1 ranged from 0 to 16 412 ng mL⁻¹ (median: 943 ng mL⁻¹; mean: 1674 ng mL⁻¹; SD ± 2333). Eight women (8.7%) had no detectable IgG to SEB, six (6.5%) had no detectable IgG to SEC and 10 (10.9%) had no detectable IgG to TSST-1.

Ninety-six cord blood samples were collected and tested for the presence of IgG to SEB, SEC and TSST-1. The median level of antibody to SEB was 4608 ng mL⁻¹ (mean: 8987 ng mL⁻¹; range: 0–46 605 ng mL⁻¹; SD ± 11 190). Antibody levels to SEC ranged from 0 to 53 886 ng mL⁻¹ (median: 4001 ng mL⁻¹; mean: 8970 ng mL⁻¹; SD ± 10 861) and those for TSST-1 ranged from 0 to 14 154 ng mL⁻¹ (median: 1170 ng mL⁻¹; mean: 2512 ng mL⁻¹; SD ± 3084). Seven samples (7.3%) had no detectable IgG to SEB, three (3.1%) had no detectable IgG to SEC and seven (7.3%) had no detectable IgG to TSST-1. There was no significant association between maternal age and antibody levels or between length of gestation and antibody levels. However, 84% of pregnant women over the age of 30, but only 68% of pregnant women under the age of 30 had antibody to all three toxins (difference not significant).

Male infants ($n = 51$) had significantly lower levels of cord blood IgG to SEC than female infants ($n = 45$) ($P = 0.05$, Mann–Whitney *U*-test, Fig. 1). For anti-SEC IgG, the mean level in males was 7440 ng mL⁻¹ and in females the mean level was 10 705 ng mL⁻¹. No significant differences were found between male and female infants for cord blood IgG levels to SEB or TSST-1.

Antenatal anti-SEB IgG correlated positively with cord blood anti-SEB IgG ($\rho = 0.815$, $P < 0.01$). Antenatal anti-SEC IgG correlated positively with cord blood anti-SEC IgG ($\rho = 0.856$, $P < 0.01$, Fig. 2). Antenatal anti-TSST-1 IgG




Fig. 1. Levels of anti-SEC IgG in the cord blood of male and female infants.




Fig. 2. Correlation between levels of IgG to SEC in antenatal and cord blood.

correlated positively with cord blood anti-TSST-1 IgG ($\rho=0.721$, $P < 0.01$). Paired sample tests showed that specific IgG levels to each of the toxins were significantly greater in the cord blood samples than in the antenatal blood samples ($P < 0.01$).

Detection of IgA to staphylococcal pyrogenic toxins in breast milk

Breast milk samples were collected from 83 women in the first 7 days following the birth of their baby. The median level of antibody to SEB was 201 ng mL^{-1} (mean: 423 ng mL^{-1} ; range: $0\text{--}10\,000 \text{ ng mL}^{-1}$; $SD \pm 1135$). Antibody levels to SEC ranged from 0 to 5370 ng mL^{-1} (median: 154 ng mL^{-1} ; mean: 402 ng mL^{-1} ; $SD \pm 787$) and those for TSST-1 ranged from 0 to 3050 ng mL^{-1} (median: 0 ng mL^{-1} ; mean: 146 ng mL^{-1} ; $SD \pm 472$).

Breast milk anti-SEB IgA showed a significant positive correlation with cord blood anti-SEB IgG ($\rho=0.233$, $P < 0.05$). Breast milk anti-SEC-1 IgA correlated positively with cord blood anti-SEC-1 IgG ($\rho=0.321$, $P < 0.01$). Breast milk anti-TSST-1 IgA showed no significant association with cord blood anti-TSST-1 IgG.

Carriage of *S. aureus* in pregnant women

Of the 96 women, 29 (30.2%) were colonized with *S. aureus*. Sixty-seven women (69.8%) had negative nasopharyngeal and stool cultures, four (4.2%) had positive nasopharyngeal and stool cultures, 16 (16.7%) were nasopharyngeal carriers only and nine (9.4%) were stool carriers only. Of the women with positive nasopharyngeal cultures, 50% were colonized with *S. aureus* on both sampling occasions. Of the women with positive stool cultures, 23% were colonized with *S. aureus* on both sampling occasions. The mean *S. aureus*

count per gram of faeces was 1.99×10^4 , (range: $1.2 \times 10^3\text{--}1.2 \times 10^5$; $SD \pm 3.1 \times 10^4$).

Colonization with *S. aureus* was not significantly associated with IgG levels to the *S. aureus* toxins in antenatal or cord blood or with IgA levels to the *S. aureus* toxins in breast milk. There was no relationship between carriage of *S. aureus* and maternal age.

Detection of toxin production by *S. aureus* isolates

Staphylococcus aureus isolates were tested for the production of SEB, SEC and TSST-1. Thirteen per cent of isolates produced SEB, 13% produced SEC and 24% produced TSST-1. Some isolates produced more than one toxin.

Women colonized with an isolate producing the toxin SEB were significantly more likely to have greater levels of anti-SEB IgG in cord blood compared with women not carrying an SEB-producing isolate ($P < 0.05$, Mann–Whitney *U*-test, Fig. 3). This trend was also found for SEC and TSST-1 but was not statistically significant.

Breast milk anti-SEC IgA levels were significantly greater if the mother was colonized with a *S. aureus* isolate producing SEC ($P < 0.05$). Mothers colonized with a TSST-1-producing isolate had greater levels of anti-TSST-1 IgA ($P=0.001$, Fig. 4). This association was also seen for SEB/anti-SEB IgA but was not statistically significant ($P=0.078$).

Discussion

An assumption of the common bacterial toxins hypothesis is that healthy infants will mount an effective immunological response when first exposed to specific toxin antigens, or they will succumb to their effects if they lack sufficient passive neutralizing antibody (Morris *et al.*, 1987). Passive




Fig. 3. Colonization with SEB-producing *Staphylococcus aureus* and its association with cord blood anti-SEB IgG levels.




Fig. 4. Colonization with TSST-1-producing *Staphylococcus aureus* and its association with breast milk anti-TSST-1 IgA levels.

immunity is conferred on the infant by placental transfer of IgG from the mother. Thus, the mother's IgG profile will determine the infant's immunological status at birth.

In this study, analysis of antenatal serum detected IgG to SEB, SEC and TSST-1 in 92%, 93.5% and 89.1% of samples, respectively. Analysis of the cord sera detected IgG to SEB, SEC and TSST in 92.7%, 96.9% and 92.7% of samples, respectively. These proportions are very similar to those found in healthy adults from Europe and the United States (Notermans *et al.*, 1983; Vergeront *et al.*, 1983; Loch *et al.*, 1986; Parsonnet *et al.*, 2005). There are a small proportion of infants who are born with no detectable passive immunity to one or more of the toxins and could be at risk of death if, on exposure to the toxins, they fail to mount an effective immune response. A study by Siarakas *et al.* (1999), comparing serum antibody levels to SEB between SIDS cases and an age-matched comparison group of infants, found that a significantly smaller proportion of the SIDS cases had specific IgG and IgM. This implies that the SIDS infants had low levels of specific maternal IgG, or that on exposure to the toxin, antibody had been consumed and the infant had no time to mount an effective immune response. Factors associated with an increased risk of SIDS are also associated with the induction of bacterial pyrogenic toxins (Harrison *et al.*, 2004).

Cord blood samples from male infants born to women in this study had significantly lower levels of anti-SEC IgG compared with female infants (Fig. 1). Male infants are at an increased risk of SIDS (Leach *et al.*, 1999). Afro-American women are reported to have significantly lower serum levels of IgG to TSST-1 (Parsonnet *et al.*, 2005) compared with white American women and the SIDS rate for infants born to Afro-American women is twice that of white American women (Mathews & MacDorman, 2007). However,

Japanese women also have lower levels of IgG to TSST-1 (Takahashi *et al.*, 2006) than European or white American women, but the SIDS rate in Japan is low (0.4/1000) (Fujita, 2002). Traditionally, Japanese infants have been placed to sleep in the supine position. At the end of 1998, only 20–30% of African American infants were placed to sleep supine. Sleeping supine greatly reduces the risk of SIDS (Leach *et al.*, 1999). Infants sleeping supine are less likely to be exposed to the factors that induce production of pyrogenic toxins in the upper respiratory tract (Blackwell *et al.*, 1999; Harrison *et al.*, 1999b; Molony *et al.*, 1999).

Transplacental transfer of IgG is upregulated in the last trimester of pregnancy (Holt & Jones, 2000), and in a full-term gestation, IgG concentration is usually higher in the cord serum than in the maternal serum (Wilmott-Pitcher *et al.*, 1980). Essery *et al.* (1999), analysing sera collected from pregnant women between 30 and 40 weeks gestation, found that levels of SEC- and TSST-1-specific IgG decreased with weeks of gestation but levels of SEB-specific IgG showed an increase. In our previous study (Harrison *et al.*, 2004), we reported that antibody levels to SEC and TSST-1 in cord blood samples showed a trend to increase with length of gestation. In this study, paired sample tests revealed that antibody levels to each of the toxins were significantly higher in cord sera than in the antenatal sera. The findings indicate active transport across the placenta, which might explain the decrease in the serum levels of some antitoxins in the latter stages of pregnancy reported by Essery *et al.* (1999).

Antenatal anti-SEC IgG showed a significant positive correlation with cord blood anti-SEC IgG (Fig. 2) and this was also the case for specific IgG to SEB and TSST-1. This relationship has also been found in a Japanese study (Takahashi *et al.*, 2006) and suggests that active immunity to the toxins is persistent or that low-level exposure to the antigens is common.

Humans become colonized with *S. aureus* soon after birth and some of the highest carriage rates are seen in the first few months of life (Blackwell *et al.*, 1999; Harrison *et al.*, 1999a; Peacock *et al.*, 2003; Lindberg *et al.*, 2004). Common sites of colonization include the upper respiratory tract and the intestinal tract. Carriage rates decline in infancy but both sites remain an ecological niche for *S. aureus* throughout life. Of the 96 women in this study, 30% were colonized with *S. aureus*, 21% had positive nasopharyngeal cultures and 14% had positive stool cultures. These carriage rates are similar to those demonstrated in previous studies (Peacock *et al.*, 2003; Harrison *et al.*, 2004; Lindberg *et al.*, 2004). Fifty per cent of the women had positive nasopharyngeal cultures on both sampling occasions and 23% had two positive stool cultures. Thirteen per cent of the isolates produced the toxin SEB, 13% produced SEC and 24% produced TSST-1. Intermittent carriage and undetected carriage of toxin-producing

S. aureus isolates from other body sites could explain as to why between 89% and 97% of the women have antibodies to one or more of the toxins even though, at any one time, carriage rates are only 30%.

There was a trend for women colonized with *S. aureus* toxin-producing isolates to have greater levels of antitoxin IgG in cord blood samples compared with women not colonized with a toxin-producing isolate. This relationship reached statistical significance ($P < 0.05$) for SEB.

Levels of passive immunity could be boosted through breast feeding. Women colonized with isolates producing TSST-1 had significantly greater levels of breast milk anti-TSST-1 IgA (Fig. 4, $P < 0.01$) and the same trend was shown for SEC/anti-SEC IgA ($P < 0.05$) and SEB/anti-SEB IgA ($P = 0.078$). Gordon *et al.* (1999) found that two-thirds of mothers colonized with *S. aureus* in the nose or throat had milk samples with levels of IgA to the toxins equal to or higher than the mean. There is conflicting evidence regarding the role of breast feeding in SIDS (Ford *et al.*, 1993; Fleming *et al.*, 1996) but some epidemiological studies report a protective, independent effect (Ford *et al.*, 1993; Alm *et al.*, 2002).

In conclusion, carriage of a *S. aureus* toxin-producing isolate in the later stages of pregnancy triggers the production of specific IgG and IgA, which in turn provides protection for the infant in the first few months of life. Experimental research is being carried out on the development of vaccines that induce IgG to the staphylococcal pyrogenic toxins (Cui *et al.*, 2005; Hu *et al.*, 2005; Narita *et al.*, 2008). Vaccination of pregnant women may be possible in the future and prevent infants being born with little or no passive immunity to the toxins.

Acknowledgements

This work was supported by a grant from the Peter Boizot Trust. We are grateful to the families who participated in this study, to Sandra Lively, Parentcraft Sister, for her help with recruitment and to the midwives for collection of cord blood samples.

References

- Alm B, Wennergren G, Norvenius SG, Skærvén R, Lagercrantz H, Helweg-Larsen K & Irgens LM (2002) Breast feeding and the sudden infant death syndrome in Scandinavia, 1992–95. *Arch Dis Child* **86**: 400–402.
- Al Madani O, Gordon AE, Weir DM, Raza MW, Busuttil A & Blackwell CC (1999) Pyrogenic toxins of *Staphylococcus aureus* in sudden unexpected nocturnal deaths in adults and older children: factors influencing the control of inflammatory responses to toxic shock syndrome toxins. *FEMS Immunol Med Microbiol* **25**: 207–219.
- Blackwell CC, Mackenzie DAC, James VS, Elton RA, Zorgani AA, Weir DM & Busuttil A (1999) Toxigenic bacteria and sudden infant death syndrome (SIDS): nasopharyngeal flora during the first year of life. *FEMS Immunol Med Microbiol* **25**: 51–55.
- Cui JC, Hu DL, Lin YC, Qian AD & Nakane A (2005) Immunization with glutathione S-transferase and mutant toxic shock syndrome toxin 1 fusion protein protects against *Staphylococcus aureus* infection. *FEMS Immunol Med Microbiol* **45**: 45–51.
- Essery SD, Raza MW, Zorgani A, MacKenzie DAC, James VS, Weir DM, Busuttil A, Hallam N & Blackwell C (1999) The protective effect of immunisation against diphtheria, pertussis and tetanus (DPT) in relation to sudden infant death syndrome. *FEMS Immunol Med Microbiol* **25**: 183–192.
- Fleming PJ, Blair PS, Bacon C, Bensley D, Smith I, Taylor E, Berry J, Golding J & Tripp J (1996) Environment of infants during sleep and risk of the sudden infant death syndrome: results of 1993–5 case-control study for confidential inquiry into stillbirths and deaths in infancy. Confidential enquiry into stillbirths and deaths regional coordinators and researchers. *Br Med J* **313**: 191–195.
- Ford RP, Taylor BJ, Mitchell EA, Enright SA, Stewart AW, Scragg R, Hassall IB, Barry DM, Allen EM & Roberts PA (1993) Breast feeding and the risk of sudden infant death syndrome. *Int J Epidemiol* **22**: 51–59.
- Fujita T (2002) Sudden infant death syndrome in Japan 1995–1998. *Forensic Sci Int* **130S**: S71–S77.
- Gordon AE, Saadi AT, MacKenzie DAC, Molony N, James VS, Weir DM, Busuttil A & Blackwell CC (1999) The protective effect of breast feeding in relation to sudden infant death syndrome (SIDS): III. Detection of IgA antibodies in human milk that bind to bacterial toxins implicated in SIDS. *FEMS Immunol Med Microbiol* **25**: 175–182.
- Harrison LM, Morris JA, Telford DR, Brown SM & Jones K (1999a) The nasopharyngeal flora in infancy: effects of age, gender, season, viral upper respiratory-tract infection and sleeping position. *FEMS Immunol Med Microbiol* **25**: 19–28.
- Harrison LM, Morris JA, Telford DR, Brown SM & Jones K (1999b) Sleeping position in infants over 6 months of age: implications for theories of sudden infant death syndrome. *FEMS Immunol Med Microbiol* **25**: 29–35.
- Harrison LM, Morris JA, Bishop LA, Lauder RM, Taylor CA & Telford DR (2004) Detection of specific antibodies in cord blood, infant and maternal saliva and breast milk to staphylococcal toxins implicated in sudden infant death syndrome (SIDS). *FEMS Immunol Med Microbiol* **42**: 94–104.
- Holt PG & Jones CA (2000) The development of the immune system during pregnancy and early life. *Allergy* **55**: 688–697.
- Hu DL, Cui JC, Omoe K, Sashinami H, Yokomizo Y, Shinagawa K & Nakane A (2005) A mutant of staphylococcal enterotoxin C devoid of bacterial superantigenic activity elicits a Th2 immune response for protection against *Staphylococcus aureus*. *Infect Immun* **73**: 174–180.
- Leach CEA, Blair PS, Fleming PJ, Smith IJ, Platt MW, Berry PJ & Golding J (the CESDI SUDI Research Group) (1999)

- Epidemiology of SIDS and explained sudden infant deaths. *Pediatrics* **140**: 143.
- Lindberg E, Adlerberth I, Hesselmar B, Saalman R, Strannegard I, Aberg N & Wold AE (2004) High rate of transfer of *Staphylococcus aureus* from parental skin to infant gut flora. *J Clin Microbiol* **42**: 530–534.
- Loch EG, Crass BA & Bergdoll MS (1986) *Staphylococcus aureus*-toxic shock syndrome toxin-1 antibody titres in serum of German women. *Arch Gynecol* **237**: 229–233.
- Malam JE, Carrick GF, Telford DR & Morris JA (1992) Staphylococcal toxins and sudden infant death syndrome. *J Clin Pathol* **45**: 716–721.
- Mathews TJ & MacDorman MF (2007) Infant mortality statistics from the 2004 period linked birth/infant death data set. *Natl Vital Stat Rep* **55**: 25.
- Molony N, Blackwell CC & Busuttil A (1999) The effect of prone posture on nasal temperature in children in relation to induction of staphylococcal toxins implicated in sudden infant death syndrome. *FEMS Immunol Med Microbiol* **25**: 109–113.
- Morris JA (1999) The common bacterial toxins hypothesis of sudden infant death syndrome. *FEMS Immunol Med Microbiol* **25**: 11–17.
- Morris JA, Haran D & Smith A (1987) Hypothesis: common bacterial toxins are a possible cause of the sudden infant death syndrome. *Med Hypotheses* **22**: 211–222.
- Murrell WG, Stewart BJ, O'Neill C, Siarakas S & Kariks S (1993) Enterotoxigenic bacteria in sudden infant death syndrome. *J Med Microbiol* **39**: 507–517.
- Narita K, Hu D, Tsuji T & Nakane A (2008) Intranasal immunisation of mutant toxic shock syndrome toxin 1 elicits systemic and mucosal immune response against *Staphylococcus aureus* infection. *FEMS Immunol Med Microbiol* **52**: 389–396.
- Newbould MJ, Malam J, McIlmurray JM, Morris JM, Telford DR & Barson AJ (1989) Immunohistological localisation of staphylococcal toxic shock syndrome toxin (TSST-1) antigen in sudden infant death syndrome. *J Clin Pathol* **42**: 935–939.
- Notermans S, van Leeuwen WJ, Dufrenne J & Tips PD (1983) Serum antibodies to enterotoxins produced by *Staphylococcus aureus* with special reference to enterotoxin F and toxic shock syndrome. *J Clin Microbiol* **18**: 1055–1060.
- Parsonnet J, Hansmann A, Delany ML et al. (2005) Prevalence of toxic shock syndrome toxin 1-producing *Staphylococcus aureus* and the presence of antibodies to this superantigen in menstruating women. *J Clin Microbiol* **43**: 4628–4634.
- Peacock SJ, Justice A, Griffiths D, de Silva GDI, Kantzanou MN, Crook D, Sleeman K & Day NP (2003) Determinants of acquisition and carriage of *Staphylococcus aureus* in infancy. *J Clin Microbiol* **41**: 5718–5725.
- Siarakas S, Brown AJ & Murrell WG (1999) Immunological evidence for a bacterial toxin aetiology in sudden infant death syndrome. *FEMS Immunol Med Microbiol* **25**: 37–50.
- Takahashi N, Hattori M, Miwa K & Nishida H (2006) Antibodies against superantigenic exotoxins produced by *Staphylococcus aureus* in sera from mothers and their infants' cord blood. *Am J Perinat* **23**: 413–419.
- Telford DR, Morris JA, Hughes P, Conway AR, Lee S, Barson AJ & Drucker DB (1989) The nasopharyngeal bacterial flora in the sudden infant death syndrome. *J Infect* **18**: 125–130.
- Vergeront JM, Stoltz SJ, Crass BA, Nelson DB, Davis JP & Bergdoll MS (1983) Prevalence of serum antibody to staphylococcal enterotoxin F among Wisconsin residents: implications for toxic shock syndrome. *J Infect Dis* **148**: 692–698.
- Wilmott-Pitcher RW, Hindocha P & Wood CB (1980) The placental transfer of IgG subclasses in human pregnancy. *Clin Exp Immunol* **41**: 303–308.
- Zorgani A, Essery SD, Al Madani O et al. (1999) Detection of pyrogenic toxins of *Staphylococcus aureus* in sudden infant death syndrome. *FEMS Immunol Med Microbiol* **25**: 103–108.

Author Query Form

Journal FEMSIM
Article 463

Dear Author,

During the copy-editing of your paper, the following queries arose. Please respond to these by marking up your proofs with the necessary changes/additions. Please write your answers clearly on the query sheet if there is insufficient space on the page proofs. If returning the proof by fax do not write too close to the paper's edge. Please remember that illegible mark-ups may delay publication.