

Search for Pair Production of Doubly Charged Higgs Bosons in the $H^{++}H^{--} \rightarrow \mu^+\mu^+\mu^-\mu^-$ Final State

V. M. Abazov,³⁶ B. Abbott,⁷⁵ M. Abolins,⁶⁵ B. S. Acharya,²⁹ M. Adams,⁵¹ T. Adams,⁴⁹ E. Aguilo,⁶ S. H. Ahn,³¹ M. Ahsan,⁵⁹ G. D. Alexeev,³⁶ G. Alkhazov,⁴⁰ A. Alton,^{64,*} G. Alverson,⁶³ G. A. Alves,² M. Anastasoae,³⁵ L. S. Ancu,³⁵ T. Andeen,⁵³ S. Anderson,⁴⁵ B. Andrieu,¹⁷ M. S. Anzels,⁵³ M. Aoki,⁵⁰ Y. Arnaud,¹⁴ M. Arov,⁶⁰ M. Arthaud,¹⁸ A. Askew,⁴⁹ B. Åsman,⁴¹ A. C. S. Assis Jesus,³ O. Atramentov,⁴⁹ C. Avila,⁸ C. Ay,²⁴ F. Badaud,¹³ A. Baden,⁶¹ L. Bagby,⁵⁰ B. Baldin,⁵⁰ D. V. Bandurin,⁵⁹ P. Banerjee,²⁹ S. Banerjee,²⁹ E. Barberis,⁶³ A.-F. Barfuss,¹⁵ P. Bargassa,⁸⁰ P. Baringer,⁵⁸ J. Barreto,² J. F. Bartlett,⁵⁰ U. Bassler,¹⁸ D. Bauer,⁴³ S. Beale,⁶ A. Bean,⁵⁸ M. Begalli,³ M. Begel,⁷³ C. Belanger-Champagne,⁴¹ L. Bellantoni,⁵⁰ A. Bellavance,⁵⁰ J. A. Benitez,⁶⁵ S. B. Beri,²⁷ G. Bernardi,¹⁷ R. Bernhard,²³ I. Bertram,⁴² M. Besançon,¹⁸ R. Beuselinck,⁴³ V. A. Bezzubov,³⁹ P. C. Bhat,⁵⁰ V. Bhatnagar,²⁷ C. Biscarat,²⁰ G. Blazey,⁵² F. Blekman,⁴³ S. Blessing,⁴⁹ D. Bloch,¹⁹ K. Bloom,⁶⁷ A. Boehnlein,⁵⁰ D. Boline,⁶² T. A. Bolton,⁵⁹ G. Borissov,⁴² T. Bose,⁷⁷ A. Brandt,⁷⁸ R. Brock,⁶⁵ G. Brooijmans,⁷⁰ A. Bross,⁵⁰ D. Brown,⁸¹ N. J. Buchanan,⁴⁹ D. Buchholz,⁵³ M. Buehler,⁸¹ V. Buescher,²² V. Bunichev,³⁸ S. Burdin,^{42,+} S. Burke,⁴⁵ T. H. Burnett,⁸² C. P. Buszello,⁴³ J. M. Butler,⁶² P. Calfayan,²⁵ S. Calvet,¹⁶ J. Cammin,⁷¹ W. Carvalho,³ B. C. K. Casey,⁵⁰ H. Castilla-Valdez,³³ S. Chakrabarti,¹⁸ D. Chakraborty,⁵² K. Chan,⁶ K. M. Chan,⁵⁵ A. Chandra,⁴⁸ F. Charles,¹⁹ E. Cheu,⁴⁵ F. Chevallier,¹⁴ D. K. Cho,⁶² S. Choi,³² B. Choudhary,²⁸ L. Christofek,⁷⁷ T. Christoudias,⁴³ S. Cihangir,⁵⁰ D. Claes,⁶⁷ Y. Coadou,⁶ M. Cooke,⁸⁰ W. E. Cooper,⁵⁰ M. Corcoran,⁸⁰ F. Couderc,¹⁸ M.-C. Cousinou,¹⁵ S. Crépe-Renaudin,¹⁴ D. Cutts,⁷⁷ M. Ćwiok,³⁰ H. da Motta,² A. Das,⁴⁵ G. Davies,⁴³ K. De,⁷⁸ S. J. de Jong,³⁵ E. De La Cruz-Burelo,⁶⁴ C. De Oliveira Martins,³ J. D. Degenhardt,⁶⁴ F. Déliot,¹⁸ M. Demarteau,⁵⁰ R. Demina,⁷¹ D. Denisov,⁵⁰ S. P. Denisov,³⁹ S. Desai,⁵⁰ H. T. Diehl,⁵⁰ M. Diesburg,⁵⁰ A. Dominguez,⁶⁷ H. Dong,⁷² L. V. Dudko,³⁸ L. Dufлот,¹⁶ S. R. Dugad,²⁹ D. Duggan,⁴⁹ A. Duperrin,¹⁵ J. Dyer,⁶⁵ A. Dyshkant,⁵² M. Eads,⁶⁷ D. Edmunds,⁶⁵ J. Ellison,⁴⁸ V. D. Elvira,⁵⁰ Y. Enari,⁷⁷ S. Eno,⁶¹ P. Ermolov,³⁸ H. Evans,⁵⁴ A. Evdokimov,⁷³ V. N. Evdokimov,³⁹ A. V. Ferapontov,⁵⁹ T. Ferbel,⁷¹ F. Fiedler,²⁴ F. Filthaut,³⁵ W. Fisher,⁵⁰ H. E. Fisk,⁵⁰ M. Fortner,⁵² H. Fox,⁴² S. Fu,⁵⁰ S. Fuess,⁵⁰ T. Gadfort,⁷⁰ C. F. Galea,³⁵ E. Gallas,⁵⁰ C. Garcia,⁷¹ A. Garcia-Bellido,⁸² V. Gavrilov,³⁷ P. Gay,¹³ W. Geist,¹⁹ D. Gelé,¹⁹ C. E. Gerber,⁵¹ Y. Gershtein,⁴⁹ D. Gillberg,⁶ G. Ginther,⁷¹ N. Gollub,⁴¹ B. Gómez,⁸ A. Goussiou,⁸² P. D. Grannis,⁷² H. Greenlee,⁵⁰ Z. D. Greenwood,⁶⁰ E. M. Gregores,⁴ G. Grenier,²⁰ Ph. Gris,¹³ J.-F. Grivaz,¹⁶ A. Grohsjean,²⁵ S. Grünendahl,⁵⁰ M. W. Grünewald,³⁰ F. Guo,⁷² J. Guo,⁷² G. Gutierrez,⁵⁰ P. Gutierrez,⁷⁵ A. Haas,⁷⁰ N. J. Hadley,⁶¹ P. Haefner,²⁵ S. Hagopian,⁴⁹ J. Haley,⁶⁸ I. Hall,⁶⁵ R. E. Hall,⁴⁷ L. Han,⁷ K. Harder,⁴⁴ A. Harel,⁷¹ R. Harrington,⁶³ J. M. Hauptman,⁵⁷ R. Hauser,⁶⁵ J. Hays,⁴³ T. Hebbeker,²¹ D. Hedin,⁵² J. G. Hegeman,³⁴ J. M. Heinmiller,⁵¹ A. P. Heinson,⁴⁸ U. Heintz,⁶² C. Hensel,⁵⁸ K. Herner,⁷² G. Hesketh,⁶³ M. D. Hildreth,⁵⁵ R. Hirosky,⁸¹ J. D. Hobbs,⁷² B. Hoeneisen,¹² H. Hoeth,²⁶ M. Hohlfield,²² S. J. Hong,³¹ S. Hossain,⁷⁵ P. Houben,³⁴ Y. Hu,⁷² Z. Hubacek,¹⁰ V. Hynek,⁹ I. Iashvili,⁶⁹ R. Illingworth,⁵⁰ A. S. Ito,⁵⁰ S. Jabeen,⁶² M. Jaffré,¹⁶ S. Jain,⁷⁵ K. Jakobs,²³ C. Jarvis,⁶¹ R. Jesik,⁴³ K. Johns,⁴⁵ C. Johnson,⁷⁰ M. Johnson,⁵⁰ A. Jonckheere,⁵⁰ P. Jonsson,⁴³ A. Juste,⁵⁰ E. Kajfasz,¹⁵ A. M. Kalinin,³⁶ J. M. Kalk,⁶⁰ S. Kappler,²¹ D. Karmanov,³⁸ P. A. Kasper,⁵⁰ I. Katsanos,⁷⁰ D. Kau,⁴⁹ V. Kaushik,⁷⁸ R. Kehoe,⁷⁹ S. Kermiche,¹⁵ N. Khalatyan,⁵⁰ A. Khanov,⁷⁶ A. Kharchilava,⁶⁹ Y. M. Kharzheev,³⁶ D. Khatidze,⁷⁰ T. J. Kim,³¹ M. H. Kirby,⁵³ M. Kirsch,²¹ B. Klima,⁵⁰ J. M. Kohli,²⁷ J.-P. Konrath,²³ V. M. Korablev,³⁹ A. V. Kozelov,³⁹ J. Kraus,⁶⁵ D. Krop,⁵⁴ T. Kuhl,²⁴ A. Kumar,⁶⁹ A. Kupco,¹¹ T. Kurča,²⁰ J. Kvita,⁹ F. Lacroix,¹³ D. Lam,⁵⁵ S. Lammers,⁷⁰ G. Landsberg,⁷⁷ P. Lebrun,²⁰ W. M. Lee,⁵⁰ A. Leflat,³⁸ J. Lellouch,¹⁷ J. Leveque,⁴⁵ J. Li,⁷⁸ L. Li,⁴⁸ Q. Z. Li,⁵⁰ S. M. Lietti,⁵ J. G. R. Lima,⁵² D. Lincoln,⁵⁰ J. Linnemann,⁶⁵ V. V. Lipaev,³⁹ R. Lipton,⁵⁰ Y. Liu,⁷ Z. Liu,⁶ A. Lobodenko,⁴⁰ M. Lokajicek,¹¹ P. Love,⁴² H. J. Lubatti,⁸² R. Luna,³ A. L. Lyon,⁵⁰ A. K. A. Maciel,² D. Mackin,⁸⁰ R. J. Madaras,⁴⁶ P. Mättig,²⁶ C. Magass,²¹ A. Magerkurth,⁶⁴ P. K. Mal,⁸² H. B. Malbouisson,³ S. Malik,⁶⁷ V. L. Malyshev,³⁶ H. S. Mao,⁵⁰ Y. Maravin,⁵⁹ B. Martin,¹⁴ R. McCarthy,⁷² A. Melnitchouk,⁶⁶ L. Mendoza,⁸ P. G. Mercadante,⁵ M. Merkin,³⁸ K. W. Merritt,⁵⁰ A. Meyer,²¹ J. Meyer,^{22,§} T. Millet,²⁰ J. Mitrevski,⁷⁰ J. Molina,³ R. K. Mommsen,⁴⁴ N. K. Mondal,²⁹ R. W. Moore,⁶ T. Moulík,⁵⁸ G. S. Muanza,²⁰ M. Mulders,⁵⁰ M. Mulhearn,⁷⁰ O. Mundal,²² L. Mundim,³ E. Nagy,¹⁵ M. Naimuddin,⁵⁰ M. Narain,⁷⁷ N. A. Naumann,³⁵ H. A. Neal,⁶⁴ J. P. Negret,⁸ P. Neustroev,⁴⁰ H. Nilsen,²³ H. Nogima,³ S. F. Novaes,⁵ T. Nunnemann,²⁵ V. O'Dell,⁵⁰ D. C. O'Neil,⁶ G. Obrant,⁴⁰ C. Ochando,¹⁶ D. Onoprienko,⁵⁹ N. Oshima,⁵⁰ N. Osman,⁴³ J. Osta,⁵⁵ R. Otec,¹⁰ G. J. Otero y Garzón,⁵⁰ M. Owen,⁴⁴ P. Padley,⁸⁰ M. Pangilinan,⁷⁷ N. Parashar,⁵⁶ S.-J. Park,⁷¹ S. K. Park,³¹ J. Parsons,⁷⁰ R. Partridge,⁷⁷ N. Parua,⁵⁴ A. Patwa,⁷³ G. Pawloski,⁸⁰ B. Penning,²³ M. Perfilov,³⁸ K. Peters,⁴⁴ Y. Peters,²⁶ P. Pétróff,¹⁶ M. Petteni,⁴³ R. Piegaia,¹ J. Piper,⁶⁵ M.-A. Pleier,²² P. L. M. Podesta-Lerma,^{33,‡} V. M. Podstavkov,⁵⁰ Y. Pogorelov,⁵⁵ M.-E. Pol,² P. Polozov,³⁷ B. G. Pope,⁶⁵ A. V. Popov,³⁹ C. Potter,⁶

W. L. Prado da Silva,³ H. B. Prosper,⁴⁹ S. Protopopescu,⁷³ J. Qian,⁶⁴ A. Quadt,^{22,§} B. Quinn,⁶⁶ A. Rakitine,⁴² M. S. Rangel,² K. Ranjan,²⁸ P. N. Ratoff,⁴² P. Renkel,⁷⁹ S. Reucroft,⁶³ P. Rich,⁴⁴ J. Rieger,⁵⁴ M. Rijssenbeek,⁷² I. Ripp-Baudot,¹⁹ F. Rizatdinova,⁷⁶ S. Robinson,⁴³ R. F. Rodrigues,³ M. Rominsky,⁷⁵ C. Royon,¹⁸ P. Rubinov,⁵⁰ R. Ruchti,⁵⁵ G. Safronov,³⁷ G. Sajot,¹⁴ A. Sánchez-Hernández,³³ M. P. Sanders,¹⁷ A. Santoro,³ G. Savage,⁵⁰ L. Sawyer,⁶⁰ T. Scanlon,⁴³ D. Schaile,²⁵ R. D. Schamberger,⁷² Y. Scheglov,⁴⁰ H. Schellman,⁵³ T. Schliephake,²⁶ C. Schwanenberger,⁴⁴ A. Schwartzman,⁶⁸ R. Schwienhorst,⁶⁵ J. Sekaric,⁴⁹ H. Severini,⁷⁵ E. Shabalina,⁵¹ M. Shamim,⁵⁹ V. Shary,¹⁸ A. A. Shchukin,³⁹ R. K. Shivpuri,²⁸ V. Siccaldi,¹⁹ V. Simak,¹⁰ V. Sirotenko,⁵⁰ P. Skubic,⁷⁵ P. Slattery,⁷¹ D. Smirnov,⁵⁵ G. R. Snow,⁶⁷ J. Snow,⁷⁴ S. Snyder,⁷³ S. Söldner-Rembold,⁴⁴ L. Sonnenschein,¹⁷ A. Sopczak,⁴² M. Sosebee,⁷⁸ K. Soustruznik,⁹ B. Spurlock,⁷⁸ J. Stark,¹⁴ J. Steele,⁶⁰ V. Stolin,³⁷ D. A. Stoyanova,³⁹ J. Strandberg,⁶⁴ S. Strandberg,⁴¹ M. A. Strang,⁶⁹ E. Strauss,⁷² M. Strauss,⁷⁵ R. Ströhmer,²⁵ D. Strom,⁵³ L. Stutte,⁵⁰ S. Sumowidagdo,⁴⁹ P. Svoisky,⁵⁵ A. Sznajder,³ P. Tamburello,⁴⁵ A. Tanasijczuk,¹ W. Taylor,⁶ J. Temple,⁴⁵ B. Tiller,²⁵ F. Tissandier,¹³ M. Titov,¹⁸ V. V. Tokmenin,³⁶ T. Toole,⁶¹ I. Torchiani,²³ T. Trefzger,²⁴ D. Tsybychev,⁷² B. Tuchming,¹⁸ C. Tully,⁶⁸ P. M. Tuts,⁷⁰ R. Unalan,⁶⁵ L. Uvarov,⁴⁰ S. Uvarov,⁴⁰ S. Uzunyan,⁵² B. Vachon,⁶ P. J. van den Berg,³⁴ R. Van Kooten,⁵⁴ W. M. van Leeuwen,³⁴ N. Varelas,⁵¹ E. W. Varnes,⁴⁵ I. A. Vasilyev,³⁹ M. Vaupel,²⁶ P. Verdier,²⁰ L. S. Vertogradov,³⁶ M. Verzocchi,⁵⁰ F. Villeneuve-Seguier,⁴³ P. Vint,⁴³ P. Vokac,¹⁰ E. Von Toerne,⁵⁹ M. Voutilainen,^{68,||} R. Wagner,⁶⁸ H. D. Wahl,⁴⁹ L. Wang,⁶¹ M. H. L. S. Wang,⁵⁰ J. Warchol,⁵⁵ G. Watts,⁸² M. Wayne,⁵⁵ G. Weber,²⁴ M. Weber,⁵⁰ L. Welty-Rieger,⁵⁴ A. Wenger,^{23,¶} N. Vermes,²² M. Wetstein,⁶¹ A. White,⁷⁸ D. Wicke,²⁶ G. W. Wilson,⁵⁸ S. J. Wimpenny,⁴⁸ M. Wobisch,⁶⁰ D. R. Wood,⁶³ T. R. Wyatt,⁴⁴ Y. Xie,⁷⁷ S. Yacoob,⁵³ R. Yamada,⁵⁰ M. Yan,⁶¹ T. Yasuda,⁵⁰ Y. A. Yatsunenko,³⁶ K. Yip,⁷³ H. D. Yoo,⁷⁷ S. W. Youn,⁵³ J. Yu,⁷⁸ A. Zatserklyaniy,⁵² C. Zeitnitz,²⁶ T. Zhao,⁸² B. Zhou,⁶⁴ J. Zhu,⁷² M. Zielinski,⁷¹ D. Zieminska,⁵⁴ A. Zieminski,^{54,**} L. Zivkovic,⁷⁰ V. Zutshi,⁵² and E. G. Zverev³⁸

(D0 Collaboration)

¹*Universidad de Buenos Aires, Buenos Aires, Argentina*²*LAFEX, Centro Brasileiro de Pesquisas Físicas, Rio de Janeiro, Brazil*³*Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil*⁴*Universidade Federal do ABC, Santo André, Brazil*⁵*Instituto de Física Teórica, Universidade Estadual Paulista, São Paulo, Brazil*⁶*University of Alberta, Edmonton, Alberta, Canada,**Simon Fraser University, Burnaby, British Columbia, Canada,**York University, Toronto, Ontario, Canada,**and McGill University, Montreal, Quebec, Canada*⁷*University of Science and Technology of China, Hefei, People's Republic of China*⁸*Universidad de los Andes, Bogotá, Colombia*⁹*Center for Particle Physics, Charles University, Prague, Czech Republic*¹⁰*Czech Technical University, Prague, Czech Republic*¹¹*Center for Particle Physics, Institute of Physics, Academy of Sciences of the Czech Republic, Prague, Czech Republic*¹²*Universidad San Francisco de Quito, Quito, Ecuador*¹³*LPC, Université Blaise Pascal, CNRS/IN2P3, Clermont, France*¹⁴*LPSC, Université Joseph Fourier Grenoble 1, CNRS/IN2P3, Institut National Polytechnique de Grenoble, France*¹⁵*CPPM, IN2P3/CNRS, Université de la Méditerranée, Marseille, France*¹⁶*LAL, Université Paris-Sud, IN2P3/CNRS, Orsay, France*¹⁷*LPNHE, IN2P3/CNRS, Universités Paris VI and VII, Paris, France*¹⁸*DAPNIA/Service de Physique des Particules, CEA, Saclay, France*¹⁹*IPHC, Université Louis Pasteur et Université de Haute Alsace, CNRS/IN2P3, Strasbourg, France*²⁰*IPNL, Université Lyon 1, CNRS/IN2P3, Villeurbanne, France, and**Université de Lyon, Lyon, France*²¹*III. Physikalisches Institut A, RWTH Aachen, Aachen, Germany*²²*Physikalisches Institut, Universität Bonn, Bonn, Germany*²³*Physikalisches Institut, Universität Freiburg, Freiburg, Germany*²⁴*Institut für Physik, Universität Mainz, Mainz, Germany*²⁵*Ludwig-Maximilians-Universität München, München, Germany*²⁶*Fachbereich Physik, University of Wuppertal, Wuppertal, Germany*²⁷*Panjab University, Chandigarh, India*²⁸*Delhi University, Delhi, India*²⁹*Tata Institute of Fundamental Research, Mumbai, India*

- ³⁰University College Dublin, Dublin, Ireland
³¹Korea Detector Laboratory, Korea University, Seoul, Korea
³²SungKyunKwan University, Suwon, Korea
³³CINVESTAV, Mexico City, Mexico
³⁴FOM-Institute NIKHEF and University of Amsterdam/NIKHEF, Amsterdam, The Netherlands
³⁵Radboud University Nijmegen/NIKHEF, Nijmegen, The Netherlands
³⁶Joint Institute for Nuclear Research, Dubna, Russia
³⁷Institute for Theoretical and Experimental Physics, Moscow, Russia
³⁸Moscow State University, Moscow, Russia
³⁹Institute for High Energy Physics, Protvino, Russia
⁴⁰Petersburg Nuclear Physics Institute, St. Petersburg, Russia
⁴¹Lund University, Lund, Sweden, Royal Institute of Technology and Stockholm University, Stockholm, Sweden, and Uppsala University, Uppsala, Sweden
⁴²Lancaster University, Lancaster, United Kingdom
⁴³Imperial College, London, United Kingdom
⁴⁴University of Manchester, Manchester, United Kingdom
⁴⁵University of Arizona, Tucson, Arizona 85721, USA
⁴⁶Lawrence Berkeley National Laboratory and University of California, Berkeley, California 94720, USA
⁴⁷California State University, Fresno, California 93740, USA
⁴⁸University of California, Riverside, California 92521, USA
⁴⁹Florida State University, Tallahassee, Florida 32306, USA
⁵⁰Fermi National Accelerator Laboratory, Batavia, Illinois 60510, USA
⁵¹University of Illinois at Chicago, Chicago, Illinois 60607, USA
⁵²Northern Illinois University, DeKalb, Illinois 60115, USA
⁵³Northwestern University, Evanston, Illinois 60208, USA
⁵⁴Indiana University, Bloomington, Indiana 47405, USA
⁵⁵University of Notre Dame, Notre Dame, Indiana 46556, USA
⁵⁶Purdue University Calumet, Hammond, Indiana 46323, USA
⁵⁷Iowa State University, Ames, Iowa 50011, USA
⁵⁸University of Kansas, Lawrence, Kansas 66045, USA
⁵⁹Kansas State University, Manhattan, Kansas 66506, USA
⁶⁰Louisiana Tech University, Ruston, Louisiana 71272, USA
⁶¹University of Maryland, College Park, Maryland 20742, USA
⁶²Boston University, Boston, Massachusetts 02215, USA
⁶³Northeastern University, Boston, Massachusetts 02115, USA
⁶⁴University of Michigan, Ann Arbor, Michigan 48109, USA
⁶⁵Michigan State University, East Lansing, Michigan 48824, USA
⁶⁶University of Mississippi, University, Mississippi 38677, USA
⁶⁷University of Nebraska, Lincoln, Nebraska 68588, USA
⁶⁸Princeton University, Princeton, New Jersey 08544, USA
⁶⁹State University of New York, Buffalo, New York 14260, USA
⁷⁰Columbia University, New York, New York 10027, USA
⁷¹University of Rochester, Rochester, New York 14627, USA
⁷²State University of New York, Stony Brook, New York 11794, USA
⁷³Brookhaven National Laboratory, Upton, New York 11973, USA
⁷⁴Langston University, Langston, Oklahoma 73050, USA
⁷⁵University of Oklahoma, Norman, Oklahoma 73019, USA
⁷⁶Oklahoma State University, Stillwater, Oklahoma 74078, USA
⁷⁷Brown University, Providence, Rhode Island 02912, USA
⁷⁸University of Texas, Arlington, Texas 76019, USA
⁷⁹Southern Methodist University, Dallas, Texas 75275, USA
⁸⁰Rice University, Houston, Texas 77005, USA
⁸¹University of Virginia, Charlottesville, Virginia 22901, USA
⁸²University of Washington, Seattle, Washington 98195, USA

(Received 11 March 2008; published 13 August 2008)

We report the results of a search for pair production of doubly charged Higgs bosons via $p\bar{p} \rightarrow H^{++}H^{--}X \rightarrow \mu^+\mu^+\mu^-\mu^-X$ at $\sqrt{s} = 1.96$ TeV. We use a data set corresponding to an integrated luminosity of 1.1 fb^{-1} collected from 2002 to 2006 by the D0 detector at the Fermilab Tevatron Collider. In the absence of an excess above the standard model background, lower mass limits of $M(H_L^{\pm\pm}) > 150 \text{ GeV}/c^2$

and $M(H_R^{\pm\pm}) > 127 \text{ GeV}/c^2$ at 95% C.L. are set, respectively, for left-handed and right-handed doubly charged Higgs bosons assuming a 100% branching ratio into muons.

DOI: 10.1103/PhysRevLett.101.071803

PACS numbers: 14.80.Cp, 13.85.Rm

In the standard model (SM) of electroweak interactions, elementary fermions and bosons acquire mass via a weak isospin scalar doublet. This mechanism results in the existence of an additional particle, the Higgs boson, which has not yet been observed. Extensions of the Higgs sector involving higher isospin multiplets predict the existence of doubly charged Higgs bosons, which can be relatively light and hence accessible at current experimental facilities. Doubly charged Higgs bosons appear in many scenarios such as left-right symmetric models [1], Higgs triplet models [2], and little Higgs models [3]. At the Fermilab Tevatron Collider, the two main production mechanisms are pair production via $p\bar{p} \rightarrow Z/\gamma^* X \rightarrow H^{++}H^{--}X$ and single production via WW fusion, $p\bar{p} \rightarrow W^\pm W^\pm X \rightarrow H^{\pm\pm}X$. However, higher isospin Higgs multiplets are generally severely constrained by $\rho \equiv m_W^2/(\cos\theta_W m_Z)^2 = 1$ at tree level. The existing phenomenological and theoretical constraints are easily satisfied when the $W^\pm W^\pm \rightarrow H^{\pm\pm}X$ coupling is vanishing [4]. If the H^{++} coupling to W boson pairs is suppressed, the dominant final states are expected to be like-sign lepton pairs. Left-handed ($H_L^{\pm\pm}$) and right-handed ($H_R^{\pm\pm}$) states are distinguished by their coupling to left-handed and right-handed leptons, respectively. While there is a prejudice that decays to heavier leptons would be favored, the relative branching ratios into different lepton flavors cannot be predicted by the theory. Therefore, it is customary to consider benchmark scenarios assuming a 100% branching ratio into a particular lepton flavor. The pair production cross section for left-handed doubly charged Higgs bosons for $100 \leq M(H^{\pm\pm}) \leq 200 \text{ GeV}/c^2$ is about a factor of 2 larger than that for the right-handed states due to different couplings to the intermediate Z boson [5]. Previous searches for $H^{\pm\pm}$ have been performed by the LEP Collaborations [6] in e^+e^- collisions and by the D0 [7] and CDF [8] Collaborations at the Tevatron $p\bar{p}$ collider. This Letter presents the results of a direct search for $p\bar{p} \rightarrow H^{++}H^{--}X$ with $H^{\pm\pm} \rightarrow \mu^\pm\mu^\pm$ by the D0 Collaboration with improved sensitivity.

The main D0 detector systems are a central tracking system, a liquid-argon and uranium calorimeter, and a muon detector [9]. The central tracking system consists of the silicon microstrip tracker (SMT) and the central fiber tracker (CFT) surrounded by a 2 T solenoidal magnet, with designs optimized for tracking and vertexing capability at pseudorapidity [10] $|\eta| < 3$ and $|\eta| < 2.5$, respectively. The liquid-argon and uranium calorimeter has a central calorimeter covering a region up to $|\eta| \approx 1.1$ and two end calorimeters extending the coverage to $|\eta| \approx 4.2$ [11]. The muon detector has layers of proportional drift tubes and scintillation counters before and after a 1.8 T iron toroid

[12]. This analysis is based on the run II data set collected with the D0 detector at $\sqrt{s} = 1.96 \text{ TeV}$ from April 2002 to February 2006 corresponding to 1.1 fb^{-1} . Events are collected using a suite of dimuon and single muon triggers.

In the previous D0 analysis [7], two like-sign muons were required in the final state. In this analysis, we require a third muon, which increases the sensitivity by decreasing backgrounds. We follow five steps to select events. In the first step (S1), events are required to have at least two muons. Each muon must have a transverse momentum $p_T > 15 \text{ GeV}/c$ and $|\eta| < 2.0$. Muons are selected using patterns of hits in the wire chambers and scintillators in the muon system. Each muon must be matched to a track in the




FIG. 1 (color online). Distributions of the dimuon invariant mass and $\Delta\phi$ between the two muons for data compared to the sum of MC backgrounds after selection S1 (a),(b), the preselection S1 with the like-sign requirement S4 (c),(d), and the final requirement S1–S5 (e),(f). The signal expected for a left-handed $H^{\pm\pm}$, with $M(H^{\pm\pm}) = 140 \text{ GeV}/c^2$, is also shown by the open histogram (e),(f).

central tracker with at least five hits in the CFT layers and at least two hits in the SMT layers. Muons from cosmic rays are removed by using a timing information on the hits in the scintillator layers.

In the second step (S_2), isolation criteria based on the calorimeter and tracking information are applied to remove the background from multijet production with muons originating from in-flight decay of pions or kaons or from semileptonic decays of B or D mesons. The sum of the transverse energies of the calorimeter cells in an annulus of radius $0.1 < \mathcal{R} < 0.4$, where $\mathcal{R} = \sqrt{(\Delta\phi)^2 + (\Delta\eta)^2}$ and ϕ is the azimuthal angle, around the muon direction is required to be less than 2.5 GeV. A similar condition is defined for the scalar sum of the p_T of all tracks, excluding the muon in a cone of radius $\mathcal{R} = 0.5$ centered around the muon, which must be less than 2.5 GeV/ c .

Selection S_3 reduces the remaining $Z \rightarrow \mu^+ \mu^-$ and multijet backgrounds. The azimuthal angle $\Delta\phi$ between at least one pair of muons is required to be less than 2.5 radians, since the two muons from Z boson decays are mostly back-to-back. This requirement also rejects a fraction of the multijet background with nearly back-to-back muons.

Selection S_4 requires at least two muons to be of like sign. The final selection (S_5) requires a third muon, satisfying the S_1 selection and the isolation selection criteria S_2 but without the minimum hit requirement on the central track.

The dominant background in this analysis arises from electroweak processes where real high p_T muons are created from W or Z boson decays as well as nonisolated muons originating from jets. The SM backgrounds and signal processes are generated with PYTHIA [13] and normalized using the theoretical cross section. The $Z/\gamma^* \rightarrow \ell^+ \ell^-$ cross section is calculated at next-to-next-to-leading order (NNLO) [14]. The $t\bar{t}$ cross section is calculated at NNLO [15] and the WW , ZZ , and WZ cross sections are calculated with MCFM [16] at next-to-leading order (NLO).

All samples are processed through the D0 detector simulation based on GEANT [17] and the same reconstruction software as for the data. The muon reconstruction and isolation efficiencies differ between Monte Carlo (MC) simulations and data, and these differences are corrected. Trigger efficiency corrections are not applied to the MC samples. Instead, the MC samples are normalized to the data using the Z boson mass peak at the selection level S_2 .

Another important background comes from multijet production, mainly $b\bar{b}$ events decaying semileptonically into muons that appear isolated. The multijet background is derived from the data sample with nonisolated muons obtained by inverting the isolation requirements for both muons after selection S_1 . The efficiency of the isolation requirement is assumed to be identical for multijet events with like-sign and opposite-sign muon pairs. It is also assumed that all like-sign events after subtracting SM backgrounds are multijet events. The SM backgrounds are subtracted in the following samples used for the multijet background determination. The total number of multijet events before the isolation requirement (4492 ± 120) is then given by the number of nonisolated events for all charge combinations multiplied by the ratio of the total number of events to the number of nonisolated events in the like-sign sample. The number of multijet events after the isolation requirement (194 ± 18) is obtained by multiplying this number with the isolation efficiency ($4.3 \pm 0.5\%$), given by the ratio of isolated to all like-sign multijet events.

A second instrumental background arises from $Z/\gamma^* \rightarrow \mu^+ \mu^-$ events in which the charge of one of the muons is misidentified. The first source of charge misidentification is due to fewer CFT layers at large η and a consequent increase in the charge misidentification probability. The second source affects very high p_T tracks for which the uncertainty on the measured curvature can cause charge misidentification. The charge misidentification rate is obtained by dividing the number of like-sign events (S_1 , S_2 , and S_4) by the number of events without the like-sign

TABLE I. The expected number of events for a signal with $M(H_L^{\pm\pm}) = 140$ GeV/ c^2 and background and the number of observed events after each selection step. The like-sign requirement is applied in the last two columns. The statistical and systematical uncertainties are combined in the table.

Selection	Preselection S_1	Isolation S_2	$\Delta\phi < 2.5$ S_3	Like sign S_4	Third muon S_5	S_1 & S_4	S_2 & S_4
$Z/\gamma^* \rightarrow \mu^+ \mu^-$	$69\,181 \pm 4642$	$58\,264 \pm 3910$	4936 ± 333	5.3 ± 1.6	<0.01	84 ± 24	42 ± 12
Multijet	4492 ± 120	194 ± 18	18 ± 2	6.3 ± 0.8	0.2 ± 0.1	1620 ± 34	70 ± 5
$Z/\gamma^* \rightarrow \tau^+ \tau^-$	328 ± 25	269 ± 21	20 ± 3	<0.01	<0.01	3.2 ± 1.3	0.2 ± 0.3
$t\bar{t}$	38 ± 3	20 ± 1	14 ± 1	0.03 ± 0.01	<0.01	6.6 ± 0.5	<0.1
WW	40 ± 3	34 ± 2	20 ± 1	<0.01	<0.01	0.08 ± 0.02	0.04 ± 0.01
WZ	19 ± 1	16 ± 1	11 ± 1	2.95 ± 0.20	1.62 ± 0.11	5.14 ± 0.35	4.25 ± 0.29
ZZ	10 ± 1	9 ± 1	5 ± 1	0.63 ± 0.05	0.47 ± 0.03	1.12 ± 0.08	0.90 ± 0.06
Total background	$74\,108 \pm 4644$	$58\,806 \pm 3910$	5024 ± 333	15.2 ± 18	2.3 ± 0.2	1720 ± 41	117 ± 13
Signal	20.5 ± 2.7	18.5 ± 2.4	16.3 ± 2.1	11.6 ± 1.5	10.1 ± 1.3	16.4 ± 2.2	13.6 ± 1.8
Data	72974	58763	4558	16	3	1678	96

requirement ($S1$ and $S2$) in the dimuon invariant mass region above $70 \text{ GeV}/c^2$, after subtracting the SM sources of background except $Z/\gamma^* \rightarrow \mu^+\mu^-$ events from the data. This mass requirement removes most multijet background events in the low mass range. From these ratios, we determine the average probability for charge misidentification in data and MC simulations to be $P_{\text{data}} = (6.2 \pm 1.1) \times 10^{-4}$ and $P_{\text{MC}} = (3.1 \pm 0.4) \times 10^{-4}$, respectively, assuming the multijet background is negligible. The uncertainties are statistical. A possible bias in such an estimate due to the potential presence of signal events in the sample has been estimated to be well below the assigned systematic uncertainty. Since the charge misidentification rate in MC simulations is underestimated, the ratio of P_{data} to P_{MC} is taken as a correction equal to 2.0 ± 0.4 . This ratio is applied to the $Z/\gamma^* \rightarrow \mu^+\mu^-$ MC sample when estimating the like-sign contribution.

The distributions of dimuon invariant mass and $\Delta\phi$ after selection $S1$ are shown in Figs. 1(a) and 1(b). The data are compared with the sum of the background contributions. For those events with more than one pair of muons fulfilling the selection criteria, the dimuon invariant mass and $\Delta\phi$ are calculated only for the pair with the highest individual momenta. The number of remaining events after each selection is shown in Table I. There is good agreement between data and the sum of the backgrounds. Figures 1(c) and 1(d) show the dimuon invariant mass and $\Delta\phi$ distributions after the $S1$ and $S4$ requirements. The excess of events at $150 \text{ GeV}/c^2$ has a significance of less than 2.6σ . The last two columns of Table I give the individual like-sign backgrounds after the various selection stages. This

demonstrates that the like-sign backgrounds are well understood.

After all five selection criteria, three data events remain, in good agreement with the SM background expectation of 2.3 ± 0.2 events. Total signal efficiencies are 32%–34% and are nearly independent of mass. The dimuon invariant mass and $\Delta\phi$ distributions for these events are compared to the sum of the backgrounds in Figs. 1(e) and 1(f).

Since no excess is observed, we use the dimuon invariant mass distribution in Fig. 1(e) to compute upper limits on the production cross section times the branching fraction as a function of $M(H^{\pm\pm})$ using the CL_S method [18] as implemented in the MCLIMIT program [19]. The expected rate for the signal as a function of $M(H^{\pm\pm})$ is determined by the NLO cross section [5] and measured luminosity and corrected for the signal efficiency.

A number of systematic uncertainties on the signal and background are taken into account in the limit calculation. The uncertainties on the correction of the muon identification are 2% and 6% for the backgrounds and signal, respectively. The uncertainty on the isolation efficiency for the multijet background is 12%. The 20% uncertainty on the correction for charge misidentification is included. The uncertainty on the luminosity for the signal is estimated to be 6.1% [20]. The uncertainty on the normalization using NNLO MC SM background production cross sections is taken to be 5%. The parton distribution function uncertainties on the cross section for backgrounds are taken to be 4% [21].

The cross section limit as a function of $M(H^{\pm\pm})$ is shown in Fig. 2 together with the theoretical cross section for left- and right-handed doubly charged Higgs bosons. Mass limits for other branching ratios can be obtained by scaling the theory cross section by the square of the branching ratio. At the 95% C.L., lower mass limits of $150 \text{ GeV}/c^2$ for left-handed and $127 \text{ GeV}/c^2$ for right-handed doubly charged Higgs bosons are obtained. This extends the previous mass limit [8] for a doubly charged Higgs boson decaying into muons.

We thank the staffs at Fermilab and collaborating institutions and acknowledge support from the DOE and NSF (USA); CEA and CNRS/IN2P3 (France); FASI, Rosatom, and RFBR (Russia); CNPq, FAPERJ, FAPESP, and FUNDUNESP (Brazil); DAE and DST (India); Colciencias (Colombia); CONACyT (Mexico); KRF and KOSEF (Korea); CONICET and UBACyT (Argentina); FOM (The Netherlands); STFC (United Kingdom); MSMT and GACR (Czech Republic); CRC Program, CFI, NSERC, and WestGrid Project (Canada); BMBF and DFG (Germany); SFI (Ireland); The Swedish Research Council (Sweden); CAS and CNSF (China); and the Alexander von Humboldt Foundation.




FIG. 2 (color online). The cross section limit as a function of the Higgs mass $M(H^{\pm\pm})$ at the 95% C.L. The mass regions excluded by LEP and CDF are also shown. The $\pm 1\sigma$ uncertainty on the expected limit is given by the shaded yellow band.

*Visitor from Augustana College, Sioux Falls, SD, USA.

- ⁺Visitor from The University of Liverpool, Liverpool, United Kingdom.
- [‡]Visitor from ICN-UNAM, Mexico City, Mexico.
- [§]Visitor from II. Physikalisches Institut, Georg-August-University, Göttingen, Germany.
- ^{||}Visitor from Helsinki Institute of Physics, Helsinki, Finland.
- [¶]Visitor from Universität Zürich, Zürich, Switzerland.
- **Deceased.
- [1] J. C. Pati and A. Salam, Phys. Rev. D **10**, 275 (1974); R. N. Mohapatra and J. C. Pati, Phys. Rev. D **11**, 566 (1975); **11**, 2558 (1975); G. Senjanovic and R. N. Mohapatra, Phys. Rev. D **12**, 1502 (1975).
- [2] G. B. Gelmini and M. Roncadelli, Phys. Lett. **99B**, 411 (1981).
- [3] N. Arkani-Hamed, A. G. Cohen, and H. Georgi, Phys. Lett. B **513**, 232 (2001).
- [4] J. F. Gunion, H. E. Haber, G. Kane, and S. Dawson, *The Higgs Hunter's Guide* (Addison-Wesley, Redwood City, CA, 1989).
- [5] M. Mühlleitner and M. Spira, Phys. Rev. D **68**, 117701 (2003).
- [6] G. Abbiendi *et al.* (OPAL Collaboration), Phys. Lett. B **526**, 221 (2002); P. D. Acton *et al.* (OPAL Collaboration), Phys. Lett. B **295**, 347 (1992); P. Achard *et al.* (L3 Collaboration), Phys. Lett. B **576**, 18 (2003); J. Abdallah *et al.* (DELPHI Collaboration), Phys. Lett. B **552**, 127 (2003).
- [7] V. M. Abazov *et al.* (D0 Collaboration), Phys. Rev. Lett. **93**, 141801 (2004).
- [8] D. Acosta *et al.* (CDF Collaboration), Phys. Rev. Lett. **93**, 221802 (2004).
- [9] V. M. Abazov *et al.* (D0 Collaboration), Nucl. Instrum. Methods Phys. Res., Sect. A **565**, 463 (2006).
- [10] The pseudorapidity is defined as $\eta = -\ln[\tan(\frac{\theta}{2})]$ as a function of the polar angle θ .
- [11] S. Abachi *et al.* (D0 Collaboration), Nucl. Instrum. Methods Phys. Res., Sect. A **338**, 185 (1994).
- [12] V. M. Abazov *et al.*, Nucl. Instrum. Methods Phys. Res., Sect. A **552**, 372 (2005).
- [13] T. Sjöstrand *et al.*, Comput. Phys. Commun. **135**, 238 (2001).
- [14] R. Hamberg, W. L. van Neerven, and T. Matsuura, Nucl. Phys. **B359**, 343 (1991); **B644**, 403(E) (2002).
- [15] N. Kidonakis and R. Vogt, Int. J. Mod. Phys. A **20**, 3171 (2005).
- [16] J. M. Campbell and R. K. Ellis, Phys. Rev. D **60**, 113006 (1999); <http://mcfm.fnal.gov/>.
- [17] R. Brun and F. Carminati, CERN Program Library Long Writeup No. W5013, 1993.
- [18] A. Read, J. Phys. G **28**, 2693 (2002).
- [19] T. Junk, Nucl. Instrum. Methods Phys. Res., Sect. A **434**, 435 (1999).
- [20] T. Andeen *et al.*, FERMILAB Report No. FERMILAB-TM-2365, 2007.
- [21] J. Pumplin *et al.*, J. High Energy Phys. 07 (2002) 12.