

MAXIMAL LEFT IDEALS OF THE BANACH ALGEBRA OF BOUNDED OPERATORS ON A BANACH SPACE

H. G. DALES, TOMASZ KANIA, TOMASZ KOCHANEK, PIOTR KOSZMIDER,
AND NIELS JAKOB LAUSTSEN

ABSTRACT. We address the following two questions regarding the maximal left ideals of the Banach algebra $\mathcal{B}(E)$ of bounded operators acting on an infinite-dimensional Banach space E :

- (I) Does $\mathcal{B}(E)$ always contain a maximal left ideal which is not finitely generated?
- (II) Is every finitely-generated, maximal left ideal of $\mathcal{B}(E)$ necessarily of the form

$$(*) \quad \{T \in \mathcal{B}(E) : Tx = 0\}$$

for some non-zero $x \in E$?

Since the two-sided ideal $\mathcal{F}(E)$ of finite-rank operators is not contained in any of the maximal left ideals given by (*), a positive answer to the second question would imply a positive answer to the first.

Our main results are: (i) Question (I) has a positive answer for most (possibly all) infinite-dimensional Banach spaces; (ii) Question (II) has a positive answer if and only if no finitely-generated, maximal left ideal of $\mathcal{B}(E)$ contains $\mathcal{F}(E)$; (iii) the answer to Question (II) is positive for many, but not all, Banach spaces.

To appear in *Studia Mathematica*.

1. INTRODUCTION AND STATEMENT OF MAIN RESULTS

The purpose of this paper is to study the maximal left ideals of the Banach algebra $\mathcal{B}(E)$ of (bounded, linear) operators acting on a Banach space E , particularly the maximal left ideals that are finitely generated. A general introduction to the Banach algebra $\mathcal{B}(E)$ can be found in [11, §2.5]. Our starting point is the elementary observation that $\mathcal{B}(E)$ always contains a large supply of singly-generated, maximal left ideals, namely

$$(1.1) \quad \mathcal{ML}_x = \{T \in \mathcal{B}(E) : Tx = 0\} \quad (x \in E \setminus \{0\})$$

(see Proposition 2.4 for details). We call the maximal left ideals of this form *fixed*, inspired by the analogous terminology for ultrafilters.

The Banach algebra $\mathcal{B}(E)$ is semisimple, as is well known (*e.g.*, see [11, Theorem 2.5.8]); that is, the intersection of its maximal left ideals is $\{0\}$. We observe that this is already true for the intersection of the fixed maximal left ideals.

2010 *Mathematics Subject Classification*. Primary 47L10, 46H10; Secondary 47L20.

Key words and phrases. Finitely-generated, maximal left ideal; Banach algebra; bounded operator; inessential operator; Banach space; Argyros–Haydon space; Sinclair–Tullo theorem.

In the case where the Banach space E is finite-dimensional, an elementary result in linear algebra states that the mapping

$$F \mapsto \{T \in \mathcal{B}(E) : F \subseteq \ker T\}$$

is an anti-isomorphism of the lattice of linear subspaces of E onto the lattice of left ideals of $\mathcal{B}(E)$ (e.g., see [24, p. 173, Exercise 3]). Hence each maximal left ideal \mathcal{L} of $\mathcal{B}(E)$ corresponds to a unique minimal, non-zero linear subspace of E , that is, a one-dimensional subspace, and therefore \mathcal{L} is fixed. This conclusion is also an easy consequence of our work, as outlined in Remark 1.2(i), below. In contrast, this statement is false whenever E is infinite-dimensional because the two-sided ideal $\mathcal{F}(E)$ of finite-rank operators is proper, so that, by Krull's theorem, it is contained in a maximal left ideal, which cannot be fixed since, for each $x \in E \setminus \{0\}$, there is a finite-rank operator T on E such that $Tx \neq 0$.

Inspired by these observations and his collaboration [12] with Żelazko, the first-named author raised the following two questions for an infinite-dimensional Banach space E :

- (I) Does $\mathcal{B}(E)$ always contain a maximal left ideal which is not finitely generated?
- (II) Is every finitely-generated, maximal left ideal of $\mathcal{B}(E)$ necessarily fixed?

In the light of the previous paragraph, we note that a positive answer to (II) would imply a positive answer to (I).

The answers to the above questions depend only on the isomorphism class of the Banach space E . This follows from the theorem of Eidelheit, which states that two Banach spaces E and F are isomorphic if and only if the corresponding Banach algebras $\mathcal{B}(E)$ and $\mathcal{B}(F)$ are isomorphic (e.g., see [11, Theorem 2.5.7]).

After presenting some preliminary material in Section 2, we shall use a counting argument in Section 3 to answer Question (I) positively for a large class of Banach spaces, including all separable Banach spaces which contain an infinite-dimensional, closed, complemented subspace with an unconditional basis and, more generally, all separable Banach spaces with an unconditional Schauder decomposition (see Corollary 3.3 for details).

We then turn our attention to Question (II). Section 4 begins with the following dichotomy, which can be viewed as the analogue of the fact that an ultrafilter on a set M is either fixed (in the sense that it consists of precisely those subsets of M which contain a fixed element $x \in M$), or it contains the Fréchet filter of all cofinite subsets of M .

Theorem 1.1 (Dichotomy for maximal left ideals). *Let E be a non-zero Banach space. Then, for each maximal left ideal \mathcal{L} of $\mathcal{B}(E)$, exactly one of the following two alternatives holds:*

- (i) \mathcal{L} is fixed; or
- (ii) \mathcal{L} contains $\mathcal{F}(E)$.

- Remark 1.2.** (i) Since $\mathcal{F}(E) = \mathcal{B}(E)$ for each finite-dimensional Banach space E , no proper left ideal of $\mathcal{B}(E)$ satisfies condition (ii) of Theorem 1.1. Hence, by this theorem, each maximal left ideal of $\mathcal{B}(E)$ is fixed whenever E is finite-dimensional.
- (ii) Another immediate consequence of Theorem 1.1 is that Question (II) has a positive answer for a Banach space E if and only if $\mathcal{F}(E)$ is not contained in any finitely-generated, maximal left ideal of $\mathcal{B}(E)$.
- (iii) In Corollary 4.1, below, we shall deduce from Theorem 1.1 a slightly stronger, but also more technical, conclusion that involves the larger ideal of inessential operators instead of $\mathcal{F}(E)$.

The other main result to be proved in Section 4 is the following dichotomy for closed left ideals of $\mathcal{B}(E)$ that are not necessarily maximal.

Theorem 1.3 (Dichotomy for closed left ideals). *Let E be a non-zero Banach space, let \mathcal{L} be a closed left ideal of $\mathcal{B}(E)$, and suppose that E is reflexive or \mathcal{L} is finitely generated. Then exactly one of the following two alternatives holds:*

- (i) \mathcal{L} is contained in a fixed maximal left ideal; or
- (ii) \mathcal{L} contains $\mathcal{F}(E)$.

We note that Theorems 1.1 and 1.3 are genuine dichotomies, in the sense that in both theorems the two alternatives (i) and (ii) are mutually exclusive because, as observed above, no fixed maximal left ideal of $\mathcal{B}(E)$ contains $\mathcal{F}(E)$.

The purpose of Sections 5 and 6 is to show that Question (II) has a positive answer for many Banach spaces, both ‘classical’ and more ‘exotic’ ones. We can summarize our results as follows, and refer to Sections 5 and 6 for full details, including precise definitions of any unexplained terminology.

Theorem 1.4. *Let E be a Banach space which satisfies one of the following six conditions:*

- (i) E has a Schauder basis and is complemented in its bidual;
- (ii) E is isomorphic to the dual space of a Banach space with a Schauder basis;
- (iii) E is an injective Banach space;
- (iv) $E = c_0(\mathbb{I})$, $E = H$, or $E = c_0(\mathbb{I}) \oplus H$, where \mathbb{I} is a non-empty index set and H is a Hilbert space;
- (v) E is a Banach space which has few operators;
- (vi) $E = C(K)$, where K is a compact Hausdorff space without isolated points, and each operator on $C(K)$ is a weak multiplication.

Then each finitely-generated, maximal left ideal of $\mathcal{B}(E)$ is fixed.

On the other hand, there is a Banach space for which the answer to Question (II) is negative; this is the main result of Section 7. Its statement involves Argyros–Haydon’s Banach space having very few operators. We denote this space by X_{AH} , and refer to Theorem 7.5 for a summary of its main properties.

Theorem 1.5. *Let $E = X_{\text{AH}} \oplus \ell_\infty$. Then the set*

$$(1.2) \quad \mathcal{K}_1 = \left\{ \begin{pmatrix} T_{1,1} & T_{1,2} \\ T_{2,1} & T_{2,2} \end{pmatrix} \in \mathcal{B}(E) : T_{1,1} \text{ is compact} \right\}$$

is a maximal two-sided ideal of codimension one in $\mathcal{B}(E)$, and hence also a maximal left ideal. Moreover, \mathcal{K}_1 is singly generated as a left ideal, and it is not fixed.

This theorem suggests that the Banach space $E = X_{\text{AH}} \oplus \ell_\infty$ is a natural candidate for providing a negative answer to Question (I). However, as we shall also show in Section 7, it does not.

Theorem 1.6. *Let $E = X_{\text{AH}} \oplus \ell_\infty$. Then the ideal \mathcal{K}_1 given by (1.2) is the unique non-fixed, finitely-generated, maximal left ideal of $\mathcal{B}(E)$. Hence*

$$(1.3) \quad \mathcal{W}_2 = \left\{ \begin{pmatrix} T_{1,1} & T_{1,2} \\ T_{2,1} & T_{2,2} \end{pmatrix} \in \mathcal{B}(E) : T_{2,2} \text{ is weakly compact} \right\},$$

which is a maximal two-sided ideal of $\mathcal{B}(E)$, is not contained in any finitely-generated, maximal left ideal of $\mathcal{B}(E)$.

To conclude this summary of our results, let us point out that Question (II) remains open in some important cases, notably for $E = C(K)$, where K is any infinite, compact metric space such that $C(K)$ is not isomorphic to c_0 .

As a final point, we shall explain how our work fits into a more general context. The main motivation behind Question (I) comes from the fact that it is the special case where $\mathcal{A} = \mathcal{B}(E)$ for a Banach space E of the following conjecture, raised and discussed in [12]:

Let \mathcal{A} be a unital Banach algebra such that every maximal left ideal of \mathcal{A} is finitely generated. Then \mathcal{A} is finite-dimensional.

A stronger form of this conjecture in the case where \mathcal{A} is commutative was proved by Ferreira and Tomassini [15]; extensions of this result are given in [12]. The conjecture is also known to be true for C^* -algebras. For a proof and a generalization of this result to the class of Hilbert C^* -modules, see [6].

The above conjecture was suggested by Sinclair and Tullo's theorem [48], which states that a Banach algebra \mathcal{A} is finite-dimensional if each closed left ideal of \mathcal{A} (not just each maximal one) is finitely generated. This result has been generalized by Boudi [7], who showed that the conclusion that \mathcal{A} is finite-dimensional remains true under the formally weaker hypothesis that each closed left ideal of \mathcal{A} is countably generated. (Boudi's theorem can in fact be deduced from Sinclair and Tullo's theorem because a closed, countably-generated left ideal is necessarily finitely generated by [12, Proposition 1.5].)

Another result that is related to our general theme, but of a different flavour from those just mentioned, is due to Grønbæk [23, Proposition 7.3], who has shown that, for a Banach space E with the approximation property, the mapping

$$F \mapsto \overline{\text{span}}\{x \otimes \lambda : x \in E, \lambda \in F\}$$

is an isomorphism of the lattice of closed linear subspaces F of the dual space of E onto the lattice of closed left ideals of the Banach algebra of compact operators on E .

2. PRELIMINARIES

Our notation is mostly standard. We write $|M|$ for the cardinality of a set M . As usual, \aleph_0 and \aleph_1 denote the first and second infinite cardinals, respectively, while $\mathfrak{c} = 2^{\aleph_0}$ is the cardinality of the continuum.

Let E be a Banach space, always supposed to be over the complex field \mathbb{C} . We denote by I_E the identity operator on E . For a non-empty set \mathbb{I} , we define

$$\ell_\infty(\mathbb{I}, E) = \{f: \mathbb{I} \rightarrow E : \|f\|_\infty < \infty\}, \quad \text{where} \quad \|f\|_\infty = \sup_{i \in \mathbb{I}} \|f(i)\|,$$

so that $\ell_\infty(\mathbb{I}, E)$ is a Banach space with respect to the norm $\|\cdot\|_\infty$. The following special conventions apply:

- $\ell_\infty(\mathbb{I}) = \ell_\infty(\mathbb{I}, \mathbb{C})$;
- $\ell_\infty = \ell_\infty(\mathbb{N})$;
- $E^n = \ell_\infty(\{1, \dots, n\}, E)$ for each $n \in \mathbb{N}$.

We write E^* for the (continuous) dual space of the Banach space E . The duality bracket between E and E^* is $\langle \cdot, \cdot \rangle$, while $\kappa_E: E \rightarrow E^{**}$ denotes the canonical embedding of E into its bidual. By an *operator* we understand a bounded, linear operator between Banach spaces; we write $\mathcal{B}(E, F)$ for the Banach space of all operators from E to another Banach space F , and denote by $T^* \in \mathcal{B}(F^*, E^*)$ the adjoint of an operator $T \in \mathcal{B}(E, F)$.

We shall require the following standard notions for $T \in \mathcal{B}(E, F)$:

- (i) T is a *finite-rank operator* if it has finite-dimensional range. We write $\mathcal{F}(E, F)$ for the set of finite-rank operators from E to F . It is well known that

$$\mathcal{F}(E, F) = \text{span}\{y \otimes \lambda : y \in F, \lambda \in E^*\},$$

where $y \otimes \lambda$ denotes the rank-one operator given by

$$y \otimes \lambda: x \mapsto \langle x, \lambda \rangle y, \quad E \rightarrow F \quad (y \in F, \lambda \in E^*).$$

The following elementary observation will be used several times:

$$(2.1) \quad R(y \otimes \lambda)S = (Ry) \otimes (S^*\lambda) \quad (y \in F, \lambda \in E^*),$$

valid for any Banach spaces D, E, F , and G and any $S \in \mathcal{B}(D, E)$ and $R \in \mathcal{B}(F, G)$.

- (ii) T is *compact* if the image under T of the unit ball of E is a relatively norm-compact subset of F . We write $\mathcal{K}(E, F)$ for the set of compact operators from E to F .
- (iii) T is *weakly compact* if the image under T of the unit ball of E is a relatively weakly compact subset of F . We write $\mathcal{W}(E, F)$ for the set of weakly compact operators from E to F .
- (iv) T is *bounded below* if, for some $\varepsilon > 0$, we have $\|Tx\| \geq \varepsilon\|x\|$ for each $x \in E$; or, equivalently, T is injective and has closed range. This

notion is dual to surjectivity in the following precise sense (*e.g.*, see [38, Theorem 3.1.22]):

$$(2.2) \quad \begin{array}{ll} T \text{ is surjective} & \iff T^* \text{ is bounded below,} \\ T \text{ is bounded below} & \iff T^* \text{ is surjective.} \end{array}$$

- (v) T is *strictly singular* if no restriction of T to an infinite-dimensional subspace of E is bounded below; that is, for each $\varepsilon > 0$, each infinite-dimensional subspace of E contains a unit vector x with $\|Tx\| \leq \varepsilon$. We write $\mathcal{S}(E, F)$ for the set of strictly singular operators from E to F .
- (vi) T is a *Fredholm operator* if its kernel is finite-dimensional and its range is finite-codimensional, in which case T has closed range.
- (vii) T is an *upper semi-Fredholm operator* if it has finite-dimensional kernel and closed range.
- (viii) T is *inessential* if $I_E - ST$ is a Fredholm operator for each $S \in \mathcal{B}(F, E)$. We write $\mathcal{E}(E, F)$ for the set of inessential operators from E to F .

The six classes \mathcal{B} , \mathcal{F} , \mathcal{H} , \mathcal{W} , \mathcal{S} , and \mathcal{E} introduced above define operator ideals in the sense of Pietsch [42], all of which except \mathcal{F} are closed. The following inclusions always hold:

$$\begin{aligned} \mathcal{F}(E, F) &\subseteq \mathcal{H}(E, F) \subseteq \mathcal{W}(E, F) \cap \mathcal{S}(E, F) \\ &\subseteq \mathcal{S}(E, F) \subseteq \mathcal{E}(E, F) \subseteq \mathcal{B}(E, F); \end{aligned}$$

no others are true in general (even in the case where $E = F$). In line with common practice, we set $\mathcal{I}(E) = \mathcal{I}(E, E)$ for each of the above operator ideals \mathcal{I} .

We remark that a left, right, or two-sided ideal of $\mathcal{B}(E)$ is proper if and only if it does not contain the identity operator I_E . This shows in particular that the two-sided ideal $\mathcal{E}(E)$ (and hence also $\mathcal{F}(E)$, $\mathcal{H}(E)$, and $\mathcal{S}(E)$) is proper whenever E is infinite-dimensional.

Remark 2.1. The ideal of inessential operators on a single Banach space E was originally introduced by Kleinecke [30] as the preimage of the Jacobson radical of the Calkin algebra $\mathcal{B}(E)/\mathcal{F}(E)$, following Yood's observation [52, p. 615] that this radical may be non-zero. The early theory of inessential operators is expounded in the monograph [8] of Caradus, Pfaffenberger, and Yood. Pietsch [42] subsequently gave the 'operator ideal' definition of $\mathcal{E}(E, F)$ stated in (viii), above, and showed that it coincides with Kleinecke's original definition in the case where $E = F$.

The following notion is central to this paper. Let Γ be a non-empty subset of $\mathcal{B}(E)$ for some Banach space E . The *left ideal generated by Γ* is the smallest left ideal \mathcal{L}_Γ of $\mathcal{B}(E)$ that contains Γ . It can be described explicitly as

$$(2.3) \quad \mathcal{L}_\Gamma = \left\{ \sum_{j=1}^n S_j T_j : S_1, \dots, S_n \in \mathcal{B}(E), T_1, \dots, T_n \in \Gamma, n \in \mathbb{N} \right\}.$$

A left ideal \mathcal{L} of $\mathcal{B}(E)$ is *singly* (respectively, *finitely*, *countably*) *generated* if $\mathcal{L} = \mathcal{L}_\Gamma$ for some singleton (respectively, non-empty and finite, countable) subset Γ of $\mathcal{B}(E)$.

In the case where Γ is a non-empty, norm-bounded subset of $\mathcal{B}(E)$, we can define an operator $\Psi_\Gamma: E \rightarrow \ell_\infty(\Gamma, E)$ by

$$(2.4) \quad (\Psi_\Gamma x)(T) = Tx \quad (x \in E, T \in \Gamma).$$

In particular, when Γ is finite, say $\Gamma = \{T_1, \dots, T_n\}$, where $n \in \mathbb{N}$ and T_1, \dots, T_n are distinct, we shall identify $\ell_\infty(\Gamma, E)$ with E^n in the natural way. Then

$$(2.5) \quad \Psi_\Gamma = \sum_{j=1}^n \iota_j T_j \in \mathcal{B}(E, E^n),$$

where $\iota_j: E \rightarrow E^n$ denotes the canonical j^{th} coordinate embedding, and (2.3) simplifies to

$$(2.6) \quad \mathcal{L}_\Gamma = \left\{ \sum_{j=1}^n S_j T_j : S_1, \dots, S_n \in \mathcal{B}(E) \right\} = \{S\Psi_\Gamma : S \in \mathcal{B}(E^n, E)\}.$$

The operator Ψ_Γ will play a key role in our work. We shall give here only one, very simple, application of Ψ_Γ , showing that each finitely-generated left ideal of operators is already singly generated for most ‘classical’ Banach spaces.

Proposition 2.2. *Let E be a Banach space which contains a complemented subspace that is isomorphic to $E \oplus E$. Then each finitely-generated left ideal of $\mathcal{B}(E)$ is singly generated.*

Proof. Let Γ be a non-empty, finite subset of $\mathcal{B}(E)$ with $n = |\Gamma| \in \mathbb{N}$. By the assumption, E contains a complemented subspace which is isomorphic to E^n , and hence there are operators $U \in \mathcal{B}(E^n, E)$ and $V \in \mathcal{B}(E, E^n)$ such that $I_{E^n} = VU$. We shall now complete the proof by showing that the left ideal \mathcal{L}_Γ is generated by the single operator $T = U\Psi_\Gamma \in \mathcal{B}(E)$.

By (2.6), we have $T \in \mathcal{L}_\Gamma$, so that $\mathcal{L}_{\{T\}} \subseteq \mathcal{L}_\Gamma$.

Conversely, each $R \in \mathcal{L}_\Gamma$ has the form $R = S\Psi_\Gamma$ for some $S \in \mathcal{B}(E^n, E)$ by (2.6), and therefore $R = S(VU)\Psi_\Gamma = (SV)T \in \mathcal{L}_{\{T\}}$. \square

Remark 2.3. Not all finitely-generated, maximal left ideals in a Banach algebra are singly generated. For instance, let

$$\mathbb{B} = \{(z, w) \in \mathbb{C}^2 : |z|^2 + |w|^2 \leq 1\}$$

be the closed unit ball in \mathbb{C}^2 , and consider the ‘polyball algebra’ \mathcal{A} on \mathbb{B} , so that by definition \mathcal{A} is the closure with respect to the uniform norm of the polynomials in two variables restricted to \mathbb{B} . It is shown in [50, Example 15.8] that the maximal ideal $\mathcal{M} = \{f \in \mathcal{A} : f(0, 0) = 0\}$ of \mathcal{A} is generated by the two coordinate functionals, but on the other hand, it is clear that \mathcal{M} is not singly generated.

It is significantly harder to find a Banach space E such that $\mathcal{B}(E)$ contains a maximal left ideal which is finitely, but not singly, generated. Such an example has, however, recently been obtained [29].

Our next result collects some basic facts about the fixed maximal left ideals of $\mathcal{B}(E)$, most of which were already stated in the Introduction.

Proposition 2.4. *Let x and y be non-zero elements of a Banach space E . Then:*

- (i) *the set \mathcal{ML}_x given by (1.1) is the left ideal of $\mathcal{B}(E)$ generated by the projection $I_E - x \otimes \lambda$, where $\lambda \in E^*$ is any functional such that $\langle x, \lambda \rangle = 1$;*
- (ii) *the left ideal \mathcal{ML}_x is maximal;*
- (iii) *$\mathcal{ML}_x = \mathcal{ML}_y$ if and only if x and y are proportional.*

In particular, $\mathcal{B}(E)$ contains $|E|$ distinct, fixed maximal left ideals whenever E is infinite-dimensional.

Proof. (i). Let $P = I_E - x \otimes \lambda$. The set \mathcal{ML}_x is clearly a left ideal which contains P , and hence $\mathcal{L}_{\{P\}} \subseteq \mathcal{ML}_x$. The reverse inclusion holds because $T(x \otimes \lambda) = 0$ for each $T \in \mathcal{ML}_x$, so that $T = TP \in \mathcal{L}_{\{P\}}$.

(ii). The left ideal \mathcal{ML}_x is evidently proper. To verify that it is maximal, suppose that $T \in \mathcal{B}(E) \setminus \mathcal{ML}_x$. Then $Tx \neq 0$, so that $\langle Tx, \mu \rangle = 1$ for some $\mu \in E^*$. The operator $S = I_E - (x \otimes \mu)T$ belongs to \mathcal{ML}_x because $(x \otimes \mu)Tx = \langle Tx, \mu \rangle x = x$, and consequently

$$I_E = S + (x \otimes \mu)T \in \mathcal{ML}_x + \mathcal{L}_{\{T\}}.$$

(iii). It is clear that $\mathcal{ML}_x = \mathcal{ML}_y$ if x and y are proportional. We prove the converse by contraposition. Suppose that x and y are linearly independent. Then we can take $\lambda \in E^*$ such that $\langle x, \lambda \rangle = 1$ and $\langle y, \lambda \rangle = 0$, and hence $x \otimes \lambda \in \mathcal{ML}_y \setminus \mathcal{ML}_x$. \square

We conclude this preliminary section with the observation that the answer to the analogue of Question (I) for two-sided ideals is negative, as the following example shows.

Example 2.5. Consider the Hilbert space $H = \ell_2(\mathbb{N}_1)$, and take a projection $P \in \mathcal{B}(H)$ with separable, infinite-dimensional range. The ideal classification of Gramsch [20] and Luft [37] implies that the ideal $\mathcal{X}(H)$ of operators with separable range is the unique maximal two-sided ideal of $\mathcal{B}(H)$. Given $T \in \mathcal{X}(H)$, let $Q \in \mathcal{B}(H)$ be the orthogonal projection onto $\overline{T(H)}$. Then $T = QT$, and also $Q = VPU$ for some operators $U, V \in \mathcal{B}(H)$, so that $T = VPUT$. Hence $\mathcal{X}(H)$ is the two-sided ideal of $\mathcal{B}(H)$ generated by the single operator P . Since $\mathcal{X}(H)$ is the only maximal two-sided ideal of $\mathcal{B}(H)$, we conclude that each maximal two-sided ideal of $\mathcal{B}(H)$ is singly generated, and therefore the analogue of Question (I) for two-sided ideals has a negative answer.

With slightly more work, we can give a similar example based on a separable Banach space. To this end, consider the p^{th} quasi-reflexive James space J_p for some $p \in (1, \infty)$. Edelstein and Mityagin [14] observed that the two-sided ideal $\mathcal{W}(J_p)$ of weakly compact operators is maximal because it has codimension one in $\mathcal{B}(J_p)$, and $\mathcal{B}(J_p)$ contains no other maximal two-sided ideals by [32, Theorem 4.16]. We shall now show that $\mathcal{W}(J_p)$ is

singly generated as a two-sided ideal. Let

$$J_p^{(n)} = \{(\alpha_j)_{j \in \mathbb{N}} \in J_p : \alpha_j = 0 \ (j > n)\} \quad (n \in \mathbb{N}).$$

Then the Banach space

$$J_p^{(\infty)} = \left(\bigoplus_{n \in \mathbb{N}} J_p^{(n)} \right)_{\ell_p}$$

is reflexive and isomorphic to a complemented subspace of J_p . (The latter observation is due to Edelstein and Mityagin [14, Lemma 6(d)]; an alternative approach can be found in [32, Proposition 4.4(iv)].) Take a projection $P \in \mathcal{B}(J_p)$ whose range is isomorphic to $J_p^{(\infty)}$. By [33, Theorem 4.3], we have

$$\begin{aligned} \mathcal{W}(J_p) &= \{TS : S \in \mathcal{B}(J_p, J_p^{(\infty)}), T \in \mathcal{B}(J_p^{(\infty)}, J_p)\} \\ &= \{VPU : U, V \in \mathcal{B}(J_p)\}, \end{aligned}$$

so that $\mathcal{W}(J_p)$ is the two-sided ideal of $\mathcal{B}(J_p)$ generated by the single operator P . On the other hand, Corollary 4.8, below, will show that $\mathcal{W}(J_p)$ is not finitely generated as a left ideal because J_p is non-reflexive.

3. COUNTING MAXIMAL LEFT IDEALS

Let E be an infinite-dimensional Banach space. An infinite family $(E_\gamma)_{\gamma \in \Gamma}$ of non-zero, closed subspaces of E is an *unconditional Schauder decomposition* of E if, for each $x \in E$, there is a unique family $(x_\gamma)_{\gamma \in \Gamma}$ with $x_\gamma \in E_\gamma$ for each $\gamma \in \Gamma$ such that the series $\sum_{\gamma \in \Gamma} x_\gamma$ converges unconditionally to x . In this case we can associate a projection $P_\Upsilon \in \mathcal{B}(E)$ with each subset Υ of Γ by the definitions

$$(3.1) \quad P_\emptyset = 0 \quad \text{and} \quad P_\Upsilon x = \sum_{\gamma \in \Upsilon} x_\gamma \quad (x \in E) \quad \text{for} \quad \Upsilon \neq \emptyset,$$

where $(x_\gamma)_{\gamma \in \Gamma}$ is related to x as above.

Using this notion, we can transfer a classical algebraic result of Rosenberg [46] to $\mathcal{B}(E)$.

Proposition 3.1. *Let E be a non-zero Banach space with an unconditional Schauder decomposition $(E_\gamma)_{\gamma \in \Gamma}$. Then the Banach algebra $\mathcal{B}(E)$ contains at least $2^{2^{|\Gamma|}}$ maximal left ideals which are not fixed.*

Proof. The power set $\mathfrak{P}(\Gamma)$ of Γ is a Boolean algebra, and

$$\mathfrak{I} = \{\Upsilon \in \mathfrak{P}(\Gamma) : |\Upsilon| < |\Gamma|\}$$

is a proper Boolean ideal of $\mathfrak{P}(\Gamma)$. Since Γ is infinite, a classical result of Pospíšil (see [44], or [9, Corollary 7.4] for an exposition) states that the collection $\mathbb{M}_{\mathfrak{I}}$ of maximal Boolean ideals of $\mathfrak{P}(\Gamma)$ containing \mathfrak{I} has cardinality $2^{2^{|\Gamma|}}$.

For each $\mathfrak{M} \in \mathbb{M}_{\mathfrak{I}}$, let $\mathcal{P}(\mathfrak{M}) = \{P_\Upsilon : \Upsilon \in \mathfrak{M}\} \subseteq \mathcal{B}(E)$, where P_Υ is the projection given by (3.1). Assume towards a contradiction that the left ideal $\mathcal{L}_{\mathcal{P}(\mathfrak{M})}$ is not proper. Then, for some $n \in \mathbb{N}$, there are operators

$T_1, \dots, T_n \in \mathcal{B}(E)$ and sets $\Upsilon_1, \dots, \Upsilon_n \in \mathfrak{M}$ such that $I_E = \sum_{j=1}^n T_j P_{\Upsilon_j}$. Right-composing both sides of this identity with the projection $P_{\Gamma \setminus \Upsilon}$, where $\Upsilon = \bigcup_{j=1}^n \Upsilon_j \in \mathfrak{M}$, we obtain $P_{\Gamma \setminus \Upsilon} = 0$, so that $\Gamma = \Upsilon \in \mathfrak{M}$, which contradicts the fact that \mathfrak{M} is a proper Boolean ideal.

We can therefore choose a maximal left ideal $\mathcal{M}_{\mathfrak{M}}$ of $\mathcal{B}(E)$ such that $\mathcal{L}_{\mathcal{P}(\mathfrak{M})} \subseteq \mathcal{M}_{\mathfrak{M}}$. This maximal left ideal $\mathcal{M}_{\mathfrak{M}}$ cannot be fixed because, for each $x \in E \setminus \{0\}$, we have $x = \sum_{\gamma \in \Gamma} P_{\{\gamma\}} x$, so that $P_{\{\gamma\}} x \neq 0$ for some $\gamma \in \Gamma$. Hence $P_{\{\gamma\}} \notin \mathcal{M}_{\mathfrak{M}}$, but on the other hand $P_{\{\gamma\}} \in \mathcal{L}_{\mathcal{P}(\mathfrak{M})} \subseteq \mathcal{M}_{\mathfrak{M}}$ since $\{\gamma\} \in \mathfrak{I} \subseteq \mathfrak{M}$.

Consequently we have a mapping $\mathfrak{M} \mapsto \mathcal{M}_{\mathfrak{M}}$ from $\mathbb{M}_{\mathfrak{I}}$ into the set of non-fixed, maximal left ideals of $\mathcal{B}(E)$. We shall complete the proof by showing that this mapping is injective. Suppose that $\mathfrak{M}, \mathfrak{N} \in \mathbb{M}_{\mathfrak{I}}$ are distinct, and take a set $\Upsilon \in \mathfrak{M} \setminus \mathfrak{N}$. The maximality of \mathfrak{N} implies that $\Gamma \setminus \Upsilon \in \mathfrak{N}$, and therefore

$$I_E = P_{\Upsilon} + P_{\Gamma \setminus \Upsilon} \in \mathcal{L}_{\mathcal{P}(\mathfrak{M})} + \mathcal{L}_{\mathcal{P}(\mathfrak{N})} \subseteq \mathcal{M}_{\mathfrak{M}} + \mathcal{M}_{\mathfrak{N}}.$$

Thus, since the left ideals $\mathcal{M}_{\mathfrak{M}}$ and $\mathcal{M}_{\mathfrak{N}}$ are proper, they are distinct. \square

Corollary 3.2. *Let E be a non-zero Banach space with an unconditional Schauder decomposition $(E_{\gamma})_{\gamma \in \Gamma}$, and suppose that E contains a dense subset D such that $2^{|D|} < 2^{2^{|\Gamma|}}$. Then $\mathcal{B}(E)$ contains at least $2^{2^{|\Gamma|}}$ maximal left ideals which are not finitely generated.*

Proof. Since each element of E is the limit point of a sequence in D , we have $|E| \leq |D|^{\aleph_0}$. Further, each operator on E is uniquely determined by its action on D , and consequently

$$(3.2) \quad |\mathcal{B}(E)| \leq |E^D| = |E|^{|D|} \leq (|D|^{\aleph_0})^{|D|} = |D|^{|D|} = 2^{|D|},$$

where the final equality follows from [25, Lemma 5.6], for example. Hence $\mathcal{B}(E)$ contains at most $(2^{|D|})^{\aleph_0} = 2^{|D|}$ countable subsets, so that $\mathcal{B}(E)$ contains at most $2^{|D|}$ countably-generated left ideals. On the other hand, Proposition 3.1 implies that there are at least $2^{2^{|\Gamma|}}$ distinct maximal left ideals of $\mathcal{B}(E)$. We have $2^{|D|} < 2^{2^{|\Gamma|}}$ by the assumption, and hence $\mathcal{B}(E)$ contains at least $2^{2^{|\Gamma|}}$ maximal left ideals which are not countably generated, and thus not finitely generated. \square

The most important case of this corollary is as follows.

Corollary 3.3. *Let E be a non-zero, separable Banach space with an unconditional Schauder decomposition $(E_{\gamma})_{\gamma \in \Gamma}$. Then $\mathcal{B}(E)$ contains precisely $2^{\mathfrak{c}}$ maximal left ideals which are not finitely generated.*

Proof. The index set Γ is necessarily countable because E is separable. Hence, by Corollary 3.2, $\mathcal{B}(E)$ contains at least $2^{\mathfrak{c}}$ maximal left ideals which are not finitely generated. On the other hand, (3.2) implies that $\mathcal{B}(E)$ has cardinality \mathfrak{c} , so that $\mathcal{B}(E)$ contains no more than $2^{\mathfrak{c}}$ distinct subsets. \square

Example 3.4. (i) Let E be a Banach space with an unconditional Schauder basis $(e_n)_{n \in \mathbb{N}}$. Then E satisfies the conditions of Corollary 3.3,

and hence $\mathcal{B}(E)$ contains 2^c maximal left ideals which are not finitely generated.

The class of Banach spaces which have an unconditional Schauder basis is large and includes for instance the classical sequence spaces c_0 and ℓ_p for $p \in [1, \infty)$, the Lebesgue spaces $L_p[0, 1]$ for $p \in (1, \infty)$, the Lorentz and Orlicz sequence spaces $d_{w,p}$ and h_M (e.g., see [35, Chapter 4]), the Tsirelson space T (e.g., see [35, Example 2.e.1]), and the Schlumprecht space S (see [47, Proposition 2]).

- (ii) Suppose that E is a Banach space containing an infinite-dimensional, closed, complemented subspace F with an unconditional Schauder decomposition $(F_\gamma)_{\gamma \in \Gamma}$. Then E also has an unconditional Schauder decomposition, obtained by adding any closed, complementary subspace of F to the collection $(F_\gamma)_{\gamma \in \Gamma}$.

In particular, generalizing (i), we see that each separable Banach space E which contains an infinite-dimensional, closed, complemented subspace with an unconditional Schauder basis satisfies the conditions of Corollary 3.3, and hence $\mathcal{B}(E)$ contains 2^c maximal left ideals that are not finitely generated. This applies for instance to $E = L_1[0, 1]$ because it contains a complemented copy of ℓ_1 (e.g., see [1, Lemma 5.1.1]); to $E = C(K)$ for any infinite, compact metric space K because E contains a complemented copy of c_0 (e.g., see [1, Proposition 4.3.11]); to $E = J_p$ for $p \in (1, \infty)$, the p^{th} quasi-reflexive James space, because J_p contains a complemented copy of ℓ_p (see [14, Lemma 6(d)] or [32, Proposition 4.4(iii)]); and to $E = \mathcal{K}(X)$, where X is any Banach space with an unconditional Schauder basis, because E contains a complemented copy of c_0 consisting of the compact operators whose matrix representation with respect to the unconditional Schauder basis is diagonal.

- (iii) There are separable Banach spaces E such that E has a unconditional Schauder decomposition $(E_n)_{n \in \mathbb{N}}$ with each E_n finite-dimensional, but E does not have an unconditional Schauder basis, notably Kalton and Peck's twisted ℓ_p -spaces Z_p for $p \in (1, \infty)$ (see [28, Corollary 9] and the remark following it). Each such Banach space E satisfies the conditions of Corollary 3.3, and hence $\mathcal{B}(E)$ contains 2^c maximal left ideals which are not finitely generated.

Remark 3.5. Corollary 3.3 is not true for all separable, infinite-dimensional Banach spaces. Indeed, we shall show in Theorem 6.2, below, that there are separable, infinite-dimensional Banach spaces E such that $\mathcal{B}(E)$ contains just one maximal left ideal which is not fixed, and this ideal is not finitely generated.

4. PROOFS OF THE DICHOTOMY THEOREMS 1.1 AND 1.3

The main purpose of this section is to prove Theorems 1.1 and 1.3. We begin with the former, whose proof is elementary.

Proof of Theorem 1.1. Let \mathcal{L} be a maximal left ideal of $\mathcal{B}(E)$ such that $\mathcal{F}(E) \not\subseteq \mathcal{L}$. Then there exist $x \in E$ and $\lambda \in E^*$ such that $x \otimes \lambda \notin \mathcal{L}$.

The maximality of \mathcal{L} implies that $I_E - T(x \otimes \lambda) \in \mathcal{L}$ for some operator $T \in \mathcal{B}(E)$, and $Tx \neq 0$ because \mathcal{L} is proper. Choose $\mu \in E^*$ such that $\langle Tx, \mu \rangle = 1$, and define $P = I_E - Tx \otimes \mu \in \mathcal{B}(E)$. We then have $PTx = 0$, so that $P = P(I_E - T(x \otimes \lambda)) \in \mathcal{L}$, and hence $\mathcal{ML}_{Tx} \subseteq \mathcal{L}$ by Proposition 2.4(i). Consequently these two maximal left ideals are equal, which shows that \mathcal{L} is fixed. \square

Theorem 1.1 easily leads to the following stronger conclusion.

Corollary 4.1 (Strong dichotomy for maximal left ideals). *Let E be a non-zero Banach space. Then, for each maximal left ideal \mathcal{L} of $\mathcal{B}(E)$, exactly one of the following two alternatives holds:*

- (i) \mathcal{L} is fixed; or
- (ii) \mathcal{L} contains $\mathcal{E}(E)$.

Proof. Let \mathcal{L} be a non-fixed, maximal left ideal of $\mathcal{B}(E)$. Then \mathcal{L} is closed and contains $\mathcal{F}(E)$ by Theorem 1.1, so that $\overline{\mathcal{F}(E)} \subseteq \mathcal{L}$, and thus $\pi(\mathcal{L})$ is a maximal left ideal of the Calkin algebra $\mathcal{B}(E)/\overline{\mathcal{F}(E)}$, where

$$\pi: \mathcal{B}(E) \rightarrow \mathcal{B}(E)/\overline{\mathcal{F}(E)}$$

denotes the quotient homomorphism. In particular $\pi(\mathcal{L})$ contains the Jacobson radical of $\mathcal{B}(E)/\overline{\mathcal{F}(E)}$, which implies that \mathcal{L} contains $\mathcal{E}(E)$ by Remark 2.1, and therefore (ii) holds. \square

Remark 4.2. Corollary 4.1 can be seen as a counter-part for maximal left ideals of [32, Proposition 6.6], which states that each maximal two-sided ideal of $\mathcal{B}(E)$ contains $\mathcal{E}(E)$ for each infinite-dimensional Banach space E .

Remark 4.3. Let \mathcal{A} be a unital C^* -algebra. We write $a \mapsto a^*$ for the involution on \mathcal{A} . (This should not be confused with the notation T^* for the adjoint of an operator T between Banach spaces used elsewhere in this paper.) A *state* on \mathcal{A} is a norm-one functional λ on \mathcal{A} which is positive, in the sense that $\langle a^*a, \lambda \rangle \geq 0$ for each $a \in \mathcal{A}$. Given a state λ on \mathcal{A} , the set

$$\mathcal{N}_\lambda = \{a \in \mathcal{A} : \langle a^*a, \lambda \rangle = 0\}$$

is a closed left ideal of \mathcal{A} by the Cauchy–Schwarz inequality (*e.g.*, see [26, Proposition 4.5.1] or [39, p. 93]). The collection of all states on \mathcal{A} forms a weak*-compact, convex subset of the dual space of \mathcal{A} , called the *state space* of \mathcal{A} . Its extreme points are the *pure states* on \mathcal{A} . Prosser [45, Theorem 6.2] has shown that the map $\lambda \mapsto \mathcal{N}_\lambda$ gives a bijective correspondence between the pure states on \mathcal{A} and the maximal left ideals of \mathcal{A} ; expositions of this result can be found in [27, Theorem 10.2.10] and [39, Theorem 5.3.5].

In the case where $\mathcal{A} = \mathcal{B}(H)$ for some Hilbert space H , the fixed maximal left ideals correspond to the vector states, which are defined as follows. Let $x \in H$ be a unit vector. Then the functional ω_x given by

$$\langle T, \omega_x \rangle = (Tx | x) \quad (T \in \mathcal{B}(H)),$$

where $(\cdot | \cdot)$ denotes the inner product on H , is a pure state on $\mathcal{B}(H)$, called the *vector state* induced by x ; and we have $\mathcal{ML}_x = \mathcal{N}_{\omega_x}$, as is easy

to check. The conclusion of Corollary 4.1 is known in this case because $\mathcal{K}(H) = \mathcal{E}(H)$, and by [27, Corollary 10.4.4] each pure state λ on $\mathcal{B}(H)$ is either a vector state, or $\mathcal{K}(H) \subseteq \ker \lambda$, in which case $\mathcal{K}(H) \subseteq \mathcal{N}_\lambda$.

Finally, suppose that the Hilbert space H is separable and infinite-dimensional. Then clearly $\mathcal{B}(H)$ has \mathfrak{c} vector states, whereas it has $2^{\mathfrak{c}}$ pure states by [27, Proposition 10.4.15]. These conclusions also follow from Proposition 2.4 and Example 3.4(i), respectively.

We shall now turn our attention to the proof of Theorem 1.3. This requires some preparation. Let E be a Banach space. For each non-empty, bounded subset Γ of $\mathcal{B}(E)$, we can define an operator Ω_Γ from the Banach space

$$\ell_1(\Gamma, E^*) = \left\{ g: \Gamma \rightarrow E^* : \sum_{T \in \Gamma} \|g(T)\| < \infty \right\}$$

into E^* by $\Omega_\Gamma g = \sum_{T \in \Gamma} T^*g(T)$ for each $g \in \ell_1(\Gamma, E^*)$; that is,

$$(4.1) \quad \langle x, \Omega_\Gamma g \rangle = \sum_{T \in \Gamma} \langle Tx, g(T) \rangle \quad (x \in E, g \in \ell_1(\Gamma, E^*)).$$

The following lemma lists some basic properties of this operator, as well as of the operator Ψ_Γ given by (2.4), and explains their relevance for our present purpose. To state it, we require the notion of the *pre-annihilator* ${}^\perp M$ of a subset M of a dual Banach space E^* :

$${}^\perp M = \{x \in E : \langle x, \lambda \rangle = 0 \ (\lambda \in M)\}.$$

Lemma 4.4. *Let E be a non-zero Banach space, and let Γ be a non-empty, bounded subset of $\mathcal{B}(E)$. Then:*

- (i) $\ker \Psi_\Gamma = {}^\perp \Omega_\Gamma(\ell_1(\Gamma, E^*))$, and a non-zero element x of E belongs to this set if and only if $\mathcal{L}_\Gamma \subseteq \mathcal{ML}_x$;
- (ii) the following three conditions are equivalent:
 - (a) no fixed maximal left ideal of $\mathcal{B}(E)$ contains \mathcal{L}_Γ ;
 - (b) the operator Ψ_Γ is injective;
 - (c) the range of the operator Ω_Γ is weak*-dense in E^* .

Now suppose either that the set Γ is finite or that the left ideal \mathcal{L}_Γ is closed. Then:

- (iii) for each $\lambda \in E^*$, the set $\mathcal{J}_\lambda = \{y \otimes \lambda : y \in E\}$ is a left ideal of $\mathcal{B}(E)$, and the following three conditions are equivalent:
 - (a) $\mathcal{J}_\lambda \subseteq \mathcal{L}_\Gamma$;
 - (b) $y \otimes \lambda \in \mathcal{L}_\Gamma$ for some $y \in E \setminus \{0\}$;
 - (c) $\lambda \in \Omega_\Gamma(\ell_1(\Gamma, E^*))$;
- (iv) the operator Ω_Γ is surjective if and only if \mathcal{L}_Γ contains $\mathcal{F}(E)$.

Proof. (i). Suppose that $x \in \ker \Psi_\Gamma$. Then $Tx = 0$ for each $T \in \Gamma$, so that (4.1) implies that $x \in {}^\perp \Omega_\Gamma(\ell_1(\Gamma, E^*))$.

Conversely, suppose that $x \in {}^\perp \Omega_\Gamma(\ell_1(\Gamma, E^*))$, and let $T \in \Gamma$ and $\lambda \in E^*$ be given. Defining $g \in \ell_1(\Gamma, E^*)$ by $g(T) = \lambda$ and $g(S) = 0$ for $S \in \Gamma \setminus \{T\}$, we have $0 = \langle x, \Omega_\Gamma g \rangle = \langle Tx, \lambda \rangle$ by (4.1). Since $\lambda \in E^*$ was arbitrary, this shows that $Tx = 0$, and hence $x \in \ker \Psi_\Gamma$.

To prove the second clause, we observe that for each $x \in E \setminus \{0\}$, we have $x \in \ker \Psi_\Gamma$ if and only if $\Gamma \subseteq \mathcal{ML}_x$, and hence if and only if $\mathcal{L}_\Gamma \subseteq \mathcal{ML}_x$.

(ii). The equivalence of (a) and (b) is immediate from (i).

To see that (b) and (c) are equivalent, we observe that, by (i) and [38, Proposition 2.6.6(c)], the weak* closure of the range of Ω_Γ is equal to the annihilator

$$(\ker \Psi_\Gamma)^\perp = \{\lambda \in E^* : \langle x, \lambda \rangle = 0 \ (x \in \ker \Psi_\Gamma)\}$$

of $\ker \Psi_\Gamma$. Hence (b) implies (c). Conversely, suppose that $(\ker \Psi_\Gamma)^\perp = E^*$. Then [38, Proposition 1.10.15(c)] implies that $\ker \Psi_\Gamma = {}^\perp(E^*) = \{0\}$.

(iii). Equation (2.1) shows that \mathcal{I}_λ is a left ideal.

The implication (a) \Rightarrow (b) is evident.

(b) \Rightarrow (c). Suppose that $y \otimes \lambda \in \mathcal{L}_\Gamma$ for some $y \in E \setminus \{0\}$. Then there are $n \in \mathbb{N}$, $S_1, \dots, S_n \in \mathcal{B}(E)$, and $T_1, \dots, T_n \in \Gamma$ such that $y \otimes \lambda = \sum_{j=1}^n S_j T_j$, where we may suppose that T_1, \dots, T_n are distinct. Choose $\mu \in E^*$ such that $\langle y, \mu \rangle = 1$, and define $g: \Gamma \rightarrow E^*$ by

$$g(T) = \begin{cases} S_j^* \mu & \text{if } T = T_j \text{ for some } j \in \{1, \dots, n\}, \\ 0 & \text{otherwise.} \end{cases}$$

Then g has finite support, so that trivially it belongs to $\ell_1(\Gamma, E^*)$, and $\Omega_\Gamma g = \lambda$ because (4.1) implies that

$$\langle x, \Omega_\Gamma g \rangle = \sum_{j=1}^n \langle T_j x, S_j^* \mu \rangle = \left\langle \sum_{j=1}^n S_j T_j x, \mu \right\rangle = \langle (y \otimes \lambda)x, \mu \rangle = \langle x, \lambda \rangle$$

for each $x \in E$. Hence $\lambda \in \Omega_\Gamma(\ell_1(\Gamma, E^*))$.

(c) \Rightarrow (a). Suppose that $\lambda = \Omega_\Gamma g$ for some $g \in \ell_1(\Gamma, E^*)$. Then, for each $y \in E$, we have

$$y \otimes \lambda = \sum_{T \in \Gamma} y \otimes T^* g(T) = \sum_{T \in \Gamma} (y \otimes g(T))T,$$

which belongs to \mathcal{L}_Γ because each term of the sum on the right-hand side does, and either this sum is finite or \mathcal{L}_Γ is closed by the assumption.

(iv). Suppose that Ω_Γ is surjective. Then (iii) implies that $\mathcal{I}_\lambda \subseteq \mathcal{L}_\Gamma$ for each $\lambda \in E^*$, and consequently $\mathcal{F}(E) = \text{span} \bigcup_{\lambda \in E^*} \mathcal{I}_\lambda \subseteq \mathcal{L}_\Gamma$.

Conversely, suppose that $\mathcal{F}(E) \subseteq \mathcal{L}_\Gamma$, and let $\lambda \in E^*$ be given. Since $\mathcal{I}_\lambda \subseteq \mathcal{F}(E)$, (iii) implies that $\lambda \in \Omega_\Gamma(\ell_1(\Gamma, E^*))$, so that the map Ω_Γ is surjective. \square

Corollary 4.5. *Let E be a Banach space, and let Γ be a non-empty, bounded subset of $\mathcal{B}(E)$ such that the left ideal \mathcal{L}_Γ is closed. Then the operator Ω_Γ has closed range.*

Proof. We may suppose that E is non-zero. Consider a sequence $(\lambda_j)_{j \in \mathbb{N}}$ in $\Omega_\Gamma(\ell_1(\Gamma, E^*))$ that converges to some element $\lambda \in E^*$. Lemma 4.4(iii) shows that $\mathcal{I}_{\lambda_j} \subseteq \mathcal{L}_\Gamma$ for each $j \in \mathbb{N}$, and hence

$$y \otimes \lambda = \lim_{j \rightarrow \infty} y \otimes \lambda_j \in \mathcal{L}_\Gamma \quad (y \in E)$$

because \mathcal{L}_Γ is closed, so that $\lambda \in \Omega_\Gamma(\ell_1(\Gamma, E^*))$ by another application of Lemma 4.4(iii). Thus Ω_Γ has closed range. \square

We can now characterize the closed left ideals of $\mathcal{B}(E)$ that contain $\mathcal{F}(E)$ as follows, provided either that E is reflexive or that we restrict our attention to the closed left ideals that are finitely generated. Note that Theorem 1.3 is simply a restatement of the equivalence of conditions (a) and (f).

Theorem 4.6. *Let E be a non-zero Banach space, let \mathcal{L} be a closed left ideal of $\mathcal{B}(E)$, and take a non-empty, bounded subset Γ of $\mathcal{B}(E)$ such that $\mathcal{L} = \mathcal{L}_\Gamma$. Suppose either that E is reflexive or that Γ is finite. Then the following six conditions are equivalent:*

- (a) *no fixed maximal left ideal of $\mathcal{B}(E)$ contains \mathcal{L} ;*
- (b) *the operator Ψ_Γ is injective;*
- (c) *the operator Ψ_Γ is bounded below;*
- (d) *the range of the operator Ω_Γ is weak*-dense in E^* ;*
- (e) *the operator Ω_Γ is surjective;*
- (f) *\mathcal{L} contains $\mathcal{F}(E)$.*

Proof. The following implications hold without supposing that E is reflexive or Γ is finite: Conditions (a), (b), and (d) are mutually equivalent by Lemma 4.4(ii), while conditions (e) and (f) are equivalent by Lemma 4.4(iv). Evidently (c) implies (b), and (e) implies (d). In fact, (e) implies (c), as we shall now show. Suppose that Ω_Γ is surjective. We can define a linear isometry $\Xi_\Gamma: \ell_1(\Gamma, E^*) \rightarrow \ell_\infty(\Gamma, E)^*$ by

$$(4.2) \quad \langle f, \Xi_\Gamma g \rangle = \sum_{T \in \Gamma} \langle f(T), g(T) \rangle \quad (f \in \ell_\infty(\Gamma, E), g \in \ell_1(\Gamma, E^*)),$$

and $\Omega_\Gamma = \Psi_\Gamma^* \Xi_\Gamma$. Hence the surjectivity of Ω_Γ implies that Ψ_Γ^* is surjective, and therefore Ψ_Γ is bounded below by (2.2).

The remaining implications do require further assumptions. We consider first the case where E is reflexive. Then the weak and weak* topologies on E^* coincide, so that the range of Ω_Γ is weak*-dense if and only if it is weakly dense, if and only if it is norm-dense by Mazur's theorem (*e.g.*, see [38, Theorem 2.5.16]). However, Ω_Γ has closed range by Corollary 4.5, and consequently (d) implies (e), which completes the proof in this case.

Secondly, suppose that the set Γ is finite. Then the isometry

$$\Xi_\Gamma: \ell_1(\Gamma, E^*) \rightarrow \ell_\infty(\Gamma, E)^*$$

defined by (4.2), above, is an isomorphism, so that Ω_Γ and Ψ_Γ^* are equal up to an isometric identification. Hence (c) and (e) are equivalent by (2.2). Moreover, Corollary 4.5 shows that Ψ_Γ^* has closed range, and therefore Ψ_Γ has closed range by the closed range theorem (*e.g.*, see [38, Theorem 3.1.21]). Thus (b) implies (c), and the proof is complete. \square

For later reference, we note that the arguments which establish the equivalence of conditions (c), (e), and (f) for finite Γ given in the first and last paragraph of the proof of Theorem 4.6, above, remain true in the case where

the left ideal $\mathcal{L} = \mathcal{L}_\Gamma$ is not necessarily closed. Hence we have the following conclusion.

Corollary 4.7. *Let E be a Banach space, and let Γ be a non-empty, finite subset of $\mathcal{B}(E)$. Then the following three conditions are equivalent:*

- (a) *the operator Ψ_Γ is bounded below;*
- (b) *the operator Ω_Γ is surjective;*
- (c) *\mathcal{L}_Γ contains $\mathcal{F}(E)$.*

As an easy consequence of these results, we obtain that the ideal of weakly compact operators is finitely generated as a left ideal only in the trivial case where it is not proper.

Corollary 4.8. *The following three conditions are equivalent for a Banach space E :*

- (a) *$\mathcal{W}(E)$ is finitely generated as a left ideal;*
- (b) *$\mathcal{W}(E) = \mathcal{B}(E)$;*
- (c) *the Banach space E is reflexive.*

Proof. The equivalence of (b) and (c) is standard, and (b) obviously implies (a).

To see that (a) implies (c), suppose that $\mathcal{W}(E) = \mathcal{L}_\Gamma$ for some non-empty, finite subset $\Gamma = \{T_1, \dots, T_n\}$ of $\mathcal{B}(E)$. It clearly suffices to consider the case where E is non-zero. Theorem 4.6 (or Corollary 4.7) implies that the operator Ψ_Γ is bounded below because $\mathcal{W}(E)$ contains $\mathcal{F}(E)$. Since the operators T_1, \dots, T_n are weakly compact, the same is true for Ψ_Γ by the definition (2.5). Hence the Davis–Figiel–Johnson–Pełczyński factorization theorem (see [13], or [36, Theorem 2.g.11] for an exposition) implies that, for some reflexive Banach space F , there are operators $R \in \mathcal{B}(E, F)$ and $S \in \mathcal{B}(F, E^n)$ such that $\Psi_\Gamma = SR$. Now R is bounded below because Ψ_Γ is, and therefore E is isomorphic to the subspace $R(E)$ of the reflexive space F , so that E is reflexive. \square

We conclude this section with an example that shows that Theorem 4.6 may not be true if we drop the assumption that either the Banach space E is reflexive or the set Γ is finite. This requires the following easy variant of Lemma 4.4(iii).

Lemma 4.9. *Let T be an operator on a Banach space E , and suppose that $y \otimes \lambda \in \overline{\mathcal{L}_{\{T\}}}$ for some $y \in E \setminus \{0\}$ and $\lambda \in E^*$. Then $\lambda \in \overline{T^*(E^*)}$.*

Proof. Let $(S_j)_{j \in \mathbb{N}}$ be a sequence in $\mathcal{B}(E)$ such that $S_j T \rightarrow y \otimes \lambda$ as $j \rightarrow \infty$, and choose $\mu \in E^*$ such that $\langle y, \mu \rangle = 1$. Then

$$T^*(S_j^* \mu) = (S_j T)^* \mu \rightarrow (y \otimes \lambda)^* \mu = \langle y, \mu \rangle \lambda = \lambda \quad \text{as } j \rightarrow \infty,$$

from which the conclusion follows. \square

Example 4.10. Let T be the operator on ℓ_∞ given by

$$(4.3) \quad T(\alpha_j)_{j \in \mathbb{N}} = \left(-\frac{\alpha_n}{2^n} + \sum_{j=n+1}^{\infty} \frac{\alpha_j}{2^j} \right)_{n \in \mathbb{N}} \quad ((\alpha_j)_{j \in \mathbb{N}} \in \ell_\infty).$$

Then T is compact and leaves the subspace c_0 invariant. Define

$$T_0: x \mapsto Tx, \quad c_0 \rightarrow c_0,$$

and consider the closed left ideal $\mathcal{L} = \overline{\mathcal{L}_{\{T_0\}}}$ of $\mathcal{B}(c_0)$. We have $\mathcal{L} \subseteq \mathcal{K}(c_0)$ because T_0 is compact. Our aim is to show that \mathcal{L} satisfies condition (a), but not condition (f), of Theorem 4.6.

We begin by verifying that $\ker T = \mathbb{C}(1, 1, \dots)$. First, it is clear that $T(1, 1, \dots) = (0, 0, \dots)$. Conversely, suppose that $(\alpha_j)_{j \in \mathbb{N}} \in \ker T$. Then

$$\frac{\alpha_n}{2^n} = \sum_{j=n+1}^{\infty} \frac{\alpha_j}{2^j} \quad (n \in \mathbb{N}),$$

so that

$$\frac{\alpha_n}{2^n} = \frac{\alpha_{n+1}}{2^{n+1}} + \sum_{j=n+2}^{\infty} \frac{\alpha_j}{2^j} = \frac{\alpha_{n+1}}{2^{n+1}} + \frac{\alpha_{n+1}}{2^{n+1}} = \frac{\alpha_{n+1}}{2^n} \quad (n \in \mathbb{N}).$$

Hence $\alpha_n = \alpha_{n+1}$ for each $n \in \mathbb{N}$, and the conclusion follows.

This shows in particular that T_0 is injective because $c_0 \cap \ker T = \{0\}$. Consequently $T_0 \notin \mathcal{ML}_x$ for each $x \in c_0 \setminus \{0\}$, and so \mathcal{L} satisfies condition (a) of Theorem 4.6.

On the other hand, identifying c_0^{**} with ℓ_∞ in the usual way, we find that $T_0^{**} = T$, which is not injective, so that T_0^{**} does not have norm-dense range by [38, Theorem 3.1.17(b)]. Take $\lambda \in c_0^* \setminus \overline{T_0^{**}(c_0^*)}$ and $y \in E \setminus \{0\}$. Then, by Lemma 4.9, $y \otimes \lambda \notin \mathcal{L}$, so that \mathcal{L} does not satisfy condition (f) of Theorem 4.6.

5. 'CLASSICAL' BANACH SPACES FOR WHICH EACH FINITELY-GENERATED, MAXIMAL LEFT IDEAL IS FIXED

The purpose of this section is to show that Question (II) has a positive answer for many standard Banach spaces E .

We begin by showing that a much stronger conclusion is true in certain cases, namely that no finitely-generated, proper left ideal of $\mathcal{B}(E)$ contains $\mathcal{F}(E)$. This result relies on the following characterization of the finite subsets Γ of $\mathcal{B}(E)$ that do *not* generate a proper left ideal in terms of standard operator-theoretic properties of Ψ_Γ .

Lemma 5.1. *Let E be a non-zero Banach space. Then the following three conditions are equivalent for each non-empty, finite subset Γ of $\mathcal{B}(E)$:*

- (a) *the operator Ψ_Γ is bounded below and its range is complemented in $E^{|\Gamma|}$;*
- (b) *the operator Ψ_Γ is left invertible;*
- (c) $\mathcal{L}_\Gamma = \mathcal{B}(E)$.

Proof. The equivalence of (a) and (b) is an easy standard result, true for any operator between Banach spaces, while the equivalence of (b) and (c) follows immediately from (2.6). \square

Proposition 5.2. *Let E be a Banach space, and let $n \in \mathbb{N}$. Then $\mathcal{F}(E)$ is contained in a proper left ideal of $\mathcal{B}(E)$ generated by n operators if and*

only if E^n contains a closed subspace which is isomorphic to E and which is not complemented in E^n .

Proof. We may suppose that E is non-zero, and prove both implications by contraposition.

\Rightarrow . Suppose that every closed subspace of E^n that is isomorphic to E is complemented in E^n , and let Γ be a subset of $\mathcal{B}(E)$ of cardinality n such that $\mathcal{F}(E) \subseteq \mathcal{L}_\Gamma$. We must prove that $\mathcal{L}_\Gamma = \mathcal{B}(E)$; that is, by Lemma 5.1, we must show that the operator Ψ_Γ is bounded below and has complemented range. Corollary 4.7 implies that Ψ_Γ is indeed bounded below, and its range is therefore a closed subspace of E^n isomorphic to E , so that it is complemented by the assumption.

\Leftarrow . Suppose that $\mathcal{B}(E)$ is the only left ideal with (at most) n generators that contains $\mathcal{F}(E)$, and let F be a closed subspace of E^n such that F is isomorphic to E . We must prove that F is complemented in E^n . Take an operator $T \in \mathcal{B}(E, E^n)$ which is bounded below and has range F , and let $T_j = \rho_j T \in \mathcal{B}(E)$, where

$$\rho_j: (x_k)_{k=1}^n \mapsto x_j, \quad E^n \rightarrow E \quad (j \in \{1, \dots, n\}).$$

Re-ordering the coordinates of E^n , we may suppose that there exists a number $m \leq n$ such that T_1, \dots, T_m are distinct and the set $\Gamma = \{T_1, \dots, T_m\}$ contains T_j for each $j \in \{1, \dots, n\}$. Then $\|\Psi_\Gamma x\| = \max_{1 \leq j \leq m} \|T_j x\| = \|Tx\|$ for each $x \in E$, so that Ψ_Γ is bounded below, and therefore \mathcal{L}_Γ contains $\mathcal{F}(E)$ by Corollary 4.7. Now the assumption implies that $\mathcal{L}_\Gamma = \mathcal{B}(E)$; that is, we can find an operator $S \in \mathcal{B}(E^m, E)$ such that

$$I_E = S\Psi_\Gamma = S\left(\sum_{j=1}^m \iota_j \rho_j\right)T.$$

Hence T has a left inverse, and consequently its range, which is equal to F , is complemented in E^n . \square

Combining this result with Theorem 1.1, we reach the following conclusion.

Corollary 5.3. *Let E be a non-zero Banach space such that, for each $n \in \mathbb{N}$, every closed subspace of E^n that is isomorphic to E is complemented in E^n . Then $\mathcal{B}(E)$ is the only finitely-generated left ideal of $\mathcal{B}(E)$ which contains $\mathcal{F}(E)$, and hence each finitely-generated, maximal left ideal of $\mathcal{B}(E)$ is fixed.*

Example 5.4. The condition of Corollary 5.3 on the Banach space E is satisfied in each of the following three cases:

- (i) E is a Hilbert space.
- (ii) E is an injective Banach space; that is, whenever a Banach space F contains a closed subspace G which is isomorphic to E , then G is complemented in F . For instance, the Banach space $E = \ell_\infty(\mathbb{I})$ is injective for each non-empty set \mathbb{I} . More generally, $C(K)$ is injective whenever the Hausdorff space K is Stonean (that is, compact and

extremely disconnected), as shown by Goodner [18] and Nachbin [40] for real scalars and generalized to the complex case by Cohen [10].

- (iii) $E = c_0(\mathbb{I})$ for a non-empty set \mathbb{I} (this follows from Sobczyk's theorem [49] for countable \mathbb{I} and from [21] (or [3, Proposition 2.8]) in the general case); here $c_0(\mathbb{I})$ denotes the closed subspace of $\ell_\infty(\mathbb{I})$ consisting of those functions $f: \mathbb{I} \rightarrow \mathbb{C}$ for which the set $\{i \in \mathbb{I} : |f(i)| \geq \varepsilon\}$ is finite for each $\varepsilon > 0$.

Thus, in each of these three cases, $\mathcal{B}(E)$ is the only finitely-generated left ideal of $\mathcal{B}(E)$ which contains $\mathcal{F}(E)$, and each finitely-generated, maximal left ideal of $\mathcal{B}(E)$ is fixed.

Our next goal is to prove a result (Theorem 5.8) which, under much less restrictive conditions on the Banach space E than Corollary 5.3, gives the slightly weaker conclusion that $\mathcal{B}(E)$ is the only finitely-generated left ideal of $\mathcal{B}(E)$ which contains $\mathcal{K}(E)$. We note in particular that Corollary 4.1 ensures that this conclusion is still strong enough to ensure that each finitely-generated, maximal left ideal of $\mathcal{B}(E)$ is fixed, thus answering Question (II) positively for a large number of Banach spaces.

Let E be a Banach space with a Schauder basis $\mathbf{e} = (e_j)_{j \in \mathbb{N}}$. For each $k \in \mathbb{N}$, we denote by P_k the k^{th} basis projection associated with \mathbf{e} . The *basis constant* of \mathbf{e} is

$$K_{\mathbf{e}} = \sup\{\|P_k\| : k \in \mathbb{N}\} \in [1, \infty).$$

The basis \mathbf{e} is *monotone* if $K_{\mathbf{e}} = 1$.

Lemma 5.5. *Let E be a Banach space with a Schauder basis $\mathbf{e} = (e_j)_{j \in \mathbb{N}}$, and let $\gamma = (\gamma_j)_{j \in \mathbb{N}}$ be a decreasing sequence of non-negative real numbers. Then*

$$(5.1) \quad \Delta_\gamma: \sum_{j=1}^{\infty} \alpha_j e_j \mapsto \sum_{j=1}^{\infty} \gamma_j \alpha_j e_j$$

defines an operator Δ_γ on E of norm at most $K_{\mathbf{e}}\gamma_1$. This operator is compact if and only if $\gamma_j \rightarrow 0$ as $j \rightarrow \infty$.

Proof. Equation (5.1) clearly defines a linear mapping Δ_γ from the dense subspace $\text{span}\{e_j : j \in \mathbb{N}\}$ of E into E , and so it suffices to show that this mapping is bounded with norm at most $K_{\mathbf{e}}\gamma_1$. Now, for each element $x = \sum_{j=1}^k \alpha_j e_j$ of $\text{span}\{e_j : j \in \mathbb{N}\}$, where $k \in \mathbb{N}$ and $\alpha_1, \dots, \alpha_k \in \mathbb{C}$, we have

$$\Delta_\gamma x = \gamma_1 P_1 x + \sum_{j=2}^k \gamma_j (P_j x - P_{j-1} x) = \sum_{j=1}^{k-1} (\gamma_j - \gamma_{j+1}) P_j x + \gamma_k x,$$

and thus

$$\|\Delta_\gamma x\| \leq \sum_{j=1}^{k-1} (\gamma_j - \gamma_{j+1}) K_{\mathbf{e}} \|x\| + \gamma_k \|x\| \leq K_{\mathbf{e}} \gamma_1 \|x\|,$$

as required.

To prove the final clause, we note that, by a standard result, $(P_j)_{j \in \mathbb{N}}$ is a bounded left approximate identity for $\mathcal{K}(E)$ (e.g., see [11, p. 318]), so that Δ_γ is compact if and only if $P_j \Delta_\gamma \rightarrow \Delta_\gamma$ as $j \rightarrow \infty$. Hence the estimates

$$\gamma_{j+1} \leq \|(I_E - P_j) \Delta_\gamma\| \leq K_{\mathbf{e}}(K_{\mathbf{e}} + 1) \gamma_{j+1} \quad (j \in \mathbb{N}),$$

which are easy to verify, give the result. \square

Corollary 5.6. *Let E be a Banach space with a Schauder basis $\mathbf{e} = (e_j)_{j \in \mathbb{N}}$, let $k \in \mathbb{N}$, and let $\eta = (\eta_j)_{j=1}^k$ be an increasing k -tuple of non-negative real numbers. Then*

$$(5.2) \quad \Theta_\eta: \sum_{j=1}^{\infty} \alpha_j e_j \mapsto \sum_{j=1}^k \eta_j \alpha_j e_j$$

defines an operator on E of norm at most $2K_{\mathbf{e}} \eta_k$.

Proof. Define

$$f_j = \begin{cases} e_{k-j+1} & \text{for } j \leq k, \\ e_j & \text{for } j > k. \end{cases}$$

Then $\mathbf{f} = (f_j)_{j \in \mathbb{N}}$ is a Schauder basis for E (because we have re-ordered only finitely many vectors of the original basis \mathbf{e}), and the m^{th} basis projection associated with \mathbf{f} is given by $P_k - P_{k-m}$ for $m < k$ and P_m for $m \geq k$, so that $K_{\mathbf{f}} \leq 2K_{\mathbf{e}}$. Now Lemma 5.5 gives the desired conclusion because

$$\Theta_\eta = \Delta_\gamma: \sum_{j=1}^{\infty} \alpha_j f_j \mapsto \sum_{j=1}^{\infty} \gamma_j \alpha_j f_j, \quad E \rightarrow E,$$

where $\gamma = (\gamma_j)_{j \in \mathbb{N}}$ denotes the decreasing sequence $(\eta_k, \eta_{k-1}, \dots, \eta_1, 0, 0, \dots)$. \square

We now come to our key lemma.

Lemma 5.7. *Let E be a Banach space with a monotone Schauder basis, and let Γ be a non-empty, finite subset of $\mathcal{B}(E)$ for which $\mathcal{F}(E) \subseteq \mathcal{L}_\Gamma$. Then the sequence $(t_j)_{j \in \mathbb{N}}$ given by*

$$(5.3) \quad t_j = \inf \{ \|T\| : T \in \mathcal{B}(E^{|\Gamma|}, E), P_j = T \Psi_\Gamma \} \in (0, \infty) \quad (j \in \mathbb{N})$$

is increasing.

Suppose that $(t_j)_{j \in \mathbb{N}}$ is unbounded, and let $\gamma = (t_j^{-1/2})_{j \in \mathbb{N}}$. Then the operator Δ_γ given by (5.1) is compact and does not belong to \mathcal{L}_Γ .

Proof. Set $n = |\Gamma| \in \mathbb{N}$. For each $j \in \mathbb{N}$, we have $P_j \in \mathcal{F}(E) \subseteq \mathcal{L}_\Gamma$, so that (2.6) ensures that the set appearing in the definition (5.3) of t_j is non-empty, and further that $t_j \geq \|\Psi_\Gamma\|^{-1} > 0$. To see that $t_{j+1} \geq t_j$, suppose that $P_{j+1} = T \Psi_\Gamma$ for some $T \in \mathcal{B}(E^n, E)$. Then $P_j = (P_j T) \Psi_\Gamma$, so that $t_j \leq \|P_j T\| \leq \|T\|$ by the monotonicity of the Schauder basis for E .

The first part of the final clause (that Δ_γ is compact if $(t_j)_{j \in \mathbb{N}}$ is unbounded) is immediate from Lemma 5.5. We shall prove the second part

by contraposition. Suppose that $\Delta_\gamma \in \mathcal{L}_\Gamma$, so that $\Delta_\gamma = S\Psi_\Gamma$ for some $S \in \mathcal{B}(E^n, E)$. Then, for each $k \in \mathbb{N}$, we have a commutative diagram

$$\begin{array}{ccc} E & \xrightarrow{P_k} & E \\ \Psi_\Gamma \downarrow & \searrow \Delta_\gamma & \uparrow \Theta_{\eta(k)} \\ E^n & \xrightarrow{S} & E, \end{array}$$

where $\eta(k) = (t_j^{1/2})_{j=1}^k$ and the operator $\Theta_{\eta(k)}$ is given by (5.2). Hence, by the definition (5.3) of t_k and Corollary 5.6, we obtain

$$t_k \leq \|\Theta_{\eta(k)}S\| \leq 2t_k^{1/2}\|S\|,$$

which implies that the sequence $(t_j)_{j \in \mathbb{N}}$ is bounded by $4\|S\|^2$. \square

Theorem 5.8. *Let E be a Banach space which is complemented in its bidual and has a Schauder basis. Then $\mathcal{B}(E)$ is the only finitely-generated left ideal of $\mathcal{B}(E)$ which contains $\mathcal{K}(E)$, and hence each finitely-generated, maximal left ideal of $\mathcal{B}(E)$ is fixed.*

Proof. Let $\mathbf{e} = (e_j)_{j \in \mathbb{N}}$ be a Schauder basis for E . By passing to an equivalent norm on E , we may suppose that \mathbf{e} is monotone. Suppose that Γ is a non-empty, finite subset of $\mathcal{B}(E)$ such that $\mathcal{K}(E) \subseteq \mathcal{L}_\Gamma$, and set $n = |\Gamma| \in \mathbb{N}$. Lemma 5.7 implies that the sequence $(t_j)_{j \in \mathbb{N}}$ given by (5.3) is bounded, so that we can find a bounded sequence $(T_j)_{j \in \mathbb{N}}$ in $\mathcal{B}(E^n, E)$ such that $P_j = T_j\Psi_\Gamma$ for each $j \in \mathbb{N}$.

We may identify $\mathcal{B}(E^n, E^{**})$ with the dual space of the projective tensor product $E^n \widehat{\otimes} E^*$; the duality bracket is given by

$$\langle x \otimes \lambda, S \rangle = \langle \lambda, Sx \rangle \quad (x \in E^n, \lambda \in E^*, S \in \mathcal{B}(E^n, E^{**}))$$

(e.g., see [11, Proposition A.3.70]). Hence $\mathcal{B}(E^n, E^{**})$ carries a weak*-topology, with respect to which its unit ball is compact, and so the sequence $(\kappa_E T_j)_{j \in \mathbb{N}}$ has a weak*-accumulation point, say $T \in \mathcal{B}(E^n, E^{**})$. Then, for each $j \in \mathbb{N}$, $\lambda \in E^*$, and $\varepsilon > 0$, we can find an integer $k \geq j$ such that


$$\begin{aligned} \varepsilon &\geq |\langle \Psi_\Gamma e_j \otimes \lambda, T - \kappa_E T_k \rangle| \\ &= |\langle \lambda, (T - \kappa_E T_k)\Psi_\Gamma e_j \rangle| = |\langle \lambda, T\Psi_\Gamma e_j - \kappa_E e_j \rangle|. \end{aligned}$$

Since $\varepsilon > 0$ and $\lambda \in E^*$ were arbitrary, we conclude that $T\Psi_\Gamma e_j = \kappa_E e_j$, and therefore $T\Psi_\Gamma = \kappa_E$. By the assumption, κ_E has a left inverse, say $\Lambda \in \mathcal{B}(E^{**}, E)$. Consequently $I_E = (\Lambda T)\Psi_\Gamma \in \mathcal{L}_\Gamma$, and so $\mathcal{L}_\Gamma = \mathcal{B}(E)$. \square

Example 5.9. Theorem 5.8 implies that, for each of the spaces $E = \ell_p$ or $E = L_p[0, 1]$, where $p \in (1, \infty)$, $\mathcal{B}(E)$ is the only finitely-generated left ideal of $\mathcal{B}(E)$ which contains $\mathcal{K}(E)$, and each finitely-generated, maximal left ideal of $\mathcal{B}(E)$ is fixed. This conclusion is also true for $p = 1$; indeed, ℓ_1 is a dual space, and therefore complemented in its bidual, while $L_1[0, 1]$ is complemented in its bidual by [1, Theorem 6.3.10].

Many other Banach spaces are known to be complemented in their biduals. The following list gives some examples.

- (i) Let E be a Banach space which is isomorphic to a complemented subspace of a dual Banach space; that is, for some Banach space F , there are operators $U \in \mathcal{B}(E, F^*)$ and $V \in \mathcal{B}(F^*, E)$ with $I_E = VU$. Then the diagram



is commutative, which implies that the operator $\kappa_E V \kappa_F^* U^{**}$ is a projection of E^{**} onto $\kappa_E(E)$, so that E is complemented in its bidual.

- (ii) As a special case of (i), suppose that E is a non-zero Banach space for which $\mathcal{B}(E)$ is isomorphic to a complemented subspace of a dual Banach space. Then, as $\mathcal{B}(E)$ contains a complemented subspace isomorphic to E , (i) implies that E is complemented in its bidual.

(It is easy to see that $\mathcal{B}(E)$ contains a complemented subspace isomorphic to E . Indeed, choose $\lambda \in E^*$ and $y \in E$ with $\langle y, \lambda \rangle = 1$, and consider the operators $U_\lambda: E \rightarrow \mathcal{B}(E)$ and $V_y: \mathcal{B}(E) \rightarrow E$ given by $U_\lambda x = x \otimes \lambda$ for $x \in E$ and $V_y(T) = Ty$ for $T \in \mathcal{B}(E)$. They satisfy $V_y U_\lambda = I_E$, so that U_λ is an isomorphism onto its range, and $U_\lambda V_y$ is a projection of $\mathcal{B}(E)$ onto the range of U_λ .)

- (iii) Let E be a Banach lattice which does not contain a subspace isomorphic to c_0 . Then E is complemented in its bidual by [36, Theorem 1.c.4].

Remark 5.10. Theorem 5.8 does not provide any new information for Banach spaces of the form $E = C(K)$, where K is a compact Hausdorff space, because the assumption that $C(K)$ is complemented in its bidual implies that $C(K)$ is injective, so that Example 5.4(ii) already applies.

A slight variation of the proof of Theorem 5.8 gives the following conclusion.

Theorem 5.11. *Let E be a non-zero Banach space with a Schauder basis. Then $\overline{\mathcal{F}(E^*)}$ is the only finitely-generated left ideal of $\mathcal{B}(E^*)$ which contains $\mathcal{F}(E^*)$, and hence each finitely-generated, maximal left ideal of $\mathcal{B}(E^*)$ is fixed.*

Proof. Suppose that $\overline{\mathcal{F}(E^*)} \subseteq \mathcal{L}_\Gamma$, where Γ is a non-empty, finite subset of $\mathcal{B}(E^*)$, and set $n = |\Gamma| \in \mathbb{N}$. As in the proof of Theorem 5.8, we may suppose that E has a monotone Schauder basis \mathbf{e} . Then, arguing as in the proof of Lemma 5.7, we see that the sequence

$$t'_j = \inf \{ \|T\| : T \in \mathcal{B}((E^*)^n, E^*), P_j^* = T\Psi_\Gamma \} \in (0, \infty) \quad (j \in \mathbb{N})$$

is increasing and bounded. (Indeed, if the sequence $(t'_j)_{j \in \mathbb{N}}$ were unbounded, then for $\gamma = ((t'_j)^{-1/2})_{j \in \mathbb{N}}$, we would have $\Delta_\gamma^* \in \overline{\mathcal{F}(E^*)} \setminus \mathcal{L}_\Gamma$, contrary to our assumption.) Consequently there exists a bounded sequence $(T_j)_{j \in \mathbb{N}}$ in $\mathcal{B}((E^*)^n, E^*)$ such that $P_j^* = T_j \Psi_\Gamma$ for each $j \in \mathbb{N}$. Let T be a weak*-accumulation point of $(T_j)_{j \in \mathbb{N}}$, where we have identified $\mathcal{B}((E^*)^n, E^*)$ with the dual space of the projective tensor product $E \widehat{\otimes} (E^*)^n$ via the duality bracket given by

$$(5.4) \quad \langle x \otimes \mu, S \rangle = \langle x, S\mu \rangle \quad (x \in E, \mu \in (E^*)^n, S \in \mathcal{B}((E^*)^n, E^*)).$$

For each $x \in E$, $\lambda \in E^*$, and $\varepsilon > 0$, we can find $j_0 \in \mathbb{N}$ such that

$$\|x - P_j x\| \leq \varepsilon(2\|\lambda\| + 1)^{-1}$$

whenever $j \geq j_0$. Choosing $j \geq j_0$ such that $|\langle x \otimes \Psi_\Gamma \lambda, T - T_j \rangle| \leq \varepsilon/2$ and applying (5.4), we then obtain

$$\begin{aligned} |\langle x, (T\Psi_\Gamma - I_{E^*})\lambda \rangle| &\leq |\langle x, (T\Psi_\Gamma - P_j^*)\lambda \rangle| + |\langle x, (I_E - P_j)^*\lambda \rangle| \\ &\leq |\langle x, (T - T_j)\Psi_\Gamma \lambda \rangle| + |\langle x - P_j x, \lambda \rangle| \leq \varepsilon. \end{aligned}$$

This implies that $T\Psi_\Gamma = I_{E^*}$, and therefore $\mathcal{L}_\Gamma = \mathcal{B}(E^*)$. \square

Example 5.12. Theorem 5.11 applies in the following two cases which have not already been resolved:

- (i) $E = X \widehat{\otimes} X^*$, where X is a Banach space with a shrinking Schauder basis (this ensures that E has a Schauder basis). Then E^* is isomorphic to $\mathcal{B}(X^*)$, so that the conclusion is that each finitely-generated, maximal left ideal of $\mathcal{B}(\mathcal{B}(X^*))$ is fixed. The most important case is where X , and hence X^* , is a separable, infinite-dimensional Hilbert space; in this case $\mathcal{B}(X^*)$ does not have the approximation property [51], which gives this example a very different flavour from Examples 5.4 and 5.9, above.
- (ii) $E = \left(\bigoplus_{n \in \mathbb{N}} E_n\right)_{\ell_1}$, where $(E_n)_{n \in \mathbb{N}}$ is a sequence of Banach spaces with Schauder bases whose basis constants are uniformly bounded. Then E^* is isomorphic to $\left(\bigoplus_{n \in \mathbb{N}} E_n^*\right)_{\ell_\infty}$, and so the conclusion is that each finitely-generated, maximal left ideal of $\mathcal{B}\left(\left(\bigoplus_{n \in \mathbb{N}} E_n^*\right)_{\ell_\infty}\right)$ is fixed.

The conditions imposed on the Banach space E in Theorems 5.8 and 5.11 are clearly preserved under the formation of finite direct sums. In contrast, this need not be the case for the condition of Corollary 5.3. For instance, c_0 and ℓ_∞ both satisfy this condition by Example 5.4(ii)–(iii), whereas their direct sum $c_0 \oplus \ell_\infty$ does not. We shall explore this situation in greater depth in Section 7. Notably, as a particular instance of Theorem 7.3, we shall see that the main conclusion of Corollary 5.3 fails for $E = c_0 \oplus \ell_\infty$ because $\mathcal{F}(c_0 \oplus \ell_\infty)$ is contained in a proper, closed, singly-generated left ideal of $\mathcal{B}(c_0 \oplus \ell_\infty)$.

We do not know the answer to Question (II) for $E = c_0 \oplus \ell_\infty$, but the following result answers this question positively for another direct sum arising

naturally from Example 5.4, with the ideal $\mathcal{S}(E)$ of strictly singular operators taking the role that was played by $\mathcal{F}(E)$ in Corollary 5.3 and $\mathcal{K}(E)$ in Theorem 5.8.

Proposition 5.13. *Let $E = c_0(\mathbb{I}) \oplus H$, where \mathbb{I} is a non-empty set and H is a Hilbert space. Then $\mathcal{B}(E)$ is the only finitely-generated left ideal of $\mathcal{B}(E)$ which contains $\mathcal{S}(E)$, and hence each finitely-generated, maximal left ideal of $\mathcal{B}(E)$ is fixed.*

Proof. Let \mathcal{L} be a finitely-generated left ideal of $\mathcal{B}(E)$ such that $\mathcal{S}(E)$ is contained in \mathcal{L} . We may suppose that \mathbb{I} is infinite and H is infinite-dimensional. Proposition 2.2 implies that \mathcal{L} is generated by a single operator $T \in \mathcal{B}(E)$, say, while Corollary 4.7 shows that T is bounded below and thus is an upper semi-Fredholm operator.

We can represent T as a matrix of operators:

$$T = \begin{pmatrix} T_{1,1}: c_0(\mathbb{I}) \rightarrow c_0(\mathbb{I}) & T_{1,2}: H \rightarrow c_0(\mathbb{I}) \\ T_{2,1}: c_0(\mathbb{I}) \rightarrow H & T_{2,2}: H \rightarrow H \end{pmatrix}.$$

Each operator from H to $c_0(\mathbb{I})$ is strictly singular because no infinite-dimensional subspace of $c_0(\mathbb{I})$ is isomorphic to a Hilbert space. Similarly, each operator from $c_0(\mathbb{I})$ to H is strictly singular. Hence, by [35, Proposition 2.c.10],

$$T - \begin{pmatrix} 0 & T_{1,2} \\ T_{2,1} & 0 \end{pmatrix} = \begin{pmatrix} T_{1,1} & 0 \\ 0 & T_{2,2} \end{pmatrix}$$

is an upper semi-Fredholm operator, which clearly implies that $T_{1,1}$ and $T_{2,2}$ are upper semi-Fredholm operators. Let $P_1 \in \mathcal{F}(c_0(\mathbb{I}))$ and $P_2 \in \mathcal{F}(H)$ be projections onto the kernels of $T_{1,1}$ and $T_{2,2}$, respectively. Then

$$\tilde{T}_{1,1}: x \mapsto T_{1,1}x, \quad \ker P_1 \rightarrow T_{1,1}(c_0(\mathbb{I})),$$

is an isomorphism, so that $T_{1,1}(c_0(\mathbb{I}))$ is isomorphic to $\ker P_1$, which in turn is isomorphic to $c_0(\mathbb{I})$ (because $\ker P_1$ has finite codimension in $c_0(\mathbb{I})$). Consequently, as in Example 5.4(iii), $T_{1,1}(c_0(\mathbb{I}))$ is complemented in $c_0(\mathbb{I})$, so that we can extend the inverse of $\tilde{T}_{1,1}$ to obtain an operator $S_1 \in \mathcal{B}(c_0(\mathbb{I}))$ which satisfies $S_1 T_{1,1} = I_{c_0(\mathbb{I})} - P_1$. Similarly, we can find an operator $S_2 \in \mathcal{B}(H)$ such that $S_2 T_{2,2} = I_H - P_2$. In conclusion, we have

$$I_E = \begin{pmatrix} S_1 & 0 \\ 0 & S_2 \end{pmatrix} T + \begin{pmatrix} P_1 & -S_1 T_{1,2} \\ -S_2 T_{2,1} & P_2 \end{pmatrix} \in \mathcal{L} + \mathcal{S}(E) = \mathcal{L},$$

and thus $\mathcal{L} = \mathcal{B}(E)$.

Since $\mathcal{S}(E) \subseteq \mathcal{E}(E)$, the final clause follows immediately from Corollary 4.1. \square

6. ‘EXOTIC’ BANACH SPACES FOR WHICH EACH FINITELY-GENERATED, MAXIMAL LEFT IDEAL IS FIXED

In this section, we shall answer Question (II) positively for two classes of custom-made Banach spaces of a distinctly non-classical nature, using an approach which is completely different from the one taken in Section 5. More precisely, for each Banach space E in either of these two classes, we are able

to describe all the maximal left ideals of $\mathcal{B}(E)$ explicitly, and it will then follow easily that only the fixed maximal left ideals are finitely generated. The reason that we can describe all the maximal left ideals of $\mathcal{B}(E)$ is, roughly speaking, that $\mathcal{B}(E)$ is ‘small’. As we shall see, in both cases each non-fixed, maximal left ideal of $\mathcal{B}(E)$ is a two-sided ideal of codimension one.

We begin with a lemma which can be viewed as a counter-part of Corollary 4.8 for left ideals of strictly singular operators.

Lemma 6.1. *Let E be a Banach space, and let \mathcal{L} be a left ideal of $\mathcal{B}(E)$ such that $\mathcal{F}(E) \subseteq \mathcal{L} \subseteq \mathcal{S}(E)$. Then the following three conditions are equivalent:*

- (a) \mathcal{L} is finitely generated;
- (b) $\mathcal{L} = \mathcal{B}(E)$;
- (c) E is finite-dimensional.

Proof. The implications (c) \Rightarrow (b) \Rightarrow (a) are clear.

To see that (a) implies (c), suppose that $\mathcal{L} = \mathcal{L}_\Gamma$ for some non-empty, finite subset Γ of $\mathcal{B}(E)$. Corollary 4.7 implies that the operator Ψ_Γ is bounded below, while (2.5) and the fact that $\Gamma \subseteq \mathcal{S}(E)$ show that Ψ_Γ is strictly singular. Hence the domain E of Ψ_Γ is finite-dimensional. \square

A Banach space E has *few operators* if E is infinite-dimensional and each operator on E is the sum of a scalar multiple of the identity operator and a strictly singular operator; that is, $\mathcal{B}(E) = \mathbb{C}I_E + \mathcal{S}(E)$. Gowers and Maurey [19] showed that each hereditarily indecomposable Banach space has few operators, and constructed the first example of such a space.

Theorem 6.2. *Let E be a Banach space which has few operators. Then $\mathcal{S}(E)$ is the unique non-fixed, maximal left ideal of $\mathcal{B}(E)$, and $\mathcal{S}(E)$ is not finitely generated as a left ideal.*

Proof. Let \mathcal{L} be a maximal left ideal of $\mathcal{B}(E)$, and suppose that \mathcal{L} is not fixed. Then, by Corollary 4.1, \mathcal{L} contains $\mathcal{E}(E)$ and hence $\mathcal{S}(E)$, which has codimension one in $\mathcal{B}(E)$, so that $\mathcal{L} = \mathcal{E}(E) = \mathcal{S}(E)$. This proves the first clause. The second clause follows from Lemma 6.1. \square

To set the scene for our second result, we begin with a short excursion into the theory of semi-direct products of Banach algebras. Let \mathcal{B} be a Banach algebra, and let \mathcal{C} and \mathcal{I} be a closed subalgebra and a closed, two-sided ideal of \mathcal{B} , respectively. Then \mathcal{B} is the *semi-direct product* of \mathcal{C} and \mathcal{I} if \mathcal{C} and \mathcal{I} are complementary subspaces of \mathcal{B} ; that is, $\mathcal{C} + \mathcal{I} = \mathcal{B}$ and $\mathcal{C} \cap \mathcal{I} = \{0\}$. In this case, we denote by $\rho: \mathcal{B} \rightarrow \mathcal{C}$ the projection of \mathcal{B} onto \mathcal{C} along \mathcal{I} . This is an algebra homomorphism, as is easy to check. It is relevant for our purposes because it induces an isomorphism between the following two lattices of closed left ideals

$$(6.1) \quad \text{Lat}_{\mathcal{I}}(\mathcal{B}) = \{ \mathcal{L} : \mathcal{L} \text{ is a closed left ideal of } \mathcal{B} \text{ such that } \mathcal{I} \subseteq \mathcal{L} \}$$

and

$$(6.2) \quad \text{Lat}(\mathcal{C}) = \{ \mathcal{N} : \mathcal{N} \text{ is a closed left ideal of } \mathcal{C} \}.$$

More precisely, for each $\mathcal{L} \in \text{Lat}_{\mathcal{I}}(\mathcal{B})$, we have $\rho(\mathcal{L}) = \mathcal{L} \cap \mathcal{C} \in \text{Lat}(\mathcal{C})$, and the mapping $\mathcal{L} \mapsto \rho(\mathcal{L})$ is a lattice isomorphism between $\text{Lat}_{\mathcal{I}}(\mathcal{B})$ and $\text{Lat}(\mathcal{C})$; its inverse is given by $\mathcal{N} \mapsto \mathcal{N} + \mathcal{I}$. Suppose that the left ideal $\mathcal{L} \in \text{Lat}_{\mathcal{I}}(\mathcal{B})$ is generated by a subset Γ of \mathcal{B} . Then evidently $\rho(\mathcal{L})$ is generated by the subset $\rho(\Gamma)$ of \mathcal{C} , so that ρ maps each closed, finitely-generated left ideal of \mathcal{B} containing \mathcal{I} to a closed, finitely-generated left ideal of \mathcal{C} .

We shall next state two classical results about $C(K)$ -spaces. The first is due to Pełczyński [41, Theorem 1], and characterizes the weakly compact operators from a $C(K)$ -space into an arbitrary Banach space.

Theorem 6.3. *Let K be a non-empty, compact Hausdorff space, and let E be a Banach space. Then the following three conditions are equivalent for each operator $T \in \mathcal{B}(C(K), E)$:*

- (a) T is weakly compact;
- (b) T is strictly singular;
- (c) T does not fix a copy of c_0 .

The second result describes the maximal ideals of the Banach algebra $C(K)$, as well as the finitely-generated ones. (Note that the notions of a left, right, and two-sided ideal coincide in $C(K)$ because $C(K)$ is commutative.) Given a point $k \in K$, we write $\varepsilon_k: C(K) \rightarrow \mathbb{C}$ for the evaluation map at k ; that is, $\varepsilon_k(f) = f(k)$ for each $f \in C(K)$. This is a surjective algebra homomorphism of norm one.

Theorem 6.4. *Let K be a compact Hausdorff space. Then:*

- (i) *each maximal ideal of $C(K)$ has the form $\ker \varepsilon_k$ for a unique point $k \in K$;*
- (ii) *the maximal ideal $\ker \varepsilon_k$ is finitely generated if and only if the point k is isolated in K .*

Proof. The first clause is folklore (e.g., see [11, Theorem 4.2.1(i)]), while the second is the complex-valued counter-part of a classical theorem of Gillman [16, Corollary 5.4]. Both clauses are also easy to verify directly. \square

We require one further notion before we can present our result. For a non-empty, compact Hausdorff space K and a function $g \in C(K)$, we denote by $M_g \in \mathcal{B}(C(K))$ the multiplication operator given by g ; that is, $M_g f = gf$ for each $f \in C(K)$. The mapping

$$(6.3) \quad \mu: g \mapsto M_g, \quad C(K) \rightarrow \mathcal{B}(C(K)),$$

is an isometric, unital algebra homomorphism. An operator $T \in \mathcal{B}(C(K))$ is a *weak multiplication* if it has the form $T = M_g + S$ for some $g \in C(K)$ and $S \in \mathcal{W}(C(K))$. The fourth-named author [31, Theorem 6.1] (assuming the continuum hypothesis) and Plebanek [43, Theorem 1.3] (without any assumptions beyond ZFC) have constructed an example of a connected, compact Hausdorff space K for which each operator on $C(K)$ is a weak multiplication. This ensures that the following theorem is not vacuous.

Theorem 6.5. *Let K be a compact Hausdorff space without isolated points and such that each operator on $C(K)$ is a weak multiplication.*

- (i) *The Banach algebra $\mathcal{B}(C(K))$ is the semi-direct product of the subalgebra $\mu(C(K))$ and the ideal $\mathcal{W}(C(K))$, where μ is the homomorphism given by (6.3).*
- (ii) *Let \mathcal{L} be a subset of $\mathcal{B}(C(K))$. Then the following four conditions are equivalent:*
 - (a) *\mathcal{L} is a non-fixed, maximal left ideal of $\mathcal{B}(C(K))$;*
 - (b) *\mathcal{L} is a maximal left ideal of $\mathcal{B}(C(K))$, and \mathcal{L} is not finitely generated;*
 - (c) *\mathcal{L} is a maximal two-sided ideal of $\mathcal{B}(C(K))$;*
 - (d) *$\mathcal{L} = \{M_g + S : S \in \mathcal{W}(C(K)) \text{ and } g \in C(K) \text{ with } g(k) = 0 \text{ for some } k \in K\}$*

In the positive case, the point $k \in K$ such that (d) holds is uniquely determined by \mathcal{L} .

Proof. (i). We have $\mathcal{B}(C(K)) = \mu(C(K)) + \mathcal{W}(C(K))$ because each operator on $C(K)$ is a weak multiplication. Theorem 6.3 allows us to replace $\mathcal{W}(C(K))$ with $\mathcal{S}(C(K))$, which we shall do in the remainder of this proof because the latter ideal suits our approach better.

To see that $\mu(C(K)) \cap \mathcal{S}(C(K)) = \{0\}$, suppose that $g \in C(K) \setminus \{0\}$. Take $k_0 \in K$ such that $g(k_0) \neq 0$, set $\varepsilon = |g(k_0)|/2 > 0$, and choose an open neighbourhood N of k_0 such that $|g(k)| \geq \varepsilon$ for each $k \in N$. Using Urysohn's lemma and the fact that k_0 is not isolated in K , we deduce that the subspace

$$F = \{f \in C(K) : f(k) = 0 \quad (k \in K \setminus N)\}$$

of $C(K)$ is infinite-dimensional. Since

$$\|M_g f\| = \sup\{|g(k)f(k)| : k \in N\} \geq \varepsilon \|f\| \quad (f \in F),$$

we conclude that M_g is not strictly singular, as required.

(ii). For each $k \in K$, let

$$\mathcal{Z}_k = \mu(\ker \varepsilon_k) + \mathcal{S}(C(K)),$$

so that

$$\mathcal{Z}_k = \{M_g + S : S \in \mathcal{S}(C(K)) \text{ and } g \in C(K) \text{ with } g(k) = 0\}.$$

By (i), \mathcal{Z}_k is a two-sided ideal of codimension one in $\mathcal{B}(C(K))$, and thus is maximal both as a left and a two-sided ideal. The implication (d) \Rightarrow (c) is now immediate, while (d) \Rightarrow (b) follows because $\rho(\mathcal{Z}_k) = \mu(\ker \varepsilon_k)$ is not finitely generated by Theorem 6.4(ii), so that \mathcal{Z}_k is not finitely generated as a left ideal, as explained in the paragraph following (6.2).

The implication (b) \Rightarrow (a) is clear because each fixed, maximal left ideal is finitely generated by Proposition 2.4(i).

(a) \Rightarrow (d). Suppose that \mathcal{L} is a non-fixed, maximal left ideal of $\mathcal{B}(C(K))$. Then, by Corollary 4.1, \mathcal{L} contains $\mathcal{E}(C(K))$ and thus $\mathcal{S}(C(K))$, so that \mathcal{L} is a maximal element of the lattice $\text{Lat}_{\mathcal{S}(C(K))}(\mathcal{B}(C(K)))$ given by (6.1).

Hence, in the notation of (6.2), there is a maximal element \mathcal{N} of the lattice $\text{Lat}(\mu(C(K)))$ such that $\mathcal{L} = \mathcal{N} + \mathcal{S}(C(K))$. Theorem 6.4(i) implies that $\mathcal{N} = \mu(\ker \varepsilon_k)$ for some $k \in K$, and consequently $\mathcal{L} = \mathcal{L}_k$.

(c) \Rightarrow (d). Suppose that \mathcal{L} is a maximal two-sided ideal of $\mathcal{B}(C(K))$. Then, as mentioned in Remark 4.2, \mathcal{L} contains $\mathcal{E}(C(K))$ and hence \mathcal{L} contains $\mathcal{S}(C(K))$, so that $\mathcal{L} = \mathcal{L}_k$ for some $k \in K$ by (i) and Theorem 6.4(i).

The final clause follows because $\ker \varepsilon_{k_1} \neq \ker \varepsilon_{k_2}$ whenever $k_1, k_2 \in K$ are distinct, and hence also $\mathcal{L}_{k_1} \neq \mathcal{L}_{k_2}$. \square

Remark 6.6. Example 5.4(ii)–(iii) and Theorem 6.5 show that there are compact Hausdorff spaces K such that Question (II) has a positive answer for $E = C(K)$. However, this question remains open for some very important $C(K)$ -spaces. Indeed, it is known that $C(K)$ contains a closed subspace which is isomorphic to $C(K)$ and which is not complemented in $C(K)$ for each of the following compact Hausdorff spaces K :

- (i) $K = [0, 1]$ (see [2]);
- (ii) $K = [0, \alpha]$ for any ordinal $\alpha \geq \omega^\omega$, where $[0, \alpha]$ denotes the set of ordinals less than or equal to α , equipped with the order topology. (Baker [5] showed this in the case where $\alpha = \omega^\omega$; the conclusion for general $\alpha \geq \omega^\omega$ follows immediately from Baker's result because $C[0, \alpha]$ is isomorphic to $C[0, \alpha] \oplus C[0, \omega^\omega]$.)

Hence, by Corollary 4.7, $\mathcal{F}(C(K))$ is contained in a singly-generated, proper left ideal of $\mathcal{B}(C(K))$ for each of these K , but we do not know whether such a left ideal can be chosen also to be maximal (or even closed).

This question cannot be answered by a variant of Theorem 5.8 because we can strengthen the above conclusion to obtain that $\mathcal{H}(C(K))$ is contained in a singly-generated, proper left ideal of $\mathcal{B}(C(K))$ for each of the above K . To see this, take an operator $U \in \mathcal{B}(C(K))$ which is bounded below and whose range $F = U(C(K))$ is not complemented in $C(K)$, and consider the isomorphism $\tilde{U}: x \mapsto Ux, C(K) \rightarrow F$. Then, for each $S \in \mathcal{H}(C(K))$, the operator $S\tilde{U}^{-1}: F \rightarrow C(K)$ has an extension $T \in \mathcal{H}(C(K))$ by a theorem of Grothendieck (see [22, pp. 559–560], or [34, Theorem 1]). Hence we have $S = TU$, and consequently $\mathcal{H}(C(K)) \subseteq \mathcal{L}_{\{U\}}$.

7. A NON-FIXED AND SINGLY-GENERATED, MAXIMAL LEFT IDEAL OF OPERATORS

The main aim of this section is to prove Theorems 1.5 and 1.6. Several parts of those theorems are special cases of more general results, which may be of independent interest, and so we shall take a more general approach, specializing only when we need to.

Recall that, for a non-empty set \mathbb{I} , we denote by $\ell_\infty(\mathbb{I})$ the Banach space of bounded, complex-valued functions defined on \mathbb{I} , and $\ell_\infty = \ell_\infty(\mathbb{N})$. Our first result collects some known facts about operators from $\ell_\infty(\mathbb{I})$ that we shall use several times.

Lemma 7.1. *Let \mathbb{I} be a non-empty set, and let X be a Banach space.*

- (i) *An operator from $\ell_\infty(\mathbb{I})$ to X is weakly compact if and only if it is strictly singular.*
- (ii) *Suppose that the set \mathbb{I} is infinite. Then each operator from $\ell_\infty(\mathbb{I})$ to X is weakly compact if and only if X does not contain a subspace isomorphic to ℓ_∞ .*

Proof. (i). This is a special case of Theorem 6.3.

(ii). The hard part is the implication \Leftarrow , which however follows immediately from [35, Proposition 2.f.4].

The forward implication is straightforward. Indeed, suppose contrapositively that X contains a subspace which is isomorphic to ℓ_∞ , and take an operator $U \in \mathcal{B}(\ell_\infty, X)$ which is bounded below. Choose an injective mapping $\theta: \mathbb{N} \rightarrow \mathbb{I}$, and define an operator $C_\theta \in \mathcal{B}(\ell_\infty(\mathbb{I}), \ell_\infty)$ by $C_\theta f = f \circ \theta$ for each $f \in \ell_\infty(\mathbb{I})$. Then UC_θ is not weakly compact, for instance because it fixes a copy of ℓ_∞ . \square

In the remainder of this section we shall consider a Banach space X such that

- (I) the bidual of X is isomorphic to $\ell_\infty(\mathbb{I}_1)$ for some infinite set \mathbb{I}_1 via a fixed isomorphism $V: X^{**} \rightarrow \ell_\infty(\mathbb{I}_1)$; and
- (II) no subspace of X is isomorphic to ℓ_∞ .

For example, $X = c_0$ satisfies both of these conditions with $\mathbb{I}_1 = \mathbb{N}$.

Let \mathbb{I}_2 be a disjoint copy of \mathbb{I}_1 (that is, \mathbb{I}_2 is a set of the same cardinality as \mathbb{I}_1 and satisfies $\mathbb{I}_1 \cap \mathbb{I}_2 = \emptyset$), and set $\mathbb{I} = \mathbb{I}_1 \cup \mathbb{I}_2$. We consider $\ell_\infty(\mathbb{I}_1)$ and $\ell_\infty(\mathbb{I}_2)$ as complementary subspaces of $\ell_\infty(\mathbb{I})$ in the natural way, and denote by P_1 and P_2 the corresponding projections of $\ell_\infty(\mathbb{I})$ onto $\ell_\infty(\mathbb{I}_1)$ and $\ell_\infty(\mathbb{I}_2)$, respectively. Moreover, we shall choose a bijection $\varphi: \mathbb{I}_2 \rightarrow \mathbb{I}_1$; we then obtain an isometric isomorphism C_φ of $\ell_\infty(\mathbb{I})$ onto the subspace $\ell_\infty(\mathbb{I}_2)$ by the definition $C_\varphi f = f \circ \varphi$ for each $f \in \ell_\infty(\mathbb{I})$.

Let $E = X \oplus \ell_\infty(\mathbb{I})$ with norm $\|(x, f)\|_E = \max\{\|x\|_X, \|f\|_\infty\}$. We identify operators T on E with (2×2) -matrices

$$\begin{pmatrix} T_{1,1}: X \rightarrow X & T_{1,2}: \ell_\infty(\mathbb{I}) \rightarrow X \\ T_{2,1}: X \rightarrow \ell_\infty(\mathbb{I}) & T_{2,2}: \ell_\infty(\mathbb{I}) \rightarrow \ell_\infty(\mathbb{I}) \end{pmatrix}.$$

Note that assumption (II) and Lemma 7.1(ii) imply that the operator $T_{1,2}$ is always weakly compact. This fact will play a key role for us.

Despite our focus on left ideals, our first result about the Banach space E is concerned with two-sided ideals.

Proposition 7.2. (i) *The set*

$$\mathcal{W}_1 = \left\{ \begin{pmatrix} T_{1,1} & T_{1,2} \\ T_{2,1} & T_{2,2} \end{pmatrix} \in \mathcal{B}(E) : T_{1,1} \in \mathcal{W}(X) \right\}$$

is a proper, closed two-sided ideal of $\mathcal{B}(E)$, and \mathcal{W}_1 is a maximal two-sided ideal of $\mathcal{B}(E)$ if and only if $\mathcal{W}(X)$ is a maximal two-sided ideal of $\mathcal{B}(X)$.

(ii) *The set*

$$\mathscr{W}_2 = \left\{ \begin{pmatrix} T_{1,1} & T_{1,2} \\ T_{2,1} & T_{2,2} \end{pmatrix} \in \mathscr{B}(E) : T_{2,2} \in \mathscr{W}(\ell_\infty(\mathbb{I})) \right\}$$

is a proper, closed two-sided ideal of $\mathscr{B}(E)$, and the following three conditions are equivalent:

- (a) \mathscr{W}_2 *is a maximal two-sided ideal of $\mathscr{B}(E)$;*
- (b) $\mathscr{W}(\ell_\infty(\mathbb{I}))$ *is a maximal two-sided ideal of $\mathscr{B}(\ell_\infty(\mathbb{I}))$;*
- (c) \mathbb{I} *is countable.*

Proof. (i). The mapping

$$T \mapsto T_{1,1} + \mathscr{W}(X), \quad \mathscr{B}(E) \rightarrow \mathscr{B}(X)/\mathscr{W}(X),$$

is a surjective algebra homomorphism of norm one. Hence its kernel, which is equal to \mathscr{W}_1 , is a closed two-sided ideal of $\mathscr{B}(E)$. This ideal is proper because X is non-reflexive by assumption (I). The fundamental isomorphism theorem implies that the Banach algebras $\mathscr{B}(E)/\mathscr{W}_1$ and $\mathscr{B}(X)/\mathscr{W}(X)$ are isomorphic, and so $\mathscr{B}(E)/\mathscr{W}_1$ is simple if and only if $\mathscr{B}(X)/\mathscr{W}(X)$ is simple. Consequently \mathscr{W}_1 is a maximal two-sided ideal of $\mathscr{B}(E)$ if and only if $\mathscr{W}(X)$ is a maximal two-sided ideal of $\mathscr{B}(X)$.

(ii). An obvious modification of the argument given above shows that \mathscr{W}_2 is a proper, closed two-sided ideal of $\mathscr{B}(E)$, and that conditions (a) and (b) are equivalent. The implication (c) \Rightarrow (b) follows from [35, Proposition 2.f.4].

Conversely, to prove that (b) \Rightarrow (c), suppose that $\mathscr{W}(\ell_\infty(\mathbb{I}))$ is a maximal two-sided ideal of $\mathscr{B}(\ell_\infty(\mathbb{I}))$, and denote by $\mathscr{G}_{\ell_\infty}(\ell_\infty(\mathbb{I}))$ the set of operators on $\ell_\infty(\mathbb{I})$ that factor through ℓ_∞ . This is a two-sided ideal of $\mathscr{B}(\ell_\infty(\mathbb{I}))$ because ℓ_∞ is isomorphic to $\ell_\infty \oplus \ell_\infty$. Hence $\mathscr{G}_{\ell_\infty}(\ell_\infty(\mathbb{I})) + \mathscr{W}(\ell_\infty(\mathbb{I}))$ is also a two-sided ideal, which is strictly greater than $\mathscr{W}(\ell_\infty(\mathbb{I}))$ because $\ell_\infty(\mathbb{I})$ contains a complemented copy of ℓ_∞ , and any projection with range isomorphic to ℓ_∞ belongs to $\mathscr{G}_{\ell_\infty}(\ell_\infty(\mathbb{I})) \setminus \mathscr{W}(\ell_\infty(\mathbb{I}))$. Consequently, by the maximality of $\mathscr{W}(\ell_\infty(\mathbb{I}))$, there are operators $R \in \mathscr{G}_{\ell_\infty}(\ell_\infty(\mathbb{I}))$ and $S \in \mathscr{W}(\ell_\infty(\mathbb{I}))$ such that $I_{\ell_\infty(\mathbb{I})} = R + S$. Then $R = I_{\ell_\infty(\mathbb{I})} - S$ is a Fredholm operator by [35, Proposition 2.c.10] and Lemma 7.1(i), and this implies that $I_{\ell_\infty(\mathbb{I})} = URT$ for some operators $T, U \in \mathscr{B}(\ell_\infty(\mathbb{I}))$ because $\ell_\infty(\mathbb{I})$ is isomorphic to its hyperplanes. Thus the identity operator on $\ell_\infty(\mathbb{I})$ factors through ℓ_∞ , which is possible only if \mathbb{I} is countable. \square

Set

$$(7.1) \quad L = \begin{pmatrix} 0 & 0 \\ V\kappa_X & C_\varphi \end{pmatrix} \in \mathscr{B}(E),$$

where the operators V and C_φ were introduced on p. 29. Since the ranges of V and C_φ are contained in the complementary subspaces $\ell_\infty(\mathbb{I}_1)$ and $\ell_\infty(\mathbb{I}_2)$ of $\ell_\infty(\mathbb{I})$, respectively, we have

$$\begin{aligned} \|L(x, f)\|_E &= \|V\kappa_X x + C_\varphi f\|_\infty = \max\{\|V\kappa_X x\|_\infty, \|C_\varphi f\|_\infty\} \\ &= \max\{\|V\kappa_X x\|_\infty, \|f\|_\infty\} \quad (x \in X, f \in \ell_\infty(\mathbb{I})), \end{aligned}$$

which shows that the operator L is bounded below because $V\kappa_X$ is bounded below. This conclusion is also immediate from our next result and Corollary 4.7.

Theorem 7.3. *The ideal \mathscr{W}_1 defined in Proposition 7.2(i) is the left ideal generated by the operator L given by (7.1); that is,*

$$\mathscr{W}_1 = \mathcal{L}_{\{L\}}.$$

Proof. We have $L \in \mathscr{W}_1$ because $L_{1,1} = 0$, and hence the inclusion \supseteq follows.

We shall prove the reverse inclusion in three steps. First, we see that

$$\begin{pmatrix} 0 & 0 \\ 0 & I_{\ell_\infty(\mathbb{I})} \end{pmatrix} = \begin{pmatrix} 0 & 0 \\ 0 & C_\varphi^{-1}P_2 \end{pmatrix} \begin{pmatrix} 0 & 0 \\ V\kappa_X & C_\varphi \end{pmatrix} \in \mathcal{L}_{\{L\}},$$

and consequently we have

$$(7.2) \quad \begin{pmatrix} 0 & T_{1,2} \\ 0 & T_{2,2} \end{pmatrix} = \begin{pmatrix} 0 & T_{1,2} \\ 0 & T_{2,2} \end{pmatrix} \begin{pmatrix} 0 & 0 \\ 0 & I_{\ell_\infty(\mathbb{I})} \end{pmatrix} \in \mathcal{L}_{\{L\}}$$

for each $T_{1,2} \in \mathcal{B}(\ell_\infty(\mathbb{I}), X)$ and $T_{2,2} \in \mathcal{B}(\ell_\infty(\mathbb{I}))$.

Second, let $T_{2,1} \in \mathcal{B}(X, \ell_\infty(\mathbb{I}))$. Being bounded below, the operator $V\kappa_X$ is an isomorphism onto its range $Y := V\kappa_X(X) \subseteq \ell_\infty(\mathbb{I}_1)$, so that it has an inverse $R \in \mathcal{B}(Y, X)$. By the injectivity of $\ell_\infty(\mathbb{I}_1)$, the composite operator $T_{2,1}R \in \mathcal{B}(Y, \ell_\infty(\mathbb{I}))$ extends to an operator $S \in \mathcal{B}(\ell_\infty(\mathbb{I}_1), \ell_\infty(\mathbb{I}))$, which then satisfies $SV\kappa_X = T_{2,1}$. Hence we have

$$(7.3) \quad \begin{pmatrix} 0 & 0 \\ T_{2,1} & 0 \end{pmatrix} = \begin{pmatrix} 0 & 0 \\ 0 & SP_1 \end{pmatrix} \begin{pmatrix} 0 & 0 \\ V\kappa_X & C_\varphi \end{pmatrix} \in \mathcal{L}_{\{L\}}.$$

Third, each operator $T_{1,1} \in \mathscr{W}(X)$ satisfies $T_{1,1}^{**}(X^{**}) \subseteq \kappa_X(X)$ (e.g., see [38, Theorem 3.5.8]). We can therefore define an operator $U \in \mathcal{B}(\ell_\infty(\mathbb{I}_1), X)$ by

$$Uf = \kappa_X^{-1}T_{1,1}^{**}V^{-1}f \quad (f \in \ell_\infty(\mathbb{I}_1)).$$

Since $\kappa_X UV\kappa_X = T_{1,1}^{**}\kappa_X = \kappa_X T_{1,1}$, we have $UV\kappa_X = T_{1,1}$, and so

$$(7.4) \quad \begin{pmatrix} T_{1,1} & 0 \\ 0 & 0 \end{pmatrix} = \begin{pmatrix} 0 & UP_1 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} 0 & 0 \\ V\kappa_X & C_\varphi \end{pmatrix} \in \mathcal{L}_{\{L\}}.$$

Combining (7.2)–(7.4), we conclude that each operator $T \in \mathscr{W}_1$ belongs to $\mathcal{L}_{\{L\}}$. \square

Remark 7.4. Since the operator L given by (7.1) is bounded below and generates a proper left ideal of $\mathcal{B}(E)$, its range is not complemented in E by Lemma 5.1. This is also easy to verify directly.

A Banach space F has *very few operators* if F is infinite-dimensional and each operator on F is the sum of a scalar multiple of the identity operator and a compact operator; that is, $\mathcal{B}(F) = \mathbb{C}I_F + \mathcal{K}(F)$. Argyros and Haydon [4] constructed the first example of a Banach space X_{AH} which has very few operators. We shall now specialize to the case where $X = X_{\text{AH}}$. The following result collects the properties of X_{AH} that we shall require.

Theorem 7.5 (Argyros and Haydon). *There is a Banach space X_{AH} with the following three properties:*

- (i) X_{AH} has very few operators;
- (ii) X_{AH} has a Schauder basis;
- (iii) the dual space of X_{AH} is isomorphic to ℓ_1 .

Using this, we can easily prove Theorem 1.5.

Proof of Theorem 1.5. We begin by checking that X_{AH} satisfies the two assumptions made on p. 29: indeed, Theorem 7.5(iii) ensures that X_{AH}^{**} is isomorphic to ℓ_∞ , while Theorem 7.5(ii) (or (iii)) implies that X_{AH} does not contain ℓ_∞ . Moreover, we see that $\mathscr{W}(X_{\text{AH}}) = \mathscr{K}(X_{\text{AH}})$ because Theorem 7.5(iii) implies that X_{AH} is non-reflexive, so that $\mathscr{W}(X_{\text{AH}})$ is a closed, non-zero, proper two-sided ideal of $\mathscr{B}(X_{\text{AH}})$, and $\mathscr{K}(X_{\text{AH}})$ is the *only* such ideal by Theorem 7.5(i)–(ii). Hence the set \mathscr{K}_1 given by (1.2) is equal to the ideal \mathscr{W}_1 defined in Proposition 7.2(i), and \mathscr{W}_1 is singly generated as a left ideal by Theorem 7.3. Theorem 7.5(i) implies that \mathscr{K}_1 has codimension one in $\mathscr{B}(E)$, so that it is trivially maximal as a left, right, and two-sided ideal. (The latter also follows from Proposition 7.2(i).) Being a non-zero, two-sided ideal, \mathscr{K}_1 contains $\mathscr{F}(E)$, and therefore \mathscr{K}_1 is not fixed. \square

Remark 7.6. (i) The Banach space $E = X_{\text{AH}} \oplus \ell_\infty$ is clearly non-separable, so the question naturally arises whether a separable Banach space E exists such that $\mathscr{B}(E)$ contains a non-fixed, finitely generated maximal left ideal. This has recently been answered affirmatively [29].

(ii) Proposition 5.13 and the discussion preceding it raise the question whether the class of Banach spaces for which Question (II) has a positive answer is closed under finite direct sums. Theorem 1.5 implies that this is not the case because X_{AH} and ℓ_∞ both belong to this class by Theorem 6.2 and Example 5.4(ii), respectively, whereas their direct sum does not.

We shall next give a characterization of the ideal \mathscr{K}_1 defined by (1.2). Theorem 1.6 will be an easy consequence of this result.

Theorem 7.7. *Let $E = X_{\text{AH}} \oplus \ell_\infty$. Then the following three conditions are equivalent for each subset \mathscr{L} of $\mathscr{B}(E)$:*

- (a) $\mathscr{L} = \mathscr{K}_1$;
- (b) \mathscr{L} is a non-fixed, finitely-generated, maximal left ideal of $\mathscr{B}(E)$;
- (c) \mathscr{L} is a maximal left ideal of $\mathscr{B}(E)$ and contains an operator which is bounded below.

Proof. (a) \Rightarrow (b). This is immediate from Theorem 1.5.

(b) \Rightarrow (c). Suppose that \mathscr{L} is a non-fixed, finitely-generated, maximal left ideal of $\mathscr{B}(E)$, so that $\mathscr{L} = \mathscr{L}_\Gamma$ for some non-empty, finite subset Γ of $\mathscr{B}(E)$. Set $n = |\Gamma| \in \mathbb{N}$. By Corollary 4.7, the operator Ψ_Γ is bounded below. Moreover, there is an operator $T \in \mathscr{B}(E^n, E)$ which is bounded below because X_{AH} embeds in ℓ_∞ , and ℓ_∞ is isomorphic to the direct sum of $2n - 1$ copies of itself. Hence the composite operator $T\Psi_\Gamma$ is bounded below, and it belongs to \mathscr{L} by (2.6).

(c) \Rightarrow (a). Suppose that \mathscr{L} is a maximal left ideal of $\mathscr{B}(E)$ and that \mathscr{L} contains an operator $R = (R_{j,k})_{j,k=1}^2$ which is bounded below. Then R

does not belong to any fixed maximal left ideal, so that $\mathcal{E}(E) \subseteq \mathcal{L}$ by Corollary 4.1. Lemma 7.1 shows that each operator from ℓ_∞ to X_{AH} is strictly singular, and thus inessential. Hence, by [17, Proposition 1], each operator from X_{AH} to ℓ_∞ is also inessential, and so we conclude that

$$(7.5) \quad \left\{ \begin{aligned} & \begin{pmatrix} T_{1,1} & T_{1,2} \\ T_{2,1} & T_{2,2} \end{pmatrix} : T_{1,1} \in \mathcal{K}(X_{\text{AH}}), T_{1,2} \in \mathcal{B}(\ell_\infty, X_{\text{AH}}), \\ & T_{2,1} \in \mathcal{B}(X_{\text{AH}}, \ell_\infty), T_{2,2} \in \mathcal{W}(\ell_\infty) \end{aligned} \right\} = \mathcal{E}(E) \subseteq \mathcal{L}.$$

Since the operator R is bounded below, its restriction $R|_{\ell_\infty} = \begin{pmatrix} R_{1,2} \\ R_{2,2} \end{pmatrix}$ is also bounded below, and is thus an upper semi-Fredholm operator. Consequently $\begin{pmatrix} 0 \\ R_{2,2} \end{pmatrix}$ is an upper semi-Fredholm operator by [35, Proposition 2.c.10] because $R_{1,2}$ is strictly singular, and therefore $R_{2,2}$ is an upper semi-Fredholm operator. Let $Q \in \mathcal{F}(\ell_\infty)$ be a projection onto $\ker R_{2,2}$. Then the restriction of $R_{2,2}$ to $\ker Q$ is an isomorphism onto its range, which is a closed subspace of ℓ_∞ . Since ℓ_∞ is injective, the inverse of this isomorphism extends to an operator $S: \ell_\infty \rightarrow \ker Q \subseteq \ell_\infty$, which then satisfies $SR_{2,2} = I_{\ell_\infty} - Q$. Hence

$$\begin{pmatrix} 0 & 0 \\ SR_{2,1} & I_{\ell_\infty} - Q \end{pmatrix} = \begin{pmatrix} 0 & 0 \\ 0 & S \end{pmatrix} \begin{pmatrix} R_{1,1} & R_{1,2} \\ R_{2,1} & R_{2,2} \end{pmatrix} \in \mathcal{L},$$

which by (7.5) implies that

$$\begin{pmatrix} 0 & 0 \\ 0 & I_{\ell_\infty} \end{pmatrix} \in \mathcal{L}.$$

Applying (7.5) once more, we see that $\mathcal{K}_1 \subseteq \mathcal{L}$, and so $\mathcal{K}_1 = \mathcal{L}$ by the maximality of \mathcal{K}_1 . \square

Proof of Theorem 1.6. The equivalence of conditions (a) and (b) in Theorem 7.7 shows that \mathcal{K}_1 is the unique non-fixed, finitely-generated, maximal left ideal of $\mathcal{B}(E)$. Proposition 7.2(ii) implies that \mathcal{W}_2 is a maximal two-sided ideal. Since $\mathcal{F}(E) \subseteq \mathcal{W}_2 \not\subseteq \mathcal{K}_1$, \mathcal{W}_2 is not contained in any finitely-generated, maximal left ideal of $\mathcal{B}(E)$. \square

One may wonder whether the conclusion of Theorem 1.5 that the ideal \mathcal{W}_1 introduced in Proposition 7.2(i) is maximal as a left ideal might be true more generally, that is, not only in the case where X is Argyros–Haydon’s Banach space. Our next result implies that this is false for $X = c_0$. Note that all weakly compact operators on c_0 are compact, so that, in this case, \mathcal{W}_1 is equal to

$$(7.6) \quad \mathcal{K}_1 = \left\{ \begin{pmatrix} T_{1,1} & T_{1,2} \\ T_{2,1} & T_{2,2} \end{pmatrix} \in \mathcal{B}(c_0 \oplus \ell_\infty) : T_{1,1} \in \mathcal{K}(c_0) \right\}.$$

Proposition 7.8. *The ideal \mathcal{K}_1 given by (7.6) is not contained in any finitely-generated, maximal left ideal of $\mathcal{B}(c_0 \oplus \ell_\infty)$.*

For clarity, we present the main technical step in the proof of Proposition 7.8 as a separate lemma.

Lemma 7.9. *Suppose that $T \in \mathcal{B}(c_0)$ is not an upper semi-Fredholm operator. Then there exist a projection $Q_0 \in \mathcal{B}(c_0)$ and a normalized basic sequence $(x_n)_{n \in \mathbb{N}}$ in c_0 such that $(x_n)_{n \in \mathbb{N}}$ is equivalent to the standard unit vector basis for c_0 and*

$$(7.7) \quad Q_0 x_{2n-1} = x_{2n-1}, \quad Q_0 x_{2n} = 0, \quad \text{and} \quad \|Tx_n\| \leq \frac{1}{n} \quad (n \in \mathbb{N}).$$

Proof. Let $(e_n)_{n \in \mathbb{N}}$ denote the standard unit vector basis for c_0 . Since T is not an upper semi-Fredholm operator, there are two cases to consider.

Case 1: $\dim \ker T = \infty$. Then $\ker T$ contains a closed subspace Y which is isomorphic to c_0 and complemented in c_0 (e.g., see [35, Proposition 2.a.2]). Let $(x_n)_{n \in \mathbb{N}}$ be a normalized Schauder basis for Y such that $(x_n)_{n \in \mathbb{N}}$ is equivalent to $(e_n)_{n \in \mathbb{N}}$. Since Y is complemented in c_0 and the basis $(x_n)_{n \in \mathbb{N}}$ is unconditional, there is a projection $Q_0 \in \mathcal{B}(c_0)$ which satisfies the first two identities in (7.7), while the third one is trivial because $x_n \in Y \subseteq \ker T$ for each $n \in \mathbb{N}$.

Case 2: $\dim \ker T < \infty$ and $T(c_0)$ is not closed. For each $n \in \mathbb{N}$, choose $\varepsilon_n \in (0, 1)$ such that $(1 + \|T\|)\varepsilon_n(1 - \varepsilon_n)^{-1} \leq n^{-1}$. By induction, we shall construct a normalized block basic sequence $(x_n)_{n \in \mathbb{N}}$ of $(e_n)_{n \in \mathbb{N}}$ such that $\|Tx_n\| \leq n^{-1}$ for each $n \in \mathbb{N}$.

To start the induction, we observe that T cannot be bounded below because its range is not closed, so that we can find a unit vector $y_1 \in c_0$ such that $\|Ty_1\| \leq \varepsilon_1$. Approximating y_1 within ε_1 by a finitely-supported vector and normalizing it, we obtain a finitely-supported unit vector $x_1 \in c_0$ such that $\|Tx_1\| \leq (1 + \|T\|)\varepsilon_1(1 - \varepsilon_1)^{-1} \leq 1$ by the choice of ε_1 .

Now assume inductively that unit vectors $x_1, \dots, x_n \in c_0$ with consecutive supports have been chosen for some $n \in \mathbb{N}$ such that $\|Tx_j\| \leq 1/j$ for each $j \in \{1, \dots, n\}$. Let $m \in \mathbb{N}$ be the maximum of the support of x_n , so that $x_1, \dots, x_n \in \text{span}\{e_1, \dots, e_m\}$, and let P_m be the m^{th} basis projection associated with $(e_j)_{j \in \mathbb{N}}$. If $T|_{\ker P_m}$ were bounded below, then it would have closed range, so that

$$T(c_0) = T(\ker P_m) + \text{span}\{Te_1, \dots, Te_m\}$$

would also be closed, being the sum of a closed subspace and a finite-dimensional one. This is false, and hence $T|_{\ker P_m}$ is not bounded below. We can therefore choose a unit vector $y_{n+1} \in \ker P_m$ such that $\|Ty_{n+1}\| \leq \varepsilon_{n+1}$. Now, as in the first step of the induction, we approximate y_{n+1} within ε_{n+1} by a finitely-supported vector in $\ker P_m$ and normalize it to obtain a finitely-supported unit vector $x_{n+1} \in \ker P_m$ such that

$$\|Tx_{n+1}\| \leq \frac{(1 + \|T\|)\varepsilon_{n+1}}{1 - \varepsilon_{n+1}} \leq \frac{1}{n+1}$$

by the choice of ε_{n+1} . Therefore the induction continues.

By [35, Proposition 2.a.1], the sequence $(x_n)_{n \in \mathbb{N}}$ is equivalent to $(e_n)_{n \in \mathbb{N}}$, and its closed linear span is complemented in c_0 . Hence, as in Case 1, we obtain a projection $Q_0 \in \mathcal{B}(c_0)$ such that the first two identities in (7.7) are satisfied, while the third one holds by the construction of $(x_n)_{n \in \mathbb{N}}$. \square

Proof of Proposition 7.8. Assume towards a contradiction that \mathcal{L} is a finitely-generated, maximal left ideal of $\mathcal{B}(c_0 \oplus \ell_\infty)$ such that $\mathcal{K}_1 \subseteq \mathcal{L}$. Proposition 2.2 implies that \mathcal{L} is generated by a single operator, say

$$T = \begin{pmatrix} T_{1,1} & T_{1,2} \\ T_{2,1} & T_{2,2} \end{pmatrix} \in \mathcal{B}(c_0 \oplus \ell_\infty).$$

We *claim* that $T_{1,1}$ is not an upper semi-Fredholm operator. Assume the contrary; that is, $\ker T_{1,1}$ is finite-dimensional, so that we can take a projection $P \in \mathcal{F}(c_0)$ onto $\ker T_{1,1}$, and $T_{1,1}(c_0)$ is closed. Then the restriction $\tilde{T}_{1,1}: x \mapsto T_{1,1}x$, $\ker P \rightarrow T_{1,1}(c_0)$, is an isomorphism. Its range is complemented in c_0 by Sobczyk's theorem [49] because it is isomorphic to $\ker P$, which is a closed subspace of finite codimension in c_0 , and hence isomorphic to c_0 . We can therefore extend the inverse of $\tilde{T}_{1,1}$ to an operator $S \in \mathcal{B}(c_0)$, which then satisfies $ST_{1,1} = I_{c_0} - P$. Since P has finite rank, we have

$$\begin{pmatrix} P & 0 \\ 0 & I_{\ell_\infty} \end{pmatrix} \in \mathcal{K}_1 \subseteq \mathcal{L}$$

and

$$\begin{pmatrix} T_{1,1} & 0 \\ 0 & 0 \end{pmatrix} = T - \begin{pmatrix} 0 & T_{1,2} \\ T_{2,1} & T_{2,2} \end{pmatrix} \in \mathcal{L} - \mathcal{K}_1 \subseteq \mathcal{L},$$

which implies that

$$\begin{pmatrix} I_{c_0} & 0 \\ 0 & I_{\ell_\infty} \end{pmatrix} = \begin{pmatrix} P & 0 \\ 0 & I_{\ell_\infty} \end{pmatrix} + \begin{pmatrix} S & 0 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} T_{1,1} & 0 \\ 0 & 0 \end{pmatrix} \in \mathcal{L}.$$

This, however, contradicts the assumption that the left ideal \mathcal{L} is proper, and thus completes the proof of our claim.

Hence, by Lemma 7.9, we obtain a projection $Q_0 \in \mathcal{B}(c_0)$ and a normalized basic sequence $(x_n)_{n \in \mathbb{N}}$ in c_0 such that $(x_n)_{n \in \mathbb{N}}$ is equivalent to the standard unit vector basis $(e_n)_{n \in \mathbb{N}}$ for c_0 and

$$(7.8) \quad Q_0 x_{2n-1} = x_{2n-1}, \quad Q_0 x_{2n} = 0, \quad \text{and} \quad \|T_{1,1} x_n\| \leq \frac{1}{n} \quad (n \in \mathbb{N}).$$

The sequence $(x_n)_{n \in \mathbb{N}}$ is weakly null because it is equivalent to the weakly null sequence $(e_n)_{n \in \mathbb{N}}$, and so the sequence $(Rx_n)_{n \in \mathbb{N}}$ is norm-null for each $R \in \mathcal{K}(c_0)$. Now let

$$Q = \begin{pmatrix} Q_0 & 0 \\ 0 & 0 \end{pmatrix} \in \mathcal{B}(c_0 \oplus \ell_\infty).$$

The maximality of the left ideal \mathcal{L} implies that either

$$(i) \quad Q \in \mathcal{L} \quad \text{or} \quad (ii) \quad \mathcal{L} + \mathcal{L}_{\{Q\}} = \mathcal{B}(c_0 \oplus \ell_\infty).$$

We shall complete the proof by showing that both of these alternatives are impossible.

In case (i), there is $S = (S_{j,k})_{j,k=1}^2 \in \mathcal{B}(c_0 \oplus \ell_\infty)$ with $Q = ST$. Defining $P_0 \in \mathcal{B}(c_0 \oplus \ell_\infty, c_0)$ by $P_0(x, f) = x$ for each $x \in c_0$ and $f \in \ell_\infty$, we have

$$x_{2n-1} = P_0 Q(x_{2n-1}, 0) = P_0 ST(x_{2n-1}, 0) = S_{1,1} T_{1,1} x_{2n-1} + S_{1,2} T_{2,1} x_{2n-1}$$

for each $n \in \mathbb{N}$. This, however, is absurd since the left-hand side is a unit vector, whereas the right-hand side norm-converges to 0 as $n \rightarrow \infty$: this holds because $\|T_{1,1}x_{2n-1}\| \rightarrow 0$ by (7.8) and $S_{1,2}T_{2,1} \in \mathcal{W}(c_0) = \mathcal{K}(c_0)$.

In case (ii), there are operators $U, V \in \mathcal{B}(c_0 \oplus \ell_\infty)$ such that

$$I_{c_0 \oplus \ell_\infty} = UT + VQ.$$

Define P_0 as above, and write $U = (U_{j,k})_{j,k=1}^2$. Then, since $Q_0x_{2n} = 0$, we have

$$x_{2n} = P_0(UT + VQ)(x_{2n}, 0) = U_{1,1}T_{1,1}x_{2n} + U_{1,2}T_{2,1}x_{2n} \quad (n \in \mathbb{N}),$$

which leads to a contradiction as in case (i) because the left-hand side is a unit vector, whereas the right-hand side norm-converges to 0 as $n \rightarrow \infty$. \square

ACKNOWLEDGEMENTS

We are grateful to David Blecher (Houston) and Mikael Rørdam (Copenhagen) for having informed us that the conjecture of the first author and Żelazko stated on p. 4 has a positive answer for C^* -algebras, and for having shown us how to deduce it from standard facts. We are also grateful to the referee for his/her careful reading of our paper which has led to simplified proofs of several results in Section 4.

Key parts of this work were carried out during a visit of Piotr Koszmider to Lancaster in February 2012. The visit was supported by a London Mathematical Society Scheme 2 grant (ref. 21101). The authors gratefully acknowledge this support.

REFERENCES

- [1] F. Albiac and N. J. Kalton, *Topics in Banach space theory*, Grad. Texts in Math. 233, Springer-Verlag, New York, 2006.
- [2] D. Amir, *Continuous function spaces with the bounded extension property*, Bull. Res. Council Israel Sect. F 101 (1962), 133–138.
- [3] S. A. Argyros, J. F. Castillo, A. S. Granero, M. Jiménez, and J. P. Moreno, *Complementation and embeddings of $c_0(I)$ in Banach spaces*, Proc. London Math. Soc. 85 (2002), 742–768.
- [4] S. A. Argyros and R. G. Haydon, *A hereditarily indecomposable \mathcal{L}_∞ -space that solves the scalar-plus-compact problem*, Acta Math. 206 (2011), 1–54.
- [5] J. W. Baker, *Some uncomplemented subspaces of $C(X)$ of the type $C(Y)$* , Studia Math. 36 (1970), 85–103.
- [6] D. Blecher and T. Kania, *Finite generation in C^* -algebras and Hilbert C^* -modules*, in preparation.
- [7] N. Boudi, *Banach algebras in which every left ideal is countably generated*, Irish Math. Soc. Bull. 48 (2002), 17–24.
- [8] S. R. Caradus, W. E. Pfaffenberger and B. Yood, *Calkin algebras and algebras of operators on Banach spaces*, Lect. Notes Pure Appl. Math. 9, Marcel Dekker, New York, 1974.
- [9] W. W. Comfort and S. Negreponis, *The theory of ultrafilters*, Grundlehren Math. Wiss. 211, Springer-Verlag, New York–Heidelberg, 1974.
- [10] H. B. Cohen, *Injective envelopes of Banach spaces*, Bull. Amer. Math. Soc. 70 (1964), 723–726.
- [11] H. G. Dales, *Banach algebras and automatic continuity*, London Math. Soc. Monogr. Ser. 24, Clarendon Press, Oxford, 2000.

- [12] H. G. Dales and W. Żelazko, *Generators of maximal left ideals in Banach algebras*, *Studia Math.* 212 (2012), 173–193.
- [13] W. J. Davis, T. Figiel, W. B. Johnson, and A. Pełczyński, *Factoring weakly compact operators*, *J. Funct. Anal.* 17 (1974), 311–327.
- [14] I. S. Edelstein and B. S. Mityagin, *Homotopy type of linear groups of two classes of Banach spaces*, *Funct. Anal. Appl.* 4 (1970), 221–231.
- [15] V. Ferreira and G. Tomassini, *Finiteness properties of topological algebras*, *Ann. Sc. Norm. Sup. Pisa Cl. Sci.* 5 (1978), 471–488.
- [16] L. Gillman, *Countably generated ideals in rings of continuous functions*, *Proc. Amer. Math. Soc.* 11 (1960), 660–666.
- [17] M. González, *On essentially incomparable Banach spaces*, *Math. Z.* 215 (1994), 621–629.
- [18] D. B. Goodner, *Projections in normed linear spaces*, *Trans. Amer. Math. Soc.* 69 (1950), 89–108.
- [19] W. T. Gowers and B. Maurey, *The unconditional basic sequence problem*, *J. Amer. Math. Soc.* 6 (1993), 851–874.
- [20] B. Gramsch, *Eine Idealstruktur Banachscher Operatoralgebren*, *J. Reine Angew. Math.* 225 (1967), 97–115.
- [21] A. S. Granero, *On complemented subspaces of $c_0(I)$* , *Atti Semin. Mat. Fis. Univ. Módena Reggio Emilia XLVI* (1998), 35–36.
- [22] A. Grothendieck, *Une caractérisation vectorielle métrique des espaces L_1* , *Canad. J. Math.* 7 (1955), 552–561.
- [23] N. Grønbæk, *Morita equivalence for Banach algebras*, *J. Pure Appl. Algebra* 99 (1995), 183–219.
- [24] N. Jacobson, *Basic Algebra II*, W. H. Freeman, San Francisco, California, 1980.
- [25] T. Jech, *Set Theory*, Third Millennium Ed., revised and expanded, Springer-Verlag, Berlin, 2003.
- [26] R. V. Kadison and J. R. Ringrose, *Fundamentals of the theory of operator algebras I: Elementary theory*, *Pure Appl. Math.* 100, Academic Press, New York, 1983.
- [27] R. V. Kadison and J. R. Ringrose, *Fundamentals of the theory of operator algebras II: Advanced theory*, *Pure Appl. Math.* 100, Academic Press, Orlando, Florida, 1986.
- [28] N. J. Kalton and N. T. Peck, *Twisted sums of sequence spaces and the three space problem*, *Trans. Amer. Math. Soc.* 255 (1979), 1–30.
- [29] T. Kania and N. J. Laustsen, *Ideal structure of the algebra of bounded operators acting on a Banach space*, in preparation.
- [30] D. Kleinecke, *Almost-finite, compact, and inessential operators*, *Proc. Amer. Math. Soc.* 14 (1963), 863–868.
- [31] P. Koszmider, *Banach spaces of continuous functions with few operators*, *Math. Ann.* 330 (2004), 151–183.
- [32] N. J. Laustsen, *Maximal ideals in the algebra of operators on certain Banach spaces*, *Proc. Edinb. Math. Soc.* 45 (2002), 523–546.
- [33] N. J. Laustsen, *Commutators of operators on Banach spaces*, *J. Operator Theory* 48 (2002), 503–514.
- [34] J. Lindenstrauss, *On the extension property for compact operators*, *Bull. Amer. Math. Soc.* 68 (1962), 484–487.
- [35] J. Lindenstrauss and L. Tzafriri, *Classical Banach spaces I*, *Ergeb. Math. Grenzgeb.* 92, Springer-Verlag, Berlin–New York, 1977.
- [36] J. Lindenstrauss and L. Tzafriri, *Classical Banach spaces II*, *Ergeb. Math. Grenzgeb.* 97, Springer-Verlag, Berlin–New York, 1979.
- [37] E. Luft, *The two-sided closed ideals of the algebra of bounded linear operators of a Hilbert space*, *Czechoslovak Math. J.* 18 (1968), 595–605.
- [38] R. E. Megginson, *An introduction to Banach space theory*, *Grad. Texts in Math.* 183, Springer-Verlag, New York, 1998.

- [39] G. J. Murphy, *C*-algebras and operator theory*, Academic Press, Boston, MA, 1990.
- [40] L. Nachbin, *On the Hahn–Banach theorem*, An. Acad. Brasil. Ciênc. 21 (1949), 151–154.
- [41] A. Pelczyński, *On strictly singular and strictly cosingular operators. I. Strictly singular and strictly cosingular operators in $C(S)$ -spaces*, Bull. Acad. Polon. Sci. Sér. Sci. Math. Astronom. Phys. 13 (1965), 31–36.
- [42] A. Pietsch, *Operator ideals*, North Holland, Amsterdam–New York–Oxford, 1980.
- [43] G. Plebanek, *A construction of a Banach space $C(K)$ with few operators*, Topology Appl. 143 (2004), 217–239.
- [44] B. Pospíšil, *On bicomact spaces*, Publ. Fac. Sci. Univ. Masaryk 270 (1939), 3–16.
- [45] R. T. Prosser, *On the ideal structure of operator algebras*, Mem. Amer. Math. Soc. 45 (1963).
- [46] A. Rosenberg, *The number of irreducible representations of simple rings with no minimal ideals*, Amer. J. Math. 75 (1953), 523–530.
- [47] Th. Schlumprecht, *An arbitrarily distortable Banach space*, Israel J. Math. 76 (1991), 81–95.
- [48] A. M. Sinclair and A. W. Tullio, *Noetherian Banach algebras are finite dimensional*, Math. Ann. 211 (1974), 151–153.
- [49] A. Sobczyk, *Projection of the space (m) on its subspace (c_0)* , Bull. Amer. Math. Soc. 47 (1941), 938–947.
- [50] E. L. Stout, *The theory of uniform algebras*, Bogden and Quigley, Tarrytown-on-Hudson, New York, 1971.
- [51] A. Szankowski, *$\mathcal{B}(H)$ does not have the approximation property*, Acta Math. 147 (1981), 89–108.
- [52] B. Yood, *Difference algebras of linear transformations on a Banach space*, Pacific J. Math. 4 (1954), 615–636.

DEPARTMENT OF MATHEMATICS AND STATISTICS, FYLDE COLLEGE, LANCASTER UNIVERSITY, LANCASTER LA1 4YF, UK

E-mail address: g.dales@lancaster.ac.uk

DEPARTMENT OF MATHEMATICS AND STATISTICS, FYLDE COLLEGE, LANCASTER UNIVERSITY, LANCASTER LA1 4YF, UK

E-mail address: t.kania@lancaster.ac.uk

INSTITUTE OF MATHEMATICS, UNIVERSITY OF SILESIA, UL. BANKOWA 14, 40-007 KATOWICE, POLAND

E-mail address: t_kochanek@wp.pl

INSTITUTE OF MATHEMATICS, POLISH ACADEMY OF SCIENCES, UL. ŚNIADECKICH 8, 00-956 WARSZAWA, POLAND

E-mail address: P.Koszmider@Impan.pl

DEPARTMENT OF MATHEMATICS AND STATISTICS, FYLDE COLLEGE, LANCASTER UNIVERSITY, LANCASTER LA1 4YF, UK

E-mail address: n.laustsen@lancaster.ac.uk