



## Fragment screening reveals salicylic hydroxamic acid as an inhibitor of *Trypanosoma brucei* GPI GlcNAc-PI de-*N*-acetylase



Michael D. Urbaniak<sup>†,‡</sup>, Amy S. Capes<sup>‡</sup>, Arthur Crossman, Sandra O'Neill, Stephen Thompson, Ian H. Gilbert<sup>\*</sup>, Michael A. J. Ferguson<sup>\*</sup>

Division of Biological Chemistry and Drug Discovery, College of Life Sciences, University of Dundee, Dow Street, Dundee DD1 5EH, UK

### ARTICLE INFO

#### Article history:

Received 5 November 2013  
Received in revised form 12 December 2013  
Accepted 18 December 2013  
Available online 30 December 2013

#### Keywords:

GPI  
*Trypanosoma brucei*  
Hydroxamic acid  
Inhibitor  
*N*-Deacetylase

### ABSTRACT

The zinc-metalloenzyme GlcNAc-PI de-*N*-acetylase is essential for the biosynthesis of mature GPI anchors and has been genetically validated in the bloodstream form of *Trypanosoma brucei*, which causes African sleeping sickness. We screened a focused library of zinc-binding fragments and identified salicylic hydroxamic acid as a GlcNAc-PI de-*N*-acetylase inhibitor with high ligand efficiency. This is the first small molecule inhibitor reported for the trypanosome GPI pathway. Investigating the structure activity relationship revealed that hydroxamic acid and 2-OH are essential for potency, and that substitution is tolerated at the 4- and 5-positions.

© 2013 Elsevier Ltd. All rights reserved.

## 1. Introduction

*Trypanosoma brucei* is a protozoan parasite that is transmitted by the bite of an infected tsetse fly and causes the fatal African sleeping sickness in humans and the related wasting disease Nagana in cattle. Current treatments are expensive, toxic, and difficult to administer, leaving an urgent need for improved therapeutic agents. The parasite is able to evade the host immune response by virtue of a dense surface coat composed of  $5 \times 10^6$  variant surface glycoprotein (VSG) dimers that prevent lysis by the innate immune response. The coat also undergoes antigenic variation to evade the adaptive immune response.<sup>1</sup> The VSG dimer is attached to the plasma membrane by a glycosylphosphatidylinositol (GPI) anchor, and GPI anchor biosynthesis has been genetically and chemically validated as essential for the survival of the clinically relevant bloodstream form of the parasite.<sup>2–5</sup>

The GPI biosynthetic pathway has been extensively studied in both *T. brucei* and mammalian systems, highlighting differences in both the order of assembly and substrate specificity that can be exploited to produce species-specific inhibitors. The GlcNAc-PI de-*N*-acetylase is a zinc metalloenzyme that catalyses the second

step in biosynthesis of GPI (Fig. 1),<sup>6</sup> and has been genetically validated as essential in bloodstream form *T. brucei*.<sup>3</sup> The use of synthetic substrate analogues has revealed that the human enzyme is more fastidious than the trypanosome enzyme, enabling the design of species-specific substrate-based inhibitors.<sup>7–9</sup> The substrate analogue approach has shown that the phosphate, 2'-NHAc and 3'-OH of the substrate GlcNAc-PI are critical for recognition by the *T. brucei* GlcNAc-PI de-*N*-acetylase.<sup>10</sup> In contrast, the diacylglycerol is not directly recognised and may be efficiently replaced with an octadecyl chain.<sup>9,11</sup> Recently, we have shown that substrate-based inhibitors containing the zinc binding moieties hydroxamic acid or carboxylic acid can act as inhibitors of the *T. brucei* de-*N*-acetylase.<sup>12,13</sup> These substrate-based inhibitors do not possess drug-like physicochemical properties and contain too much stereochemical complexity for tractable synthesis. Here, we report our efforts to identify alternative scaffolds for GlcNAc-PI de-*N*-acetylase inhibitors that resulted in the identification of salicylic hydroxamic acid as an inhibitor with high ligand efficiency.

## 2. Materials and methods

### 2.1. Materials


The synthesis of D-glucosamine- $\alpha$ -(1-6)-D-myo-inositol-1-octadecyl phosphate (GlcN-IPC<sub>18</sub>) has been described previously.<sup>14</sup> The corresponding *N*-acetyl derivative (GlcNAc-IPC<sub>18</sub>) was prepared by treatment with acetic anhydride, and the concentration of stock

\* Corresponding authors. Tel.: +44 (0) 1382 384219.

E-mail addresses: [i.h.gilbert@dundee.ac.uk](mailto:i.h.gilbert@dundee.ac.uk) (I.H. Gilbert), [m.a.j.ferguson@dundee.ac.uk](mailto:m.a.j.ferguson@dundee.ac.uk) (M.A.J. Ferguson).

<sup>†</sup> These authors contributed equally to the work.

<sup>‡</sup> Present address: Division of Biomedical and Life Sciences, Faculty of Health and Medicine, Lancaster University, Lancaster LA1 4GQ, UK.



**Figure 1.** GlcNAc-PI de-N-acetylase catalyses the second step in the GPI biosynthetic pathway. The zinc-dependent metalloenzyme is inhibited by zinc chelators.

solutions determined by measurement of the inositol content by selected ion-monitoring GC–MS.<sup>13</sup> Bloodstream form *Trypanosome brucei* (variant MITat1.4) was isolated and membranes (cell-free system) prepared as described previously and stored at  $-80^{\circ}\text{C}$ .<sup>15</sup> Recombinant GST-tagged *T. brucei* de-N-acetylase (GST-TbGPI12) was expressed and purified as described previously and stored at  $-80^{\circ}\text{C}$ .<sup>13</sup>

## 2.2. Mass spectrometry based activity assays

Inhibition assays were performed in 100  $\mu\text{L}$  final volume, with 1% v/v DMSO with or without inhibitor. Recombinant GST-TbGPI12 (10  $\mu\text{g}$  per assay) or trypanosome cell-free system ( $2.5 \times 10^6$  cell equivalents per assay) in incorporation buffer (25 mM Tris pH 8.0, 50 mM KCl, 5 mM  $\text{MnCl}_2$ ) was added to wells containing 500 pmol GlcNAc-IPC<sub>18</sub> with or without the inhibitor and incubated at  $37^{\circ}\text{C}$  for 1 h. The reactions were quenched by the addition of 200  $\mu\text{L}$  of 5% propan-1-ol, 5 mM  $\text{NH}_4\text{OAc}$ , and the glycolipids were bound to C<sub>18</sub> resin (50 mg Isolute cartridge), washed three times with 200  $\mu\text{L}$  5% propan-1-ol, 5 mM  $\text{NH}_4\text{OAc}$  and eluted with 100  $\mu\text{L}$  40% propan-1-ol, 5 mM  $\text{NH}_4\text{OAc}$ . Enriched glycolipids were analysed by liquid chromatography coupled to an electrospray tandem mass spectrometer (LC–MS/MS). Samples (40  $\mu\text{L}$ ) were injected on to a  $10 \times 1$  mm C<sub>18</sub> column (ACE, 5  $\mu\text{M}$ ) and eluted using a binary gradient of 5–80% propan-1-ol in 5 mM  $\text{NH}_4\text{OAc}$  (Dionex Ultimate 3000). The gradient consisted of 2 min 0% B, 2–4 min 0–100% B, 4–8 min 100%, 8–9 min 100–0% B, 9–10 min 0% B where buffer A consisted of 5% propan-1-ol, 5 mM  $\text{NH}_4\text{OAc}$  and buffer B 80% propan-1-ol, 5 mM  $\text{NH}_4\text{OAc}$ . The glycolipids were analysed on an electrospray triple quadrupole mass spectrometer (Micromass Quattro Ultima) in multiple reaction monitoring mode. The turnover of the substrate GlcNAc-IPC<sub>18</sub> ( $m/z$  715 > 223) to GlcN-IPC<sub>18</sub> ( $m/z$  672 > 223) was used to calculate the percentage of substrate conversion to product in a given sample.<sup>6</sup> Inhibitor IC<sub>50</sub> values were calculated using a four-parameter fit of eight-point potency curves derived from three independent experiments, and are quoted with standard deviation.

## 2.3. Trypanosome cell-free system assay

Trypanosome cell-free system assays, where the formation of GPI precursors is monitored by following the incorporation of [<sup>3</sup>H]-mannose were analysed using high-performance liquid chromatography and fluorography as described previously.<sup>13</sup>

## 2.4. Compound synthesis

### 2.4.1. Synthesis of 5-((tert-butoxycarbonyl)amino)-2-hydroxybenzoic acid (9)

5-Aminosalicylic acid **8** (6.89 g, 45.0 mmol) was suspended in water (60 mL), and TEA (12.4 mL, 90.0 mmol) was added. A solution of Boc<sub>2</sub>O (10.8 g, 90.0 mmol) in dioxane (120 mL) was added, and the reaction was stirred at room temperature overnight. The solvent was removed, and the solid residue suspended in water (20 mL). HCl (3 M) was added until the greyish pink precipitate stopped forming. The precipitate was filtered off and washed with water. The precipitate was dissolved in boiling acetone and hot filtered before being recrystallized twice from acetone to afford the product (8.61 g, 76%) as a white powder, mp  $278^{\circ}\text{C}$ . <sup>1</sup>H NMR (500 MHz, MeOD)  $\delta$  7.93 (1H, s), 7.40 (1H, d,  $J = 7.6$  Hz), 6.85 (1H, d,  $J = 8.9$  Hz), 1.51 (9H, s); <sup>13</sup>C NMR (125 MHz, MeOD)  $\delta$  173.3, 159.0, 155.7, 132.1, 128.7, 121.8, 118.2, 113.5, 80.8, 28.7; HRMS, calcd mass for C<sub>12</sub>H<sub>16</sub>NO<sub>5</sub><sup>+</sup> [ $M+H^+$ ]: 254.1023. Found: 254.1027 (–1.4 ppm).

### 2.4.2. Synthesis of tert-butyl (3-((benzyloxy)carbamoyl)-4-hydroxyphenyl)carbamate (10)


O-Benzylhydroxylamine hydrochloride (3.54 g, 22.2 mmol) was suspended in  $\text{CHCl}_3$  (85 mL) and cooled to  $0^{\circ}\text{C}$ , and then TEA (3.21 mL, 23.1 mmol) was added. A solution of compound **9** (5.62 g, 22.2 mmol) in THF (140 mL), DMAP (169 mg, 1.12 mmol) and DCC (5.06 g, 24.4 mmol) was added, and the reaction was stirred for 16 h. The solvent was removed and the solids were resuspended in EtOAc, filtered, and then washed with HCl (1 M, 50 mL), water (50 mL), ammonia (1 M,  $2 \times 50$  mL) and filtered through cotton wool. The solvent was removed and the crude was dried on to silica then purified by column chromatography (EtOAc/hexane, 0:100 to 1:1) to afford the product (3.63 g, 46%) as a pale yellow powder, mp  $102$ – $103^{\circ}\text{C}$ . <sup>1</sup>H NMR (500 MHz,  $\text{CDCl}_3$ )  $\delta$  11.33 (1H, br s), 9.42 (1H, br s), 7.63 (1H, br s), 7.44–7.41 (2H, m), 7.39–7.34 (3H, m), 7.13 (1H, dd,  $J = 8.9$ , 2.4 Hz), 6.90 (1H, d,  $J = 8.9$  Hz), 6.45 (1H, br s), 4.97 (2H, s), 1.49 (9H, s); <sup>13</sup>C NMR (125 MHz,  $\text{CDCl}_3$ )  $\delta$  157.1, 153.3, 134.9, 129.6, 129.3, 128.9, 128.7, 126.1, 118.8, 116.0, 112.1, 80.8, 78.6, 28.3; LRMS (ES<sup>+</sup>),  $m/z$  359.2 [ $M+H^+$ ]; HRMS, calcd mass for C<sub>19</sub>H<sub>23</sub>N<sub>2</sub>O<sub>5</sub><sup>+</sup> [ $M+H^+$ ]: 359.1601. Found: 359.1591 (2.8 ppm).

### 2.4.3. Synthesis of 5-amino-N-(benzyloxy)-2-hydroxybenzamide TFA salt (11)

Compound **10** (1.99 g, 5.56 mmol) was dissolved in wet TFA (5.3 mL), and the mixture was stirred for 1 h. The TFA was removed in vacuo; the residue was suspended in Et<sub>2</sub>O and filtered to afford the product (1.76 g, 85%) as a tan powder, mp  $169$ – $170^{\circ}\text{C}$ . <sup>1</sup>H NMR (500 MHz, DMSO-*d*<sub>6</sub>)  $\delta$  11.48 (1H, br s), 9.96 (2H, br s), 7.64 (1H, d,  $J = 2.6$  Hz), 7.47 (2H, d,  $J = 7$  Hz), 7.42–7.36 (3H, m), 7.31 (1H, dd,  $J = 8.7$ , 2.8 Hz), 7.03 (1H, d,  $J = 8.8$  Hz), 4.96 (2H, s); <sup>13</sup>C NMR (125 MHz, DMSO-*d*<sub>6</sub>)  $\delta$  163.4, 155.6, 135.7, 128.8, 128.29, 128.27, 126.9, 124.5, 122.7, 117.87, 115.5, 113.2, 77.0; LRMS (ES<sup>+</sup>),  $m/z$  259.1 [ $M+H^+$ ]; HRMS, calcd mass for C<sub>14</sub>H<sub>15</sub>N<sub>2</sub>O<sub>3</sub><sup>+</sup> [ $M+H^+$ ]: 259.1077. Found: 259.1069 (3.3 ppm).

### 2.4.4. General method for the synthesis of the amide series

Compound **11** (200 mg, 0.592 mmol), DMAP (cat.) and the acyl chloride (0.592 mmol) were dissolved in THF (2 mL) and DCE



**Figure 2.** *T. brucei* GlcNAc-PI de-*N*-acetylase inhibitors. The activity of the recombinant enzyme against the synthetic substrate *D*-glucosamine- $\alpha$ -(1-6)-*D*-myo-inositol-1-octadecyl phosphate was measured using a mass spectrometry based assay (A) Fragments were screened at 1 mM in duplicate. (B) The six most potent fragments showed dose–response. (C) For the most potent compound **1** an eight-point potency curve was determined in triplicate giving an  $IC_{50} = 63 \pm 4 \mu M$ . (D) Structures of the six most potent fragments.

(0.5 mL). Pyridine (96  $\mu L$ , 1.184 mmol) was added, and the reaction stirred for 24 h. The mixture was diluted with DCM (10 mL) and water (10 mL) then stirred vigorously. The mixture was passed through a phase separator, and then the solvent was removed. The crude was recrystallized from MeOH/Et<sub>2</sub>O. Characterisation of compounds **12–19** is reported in the [Supplementary data](#).

#### 2.4.5. General method for amide deprotection

A 0.05 M solution of benzyl-protected amide (**12–19**) and AcOH (2 equiv) in MeOH/THF, MeOH or THF, was passed through a H<sub>2</sub> flow reactor (1 mL/min, 30 °C, 1 atm) using a 20% Pd(OH)<sub>2</sub> catalyst cartridge. The solvent was removed and the product was recrystallized from THF/MeOH, or washed with THF or MeOH. Characterisation of compounds **20–27** is reported in the [Supplementary data](#).

### 3. Results and discussion

#### 3.1. Fragment screening

To discover more drug-like scaffolds for zinc-binding de-*N*-acetylase inhibitors we screened a small focused library of zinc binding fragments. Because we have previously shown that the enzyme may be inhibited by substrate analogues containing hydroxamic acid, *N*-hydroxyurea or carboxylic acid groups,<sup>12,13</sup> we selected 26 commercially available fragments, which contained these zinc binding moieties, with an average molecular weight of 220 ([Supplemental Table S1](#)). We screened the inhibition of recombinant *T. brucei* de-*N*-acetylase by the fragments using a mass spectrometry-based activity assay to directly monitor the conversion of the synthetic substrate *N*-acetyl-*D*-glucosamine- $\alpha$ -(1-6)-*D*-myo-inositol-1-octadecyl phosphate (GlcNAc-IPC<sub>18</sub>) to *D*-glucosamine- $\alpha$ -(1-6)-*D*-myo-inositol-1-octadecyl phosphate (GlcNAc-IPC<sub>18</sub>). The replacement of the diacylglycerol moiety of GlcNAc-PI with a simple C<sub>18</sub> alkyl chain in GlcNAc-IPC<sub>18</sub> has no effect on the substrate recognition, and enables the use of a unique mass-transition to follow the reaction.<sup>6</sup> Initial screening of the fragments at 1 mM in duplicate revealed six compounds with >50% inhibition ([Fig. 2A + D](#)). However, only one compound showed >50% inhibition at 100  $\mu M$  ([Fig. 2B](#)). The most potent compound **1** ([Fig. 2C](#)) had an  $IC_{50}$  of  $63 \pm 4 \mu M$  and a molecular weight of 153, giving an impressive ligand efficiency ( $-RT \cdot \ln IC_{50} / N_{\text{non-H atoms}}$ )<sup>16</sup> of  $-0.57 \text{ kcal mol}^{-1}$  per non-hydrogen atom, which is over one third of the theoretical maximum affinity for organic compounds of


$-1.5 \text{ kcal mol}^{-1}$  per non-hydrogen atom.<sup>17</sup> By way of comparison, the previously reported glucocyclitol-phospholipid substrate analogue 1R,2R-1-O-[2-C-(carboxymethyl *N*-hydroxyamide)-2-deoxy- $\beta$ -*D*-glucopyranosyl]-cyclohexanediol 2-(*n*-octadecylphosphate) has an  $IC_{50}$  of  $19 \pm 0.5 \mu M$  and a molecular weight of 666.4,<sup>13</sup> giving a substantially lower ligand efficiency of only  $-0.13 \text{ kcal mol}^{-1}$  per non-hydrogen atom.

The result of the fragment screen compared favourably to our screening of 16,037 lead-like compounds from the Dundee Drug Discovery Unit's compound collection<sup>18</sup> against the *T. brucei* de-*N*-acetylase in a cell-based functional complementation assay,<sup>1</sup> where no inhibitors were identified (data not shown). Interestingly, compound **1** (also known as SHAM) has previously been identified as an inhibitor of trypanosome alternative oxidase (TAO), and is toxic in vivo to the clinically relevant bloodstream form *T. brucei*.<sup>19,20</sup>

#### 3.2. Inhibition of the *T. brucei* cell-free system

The ability of **1** to act as an inhibitor of the *T. brucei* GPI pathway was confirmed using the trypanosome cell-free system ([Fig. 3](#)). Priming the cell-free system with GlcAc-IPC<sub>18</sub> in the presence of GDP-[<sup>3</sup>H]-mannose stimulates the production of radiolabelled mannosylated GPI intermediates that can be separated by high performance thin-layer chromatography and visualised by fluorography. Priming the cell-free system with GlcNAc-IPC<sub>18</sub> (10  $\mu M$ ) was prevented by incubation with >300  $\mu M$  of **1** ([Fig. 3B](#)), a five-fold increase over the  $IC_{50}$  value against the recombinant enzyme. To verify that the decrease in potency was not due to a reduction in potency against the endogenous enzyme compared with the recombinant enzyme, we repeated the mass spectrometry based activity assay with the *T. brucei* cell-free system. We found that it was necessary to reduce the amount of cell-free system by 40-fold in the mass spectrometry-based assay compared with the radiometric assay to achieve measurements in the linear range for the turnover of GlcNAc-IPC<sub>18</sub>. The potency of compound **1** against the cell-free system with an  $IC_{50} = 66 \pm 8 \mu M$  ([Fig. 3B](#)) was comparable to that against the recombinant enzyme.

<sup>1</sup> The assay used the CHO MS2S cell line, which has a defective de-*N*-acetylase and expresses the GPI anchored human CD55, functionally complemented by transfection with *T. brucei* de-*N*-acetylase. Cells were treated with trypsin, plated onto compound, and GPI biosynthesis monitored by  $\alpha$ CD55 ELISA at 48 h.



**Figure 3.** Trypanosome GPI biosynthesis in the cell-free system. (A) The *T. brucei* cell-free system was incubated with GlcNAc-PI (10  $\mu$ M) in the presence of 0, 100, 300, or 1000  $\mu$ M of **1** and 0.5  $\mu$ Ci of GDP-[ $^3$ H]-mannose to stimulate the production of radiolabelled mannosylated GPI intermediates. Glycolipid products were extracted, separated by high-performance thin-layer chromatography, and visualised by fluorography. DPM—dolichol-phosphate-mannose, M1—Man<sub>1</sub>GlcN-IPC<sub>18</sub>, M2—Man<sub>2</sub>GlcN-IPC<sub>18</sub>, M3—Man<sub>3</sub>GlcN-IPC<sub>18</sub>, A'—EtNPM<sub>3</sub>GlcN-IPC<sub>18</sub>. (B) Inhibition of the turnover of GlcNAc-IPC<sub>18</sub> (10  $\mu$ M) by the presence of **1** measured in the *T. brucei* cell-free system in the LC-MS/MS assay.

**Table 1**  
Inhibitory activity of commercially available analogues of **1**

ID	R <sup>1</sup>	R <sup>2</sup>	R <sup>4</sup>	Inhibition at 1 mM <sup>a</sup> (%)
1	NHOH	OH	H	97 ± 2
2	OH	OH	H	25 ± 1
3	NHOH	H	H	-6 ± 14
4	NHOH	Br	H	-4 ± 15
5	NHOH	NH <sub>2</sub>	H	-5 ± 2
6	NHOH	OH	Br	86 ± 2
7	NHOH	H	Br	9.6 ± 15

<sup>a</sup> Inhibition of *T. brucei* de-*N*-acetylase (cell-free system).


### 3.3. Structure–activity relationship

Substructure searching identified commercially available analogues of **1** that were screened for activity against in the mass spectrometry based assay against *T. brucei* cell-free system. Replacement of the hydroxamic acid with carboxylic acid decreased the potency of inhibition, potentially due to the carboxylic acid acting as a less efficient zinc chelator. Removal of the

2-hydroxyl group or substitution with bromide or amine completely abrogated the inhibitory activity, whereas substitution at the 4' position with bromine was tolerated (Table 1).






To further explore the available chemical space, a small array of analogues of compound **1** were synthesised from the literature compound **11**<sup>21</sup> by coupling to the 5-NH<sub>2</sub> position with commercially available acid chlorides (Scheme 1). After deprotection of the hydroxamic acid, the majority of compounds showed decreased inhibitory activity compared with **1** (Table 2). The two most potent compounds **20** and **21** had IC<sub>50</sub> values of 45 ± 11  $\mu$ M and 69 ± 6  $\mu$ M respectively (Supplementary Fig. 1), comparable to the parent compound albeit with reduced ligand efficiency (-0.30 and -0.29 kcal mol<sup>-1</sup> per non-hydrogen atom respectively), suggesting that the 5'-amide substitution was tolerated but contributed little to binding.

The structure–activity relationship derived from the data presented here suggest that the 2-OH group of **1** is essential for activity and substitution at the 4- and 5- position is tolerated (Fig. 4). Comparing **1** with the natural substrate GlcNAc-PI could suggest that **1** is mimicking the GlcNAc moiety, with the 2-OH of **1** mimicking the 3'-OH that is essential for substrate recognition. In the proposed reaction mechanism for the de-*N*-acetylase, the catalytic base D43 assists the nucleophilic attack of a zinc-bound water molecule on the activated acetamidocarbonyl, and the subsequent protonation of the tetrahedral intermediate promotes loss of acetic




**Scheme 1.** Synthesis of a small array of 5' amide derivatives of **1**. Reagents and conditions: (a) TEA, Boc<sub>2</sub>O, dioxane, RT, 18 h, 76%. (b) BnONH<sub>2</sub>·HCl, DCC, DMAP, TEA, CHCl<sub>3</sub>/THF, 0 °C, 16 h to RT, 24 h, 46%. (c) TFA (wet), RT, 1 h, 85%. (d) R<sub>1</sub>-COCl, pyridine, DMAP (cat.), THF/DCE 4:1, RT, 24 h, 21–79%. (e) AcOH, H<sub>2</sub>, 20% Pd(OH)<sub>2</sub>, MeOH/THF, 0.5 mL/min, 1 atm, 45–100%.

**Table 2**  
Inhibitory activity of novel derivatives of **1**

ID	R <sub>1</sub>	Inhibition at 1 mM <sup>a</sup> (%)
20		99 ± 1.6
21		96 ± 0.4
22	Me- 	62 ± 24
23	CH <sub>3</sub> (CH <sub>2</sub> ) <sub>16</sub>	25 ± 14
24	AcN- 	40 ± 17
25		75 ± 7.0
26		56 ± 8.3
27		86 ± 11

<sup>a</sup> Inhibition of *T. brucei* de-*N*-acetylase (cell-free system).



**Figure 4.** Proposed mode of action. (A) Structure–activity relationships of **1**. (B) Components of the natural substrate GlcNAc-PI recognised by the enzyme. R<sub>1</sub> =  $\alpha$ -(1-6)-*D*-myo-inositol-1-phosphate-dimyristoylglycerol. (C) The GlcNAc-PI de-*N*-acetylase contains catalytic zinc chelated by the residues H49, D52 and H157, and the catalytic base D43. (D) Compound **1** could inhibit the enzyme through binding of the hydroxamic acid to the catalytic zinc, replacing both the acetamidocarbonyl and activated water.

### 3.4. Conclusions

We have identified salicylic hydroxamic acid **1** as an inhibitor of *T. brucei* GlcNAc-PI de-*N*-acetylase with high ligand efficiency, representing the first non-substrate analogue inhibitor of the trypanosome GPI pathway. Although **1** shows modest potency with an IC<sub>50</sub> of 63 ± 4 μM, it has impressive ligand efficiency of 0.57 kcal mol<sup>-1</sup> per non-hydrogen atom, and thus may be a useful starting point to develop more potent inhibitors.

### Acknowledgments

This work was funded by the Medical Research Council (studentship to ASC) and the Wellcome Trust (Programme Grant 085622 and 077705, and Strategic Awards 083481 and 100476). The funders had no further role in the study design.

### Supplementary data

Supplementary data associated with this article can be found, in the online version, at <http://dx.doi.org/10.1016/j.carres.2013.12.016>.

### References

- Ferguson, M. A. J. *J. Cell Sci.* **1999**, *112*, 2799–2809.
- Nagamune, K.; Nozaki, T.; Maeda, Y.; Ohishi, K.; Fukuma, T.; Hara, T.; Schwarz, R. T.; Sutterlin, C.; Brun, R.; Reizman, H.; Kinoshita, T. *Proc. Natl. Acad. Sci. U.S.A.* **2000**, *97*, 10336–10341.
- Chang, T.; Milne, K. G.; Guther, M. L. S.; Smith, T. K.; Ferguson, M. A. J. *J. Biol. Chem.* **2002**, *277*, 50176–50182.
- Lillico, S.; Field, M. C.; Blundell, P.; Coombs, G. H.; Mottram, J. C. *Mol. Biol. Cell* **2003**, *14*, 1182–1194.
- Smith, T. K.; Crossman, A.; Brimacombe, J. S.; Ferguson, M. A. J. *EMBO J.* **2004**, *23*, 4701–4708.
- Urbaniak, M. D.; Crossman, A.; Chang, T.; Smith, T. K.; Aalten, D. M. F. v.; Ferguson, M. A. J. *J. Biol. Chem.* **2005**, *280*, 22831–22838.
- Smith, T. K.; Sharma, D. K.; Crossman, A.; Dix, A.; Brimacombe, J. S.; Ferguson, M. A. J. *EMBO J.* **1997**, *16*, 6667–6675.
- Smith, T. K.; Patterson, M. J.; Crossman, A.; Brimacombe, J. S.; Ferguson, M. A. J. *Biochemistry* **2000**, *39*, 11801–11807.
- Smith, T. K.; Crossman, A.; Borrissow, C. N.; Paterson, M. J.; Dix, A.; Brimacombe, J. S.; Ferguson, M. A. J. *EMBO J.* **2001**, *20*, 3322–3332.
- Urbaniak, M. D.; Ferguson, M. A. J. *Enzymes* **2009**, 49–64.
- Urbaniak, M. D.; Crossman, A.; Ferguson, M. A. J. *Chem. Biol. Drug Des.* **2008**, *72*, 127–132.
- Abdelwahab, N. Z.; Crossman, A. T.; Urbaniak, M. D.; Ferguson, M. A. J. *Carbohydr. Res.* **2011**, *346*, 708.
- Abdelwahab, N. Z.; Crossman, A. T.; Sullivan, L.; Ferguson, M. A. J.; Urbaniak, M. D. *Chem. Biol. Drug Des.* **2012**, *79*, 270–278.
- Crossman, A.; Patterson, M. J.; Ferguson, M. A. J.; Smith, T. K.; Brimacombe, J. S. *Carbohydr. Res.* **2002**, *337*, 2049–2059.
- Masterson, W. J.; Doering, T. L.; Hart, G. W.; Englund, P. W. *Cell* **1989**, *62*, 73–80.
- Hopkins, A. L.; Groom, C. R.; Alex, A. *Drug Discovery Today* **2004**, *9*, 430–431.
- Kuntz, I. D.; Chen, K.; Sharp, K. A.; Kollman, P. A. *Proc. Natl. Acad. Sci. U.S.A.* **1999**, *96*, 9997–10002.
- Brenk, R.; Schipani, A.; James, D.; Krasowski, A.; Gilbert, I. H.; Frearson, J. A.; Wyatt, P. G. *ChemMedChem* **2008**, *3*, 435–444.
- Clarkson, A. B.; Brohn, F. H. *Science* **1976**, *194*, 204–206.
- Robert Ott; Kelly Chibale; Sederick Anderson; Alex Chipeleme; Minu chaudhuri; Abdelmadjid Guerrah; Nancy Colowick; Hill, G. C. *Acta Trop.* **2006**, *100*, 172–184.
- Chen, H.; Boiziau, J.; Parker, F.; Mailliet, P.; Commercon, A.; Tocque, B.; Le Pecq, J. B.; Roques, B. P.; Garbay, C. J. *Med. Chem.* **1994**, *37*, 845–859.

acid. We propose that **1** could inhibit the enzyme through the hydroxamic acid binding to the catalytic zinc, and replacing both the acetamidocarbonyl and the activated water (Fig. 4).