

A CHARACTERISATION OF GENERICALLY RIGID FRAMEWORKS ON SURFACES OF REVOLUTION

A. NIXON*, J.C. OWEN, AND S.C. POWER†

ABSTRACT. A foundational theorem of Laman provides a counting characterisation of the finite simple graphs whose generic bar-joint frameworks in two dimensions are infinitesimally rigid. Recently a Laman-type characterisation was obtained for frameworks in three dimensions whose vertices are constrained to concentric spheres or to concentric cylinders. Noting that the plane and the sphere have 3 independent locally tangential infinitesimal motions while the cylinder has 2, we obtain here a Laman-type theorem for frameworks on algebraic surfaces with a 1-dimensional space of tangential motions. Such surfaces include the torus, helicoids and surfaces of revolution. The relevant class of graphs are the $(2, 1)$ -tight graphs, in contrast to $(2, 3)$ -tightness for the plane/sphere and $(2, 2)$ -tightness for the cylinder. The proof uses a new characterisation of simple $(2, 1)$ -tight graphs and an inductive construction requiring generic rigidity preservation for 5 graph moves, including the two Henneberg moves, an edge joining move and various vertex surgery moves.

1. INTRODUCTION

A bar-joint framework in real Euclidean space \mathbb{R}^d is a geometric realisation of the vertices of a graph with the edges considered as inextensible bars between them. Such a framework is said to be rigid if there is no non-trivial continuous motion of the framework vertices which maintains bar-lengths, and is said to be flexible if it is not rigid. A foundational theorem of Laman, obtained in 1970, asserts that the rigidity of a generically positioned framework in the plane depends only on the underlying graph and furthermore these graphs are characterised in terms of a simple counting condition. There is also an elegant recursive construction of the minimally rigid frameworks going back to Henneberg [6, 9] in which each framework may be derived from a single edge framework by repeated application of two simple construction moves, namely the Henneberg 1 move and the Henneberg 2 move. Analogous characterisations for frameworks in \mathbb{R}^3 remain open problems and no combinatorial characterisation of generic rigidity is known. We note however that a number of partial and related results are given in Whiteley [22] and that the longstanding molecular conjecture has been resolved by Katoh and Tanigawa [8].

Attention has also been given to frameworks in 3-dimensional space whose vertices are constrained to 2-dimensional surfaces. In [12] we obtained Laman-Henneberg-type theorems for the case of constraint to parallel planes, concentric spheres and concentric

*part of this research was carried out at the Fields Institute, Toronto.

†supported by EPSRC grant EP/J008648/1.

2010 *Mathematics Subject Classification*. 52C25, 05B35, 05C10, 53A05

Key words and phrases: bar-joint framework, infinitesimally rigid, framework on a surface.

cylinders. The cylinder case presents new complications both for the purely graph theoretical analysis and for the preservation of rigidity under the Henneberg moves and the further construction moves that are needed.

A fixed plane surface in \mathbb{R}^3 supports a three-dimensional vector space of internal infinitesimal motions, coming from translations and rotations, while the (infinite circular) cylinder has two such independent motions. More formally, in Definition 2.2 we define the type k of an irreducible algebraic surface where k takes values 3, 2, 1 or 0. (See also Definition 8.1.) The type of a surface is reflected in the graph counting conditions for Laman type theorems; fewer independent infinitesimal motions for the surface imply a richer set of graphs which in turn require more constructive moves and more refined proof techniques.

The main result in the present paper is the following theorem for bar-joint frameworks in \mathbb{R}^3 whose vertices are constrained to an algebraic surface \mathcal{M} of type 1. These surfaces include a range of fundamental algebraic surfaces such as the elliptical cylinder, the cone, the torus, surfaces of revolution, and various helical glide-translation surfaces.

Theorem 1.1. *Let $G = (V, E)$ be a simple graph, let \mathcal{M} be an irreducible algebraic surface in \mathbb{R}^3 of type 1 and let (G, p) be a generic framework on \mathcal{M} . Then (G, p) is isostatic on \mathcal{M} if and only if G is K_1, K_2, K_3, K_4 or is $(2, 1)$ -tight.*

For a plane surface the required graphs are the Laman graphs, corresponding to the top count $|E| = 2|V| - 3$ and the inequality $|E'| \leq 2|V'| - 3$ for every subgraph (V', E') . In fact these are the $(2, 3)$ -tight graphs, in the sense of Definition 2.8 and are necessarily simple. Observe that these graphs are necessarily simple. For the cylinder the appropriate graphs are the simple $(2, 2)$ -tight graphs. In this paper the key class of graphs are the simple $(2, 1)$ -tight graphs which were characterised recently in Nixon and Owen [13]. We shall obtain here the following alternative characterisation which turns out to be efficient for our purposes. The methods for this also lead to a new analogous characterisation of simple $(2, 2)$ -tight graphs given in Section 3.

Theorem 1.2. *A simple finite graph is $(2, 1)$ -tight if and only if it is equal to $K_5 \setminus e$ or can be obtained from this graph by the sequential application of moves of 5 types, namely the Henneberg 1 and 2 moves, the vertex-to- K_4 move, the vertex-to-4-cycle move, and the edge joining move.*

The proof of the main theorem is principally concerned with the sufficiency for rigidity of the combinatorial condition and we obtain this by showing that each of the moves in the constructive sequence for the graph preserves generic rigidity. We introduce some new methods for this and there are two moves that present particular challenges, namely the Henneberg 2 move and the vertex-to- K_4 move. For the Henneberg 2 move we give two quite different proofs which also give new proofs in the case of the circular cylinder. The first of these uses a convergence argument involving a sequence of generic realisations of G' which converge to a degenerate non-generic realisation of G' which in a natural sense covers a generic realisation of G . On the other hand in Section 6 we adopt the traditional approach of algebraic specialisation to obtain a direct entirely algebraic proof.

For the vertex-to- K_4 move we show that proper flexes are inherited under the inverse move $G' \rightarrow G$ corresponding to K_4 contraction to a vertex. This is achieved through the

consideration of a sequence (G', p^k) in which the K_4 -subframeworks contract in a manner which is *well-behaved* with respect to the distinct principal curvatures of the surface.

In all our considerations the framework vertices are constrained to a surface while the edges are straight Euclidean edges measured by distances in \mathbb{R}^3 . We note that Whiteley [20] has considered the topic of frameworks on surfaces with geodesic edges and there the appropriate combinatorial objects are looped multigraphs. Moreover, rigidity is expressed in terms of the rank of the k -frame matrix rather than the rigidity matrix. We note also that frameworks on surfaces are also implicit in the context of periodic frameworks, considered, for example, by Borcea and Streinu [3], Malestein and Theran [10, 11], Nixon and Ross [15], Owen and Power [16] and Ross [17].

The paper is structured as follows. In Section 2 we recall basic definitions and key results from [12] for generic frameworks on surfaces. In Section 3 we detail the inductive moves on graphs and obtain inductive characterisations of simple $(2, k)$ -tight graphs for $k = 1$ and 2. In Section 4 we prove the preservation of generic independence for the two Henneberg moves as moves on frameworks on algebraic surfaces. In Section 5 we prove the preservation of generic independence for various vertex surgery moves, including vertex splitting, vertex-to-4-cycle and vertex-to- K_4 . In Section 6 we provide an alternative proof of generic independence under Henneberg 2 moves that we believe could be of independent interest. In Section 7 we prove the main theorem and in the final section we discuss the difficulty of extending our results to other surfaces and comment on higher dimensional contexts.

2. FRAMEWORKS ON SURFACES

Let \mathcal{M} be a subset of \mathbb{R}^3 which is a smooth surface in the sense of being a 2-dimensional embedded differentiable manifold. The main examples we have in mind are defined as disjoint unions of parts of elementary algebraic surfaces. Accordingly we assume smoothness in the sense that local coordinate maps exist for \mathcal{M} which are analytic. In particular for every point of \mathcal{M} there is a continuous choice of normal vectors in a neighbourhood of the point and a Taylor series expansion, as in Equation 5.1, for points of \mathcal{M} in this neighbourhood.

A framework (G, p) on \mathcal{M} is a finite bar-joint framework in \mathbb{R}^3 , for a simple graph $G = (V, E)$, with framework points $p(v), v \in V$, which lie on \mathcal{M} .

An *infinitesimal flex* of (G, p) on \mathcal{M} is a sequence or vector u of velocity vectors $u_1, \dots, u_{|V|}$, considered as acting at the framework points, which are tangential to the surface and satisfy the infinitesimal flex requirement in \mathbb{R}^3 , namely

$$u_i \cdot (p_i - p_j) = u_j \cdot (p_i - p_j),$$

for each edge $v_i v_j$. It is elementary to show that u is an infinitesimal flex if and only if u lies in the nullspace (kernel) of the rigidity matrix $R_{\mathcal{M}}(G, p)$ given in the following definition. The submatrix of $R_{\mathcal{M}}(G, p)$ given by the first $|E|$ rows provides the usual rigidity matrix, $R_3(G, p)$ say, for the unrestricted framework (G, p) . The tangentiality condition corresponds to u lying in the nullspace of the matrix formed by the last $|V|$ rows.

Definition 2.1. *The rigidity matrix $R_{\mathcal{M}}(G, p)$ of (G, p) on \mathcal{M} is an $|E| + |V|$ by $3|V|$ matrix in which consecutive triples of columns correspond to framework points. The first $|E|$ rows correspond to the edges and the entries in row $e = uv$ are zero except possibly in*

the column triples for $p(u)$ and $p(v)$, where the entries are the coordinates of $p(u) - p(v)$ and $p(v) - p(u)$ respectively. The final $|V|$ rows correspond to the vertices and the entries in the row for vertex v are zero except in the columns for v where the entries are the coordinates of a normal vector $N(p(v))$ to \mathcal{M} at $p(v)$.

The case of a surface \mathcal{M} which is a subset of the nonsingular points of a polynomial equation $m(x, y, z) = 0$ is of particular interest, especially when $m(x, y, z)$ is irreducible over some coefficient field. In what follows we confine attention to the rational field and refer to such surfaces simply as *irreducible surfaces*. In this case we may take the derivative of $m(x, y, z)$ at $p(v)$ for the choice of normal $N(p(v))$. Furthermore, the rigidity matrix arises from the derivative of the augmented edge-function \tilde{f}_G , with

$$2R_{\mathcal{M}}(G, p) = (D\tilde{f}_G)(p),$$

where $\tilde{f}_G : \mathbb{R}^{3|V|} \rightarrow \mathbb{R}^{|E|+|V|}$ is given by $\tilde{f}_G(q) = (f_G(q), m(q_1), \dots, m(q_{|V|}))$ with

$$f_G(q) = (\|q_i - q_j\|^2)_{v_i v_j \in E}$$

the usual edge function for G associated with framework realisations in \mathbb{R}^{3n} , where $\|\cdot\|$ is the usual Euclidean norm.

As is well-known, for $n \geq 4$ a complete graph framework (K_n, p) in \mathbb{R}^3 , not lying in a hyperplane, has a 6-dimensional vector space of infinitesimal flexes, a basis for which may be provided by a set of linearly independent infinitesimal flexes associated with translations and rotations. When the vertices of (K_n, p) are constrained to a surface \mathcal{M} then the dimension is reduced to $\dim \ker R_{\mathcal{M}}(K_n, p) = k$ where $k = 3, 2, 1$ or 0 .

We now define smooth surfaces of type k for $k = 3, 2, 1, 0$. The type number reflects the number of independent infinitesimal motions of a typical framework on \mathcal{M} that arise from isometries of \mathbb{R}^3 that act tangentially at every point on \mathcal{M} (not just the framework joints). For the sphere, cylinder and cone the types are 3, 2 and 1 respectively, while the ellipsoid, defined by $x^2 + 2y^2 + 3z^2 = 1$, has type 0.

Definition 2.2. *A surface \mathcal{M} is said to be of type k , or to have freedom number k , if $\dim \ker R_{\mathcal{M}}(K_n, p) \geq k$ for all complete graph frameworks (K_n, p) on \mathcal{M} and k is the largest such number.*

Apart from the type 3 surfaces, which arise from concentric spheres or parallel planes, a typical K_4 framework on a surface \mathcal{M} has a two-dimensional space of infinitesimal flexes. This follows on consideration of the 10 by 12 rigidity matrix. For K_3 and K_2 frameworks the space is three-dimensional and includes rotational flexes not derivable from (tangentially acting) isometries. For the cylinder the flexes of K_4 frameworks are all associated with isometries whereas on the cone (resp. ellipsoid) there is a one-dimensional (resp. zero-dimensional) subspace determined by tangential isometries.

Definition 2.3. *Let \mathcal{M} be a smooth surface and $p = (p_1, \dots, p_n)$ a vector of points on \mathcal{M} . Then the framework (G, p) on \mathcal{M} is said to be infinitesimally rigid if every infinitesimal flex of (G, p) on \mathcal{M} corresponds to a rigid motion flex of \mathcal{M} .*

A framework (G, p) on \mathcal{M} is *independent* if $R_{\mathcal{M}}(G, p)$ has linearly independent rows and is *minimally infinitesimally rigid* on \mathcal{M} , or *isostatic* on \mathcal{M} if it is independent and infinitesimally rigid on \mathcal{M} .

From the point of view of the infinitesimal rigidity it is only the nature of $\mathcal{M}^{|V|}$ in a neighbourhood of p which is of significance. On the other hand for irreducible surfaces one can establish generic properties for the pair G, \mathcal{M} as we shall see.

Following Asimow and Roth we say that a framework (G, p) on \mathcal{M} is *regular* if the rank of $R_{\mathcal{M}}(G, p)$ takes its maximum value throughout a neighbourhood of p in $\mathcal{M}^{|V|}$. In the case that \mathcal{M} is an algebraic surface determined by an irreducible polynomial $m(x, y, z)$ over \mathbb{Q} , the framework (G, p) is said to be *generic* on \mathcal{M} if an algebraic dependency $h(\{x_i\}, \{y_i\}, \{z_i\}) = 0$ holds between the coordinates x_i, y_i, z_i of all the points p_i only when the polynomial $h(\{X_i\}, \{Y_i\}, \{Z_i\})$ lies in the ideal generated by the polynomials $m(X_i, Y_i, Z_i), 1 \leq i \leq |V|$. It is a standard exercise to show that such generic frameworks are regular.

Note that in contrast to type 3 and 2 there are diverse classical surfaces of type 1, including spheroids, with isometry group S^1 , elliptical cylinders and other noncircular cylinders, with translational isometry group \mathbb{R}^1 , and circular hyperboloids and other diverse surfaces with glide-rotation isometry group \mathbb{R}^1 .

Definition 2.4. *A simple graph G is independent for the irreducible surface \mathcal{M} if every generic framework (G, p) on \mathcal{M} is independent.*

In particular, K_4 is dependent for the sphere but independent for the cylinder. On the other hand $K_5 \setminus e$ is dependent for the cylinder but independent for the cone. Note that $K_n \setminus e$ denotes the unique graph formed by deleting any single edge from K_n .

The determination of combinatorial conditions for the generic independence of classes of frameworks is one of the fundamental problems in constraint system rigidity theory. See for example Whiteley [21, 22] and Jackson and Jordan [7]. Our main result can be viewed in this spirit. Also we note that there is the following matroidal interpretation of our main result. Let $\mathcal{L}(K_n, \mathcal{M})$ be the linear matroid for the rigidity matrix $R_{\mathcal{M}}(K_n, p)$ associated with a generic n -tuple and the irreducible surface \mathcal{M} . Then by Theorem 1.1 the bases of $\mathcal{L}(K_n, \mathcal{M})$ correspond to sets of rows determined by the $(2, 1)$ -tight subgraphs.

The notions of continuous rigidity and minimal continuous rigidity are also naturally defined in the surface setting and the following equivalence is an analogue of a theorem of Gluck [5].

Theorem 2.5. [12] *A generic framework (G, p) on an algebraic surface \mathcal{M} is infinitesimally rigid if and only if it is continuously rigid on \mathcal{M} .*

In this paper the infinitesimal rigidity perspective will be more direct and we make use of the following two results from [12], namely a version of the Maxwell counting condition and an isostatic characterisation in the spirit of Asimow and Roth [1].

Theorem 2.6. [12] *Let (G, p) be an isostatic generic framework on the algebraic surface \mathcal{M} of type $k, 0 \leq k \leq 3$, with G not equal to K_1, K_2, K_3 or K_4 . Then $|E| = 2|V| - k$ and for every subgraph H of G with at least one edge, $|E(H)| \leq 2|V(H)| - k$.*

Theorem 2.7. [12] *Let (G, p) be a generic framework on a surface \mathcal{M} of type k . Then (G, p) is isostatic on \mathcal{M} if and only if*

- (1) $\text{rank } R_{\mathcal{M}}(G, p) = 3|V| - k$ and
- (2) $2|V| - |E| = k$.

The classes of graphs in Theorem 2.6 are the simple graphs that are $(2, k)$ -tight, with $k = 0, 1, 2, 3$, in the following sense.

Definition 2.8. *A graph $G = (V, E)$ is $(2, k)$ -sparse if for all subgraphs H , with at least one edge, the inequalities $|E(H)| \leq 2|V(H)| - k$ hold. Moreover G is $(2, k)$ -tight if G is $(2, k)$ -sparse and $|E| = 2|V| - k$.*

The inductive characterisations of these classes of simple graphs for $k = 3, 2$ and 1 form a key part of our approach to proving the sufficiency of the necessary counting conditions for generic infinitesimal rigidity. We describe the various construction moves in the next section.

Theorem 2.9 (Henneberg [6] and Laman [9]). *A simple graph G is $(2, 3)$ -tight if and only if it can be generated from K_2 by Henneberg 1 and 2 moves.*

This characterisation plays a role in the following extension of Laman's theorem.

Theorem 2.10. [12] *Let $G = (V, E)$, let \mathcal{M} be a union of parallel planes or a union of concentric spheres and let p be generic on \mathcal{M} . Then (G, p) is isostatic on \mathcal{M} if and only if G is K_1, K_2 or $(2, 3)$ -tight.*

Theorem 2.11. [12, 13] *For a simple graph G the following are equivalent:*

- (1) G is $(2, 2)$ -tight,
- (2) G can be generated from K_1 by Henneberg 1, Henneberg 2 and graph extension moves,
- (3) G can be generated from K_1 by Henneberg 1, Henneberg 2, vertex-to- K_4 and vertex splitting moves.

Theorem 2.12. [12] *Let $G = (V, E)$, let \mathcal{M} be a cylinder or a union of concentric cylinders and let p be generic on \mathcal{M} . Then (G, p) is isostatic on \mathcal{M} if and only if G is K_1, K_2, K_3 or is $(2, 2)$ -tight.*

We observe that this theorem can also be proven, using the methods of this paper, by applying the equivalence of (1) and (3) in Theorem 2.11 or by applying Theorem 3.1.

We also note the following alternative characterisation of $(2, 1)$ -tight simple graphs to that which is given in Theorem 1.2. This will not be needed for the proofs in this paper but we note that at the expense of introducing the new base graph $K_4 \sqcup K_4$ one may replace the vertex-to-4-cycle move by the more localised vertex splitting move. ($K_4 \sqcup K_4$ consists of two copies of K_4 sharing exactly one edge.)

Theorem 2.13. [13] *A simple graph G is $(2, 1)$ -tight if and only if it can be generated from $K_5 \setminus e$ or $K_4 \sqcup K_4$ by Henneberg 1, Henneberg 2, vertex-to- K_4 , vertex splitting and edge joining moves.*

3. SIMPLE $(2, 1)$ -TIGHT GRAPHS

In this section we prove Theorem 1.2 and in Theorem 3.1 we obtain an analogous characterisation of $(2, 2)$ -tight simple graphs. We remark that the insistence on simplicity of the graph makes the construction problematic. Indeed if we permit loops and parallel edges then a general construction theorem of Fekete and Szegő [4] applies. In the case of $(2, 1)$ -tight graphs it ensures that the only operations required are Henneberg 1 and 2

type operations. Fekete and Szegő’s result extended work of Tay on (k, k) -tight graphs for the characterisation of the generic rigidity of body-bar frameworks in arbitrary dimension [18].

The inductive characterisation of $(2, 3)$ -tight graphs of Henneberg and Laman starts with the elementary counting observation that for $k \geq 1$ the average degree in a $(2, k)$ -tight graph is less than 4, while there can be no vertices of degree less than 2. Thus degree 2 or degree 3 vertices exist. Henneberg introduced the following two operations on graphs which maintain the sparsity count, increase the vertex count by 1, and add a new vertex of degree 2 or degree 3 respectively. See also the discussions in [9, 14, 19].

The *Henneberg 1* move augments a graph G by adding a vertex v of degree 2 and two edges vv_1, vv_2 from it to distinct neighbours v_1, v_2 in G .

The *Henneberg 2* move removes an edge v_1v_2 from a graph and adds a vertex v of degree 3 with distinct neighbours v_1, v_2, v_3 for some vertex v_3 . This is also referred to as edge splitting or 1-extension in the literature.




FIGURE 1. The Henneberg 2 move.

Laman proved that every $(2, 3)$ -tight graph can be generated recursively from K_2 by a sequence of these operations. The key step is to show that an inverse Henneberg 2 operation is possible on a degree 3 vertex v , by virtue of the fact that at least one of the three choices for the new edge (v_1v_2, v_2v_3 or v_3v_1) can be made without violating any subgraph count.

On the other hand, for a $(2, 2)$ -tight graph it is easy to see that degree 3 vertices may be contained within subgraphs isomorphic to K_4 and so admit no inverse Henneberg 2 move. Indeed there are countably many $(2, 2)$ -tight graphs for which every vertex of degree less than 4 is contained in a copy of K_4 . In view of this obstruction, in [12, 13] we considered additional graph moves preserving $(2, 2)$ -tightness, including those indicated in Figure 2 and Figure 3. These are the *vertex-to- K_4 move* and the *vertex splitting move*.

The *vertex-to- K_4 move* substitutes a copy of K_4 in place of a vertex v , with an arbitrary replacement of edges xv by edges xw with w in $V(K_4)$. The inverse operation, contracting a copy of K_4 to a single vertex will be called the *K_4 -to-vertex move*.

The *vertex splitting* operation removes an edge uv and vertex v and inserts a copy of K_3 on vertices u, v_1, v_2 and assigns all edges xv into either an edge xv_1 or an edge xv_2 . That is, let $G = (V, E)$, let $V^* := V \setminus v$, let $E^* := E(G[V^*])$ and let $E' := \{xy \in E : y = v\}$.




FIGURE 2. The vertex-to- K_4 move.




FIGURE 3. The vertex splitting move.

Thus $G = (V, E) = (V^* + v, E^* + E' + uv)$ and if G' is the result of a vertex split on the edge uv then $G' = (V', E') = (V^* + \{v_1, v_2\}, E^* + E'_1 + E'_2 + \{uv_1, uv_2, v_1v_2\})$ where $E'_i := \{xy \in E' : x = v_i\}$ is an arbitrary partition of E' . We refer to the inverse move as an edge contraction.

We remark that the vertex-to- K_4 move is a special case of the more elaborate *graph extension move* employed in [14]. We also remark that the vertex splitting move is not needed for the proof of our main theorem.

Finally, an *edge-joining* move [13] combines two graphs G, H to form a new graph with the vertices and edges of these graphs together with an additional connecting edge $e = gh$ with g in G and h in H .

The *vertex-to-4-cycle* move is a certain vertex splitting operation, as in Figure 4. The vertex v_1 is split to two vertices v_1 and v_0 and the edges v_1v_2 and v_1v_3 are duplicated as v_0v_2 and v_0v_3 . Other edges of the form vv_1 are either left or are replaced by vv_0 . The move preserves $(2, k)$ -tightness and after the vertex split there will be no edge pair wv_1, wv_0 with $w \neq v_2, v_3$ and no edge v_0v_1 .




FIGURE 4. The vertex-to-4-cycle move.

The inverse of such a move on a simple $(2, 1)$ -tight graph will be referred to as a 4-cycle contraction.

For each of our moves we use the term *admissible*, if whenever G is simple and $(2, k)$ -tight, the result of the move is also simple and $(2, k)$ -tight. It is evident that inverse Henneberg 1 moves and, when $k = 1$, inverse edge joining moves are always admissible. For each of the other moves it is easy to observe examples when the inverse moves are non-admissible.

We now prove Theorem 1.2.

Proof of Theorem 1.2. A basic incidence degree counting argument shows that if G is $(2, 1)$ -tight with no possible inverse Henneberg 1 moves then there are at least two degree 3 vertices. In fact these vertices must be in copies of K_4 graphs. If not then there is a potential inverse Henneberg 2 move and a counting argument shows that these inverse moves are admissible. (This counting argument is given explicitly in [13, Lemma 3.1].)

Suppose now that one of these K_4 graphs does not have an admissible contraction to a vertex. Then there is a vertex w outside this K_4 with two edges to distinct vertices a, b in this K_4 , neither of which is of degree 3 in G . Thus there is a 4-cycle including w, a, b and a degree 3 vertex v of the K_4 graph. The edge vw is absent from G and so there is a potential inverse 4-cycle contraction, with $v \rightarrow w$, as long as the edge wc is absent, where c is the fourth K_4 vertex, since the move creates this edge.

Suppose first that wc is absent. We claim that the 4-cycle contraction is admissible.

To check this we need only consider the count change for subgraphs Y of G , with $f(Y) = 2|V(Y)| - |E(Y)| = 1$, which include w and c (and so receive wc) but not v (since otherwise the count change is zero). There are three cases:

- (i) a and b are in Y . However, adding v and 3 edges would create Y^+ with $f(Y^+) = 0$, so Y cannot have $f(Y) = 1$.
- (ii) a and b are not in Y . However, adding a and b and their 5 edges aw, ac, bc, bw, ab to Y creates Y^+ with $f(Y^+) = 0$.
- (iii) Just b (or a) is in Y . However, adding a and the 3 edges ac, aw, ab creates Y^+ with $f(Y^+) = 0$.

Thus this 4-cycle contraction is admissible.

Suppose now that the edge wc is present. Then G contains a copy of $K_5 \setminus e$ supported by the vertices v, w, a, b, c . Moreover assume that this is true for every degree 3 vertex, that is, there is no admissible inverse Henneberg 2, admissible K_4 -to-vertex or admissible 4-cycle contraction. We show that if G is not isomorphic to $K_5 \setminus e$ then there is an inverse edge-joining move. This completes the proof of the theorem.

We argue as in Lemma 4.10 of [13]. Let $Y = \{Y_1, \dots, Y_n\}$ be the subgraphs which are copies of $K_5 \setminus e$. They are necessarily vertex disjoint since $f(Y_i \cup Y_j) = 2 - f(Y_i \cap Y_j)$ and every proper subgraph X of $K_5 \setminus e$ has $f(X) \geq 2$. Let V_0 and E_0 be the sets of edges in G which are in none of the Y_i . Then

$$f(G) = \sum_{i=1}^n f(Y_i) + 2|V_0| - |E_0|$$

so $|E_0| = 2|V_0| + n - 1$. Since every vertex of degree less than 4 is contained in some Y_i , each vertex in V_0 is incident to at least 4 edges. If every Y_i is incident to at least 2 edges in E_0 then there are at least $4|V_0| + 2n$ edge/vertex incidences in E_0 . This implies

$|E_0| \geq 2|V_0| + n$, a contradiction. Thus either there is a copy Y_i with no incidences, which would imply $G = Y_i$, since G is connected, contrary to our assumption, or there is a copy with one incidence. In this case there is an inverse edge joining move, as desired. \square

As mentioned in the introduction, a simpler version of the same proof scheme gives a construction for $(2, 2)$ -tight graphs as follows.

Theorem 3.1. *A simple graph G is $(2, 2)$ -tight if and only if it can be obtained from K_1 by the sequential application of moves of 4 types, namely the Henneberg 1 and 2 moves, the vertex-to- K_4 move and the vertex-to-4-cycle move.*

Proof. Suppose that G is simple and $(2, 2)$ -tight with at least one edge. Then, by simple counting, there exist vertices of degree 2 or 3. As in the last proof, if there are no inverse Henneberg moves then it follows that the minimum degree is 3 and any such vertex v must lie in a copy of K_4 . As above if there is no admissible contraction on this K_4 then there is a vertex w adjacent to two vertices a, b in that copy of K_4 . Since G is $(2, 2)$ -sparse there is no edge wc , with c the final K_4 vertex, and hence, the argument above, shows there is an admissible 4-cycle contraction move. \square

4. HENNEBERG MOVES ON FRAMEWORKS ON SURFACES

The role played by the Henneberg moves in rigidity theory is extensive and well studied (see for example [2, 6, 7, 9, 18, 22]) and we now consider such moves on frameworks on surfaces. The case of the Henneberg 1 move is elementary. Recall that an isostatic framework is one which is independent and infinitesimally rigid.

Lemma 4.1. *Let (G, p) be a generic framework on an algebraic surface \mathcal{M} , let G' be a graph obtained from G by a Henneberg 1 move with new vertex v , and let $p' = (p, p_v)$ be generic. Then (G, p) is isostatic on \mathcal{M} if and only if (G', p') is isostatic on \mathcal{M} .*

Proof. The rigidity matrix $R_{\mathcal{M}}(G', p')$ contains 3 new rows and 3 new columns and those columns are zero everywhere in the $|E| + |V|$ rows for G . Reorder the rows and columns so that the first three rows and columns are the new ones. By the generic location of p_v and the block upper triangular structure the first three rows are independent of the rest. Thus there is a row dependency in $R_{\mathcal{M}}(G', p')$ if and only if there is a row dependency in $R_{\mathcal{M}}(G, p)$. \square

The preservation of independence and isostaticity under the Henneberg 2 move is considerably more subtle. To see an aspect of this let (G, p) be generic on \mathcal{M} , where \mathcal{M} is the cylinder surface $x^2 + y^2 = 1$ in \mathbb{R}^3 . Examining the form of $R_{\mathcal{M}}(G, p)$, it is evident that the addition of a degree 0 vertex increases the rank by 1. Elementary linear algebra also shows that the addition of a degree 1 framework point (x_1, y_1, z_1) incident to a point (x, y, z) increases the rank by 2 if and only if (x_1, y_1, z_1) is not equal to (x, y, z) or $(-x, -y, z)$. More surprisingly, similar considerations show that there are only four possible points where the rank does not fully increase for a new degree 2 vertex. This is contrary to the situation in the plane where any point on a line through the existing edge will create a copy of K_3 whose rows give a minimal linear dependency. In view of this, adapting a typical Henneberg 2 argument for rigidity preservation would require placing the new degree 3 vertex at one of a finite number of points. However, the dependencies created by each of these points are not amenable to such an argument.

Such difficulties motivated us in [12, Section 4], to show that continuous rigidity was preserved. The proof there makes particular use of the two trivial motions of the cylinder and it is not clear how one might generalise this method to surfaces with fewer trivial motions. Our first new approach below is based on the convergence of specialised frameworks. A direct algebraic proof is given in Section 6.

The following notation and observations will be useful.

Let \mathcal{T}_{q_i} denote the tangent space for the point q_i on the surface \mathcal{M} and for a framework vector $q = (q_1, q_2, \dots, q_n)$ let

$$\mathcal{T}_q = \mathcal{T}_{q_1} \oplus \mathcal{T}_{q_2} \oplus \dots \oplus \mathcal{T}_{q_n} \subseteq \mathbb{R}^3 \oplus \dots \oplus \mathbb{R}^3 = \mathbb{R}^{3n}$$

denote the joint tangent space. This is the space of infinitesimal velocities of any framework with framework vector q . With the usual Euclidean structure we have orthogonal projections

$$P_q : \mathbb{R}^{3n} \rightarrow \mathcal{T}_q, \quad F_q : \mathbb{R}^{3n} \rightarrow \mathcal{F}(G, q), \quad Q_q : \mathbb{R}^{3n} \rightarrow \mathcal{R}_q,$$

where $\mathcal{F}(G, q)$ is the vector space of infinitesimal flexes of the framework (G, q) and where \mathcal{R}_q is the subspace of \mathcal{T}_q consisting of rigid motion infinitesimal flexes. We have $\mathcal{R}_q = \ker R_{\mathcal{M}}(K_n, q)$ for $n \geq 6 - k$, where k is the type of \mathcal{M} . Since \mathcal{M} is smooth the function $q \rightarrow P_q$ is continuous in a neighbourhood of p . This also holds in the case of a *degenerate framework vector* p , in the sense that some or all of the vectors p_i may agree. Similarly, if the spaces \mathcal{R}_q have dimension k throughout a neighbourhood of a (possibly degenerate) framework vector p then the function $q \rightarrow Q_q$ is continuous, as long as the degeneracy includes three non-collinear points. In general the function $q \rightarrow \mathcal{F}(G, q)$ is lower semi-continuous.

Lemma 4.2. *Let G be a simple graph, let G' be derived from G by a Henneberg 2 move and let \mathcal{M} be an irreducible surface. If G is minimally infinitesimally rigid on \mathcal{M} then G' is minimally infinitesimally rigid on \mathcal{M} .*

Proof. Let (G, p) be generic on \mathcal{M} with $p = (p_1, \dots, p_n)$ and let $p' = (p_0, p)$ where (G', p') is generic on \mathcal{M} . We let $v_1 v_2$ denote the edge involved in the Henneberg move and write v_0 for the new vertex. Suppose that (G', p') is not infinitesimally rigid on \mathcal{M} . Then it follows that every specialised framework on \mathcal{M} with graph G' is infinitesimally flexible. Figure 5 indicates a sequence of specialisations (G', p^k) in which only the p_0 framework point is specialised to a point p_0^k . Also p_0^k tends to p_2 in the direction a where a is a tangent vector at p_2 which is orthogonal to a tangent vector b at p_2 where b is orthogonal to $p_2 - p_1$. More precisely, the normalised vector $(p_2^k - p_0^k) / \|p_2^k - p_0^k\|$ converges to a , as $k \rightarrow \infty$. Each of the frameworks (G', p^k) has a unit norm flex u^k which is orthogonal to the space of rigid motion infinitesimal flexes of its framework. By the Bolzano-Weierstrass theorem there is a subsequence of the sequence u^k which converges to a vector, u^∞ say, of unit norm. Discarding framework points and relabelling we may assume this holds for the original sequence. The limit flex of the degenerate framework (G', p^∞) has the form

$$u^\infty = (u_0^\infty, u_1, u_2, \dots, u_n),$$

while the degenerate framework vector is

$$p^\infty = (p_0^\infty, p_1, p_2, p_3, \dots, p_n),$$

where $p_0^\infty = p_2$.

We claim that the velocities u_1, u_2 give an infinitesimal flex of p_1p_2 (as a single edge framework on \mathcal{M}). To see this note that in view of the well-behaved convergence of p_0^k to p_2 (in the a direction) it follows that the velocities u_2 and u_0^∞ have the same component in the a direction, and so $(u_2 - u_0^\infty) \cdot a = 0$. Since $u_2 - u_0^\infty$ is tangential to \mathcal{M} it follows from the choice of a that $u_2 - u_0^\infty$ is orthogonal to $p_2 - p_1$. On the other hand $u_1 - u_0^\infty$ is orthogonal to $p_2 - p_1$ and so taking differences $u_2 - u_1$ is orthogonal to $p_2 - p_1$ as desired.

It now follows, by the rigidity of (G, p) that the velocity vector

$$u_{\text{res}}^\infty = (u_1, u_2, \dots, u_n),$$

is a rigid motion flex. However u_0^∞ is determined by u_1 and u_3 , since the bars for the pairs p_0^∞, p_1 and p_0^∞, p_3 are present. Thus u_0^∞ agrees with u_2 and u_0^∞ is a rigid motion flex for (G', p^∞) . This is a contradiction since the flex has unit norm and is orthogonal to the rigid motion flexes. \square




FIGURE 5. Geometry for the sequential specialisation proof of rigidity preservation under the Henneberg 2 move.

5. VERTEX SURGERY MOVES ON FRAMEWORKS

Let \mathcal{M} be an irreducible surface of type 2, 1 or 0. We now show the preservation of independence of G on \mathcal{M} under the vertex splitting move, the vertex-to- K_4 move and the vertex-to-4-cycle move. The form of argument in the proofs is in terms of flexibility preservation for the inverse move and this is obtained once again by the consideration of certain well-behaved contraction sequences $(G', p^k) \rightarrow (G', p^\infty)$.

For completeness and comparison we include Lemma 5.1 although it will not be needed for the proof of the main theorem.

5.1. Vertex Splitting. Recall that in a vertex-splitting move $G \rightarrow G'$ a vertex v is doubled to v_1 and v_2 with a doubling of an edge vu to v_1u, v_2u , with the remaining edges to v distributed arbitrarily between v_1 and v_2 . The reverse move $G' \rightarrow G$ is an admissible triangle contraction in the sense that the result of collapsing an edge (of a triangle) to coincident endpoints results in a simple $(2, k)$ -tight graph.

The preservation of independence under vertex splitting for bar-joint frameworks in various dimensions was shown by Whiteley [21]. The arguments there were based on self-stresses and k -frames. Here we argue somewhat more directly in terms of infinitesimal flexes.

Lemma 5.1. *Let \mathcal{M} be an irreducible surface of type 2, 1 or 0 and let $G \rightarrow G'$ be a vertex splitting move. If G is independent on \mathcal{M} then G' is independent on \mathcal{M} .*

Proof. To fix notation, let G have $n - 1$ vertices v, v_3, v_4, \dots, v_n with vertex v to be split and v_3 featuring as the vertex u in the vertex-splitting move. Let p, p' be generic vectors for G, G' respectively, with $p = (p(v), p(v_3), \dots, p(v_n)) = (p_1, p_3, \dots, p_n)$ and $p' = (p(v_1)', p(v_2)', \dots, p(v_n)') = (p_1, p_2, \dots, p_n)$.

It will be sufficient to show that if G' is dependent on \mathcal{M} then so too is G . Accordingly we assume that (G', p') is a generic framework which is infinitesimally flexible. Let p_2^k be a sequence of generic points on \mathcal{M} which converges to p_1 in the following well-behaved manner, namely the unit vectors a_k in the directions $p_2^k - p_1$ converge to a unit vector a with $a \cdot (p_3 - p_1) \neq 0$. Also, let $p^k = (p_1, p_2^k, p_3, \dots, p_n)$ and let $p^\infty = (p_1, p_1, p_3, \dots, p_n)$. By the assumption, for each k there exists a unit vector $u^k = (u_1^k, \dots, u_n^k)$ in \mathcal{J}_{p^k} which is an infinitesimal flex of (G', p^k) and which has no rigid motion flex component, in the sense that $Q_{p^k} u^k = 0$. Furthermore, taking subsequences, we may assume that u^k converges to some unit norm flex $u' = (u_1, \dots, u_n)$ as k tends to infinity. By the flex condition we have

$$u_1 \cdot (p_1 - p_3) = u_3 \cdot (p_1 - p_3)$$

and

$$u_2 \cdot (p_1 - p_3) = \lim_{k \rightarrow \infty} u_2^k \cdot (p_2^k - p_3) = \lim_{k \rightarrow \infty} u_3^k \cdot (p_2^k - p_3) = u_3 \cdot (p_1 - p_3).$$

On the other hand, since a_k is a scalar multiple of $p_2^k - p_1$,

$$(u_2 - u_1) \cdot a = \lim_{k \rightarrow \infty} (u_2^k - u_1^k) \cdot a_k = \lim_{k \rightarrow \infty} (u_2^k - u_1^k) \cdot (p_2^k - p_1) = 0.$$

Thus $u_1 = u_2$ since both vectors lie in the same tangent plane and their difference is orthogonal to the linearly independent vectors a and $p_1 - p_2$.

Note that there is always a natural injective map

$$\iota : \mathcal{F}(G, p) \rightarrow \mathcal{F}(G', p^\infty),$$

provided by the map $(w_1, w_3, w_4, \dots, w_n) \rightarrow (w_1, w_1, w_3, w_4, \dots, w_n)$. Also a flex of (G', p^∞) of the form $(w_1, w_1, w_3, w_4, \dots, w_n)$ determines the flex $(w_1, w_3, w_4, \dots, w_n)$ of (G, p) . Thus the flex u' gives rise to a flex, u say, of (G, p) . From the continuity of $q \rightarrow Q_q$ it follows that $Q_{p^\infty} u' = 0$. Thus $Q_p u = 0$ and so u is a proper flex and G is dependent, as required. \square

5.2. Vertex-to- K_4 move. We now show preservation of independence under the vertex-to- K_4 move $G \rightarrow G'$. Once again we will argue in terms of a *well-behaved contraction* $(G', p^k) \rightarrow (G', p^\infty)$, this time with respect to a particular sequence of subframeworks $(K_4, (p_1, p_2^k, p_3^k, p_4^k))$ with $p_i^k \rightarrow p_i$ as $k \rightarrow \infty$, for each $i = 2, 3, 4$.

First we consider separately the case of a cylinder since here we may exploit the two trivial infinitesimal motions to obtain a simple direct proof.

Lemma 5.2. *Let \mathcal{M} be the cylinder, let G' be $(2, 2)$ -sparse with proper subgraph K_4 and let $p = (p_1, \dots, p_n)$ be a framework vector in \mathcal{M}^n for G' , with p_1, \dots, p_4 the placement of the vertices for K_4 . Let*

$$p^k = (p_1, p_2^k, p_3^k, p_4^k, p_5, \dots, p_n)$$

be a sequence in \mathcal{M}^n with $p_i^k \rightarrow p_i$ as $k \rightarrow \infty$, for $i = 2, 3, 4$, and suppose that

- (i) each framework $(K_4, (p_1, p_2^k, p_3^k, p_4^k))$ has 2 independent infinitesimal flexes on \mathcal{M} ,
 - (ii) the dimension of the infinitesimal flex space of (G', p^k) is greater than 2 for each k .
- Then there is a unit norm infinitesimal flex u' of (G', p^∞) of the form

$$u' = (0, 0, 0, 0, u_5, \dots, u_n).$$

Proof. In view of (ii) for each k there exists an infinitesimal flex $u^k = (u_1^k, \dots, u_n^k)$ of (G, p^k) in \mathbb{R}^{3n} with norm $\|u^k\| = 1$, and with $u_1^k = 0$ for all k . Thus, from (i) it follows that $u_i^k = 0$ for $i = 2, 3, 4$. Taking subsequences we may assume that u^k converges to a unit norm flex u' of (G', p^∞) and this has the desired form. \square

Corollary 5.3. *The vertex-to- K_4 move on the cylinder preserves generic infinitesimal rigidity.*

Proof. Let $G \rightarrow G'$ be a vertex-to- K_4 move. Suppose that G' is dependent. Let p^k be a sequence of generic framework vectors for G' as specified in Lemma 5.2, with $p_i^k \rightarrow p_i$ for $i = 1, \dots, 4$. Since p^k is generic (i) holds, and (ii) holds since G' is dependent. Thus, by the lemma (G', p^∞) has an infinitesimal flex of the indicated form. This in turn gives an infinitesimal flex u of (G, p) with $u_1 = 0$ and so G is dependent, as required. \square

We now consider the much more subtle case of irreducible surfaces which are of type 1 or 0. In this case we exploit the characteristic property that at a generic point p_1 on \mathcal{M} the principal curvatures are distinct.

Let $(K_4, (p_1, \dots, p_4))$ be a bar-joint framework on the irreducible surface \mathcal{M} and assume that the principal curvatures κ_s, κ_t at p_1 are well-defined, with $\kappa_s \neq \kappa_t$, and that \hat{s} and \hat{t} are associated orthonormal vectors in the tangent plane at p_1 . For definiteness let \hat{n} be the unit normal at p_1 with $\hat{s}, \hat{t}, \hat{n}$ a right-handed orthonormal triple. By Taylor's theorem, in a neighbourhood of p_1 the points p on \mathcal{M} take the form

$$p(s, t) = p_1 + (s\hat{s} + t\hat{t}) + 1/2(\kappa_s s^2 + \kappa_t t^2)\hat{n} + r(s, t) \quad (5.1)$$

with $\|r(s, t)\| = O(\|(s, t)\|^3)$, for $\|(s, t)\| \leq R$, say. In particular

$$\frac{dp}{ds} = \hat{s} + \kappa_s s \hat{n} + r_s, \quad \frac{dp}{dt} = \hat{t} + \kappa_t t \hat{n} + r_t$$

where $\|r_s\|$ and $\|r_t\|$ are of order $\|(s, t)\|^2$. Also the vectors

$$n(s, t) = dp/ds(s, t) \times dp/dt(s, t) \quad (5.2)$$

give a continuous choice of normal vectors in a neighbourhood of p_1 , with

$$\begin{aligned} n(s, t) &= \hat{s} \times \hat{t} + \kappa_t t \hat{s} \times \hat{n} + \kappa_s s \hat{n} \times \hat{t} + \underline{r} \\ &= \hat{n} - (\kappa_t t \hat{t} + \kappa_s s \hat{s}) + \underline{r} \end{aligned}$$

where $\|\underline{r}\|$ is of order $\|(s, t)\|^2$ in this neighbourhood.

Assume now that the triple p_2, p_3, p_4 is generic and lies in this neighbourhood with $p_i = p(s_i, t_i)$. For $\epsilon \leq 1$ let $p_i^\epsilon = p(\epsilon s_i, \epsilon t_i)$. By a *well-behaved K_4 contraction* of (G, p) over

the subgraph $K_4 \subseteq G$, with vertices v_1, \dots, v_4 , we mean a framework sequence (G, p^k) with

$$p^k = (p_1, p_2^{\epsilon_k}, p_3^{\epsilon_k}, p_4^{\epsilon_k}, p_5, \dots, p_n),$$

where $\epsilon_k \rightarrow 0$ as $k \rightarrow \infty$ and where the local coordinates $s_2, t_2, s_3, t_3, s_4, t_4$ satisfy the determinant condition

$$\begin{vmatrix} s_2 & t_2 & s_2 t_2 \\ s_3 & t_3 & s_3 t_3 \\ s_4 & t_4 & s_4 t_4 \end{vmatrix} \neq 0.$$

It is straightforward to see that we can choose a well-behaved K_4 contraction. For example if $s_i = i, t_i = i^2$ for $i = 2, 3, 4$ then the determinant has the value 48.

Lemma 5.4. *Let \mathcal{M} be an irreducible surface of type 1 or 0. Let $(K_4, (p_1, p_2^k, p_3^k, p_4^k)), k = 1, 2, \dots$, be a well-behaved contraction of frameworks on \mathcal{M} and let $u_k, k = 1, 2, \dots$, be an associated sequence of infinitesimal flexes which forms a convergent sequence in \mathbb{R}^{12} . Then the limit vector has the form (u_1, u_1, u_1, u_1) .*

Proof. Let $u = (u_1, \dots, u_4)$ be an infinitesimal flex of (K_4, p) . Equivalently, $u_i \cdot n_i = 0$ where n_i is the unit normal at p_i and $(p_i - p_j) \cdot (u_i - u_j) = 0$ for $1 \leq i < j \leq 4$. Since (K_4, p) is infinitesimally rigid in \mathbb{R}^3 the flex u is equal to $u_a + u_b$ where u_b is determined by translation by the vector b and where u_a corresponds to an infinitesimal rotation about a line through p_1 with direction vector a . Thus $u_1 = b$ and we may choose the magnitude and direction of a so that $u_i - u_1 = (p_i - p_1) \times a$, for $i = 2, 3, 4$. Substituting gives

$$(a \times (p_i - p_1)) \cdot n_i = u_1 \cdot n_i,$$

or equivalently,

$$a \cdot (n_i \times (p_i - p_1)) + b \cdot n_i = 0,$$

for $i = 2, 3, 4$.

We have the normal vectors $n(s, t) = dp/ds(s_i, t_i) \times dp/dt(s_i, t_i)$ as in Equation 5.2 above. At the point $p_i^\epsilon = p(\epsilon s_i, \epsilon t_i)$ these normals take the form

$$n_i^\epsilon = n(\epsilon s_i, \epsilon t_i) = \hat{n} - \epsilon(\kappa_t t_i \hat{t} + \kappa_s s_i \hat{s}) + \underline{r}_i^\epsilon$$

where $\|\underline{r}_i^\epsilon\| = O(\epsilon^2)$.

Consider now an infinitesimal flex u^ϵ of the framework (K_4, p^ϵ) on \mathcal{M} . The associated equations are

$$a^\epsilon \cdot (n_i^\epsilon \times (p_i^\epsilon - p_1)) + b^\epsilon \cdot n_i^\epsilon = 0, \tag{5.3}$$

for $i = 2, 3, 4$, and we may identify the crossed product here as

$$n_i^\epsilon \times (p_i^\epsilon - p_1) = (\hat{n} - \epsilon(\kappa_t t_i \hat{t} + \kappa_s s_i \hat{s})) \times (\epsilon(s_i \hat{s} + t_i \hat{t}) + 1/2\epsilon^2(\kappa_s s_i^2 + \kappa_t t_i^2)\hat{n}) + R_i^\epsilon$$

with $\|R_i^\epsilon\| = O(\epsilon^3)$.

We may assume, by passing to a subsequence, that ϵ runs through a sequence ϵ_k tending to zero and that the associated unit norm flexes u^ϵ converge to a limit flex u^0 of the degenerate framework $(K_4, (p_1, p_1, p_1, p_1))$ on \mathcal{M} . Let $b^0 = u_1^0$ and let b^ϵ and a^ϵ be the associated vectors. While $b^\epsilon = u_1^\epsilon$ converges to b^0 , as $\epsilon = \epsilon_k \rightarrow 0$, the sequence (a^{ϵ_k}) may be unbounded. However, in view of the three equations

$$u_i^\epsilon - u_1^\epsilon = (p_i^\epsilon - p_1) \times a^\epsilon$$

and the definition of p_i^ϵ it follows that $\|a^{\epsilon k}\|$ is at worst of order $1/\epsilon_k$. We shall show that $\|a^{\epsilon k}\|$ is in fact bounded and so, from the equation above, the desired conclusion follows.

Returning to the three equations, see Equation (5.3), which determine a^ϵ from b^ϵ we have

$$a^\epsilon \cdot (\epsilon s_i \hat{t} - \epsilon t_i \hat{s} - \epsilon^2 s_i t_i (\kappa_s - \kappa_t) \hat{n} + R_i^\epsilon) - \kappa_s (b^\epsilon \cdot \hat{s}) \epsilon s_i - \kappa_t (b^\epsilon \cdot \hat{t}) \epsilon t_i + r_i^\epsilon = 0,$$

where $r_i^\epsilon = \|b^\epsilon \cdot r_i^\epsilon\| = O(\epsilon^2)$. Note that $\|a^\epsilon \cdot R_i^\epsilon\| = O(\epsilon^2)$ and so it follows, introducing coordinates for a^ϵ , and cancelling a factor of ϵ , that

$$(a_s^\epsilon \hat{s} + a_t^\epsilon \hat{t} + a_n^\epsilon \hat{n}) \cdot (s_i \hat{t} - t_i \hat{s} - \epsilon s_i t_i (\kappa_s - \kappa_t) \hat{n}) - \kappa_s (b^\epsilon \cdot \hat{s}) s_i - \kappa_t (b^\epsilon \cdot \hat{t}) t_i = O(\epsilon),$$

for $i = 2, 3, 4$. Thus

$$-a_s^\epsilon t_i + a_t^\epsilon s_i - a_n^\epsilon \epsilon s_i t_i (\kappa_s - \kappa_t) = d_i^\epsilon, \quad \text{for } i = 2, 3, 4,$$

where

$$d_i^\epsilon = b^\epsilon \cdot (\kappa_s s_i \hat{s} + \kappa_t t_i \hat{t}) + X_i^\epsilon,$$

with $X_i^\epsilon = O(\epsilon)$.

Let $\eta = \epsilon(\kappa_s - \kappa_t)$ for $i = 2, 3, 4$, let A_ϵ be the matrix

$$\begin{bmatrix} -t_2 & s_2 & -s_2 t_2 \eta \\ -t_3 & s_3 & -s_3 t_3 \eta \\ -t_4 & s_4 & -s_4 t_4 \eta \end{bmatrix},$$

and note that $\det A_\epsilon = C\epsilon$ for some nonzero constant C . By Cramer's rule we have

$$a_n^\epsilon = (\det A_\epsilon)^{-1} \begin{vmatrix} -t_2 & s_2 & d_2^\epsilon \\ -t_3 & s_3 & d_3^\epsilon \\ -t_4 & s_4 & d_4^\epsilon \end{vmatrix} = (\det A_\epsilon)^{-1} \begin{vmatrix} -t_2 & s_2 & X_2^\epsilon \\ -t_3 & s_3 & X_3^\epsilon \\ -t_4 & s_4 & X_4^\epsilon \end{vmatrix},$$

since the column for $d_i^\epsilon - X_i^\epsilon$ is a linear combination of the first two columns. It follows that the sequence $a_n^{\epsilon k}$ is bounded.

The boundedness of $(a_s^{\epsilon k})$, and similarly $(a_t^{\epsilon k})$, follows more readily, since

$$a_s^\epsilon = (\det A_\epsilon)^{-1} \begin{vmatrix} d_2^\epsilon & s_2 & -s_2 t_2 \eta \\ d_3^\epsilon & s_3 & -s_3 t_3 \eta \\ d_4^\epsilon & s_4 & -s_4 t_4 \eta \end{vmatrix}$$

and the ϵ factors cancel. Thus, the sequence of vectors $a^{\epsilon k}$ is bounded, as desired. \square

Lemma 5.5. *Let \mathcal{M} be an irreducible surface of type 1 or 0, let G' be $(2, 1)$ -sparse with v_1, \dots, v_4 inducing a K_4 subgraph, and let $p = (p_1, \dots, p_n)$ be a generic framework vector in \mathcal{M}^n . Let*

$$p^k = (p_1, p_2^k, p_3^k, p_4^k, p_5, \dots, p_n)$$

be sequence in \mathcal{M}^n , with $p_i^k \rightarrow p_i$ as $k \rightarrow \infty$, for $i = 2, 3, 4$, such that (G, p^k) is a well-behaved contraction with limit (G', p^∞) . If the rigid infinitesimal motion spaces \mathcal{R}_{p^k} and \mathcal{R}_{p^∞} are one-dimensional and the dimension of the flex space $\mathcal{F}(G', p^k)$ is greater than 1 for all k , then there is a unit norm flex u in $\mathcal{F}(G', p^\infty)$ which is orthogonal to \mathcal{R}_{p^∞} and satisfies $u_1 = u_2 = u_3 = u_4$.

Proof. By the hypotheses for each k there exists an infinitesimal flex $u^k = (u_1^k, \dots, u_n^k)$ of (G, p^k) lying in the multiple tangent space \mathcal{T}_{p^k} such that the Euclidean norm of u^k is unity and u^k is orthogonal to the subspace \mathcal{R}_{p^k} . Taking a subsequence if necessary we may assume that u^k converges to u as $k \rightarrow \infty$. By Lemma 5.4 the velocities u_1, \dots, u_n agree. By the hypotheses, the orthogonal projections Q_k onto \mathcal{R}_{p^k} converge to the projection Q_∞ onto \mathcal{R}_{p^∞} and so u is orthogonal to \mathcal{R}_{p^∞} , as desired. \square

Corollary 5.6. *The vertex-to- K_4 move for an irreducible surface of type 1 or 0 preserves generic infinitesimal rigidity.*

Proof. This follows from the previous lemma in the same manner as the proof of Corollary 5.3. \square

5.3. The vertex-to-4-cycle move.

Lemma 5.7. *Let \mathcal{M} be an irreducible surface of type k and let $G \rightarrow G'$ be a vertex-to-4-cycle move. If G is minimally infinitesimally rigid on \mathcal{M} then G' is minimally infinitesimally rigid on \mathcal{M} .*

Proof. Once again we use a sequential contraction argument. Let G have n vertices v_1, v_2, \dots, v_n and edges v_1v_2, v_1v_3 and let $G \rightarrow G'$ be the move in question, with new vertex v_0 and edges v_0v_2, v_0v_3 . It will be sufficient to show that if G' is dependent on \mathcal{M} then so too is G .

Let p, p' be the generic framework vectors for G, G' respectively, with $p' = (p_0, p_1, \dots, p_n)$. Also let $p^k = (p_0^k, p_1, \dots, p_n)$ be generic, with p_0^k converging p_0 . By the assumption for each k there exists a unit vector $u^k = (u_0^k, u_1^k, \dots, u_n^k)$ in the joint tangent space \mathcal{T}_{p^k} which is an infinitesimal flex of (G', p^k) and which is orthogonal to the rigid motion flexes. In earlier notation, $Q_{p^k}u^k = 0$. Taking subsequences, we may assume that u^k converges to some unit norm flex $u' = (u_0, u_1, \dots, u_n)$ of the degenerate framework (G', p^∞) , as k tends to infinity, where $p^\infty = (p_1, p_1, p_2, \dots, p_n)$. Also, by the assumption on G , this degenerate framework (for G') has a space of rigid motion flexes which is naturally identifiable with the space of rigid motion flexes of (G, p) . It remains to show that $u_0 = u_1$ so that we may conclude that (u_1, \dots, u_n) is a proper flex of (G, p) , completing the proof.

It follows from the flex conditions and taking limits that $u_0 - u_2$ is orthogonal to $p_1 - p_2$, and $u_0 - u_3$ is orthogonal to $p_1 - p_3$. Also $u_1 - u_2$ is orthogonal to $p_1 - p_2$, and $u_1 - u_3$ is orthogonal to $p_1 - p_3$. It follows, subtracting, that $u_0 - u_1$ is orthogonal to $p_1 - p_2$ and to $p_2 - p_3$. At the same time $u_0 - u_1$ lies in the tangent plane at p_1 and we may choose p_2, p_3 so that 0 is the only tangent vector orthogonal to $p_1 - p_2$ and to $p_2 - p_3$. \square

6. THE ALGEBRAIC APPROACH

We now give a direct algebraic proof of the preservation of infinitesimal rigidity under the Henneberg 2 move on an irreducible surface. We expect this approach to be more widely useful in the analysis of bar-joint frameworks in higher dimensions.

Assume that \mathcal{M} is an irreducible surface of type k which is defined by the irreducible polynomial $m(x, y, z) = 0$ where the coefficients of m are in \mathbb{Q} . Suppose that G is a $(2, k)$ -tight graph and that (G, p) is a generic framework on \mathcal{M} with $p = (p_1, \dots, p_n)$. Also let

$p^+ = (p, p_v)$ where (G^+, p^+) is generic on \mathcal{M} and G^+ derives from G through a Henneberg 2 move. We write v_1v_2 for the edge involved in the Henneberg move and v for the new vertex.

Since G is independent the rigidity matrix $R_{\mathcal{M}}(G \setminus v_1v_2, p)$ has a flex vector $u = (u_1, \dots, u_n)$ in the nullspace which is not a flex of (G, p) . In particular $(p_1 - p_2) \cdot (u_1 - u_2) \neq 0$. Moreover we may choose u as a solution of the equations $R_{\mathcal{M}}(G, p)u = A$ where A is a column vector with all entries zero except for an entry of unity in the row representing the edge v_1v_2 . This gives a set of linear equations with coefficients in $\mathbb{Q}(p)$ and we can select a solution for which all coordinates of the velocities u_i lie in $\mathbb{Q}(p)$.

We show first that u does not extend to a flex of (G^+, p^+) .

Suppose by way of contradiction that $u^+ = (u, u_v)$ is an extension of u to a flex of (G^+, p^+) with component u_v acting at p_v . Introducing the notation $p_{i,j} = p_i - p_j$, $p_{v,i} = p_v - p_i$ and similarly $u_{i,j} = u_i - u_j$, $u_{v,i} = u_v - u_i$ the flex u_v satisfies four equations

$$p_{v,i} \cdot u_{v,i} = 0, \quad 1 \leq i \leq 3, \quad u_v \cdot N(p_v) = 0,$$

where $N(p_v)$ is the normal to the surface \mathcal{M} at p_v given by

$$N(p_v) = (\nabla m)(p_v) = (\partial m / \partial x, \partial m / \partial y, \partial m / \partial z)|_{p_v}.$$

Introducing the coordinate notation $(p_{v,1}^x, p_{v,1}^y, p_{v,1}^z)$ for $p_{v,1}$ these four equations for the three components of $u_{v,1}$ have a consistent solution if and only if $\det(D) = 0$, where

$$D = \begin{bmatrix} p_{v,1}^x & p_{v,1}^y & p_{v,1}^z & 0 \\ p_{2,1}^x & p_{2,1}^y & p_{2,1}^z & -u_{2,1} \cdot p_{v,2} \\ p_{3,1}^x & p_{3,1}^y & p_{3,1}^z & -u_{3,1} \cdot p_{v,3} \\ N(p_v)^x & N(p_v)^y & N(p_v)^z & u_1 \cdot N(p_v) \end{bmatrix}.$$

Let $P_v = (x, y, z)$ be the vector of indeterminates corresponding to p_v , let $P_{v,i} = P_v - p_i$, $i = 1, 2, 3$, and let

$$D(P_v) = D(x, y, z) = \begin{bmatrix} P_{v,1}^x & P_{v,1}^y & P_{v,1}^z & 0 \\ P_{2,1}^x & P_{2,1}^y & P_{2,1}^z & -u_{2,1} \cdot p_{v,2} \\ P_{3,1}^x & P_{3,1}^y & P_{3,1}^z & -u_{3,1} \cdot p_{v,3} \\ N(P_v)^x & N(P_v)^y & N(P_v)^z & u_1 \cdot N(P_v) \end{bmatrix}.$$

Then the polynomial $\det(D(P_v))$ lies in the ring $\mathbb{Q}(p)[P_v]$. Since

$$0 = \det(D) = \det(D(P_v))|_{p_v}$$

the polynomial $\det(D(P_v))$ evaluates to zero under the substitution $P_v = p_v$. Since p_v is generic on \mathcal{M} this implies that $\det(D(P_v))$ is in the ideal of $\mathbb{Q}(p)[P_v]$ generated by the surface polynomial $m(x, y, z)$. Thus $\det(D(P_v)) = h(P_v)m(P_v)$ for some polynomial $h(P_v)$ in $\mathbb{Q}(p)[P_v]$.

Since $\det(D(P_v)) = h(P_v)m(P_v)$ and $\nabla(m(P_v)) = N(P_v)$ we have

$$\nabla(\det(D(P_v))) = h(P_v)N(P_v) + \nabla(h(P_v))m(P_v)$$

and so

$$\nabla(\det(D(P_v)))|_{p'_v} = h(p'_v)N(p'_v)$$

for any point p'_v satisfying $m(p'_v) = 0$. This implies $a \cdot \nabla(\det(D(P_v)))|_{p'_v} = 0$ for any $a \in \mathbb{R}^3$ satisfying $a \cdot N(p'_v) = 0$ and any point p'_v satisfying $m(p'_v) = 0$. We consider $p'_v = p_1$ which satisfies this property.

We have $u_1 \cdot N(p_1) = 0$. Also, since the first row of the matrix $D(P_v)|_{p_1}$ is zero we get a non-zero contribution to $\nabla(\det(D(P_v)))|_{p_1}$ only from the action of the ∇ operator on the first row of $D(P_v)$. Thus, in vector form, $\nabla(\det(D(P_v)))|_{p_1}$ is the determinant of the matrix

$$\begin{bmatrix} i & j & k & 0 \\ p_{2,1}^x & p_{2,1}^y & p_{2,1}^z & -u_{2,1} \cdot p_{1,2} \\ p_{3,1}^x & p_{3,1}^y & p_{3,1}^z & -u_{3,1} \cdot p_{1,3} \\ N(p_1)^x & N(p_1)^y & N(p_1)^z & u_1 \cdot N(p_1) \end{bmatrix}.$$

Expanding the determinant along the final column gives

$$\nabla(\det(D(P_v)))|_{p_1} = ((p_{2,1} \cdot u_{2,1})p_{3,1} \times N(p_1) - (p_{3,1} \cdot u_{3,1})p_{2,1} \times N(p_1))$$

and so from the above

$$a \cdot ((p_{2,1} \cdot u_{2,1})p_{3,1} \times N(p_1) - (p_{3,1} \cdot u_{3,1})p_{2,1} \times N(p_1)) = 0$$

for all a with the property that $a \cdot N(p_1) = 0$.

The vector $a = N(p_1) \times (p_{2,1} \times N(p_1))$ satisfies $a \cdot N(p_1) = 0$ and $a \cdot \nabla(\det(D(P_v)))|_{p_1} = 0$ gives the condition $(p_{2,1} \cdot u_{2,1})b = 0$ where

$$b = (N(p_1) \times (p_{2,1} \times N(p_1))) \cdot (p_{3,1} \times N(p_1)) = N(p_1) \cdot (p_{3,1} \times p_{2,1}).$$

We have $b \neq 0$ because the condition that $N(p_1) \cdot (p_{3,1} \times p_{2,1}) = 0$ for all p_2, p_3 on \mathcal{M} contradicts the smoothness requirement that $p_{3,1} \times p_{2,1}$ becomes parallel to $N(p_1)$ for p_2 and p_3 close to p_1 . Thus $p_{2,1} \cdot u_{2,1} = 0$ which is contrary to our original choice of u and so we conclude that u does not extend to a flex of (G^+, p^+) .

On the other hand, suppose that a flex $u = (u_1, \dots, u_n)$ of $(G \setminus v_1 v_2, p)$ on \mathcal{M} does extend to a flex (u, u_v) of (G^+, p^+) on \mathcal{M} . Then u_v is the solution of the three equations

$$u_v \cdot (p_v - p_1) = u_1 \cdot (p_v - p_1), u_v \cdot (p_v - p_2) = u_2 \cdot (p_v - p_2) \text{ and } u_v \cdot N(p_v) = 0$$

and the solution is unique because $(p_v - p_1) \times (p_v - p_2) \cdot N(p_v) \neq 0$ for generic p_v, p_1, p_2 for the same reason given above that $b \neq 0$. Also if u is zero then (u, u_v) is zero and so every flex in the nullspace of $R_M(G^+, p^+)$ is the extension of a flex of $R_M(G \setminus v_1 v_2, p)$.

Finally, consider the matrix $R' = R_M(G \setminus v_1 v_2, p)$ of size $m' \times n'$. Since G is independent we have $m' + 1 \leq n'$ and $\text{rank}(R') = m'$. For the matrix $R = R_M(G^+, p^+)$ of size $m \times n$ we have $m = m' + 4$ and $n = n' + 3$ and so $m \leq n$ and $n - m = n' - m' - 1$. Every flex of R' either does not extend to a flex of R or extends to a unique flex of R and every flex of R is the extension of some flex of R' . Thus $|\text{null}(R')| > |\text{null}(R)|$. By Lemma 6.1 the rank of R is m which means that G^+ is independent on \mathcal{M} , as required.

Lemma 6.1. *Let R be an $m \times n$ matrix with $m \leq n$ and R' an $m' \times n'$ matrix with $m' \leq n'$ and $\text{rank}(R') = m'$. If $n - m = n' - m' - 1$ and $|\text{null}(R)| < |\text{null}(R')|$ then $\text{rank}(R) = m$.*

Proof. $\text{rank}(R) = n - |\text{null}(R)| \geq n - |\text{null}(R')| + 1$ and $|\text{null}(R')| = n' - m'$ so $\text{rank}(R) \geq m$. \square

7. MINIMAL RIGIDITY ON TYPE 1 SURFACES

There is a final independence preserving move that we need for the proof of the main result. Recall that if G and H are graphs with vertices $g \in G, h \in H$ then the *edge joining move* combines G and H by adding the edge gh .

Lemma 7.1. *Let \mathcal{M} be an irreducible surface of type 1. Let (G, p) and (H, q) be minimally infinitesimally rigid on \mathcal{M} , and let G' be an edge join of G and H through an edge gh . If $|V(G)|$ and $|V(H)|$ are greater than 4 and (p_g, p_h) is generic on \mathcal{M} then $(G', (p, q))$ is generically minimally infinitesimally rigid on \mathcal{M} .*

Proof. Consider the block matrix form

$$R_{\mathcal{M}}(G', (p, q)) = \begin{bmatrix} R_{\mathcal{M}}(G, p) & 0 \\ * & * \\ 0 & R_{\mathcal{M}}(H, q) \end{bmatrix}.$$

By the hypotheses the nullspaces of the rigidity matrices $R_{\mathcal{M}}(G, p)$ and $R_{\mathcal{M}}(H, q)$ are one-dimensional. Let $u = (u_p, u_q)$ be an infinitesimal flex of the edge-joined framework and let gh be the joining edge for G' . Subtracting a tangential rigid motion infinitesimal flex we may assume that u_p assigns a zero velocity to the framework joint p_g . Since u_p is a flex of (G, p) on \mathcal{M} it follows that $u_p = 0$. Also in view of the generic row in $R_{\mathcal{M}}(G', (p, q))$ for the joining edge it follows that u_q assigns a velocity to the framework vertex q_h which is linearly independent from the one-dimensional space of velocity vectors of q_h obtained from infinitesimal flexes of (H, q) . It follows that this velocity on q_h is zero and hence that u_q is zero. Thus $u = 0$ and the nullspace of $R_{\mathcal{M}}(G', (p, q))$ has dimension one, as desired. \square

We now arrive at the proof of our main result, Theorem 1.1. For the reader's convenience we first re-state the theorem.

Theorem 7.2. *Let $G = (V, E)$ be a simple graph and let \mathcal{M} be an irreducible surface of type 1. Then a generic framework (G, p) on \mathcal{M} is isostatic if and only if G is K_1, K_2, K_3, K_4 or is $(2, 1)$ -tight.*

Proof. That the underlying graph of an isostatic framework on \mathcal{M} is $(2, 1)$ -tight or is a small complete graph follows from Theorem 2.6. For the sufficiency direction one can check that the minimal graph $K_5 \setminus e$ in the inductive characterisation of $(2, 1)$ -tight graphs is isostatic on \mathcal{M} . The sufficiency of $(2, 1)$ -tightness now follows from Theorem 1.2 if minimal generic rigidity is preserved by Henneberg 1 and 2 moves, the vertex-to- K_4 move, the vertex-to-4-cycle move and the edge joining move. This is the content of Lemma 4.1, Lemma 4.2, Corollary 5.6, Lemma 5.7 and Lemma 7.1. \square

Note that we could also have used Theorem 2.13, applying Lemma 5.1, to prove the theorem.

8. EXTENSIONS

We finish by noting some further natural considerations for frameworks constrained to surfaces.

The assumption, in Theorem 1.1, that \mathcal{M} is irreducible merits two comments. Firstly, it is required to avoid to surfaces composed as unions of surfaces of differing numbers of internal motions. For example if \mathcal{M} was the union of two cylinders with distinct but parallel axes then consideration of Theorem 2.12 instantly shows that simply being $(2, 1)$ -tight is not the correct characterisation. Secondly, if \mathcal{M} is reducible but each component is irreducible then we do not expect any great difficulty in extending our results. For example we expect that Theorem 1.1 is true for concentric cones, torii or elliptical cylinders.

The usual two-dimensional torus embedded in 3-dimensional space has freedom type 1 and so isostatic frameworks on this surface are characterised as in the previous theorem. When the torus is realised in \mathbb{R}^4 one may also consider the *Clifford torus* \mathcal{T} , that is, the real algebraic variety and smooth manifold defined by the polynomial equations $x^2 + y^2 = 1$ and $z^2 + w^2 = 1$. The definition of type (freedom number) given in Definition 2.2 extends without change to an algebraic surface \mathcal{M} in \mathbb{R}^d .

Definition 8.1. *An embedded manifold \mathcal{M} in \mathbb{R}^d is of type k if $\dim \ker R_{\mathcal{M}}(K_n, q) \geq k$ for all frameworks (K_n, p) on \mathcal{M} , for $n = 2, 3, \dots$, and k is the largest such integer.*

In particular the Clifford torus has freedom type 2. The rigidity analysis in this setting requires us to adapt the definition of the rigidity matrix. The details are similar to those in Definition 2.1 with the following changes. There are now 4 columns per vertex and 2 rows per vertex where the rows for vertex i (and corresponding framework point (x_i, y_i, z_i, w_i)) are zero except in the 4-tuple corresponding to i where the entries in the first row are $x_i, y_i, 0, 0$ and the second are $0, 0, z_i, w_i$.

On the other hand take the product of a circle and an ellipse. This is the algebraic variety \mathcal{S} defined by, say, $x^2 + y^2 = 1$ and $z^2 + w^2/2 = 1$ in \mathbb{R}^4 . \mathcal{S} admits exactly 1 trivial motion. Adapting the methods of the last section would lead to the $(2, 2)$ -tight and the $(2, 1)$ -tight characterisations of frameworks on \mathcal{T} and \mathcal{S} respectively.

It is natural to seek a similar characterisation of our main result in the case of frameworks with vertices constrained to an irreducible surface of type 0. There are a variety of such surfaces that a characterisation could apply to including an elliptical cone, a mobius strip, a hyperboloid and a hyperbolic paraboloid. Note that the graphs of rigid frameworks need not be connected in this setting. As a starting point [12, Proposition 3.4] gives the necessity of the graph being simple and $(2, 0)$ -tight and we expect that the rigidity preservation methods in this paper will be useful in deriving a characterisation. However, there are immediate additional complications to establishing sufficient conditions, not least since any simple $(2, 0)$ -tight graph containing a subgraph isomorphic to K_5 has a dependent rigidity matrix on any surface. This can easily be seen by noting that K_5 is not $(3, 6)$ -sparse and hence is dependent in \mathbb{R}^3 . We also remark that $(2, 0)$ -tight graphs may be 4-regular so additional degree 4 operations seem to be necessary (such as X -replacement, see [14, 19]). This fact, together with the fact that an inductive scheme would have to avoid creating K_5 subgraphs, and the fact that there are many simple $(2, 0)$ -tight graphs (even on small vertex sets) that cannot be generated using the operations in this paper, with X -replacement, all suggest that the analogue of Theorem 1.2 for simple $(2, 0)$ -tight graphs, without K_5 subgraphs, will be significantly more challenging to establish.

Acknowledgement. We would like to thank Bill Jackson, for discussions relating to the Henneberg 2 move on manifolds.

REFERENCES

- [1] L. Asimow and B. Roth, The Rigidity of Graphs, *Trans. Amer. Math. Soc.*, 245, (1978), 279-289.
- [2] A. Berg and T. Jordan, A Proof of Connelly's Conjecture on 3-connected Circuits of the Rigidity Matroid, *Journal of Combinatorial Theory, Series B* 88 (2003) 77-97.
- [3] C. S. Borcea and I. Streinu, Periodic Frameworks and Flexibility, *Proc. R. Soc. A* (2010) 466, 2633-2649.
- [4] Z. Fekete and L. Szegő, A Note on $[k, l]$ -sparse Graphs, in *Graph Theory in Paris*, Edited by A. Bondy et al, (2007), 169-177.
- [5] H. Gluck, Almost all Simply Connected Closed Surfaces are Rigid, in *Geometric Topology*, Lecture Notes in Math., no. 438, Springer-Verlag, Berlin, 1975, pp. 225-239.
- [6] L. Henneberg, *Die graphische Statik der starren Systeme*, Leipzig (1911).
- [7] B. Jackson and T. Jordan, Connected Rigidity Matroids and Unique Realisations of Graphs, *J. Combinatorial Theory B*, vol. 94, (2005), 1-29.
- [8] N. Katoh and S. Tanigawa, A Proof of the Molecular Conjecture, *Disc. and Comp. Geom.*, 45(4), (2011), 647-700.
- [9] G. Laman, On Graphs and the Rigidity of Plane Skeletal Structures, *J. Engineering Mathematics*, 4 (1970), 331-340.
- [10] J. Malestein and L. Theran, Generic Combinatorial Rigidity of Periodic Frameworks, *Advances in Math*, 233, 291-331, (2013).
- [11] J. Malestein and L. Theran, Generic Rigidity with Forced Symmetry and Sparse Coloured Graphs, <http://arxiv.org/abs/1203.0772v1>, (2012).
- [12] A. Nixon, J.C. Owen and S.C. Power, Rigidity of Frameworks Supported on Surfaces, *SIAM J. Discrete Math.* 26, 4, 1733-1757, (2012).
- [13] A. Nixon and J.C. Owen, An Inductive Construction of $(2, 1)$ -tight Graphs, <http://arxiv.org/abs/1103.2967v2>, (2012).
- [14] A. Nixon and E. Ross, One Brick at a Time: a Survey of Inductive Constructions in Rigidity Theory, <http://arxiv.org/abs/1203.6623>, to appear in the Proceedings of the Fields Institute, edited by R. Connelly, A. Weiss and W. Whiteley.
- [15] A. Nixon and E. Ross, Periodic Rigidity on a Variable Torus Using Inductive Constructions, <http://arxiv.org/abs/1204.1349>, (2012).
- [16] J. C. Owen and S. C. Power, Infinite Bar-joint Frameworks, Crystals and Operator theory, *New York J. Math.*, 17 (2011) 445-490.
- [17] E. Ross, Inductive Constructions for Frameworks on a two-dimensional Fixed Torus, preprint, (2011).
- [18] T-S. Tay, Henneberg's Method for Bar and Body Frameworks. *Structural Topology* 17, (1991), 53-58.
- [19] T-S. Tay and W. Whiteley, Generating Isostatic Frameworks, *Structural Topology*, 11, (1985), 21-69.
- [20] W. Whiteley, The Union of Matroids and the Rigidity of Frameworks, *SIAM J. Disc. Math.*, 1 (1988), 237-255.
- [21] W. Whiteley, Vertex Splitting in Isostatic Frameworks, *Structural Topology*, 16, (1991), 23-30.
- [22] W. Whiteley, Matroids and rigid structures, in "Matroid Applications" ed. N. White, *Encyclopedia of Mathematics and its applications* 40 (1992), 1-51.

HEILBRONN INSTITUTE FOR MATH. RESEARCH, SCHOOL OF MATH., UNIVERSITY OF BRISTOL, BRISTOL, BS8 1TW, U.K.

E-mail address: tony.nixon@bristol.ac.uk

D-CUBED, SIEMENS PLM SOFTWARE, FRANCIS HOUSE, 112 HILLS ROAD, CAMBRIDGE, CB2 1PH, U.K.

E-mail address: owen.john.ext@siemens.com

DEPT. MATH. STATS., LANCASTER UNIVERSITY, LANCASTER LA1 4YF, U.K.

E-mail address: s.power@lancaster.ac.uk