

A CHAIN CONDITION FOR OPERATORS FROM $C(K)$ -SPACES

KLAAS PIETER HART, TOMASZ KANIA AND TOMASZ KOCHANEK

ABSTRACT. We introduce a chain condition (\mathfrak{A}) , defined for operators acting on $C(K)$ -spaces, which is intermediate between weak compactness and having weakly compactly generated range. It is motivated by Pełczyński's characterisation of weakly compact operators on $C(K)$ -spaces. We prove that if K is extremally disconnected and X is a Banach space then, for an operator $T: C(K) \rightarrow X$, T is weakly compact if and only if T satisfies (\mathfrak{A}) if and only if the representing vector measure of T satisfies an analogous chain condition. As a tool for proving the above-mentioned result, we derive a topological counterpart of Rosenthal's lemma. We exhibit several compact Hausdorff spaces K for which the identity operator on $C(K)$ satisfies (\mathfrak{A}) , for example both locally connected compact spaces having countable cellularity and ladder system spaces have this property. Using a Ramsey-type theorem, due to Dushnik and Miller, we prove that the collection of operators on a $C(K)$ -space satisfying (\mathfrak{A}) forms a closed left ideal of $\mathcal{B}(C(K))$.

1. INTRODUCTION

The aim of this paper is to study a certain chain condition, denoted (\mathfrak{A}) , defined for (bounded, linear) operators $T: C(K) \rightarrow X$, where K is a compact Hausdorff space, X is an arbitrary Banach space and $C(K)$ is the Banach space of all scalar-valued continuous functions on K , equipped with the supremum norm. The main motivation behind our work comes from a theorem of Pełczyński which asserts that an operator $T: C(K) \rightarrow X$ is weakly compact if and only if there is no isomorphic copy of c_0 in $C(K)$ on which T is bounded below. In other words, T fails to be weakly compact provided there is a sequence of pairwise disjoint open sets $\{O_n\}_{n=1}^\infty$ in K and a uniformly bounded family of functions $\{f_n\}_{n=1}^\infty$ in $C(K)$ such that the support of f_n is contained in O_n ($n \in \mathbb{N}$) and $\inf_n \|T(f_n)\| > 0$. Loosely speaking, Pełczyński's theorem characterises weakly compact operators from $C(K)$ -spaces as precisely those which annihilate the sequences of disjointly supported and uniformly bounded functions in $C(K)$. Our condition (\mathfrak{A}) is similar in nature to this requirement.

Let K be a compact Hausdorff space and X be a Banach space. The *support*, $\text{supp}(f)$, of a function $f \in C(K)$ is defined as the closure of the set $\{x \in K: f(x) \neq 0\}$. For any pair of functions $f, g \in C(K)$ we write $f \prec g$ whenever $f \neq g$ and $f(x) = g(x)$ for each $x \in \text{supp}(f)$. The relation \prec is a strict ordering on $C(K)$. For a bounded and linear operator $T: C(K) \rightarrow X$, the chain condition (\mathfrak{A}) is defined as follows:

2010 *Mathematics Subject Classification*. Primary 46B50, 46E15, 37F20; Secondary 54F05, 46B26.

Key words and phrases. Weakly compact operator, chain condition, extremally disconnected space, vector measure, Rosenthal's lemma, partition lemma.

(\clubsuit) for each uncountable \prec -chain F in $C(K)$ with $\sup_{f \in F} \|f\| < \infty$,

$$\inf\{\|Tf - Tg\| : f, g \in F, f \neq g\} = 0.$$

Any uniformly bounded \prec -chain $\{f_i\}_{i \in I}$ such that $\|f_i - f_j\| \geq \delta$ for $i, j \in I$, $i \neq j$, and some positive δ , will be called a δ - \prec -chain. We also say that a compact Hausdorff space K satisfies (\clubsuit) whenever the identity operator on $C(K)$ satisfies (\clubsuit), equivalently, for each $\delta > 0$ there is no uncountable δ - \prec -chain in $C(K)$.

We start Section 2 with the observations that, for each operator $T: C(K) \rightarrow X$, the condition (\clubsuit) is weaker than weak compactness (Proposition 2.1), and that these two properties coincide provided that K is extremally disconnected (i.e. open sets have open closures), in which case weak compactness may also be characterised by a counterpart of condition (\clubsuit) for the representing measure of T (Theorem 2.5). This result is obtained by refining a classical lemma due to Rosenthal to the extremally disconnected setting (Lemma 2.2) and by applying some vector-measure techniques (Proposition 2.4).

In Section 3 we give examples of compact Hausdorff spaces that satisfy (\clubsuit). They include compact Hausdorff spaces which are countable, one-point compactifications of discrete sets (Proposition 3.3), compact Hausdorff spaces which are locally connected and have countable cellularity (Theorem 3.5) and ladder system spaces (Corollary 3.8).

Like other classical chain conditions for topological spaces or Boolean algebras, this one carries some combinatorial insight. Using Ramsey-theoretic techniques we prove that the family of operators on a given $C(K)$ -space satisfying (\clubsuit) forms a closed left ideal of the Banach algebra $\mathcal{B}(C(K))$ of all operators on $C(K)$ (Theorem 4.2).

All Banach spaces are assumed to be over either real or complex scalars. By an *operator* we understand a bounded linear operator acting between Banach spaces. An operator $T: E \rightarrow F$ is *bounded below* whenever there exists $\gamma > 0$ such that $\|Tx\| \geq \gamma\|x\|$ for each $x \in E$; an operator which is bounded below is injective and has closed range. An operator $T: E \rightarrow F$ *fixes a copy* of a Banach space X , if there is a subspace E_0 of E isomorphic to X such that $T|_{E_0}$ is bounded below. The identity operator on a Banach space X is denoted I_X . For a topological space K , we denote its cardinal number by $|K|$, and define the *cellularity* of K , $c(K)$, to be the supremum of the cardinalities of families consisting of pairwise disjoint open sets in K . A topological space with countable cellularity is said to satisfy *c.c.c.* The symbol $\mathbf{1}_A$ denotes the indicator function of a set A .

2. (\clubsuit) AND WEAKLY COMPACT OPERATORS FROM $C(K)$ -SPACES

By a classical result of Pełczyński (see, *e.g.*, [3, Theorem VI.2.15]), every non-weakly compact operator from a $C(K)$ -space into an arbitrary Banach space X fixes a copy of c_0 . In the case where K is extremally disconnected the Goodner–Nachbin theorem ([8], [13]) says that $C(K)$ is (isometrically) injective, and then Rosenthal’s theorem [15, Theorem 1.3] asserts that any non-weakly compact operator $T: C(K) \rightarrow X$ fixes a copy of ℓ_∞ . The main result of this section, Theorem 2.5, is in the spirit of these facts. It asserts that weak compactness of an operator from a $C(K)$ -space, for K extremally disconnected, may be characterised in terms of the chain conditions, imposed both upon the operator itself and its representing measure.

Let us begin with the following refinement of Pełczyński's theorem.

Proposition 2.1. *Let K be a compact Hausdorff space, X be a Banach space and let $T: C(K) \rightarrow X$ be an operator. Then the following assertions are equivalent*

- (i) *T is weakly compact;*
- (ii) *for each infinite \prec -chain $F \subseteq C(K)$,*

$$\inf\{\|Tf - Tg\|: f, g \in F, f \neq g\} = 0.$$

In particular, every weakly compact operator $T: C(K) \rightarrow X$ satisfies (\clubsuit) .

Proof. For the implication (i) \Rightarrow (ii) assume contrapositively that

$$\delta = \inf\{\|Tf - Tg\|: f, g \in F, f \neq g\} > 0.$$

Let $F_0 = (f_n)_{n \in \mathbb{N}}$ be a \prec -monotone subsequence of F . For each $n \in \mathbb{N}$ set $g_n = |f_{n+1} - f_n|$. Then $Y = \overline{\text{span}}\{g_n: n \in \mathbb{N}\}$ is an isomorphic copy of c_0 and $\|Tg_n\| > \delta$ ($n \in \mathbb{N}$). Consequently, T is bounded below on Y . So T cannot be weakly compact.

To prove (ii) \Rightarrow (i) assume, for a contradiction, that T is not weakly compact. Appealing to the proof of [3, Theorem VI.2.15], we may construct a sequence $(f_n)_{n=1}^\infty$ of disjointly supported functions in $C(K)$ of norm at most one such that $\inf_{n \in \mathbb{N}} \|Tf_n\| > 0$. The family $F = \{\sum_{k=1}^n f_k\}_{n=1}^\infty$ is then a uniformly bounded infinite \prec -chain for which $\inf\{\|Tf - Tg\|: f, g \in F, f \neq g\} > 0$. \square

Given a compact Hausdorff space K and a Banach space X , every operator $T: C(K) \rightarrow X$ admits a Riesz-type representation (cf. [3, Chapter 6]) in the following precise sense: there exists a w^* -countably additive vector measure $\mu: \Sigma \rightarrow X^{**}$ (called the *representing measure* for T) on the σ -algebra Σ of all Borel subsets of K such that:

- (i) for each $x^* \in X^*$ the map $\Sigma \ni A \mapsto \mu(A)x^*$ is a regular countably additive scalar measure (and will be denoted $x^* \circ \mu$);
- (ii) the map $x^* \mapsto x^* \circ \mu$ from X^* into $C(K)^*$ is w^* -to- w^* continuous;
- (iii) $x^*T(f) = \int_K f d(x^* \circ \mu)$ for each $x^* \in X^*$ and $f \in C(K)$;
- (iv) $\|T\| = \|\mu\|(K)$.

The representing measure μ may be expressed explicitly by the formula $\mu(A) = T^{**}\varphi_A$, where $\varphi_A \in C(K)^{**}$ acts as $\varphi_A(\nu) = \nu(A)$ ($A \in \Sigma, \nu \in C(K)^*$). Equivalently, it may be defined by the prescription $\mu(A)x^* = \mu_{x^*}(A)$, where $\mu_{x^*} = T^*x^*$ is the scalar measure produced by the Riesz theorem applied to the functional x^*T .

We will need the following topological counterpart of Rosenthal's lemma (cf. [3, Lemma I.4.1]). Although its proof is almost the same as the original one, we present it for completeness and to demonstrate the rôle played by extremal disconnectedness.

Lemma 2.2. *Let K be a compact Hausdorff space which is extremally disconnected, and let $(V_n)_{n=1}^\infty$ be a sequence of pairwise disjoint open subsets of K . Suppose that $(\mu_n)_{n=1}^\infty$ is a sequence of scalar Borel measures on K having uniformly bounded variations. Then, for every $\varepsilon > 0$ there exists a strictly increasing sequence $(n_k)_{k=1}^\infty$ of natural numbers such that,*

for each $k \in \mathbb{N}$,

$$|\mu_{n_k}| \left(\overline{\bigcup_{j \neq k} V_{n_j}} \right) < \varepsilon.$$

Proof. Let us suppose, with no loss of generality, that $|\mu_n|(K) \leq 1$ for each $n \in \mathbb{N}$. Consider any sequence $(M_p)_{p=1}^\infty$ of pairwise disjoint infinite subsets of \mathbb{N} such that $\mathbb{N} = \bigcup_p M_p$. We consider two cases.

Case 1. First, if there is some $p \in \mathbb{N}$ for which

$$|\mu_k| \left(\overline{\bigcup_{\substack{j \in M_p \\ j \neq k}} V_j} \right) < \varepsilon \quad \text{for each } k \in M_p,$$

then we get the assertion from the induced enumeration of M_p : $\{n_1 < n_2 < \dots\}$.

Case 2. Now suppose that for every $p \in \mathbb{N}$ there is $k_p \in M_p$ such that

$$(2.1) \quad |\mu_{k_p}| \left(\overline{\bigcup_{\substack{j \in M_p \\ j \neq k_p}} V_j} \right) \geq \varepsilon.$$

Fix, for a moment, any $p \in \mathbb{N}$. Since $\bigcup_q V_{k_q}$ is disjoint from the open set $\bigcup_{j \in M_p, j \neq k_p} V_j$, so is its closure. Hence,

$$(2.2) \quad \bigcup_{j \in M_p, j \neq k_p} V_j \subset \bigcup_{n \in \mathbb{N}} V_n \setminus \overline{\bigcup_{q \in \mathbb{N}} V_{k_q}}.$$

Observe that $\overline{\bigcup_q V_{k_q}}$ is open, as K is extremally disconnected, whence $\overline{\bigcup_n V_n} \setminus \overline{\bigcup_q V_{k_q}}$ is closed. Therefore, by (2.2), we get

$$(2.3) \quad \overline{\bigcup_{\substack{j \in M_p \\ j \neq k_p}} V_j} \subset \overline{\bigcup_{n \in \mathbb{N}} V_n} \setminus \overline{\bigcup_{q \in \mathbb{N}} V_{k_q}}.$$

Obviously, we have

$$|\mu_{k_p}| \left(\overline{\bigcup_{q \in \mathbb{N}} V_{k_q}} \right) + |\mu_{k_p}| \left(\overline{\bigcup_{n \in \mathbb{N}} V_n} \setminus \overline{\bigcup_{q \in \mathbb{N}} V_{k_q}} \right) \leq 1,$$

so (2.3) and (2.1) imply

$$|\mu_{k_p}| \left(\overline{\bigcup_{q \in \mathbb{N}} V_{k_q}} \right) \leq 1 - \varepsilon,$$

and this inequality is valid for every $p \in \mathbb{N}$.

Consequently, we may repeat the same argument replacing the space K with the clopen subspace $\overline{\bigcup_{q \in \mathbb{N}} V_{k_q}}$ (which is extremally disconnected as this property is inherited by open subspaces) and the sequences $(\mu_n)_{n=1}^\infty$ and $(V_n)_{n=1}^\infty$ with $(\mu_{k_p})_{p=1}^\infty$ and $(V_{k_p})_{p=1}^\infty$, respectively. By continuing this we would get subsequent upper bounds $1 - 2\varepsilon, 1 - 3\varepsilon, \dots$ for some of

the variations $|\mu_n|$. Since this process has to terminate, we will end up with *Case 1*, where the assertion has been proved. \square

Proposition 2.3. *Let K be a compact Hausdorff space which is extremally disconnected, and let X be a Banach space. Every operator from $C(K)$ into X which satisfies (\mathfrak{H}) is weakly compact.*

Proof. It suffices to consider the real space $C_{\mathbb{R}}(K)$ since, we prove (arguing by contraposition) that there is a δ - \prec -chain in $C_{\mathbb{R}}(K)$ which will also do the job in $C_{\mathbb{C}}(K)$.

Assume that $T: C_{\mathbb{R}}(K) \rightarrow X$ is a non-weakly compact operator, and let μ be its representing measure. Then (see the proof of Theorem 5.5.3 in [1]) there exist a number $\varepsilon > 0$, a sequence $(O_n)_{n=1}^{\infty}$ of pairwise disjoint open subsets of K and a sequence $(x_n^*)_{n=1}^{\infty}$ in the unit ball of X^* such that $(x_n^* \circ \mu)(O_n) > \varepsilon$ for each $n \in \mathbb{N}$ (recall that $x_n^* \circ \mu = T^*x_n^*$). Since $x_n^* \circ \mu$ is a regular measure and K is extremally disconnected, we may assume that each O_n is clopen. By Lemma 2.2, we may also assume that

$$|x_n^* \circ \mu| \left(\overline{\bigcup_{j \neq n} O_j} \right) < \frac{\varepsilon}{2} \quad \text{for each } n \in \mathbb{N}.$$

Let $\{q_n: n \in \mathbb{N}\}$ be an enumeration of the rational numbers and define $A_t = \{n: q_n < t\}$ for $t \in \mathbb{R}$. Then $\{A_t: t \in \mathbb{R}\}$ is an increasing chain of infinite subsets of \mathbb{N} . For each t , the set $C_t = \overline{\bigcup_{n \in A_t} O_n}$ is clopen, hence its characteristic function, call it f_t , is continuous. The family $\{C_t: t \in \mathbb{R}\}$ is a strictly increasing chain of clopen sets, so that $s < t$ readily implies $f_s \prec f_t$. Clearly $\|f_t\| = 1$ for all t .

Now, let $s < t$. Then, there is $n \in A_t \setminus A_s$ (in fact the set is infinite). Since

$$\text{supp}(f_t - f_s) \subset \overline{\bigcup_{k \in \mathbb{N}} O_k} = O_n \cup \overline{\bigcup_{k \neq n} O_k}$$

and $f_t - f_s$ equals 1 on O_n , we get

$$\begin{aligned} \|T(f_t) - T(f_s)\| &\geq x_n^* T(f_t - f_s) = \int_K (f_t - f_s) d(x_n^* \circ \mu) \\ &= \left(\int_{O_n} + \int_{\overline{\bigcup_{k \neq n} O_k}} \right) (f_t - f_s) d(x_n^* \circ \mu) \\ &\geq (x_n^* \circ \mu)(O_n) - |x_n^* \circ \mu| \left(\overline{\bigcup_{k \neq n} O_k} \right) \\ &> \varepsilon - \frac{\varepsilon}{2} = \frac{\varepsilon}{2} \quad (i, j \in \mathbb{N}, i \neq j) \end{aligned}$$

which proves that T does not satisfy (\mathfrak{H}) . \square

We now introduce a counterpart of condition (\mathfrak{H}) for vector measures. Namely, for a set algebra Σ , Banach space X , and (finitely) additive vector measure $\mu: \Sigma \rightarrow X$, consider the following property:

(\mathfrak{B}) for each uncountable chain (with respect to inclusion) $\{E_i\}_{i \in I}$ in Σ ,

$$\inf\{\|\mu(E_i) - \mu(E_j)\| : i, j \in I, i \neq j\} = 0.$$

A vector measure $\mu: \Sigma \rightarrow X$ is called *strongly additive* provided that for every sequence $(E_n)_{n=1}^\infty \subset \Sigma$ of pairwise disjoint sets, the series $\sum_{n=1}^\infty \mu(E_n)$ is unconditionally convergent in X .

Proposition 2.4. *Let Σ be a σ -algebra of sets and X be a Banach space. A bounded vector measure $\mu: \Sigma \rightarrow X$ satisfies (\mathfrak{B}) if and only if it is strongly additive.*

Proof. (\Leftarrow) Suppose that μ does not satisfy (\mathfrak{B}). Then there exists a monotone (with respect to inclusion) sequence $(E_n)_{n=1}^\infty \subset \Sigma$ such that $\|\mu(E_m) - \mu(E_n)\| > \delta$ for $m, n \in \mathbb{N}$, $m \neq n$, and for some $\delta > 0$. Put $D_n = E_{n+1} \setminus E_n$ provided $(E_n)_{n=1}^\infty$ is increasing and $D_n = E_n \setminus E_{n+1}$ otherwise. In any case $(D_n)_{n=1}^\infty$ forms a sequence of pairwise disjoint elements from Σ such that $\|\mu(D_n)\| > \delta$ for each $n \in \mathbb{N}$. Consequently, μ is not strongly additive.

(\Rightarrow) Now, suppose that μ fails to be strongly additive. Then the Diestel–Faires theorem (cf. [3, Theorem I.4.2]) produces a closed subspace Y of X and an isomorphism $T: \ell_\infty \rightarrow Y$ such that $T(e_n) = \mu(A_n)$ for some pairwise disjoint sets $(A_n)_{n=1}^\infty \subset \Sigma$. For any $n \in \mathbb{N}$ we have $1 = \|e_n\| \leq \|T^{-1}\| \cdot \|\mu(A_n)\|$, hence for some x_n^* in the unit ball of X^* we have $x_n^* \mu(A_n) \geq \|T^{-1}\|^{-1}$. Since $(x_n^* \mu)_{n=1}^\infty$ is a uniformly bounded sequence of scalar measures, Rosenthal's lemma produces a subsequence $(x_{n_k}^* \mu)_{k=1}^\infty$ such that

$$|x_{n_k}^* \mu\left(\bigcup_{j \neq k} A_{n_j}\right)| < \frac{1}{2} \|T^{-1}\|^{-1} \quad \text{for each } k \in \mathbb{N}.$$

Let \mathcal{C} be an uncountable chain of subsets of \mathbb{N} and for each $C \in \mathcal{C}$ define $E(C) = \bigcup_{j \in C} A_{n_j}$. Plainly, $\{E(C)\}_{C \in \mathcal{C}}$ is an uncountable chain of members of Σ . Moreover, for any $C_1, C_2 \in \mathcal{C}$ with $C_1 \subsetneq C_2$, and any $k \in C_2 \setminus C_1$, we have

$$\begin{aligned} \|\mu(E(C_2)) - \mu(E(C_1))\| &= \|\mu(E(C_2 \setminus C_1))\| = \left\| \mu\left(\bigcup_{j \in C_2 \setminus C_1} A_{n_j}\right) \right\| \\ (2.4) \quad &\geq \left| x_{n_k}^* \mu\left(\bigcup_{j \in C_2 \setminus C_1} A_{n_j}\right) \right| \geq |x_{n_k}^* \mu(A_{n_k})| - |x_{n_k}^* \mu\left(\bigcup_{j \neq k} A_{n_j}\right)| \\ &> \|T^{-1}\|^{-1} - \frac{1}{2} \|T^{-1}\|^{-1} = \frac{1}{2} \|T^{-1}\|^{-1}. \end{aligned}$$

This shows that μ fails to satisfy (\mathfrak{B}), so the proof is complete. \square

We are now prepared to proceed to the main result of this section.

Theorem 2.5. *Let K be a compact Hausdorff space which is extremally disconnected, and X a Banach space, let Σ be the Borel σ -algebra of K and let $T: C(K) \rightarrow X$ be an operator with representing measure $\mu: \Sigma \rightarrow X^{**}$. Then the following assertions are equivalent:*

- (i) T is weakly compact;
- (ii) μ is strongly additive;

- (iii) T satisfies (\mathfrak{H}) ;
- (iv) μ satisfies (\mathfrak{M}) .

Proof. The equivalence of clauses (i) and (ii) is the Bartle–Dunford–Schwartz theorem (cf. [3, Theorem VI.2.5]) and is valid for any compact Hausdorff space K . The equivalence of clauses (i) and (iii) follows from Propositions 2.1 and 2.3. The equivalence of (ii) and (iv) is immediate from Proposition 2.4. \square

It is easily seen that the implication (\Leftarrow) of Proposition 2.4 holds true for any set algebra Σ , not necessarily a σ -algebra. However, the situation is not so clear for the implication (\Rightarrow) , so the natural question arises: how can one characterise the class of set algebras Σ for which every vector measure $\mu: \Sigma \rightarrow X$ satisfying (\mathfrak{M}) is strongly additive? An inspection of the proof of Proposition 2.4 suggests that some kind of interpolation property of Σ could do the job, so we may make our problem more precise. A set algebra Σ has the *subsequential completeness property* (SCP) whenever for every sequence $(E_n)_{n=1}^\infty$ of pairwise disjoint sets from Σ there is a subsequence $(E_{n_k})_{k=1}^\infty$ with $\bigcup_k E_{n_k} \in \Sigma$. (Haydon constructed a set algebra with (SCP) which is not a σ -algebra; see [11, Proposition 1E].)

Question 2.6. Suppose that a set algebra Σ has (SCP). Is it true that, for any Banach space X , every vector measure $\mu: \Sigma \rightarrow X$ satisfying (\mathfrak{M}) is necessarily strongly additive?

Remark 2.7. The second and third-named authors ([12]) have recently studied the operator ideal \mathcal{WCG} of *weakly compactly generated* operators, that is, operators whose range is contained in a weakly compactly generated subspace of their codomain. (A Banach space is *weakly compactly generated* whenever it contains a linearly dense weakly compact subset.) The class \mathcal{WCG} contains all weakly compact operators and all operators having separable range, but in contrast, a weakly compactly generated operator defined on a $C(K)$ -space need not satisfy (\mathfrak{H}) (cf. Proposition 2.1).

To see this, consider the ordinal interval $K = [0, \omega_1]$ equipped with the order topology which makes it a compact Hausdorff space. Let $D = \{0\} \cup \{\alpha + 1: \alpha < \omega_1\}$. Define a mapping $\varphi: [0, \omega_1] \rightarrow [0, \omega_1]$ by

$$\varphi(\alpha) = \begin{cases} \alpha + 1, & \text{if } \alpha \in D, \\ \alpha, & \text{if } \alpha \in [0, \omega_1] \setminus D. \end{cases}$$

Plainly, φ is continuous, hence the operator $C_\varphi: C[0, \omega_1] \rightarrow C[0, \omega_1]$ defined by $C_\varphi f = f \circ \varphi$ is bounded.

We note that $T = I_{C[0, \omega_1]} - C_\varphi$ maps the Schauder basis $\{\mathbf{1}_{[0, \alpha]}\}_{0 \leq \alpha \leq \omega_1}$ of $C[0, \omega_1]$ onto the set $\{\mathbf{1}_{\{\alpha\}}\}_{\alpha \in D} \cup \{0\}$. Consequently, the range of T is isomorphic to $c_0(\omega_1)$, which is a weakly compactly generated Banach space, so that $T \in \mathcal{WCG}(C[0, \omega_1])$. On the other hand, T maps the 1- \prec -chain $\{\mathbf{1}_{[0, \alpha]}\}_{\alpha \in D}$ onto $\{\mathbf{1}_{\{\alpha\}}\}_{\alpha \in D}$, so T fails (\mathfrak{H}) .

It is natural to ask whether the condition (\mathfrak{H}) is the same as a seemingly weaker similar statement $(\mathfrak{H}_{\text{wo}})$ in which the δ - \prec -chains are assumed to be well-ordered. This is, however, not the case, as we shall see using the following observation.

Proposition 2.8. *Let K be a compact Hausdorff space satisfying c.c.c. Then the identity map $I_{C(K)}$ satisfies $(\mathfrak{H}_{\text{wo}})$.*

Proof. Let K be a compact Hausdorff space which does not satisfy the condition $(\mathfrak{H}_{\text{wo}})$. With no loss of generality, we may assume that, for some $\delta > 0$, there exists an uncountable, well-ordered δ - \prec -chain $\{f_\alpha\}_{\alpha < \omega_1} \subset C(K)$. For each ordinal $\alpha < \omega_1$ set $g_\alpha = |f_{\alpha+1} - f_\alpha|$. The family $\{g_\alpha^{-1}((\delta, \infty)) : \alpha < \omega_1\}$ consists of uncountably many pairwise disjoint open sets, so K fails the c.c.c. condition. \square

Corollary 2.9. *The condition $(\mathfrak{H}_{\text{wo}})$ is strictly weaker than (\mathfrak{H}) .*

Proof. In the light of Proposition 2.8, it is sufficient to exhibit an example of a compact Hausdorff space which is c.c.c., yet fails (\mathfrak{H}) .

Let K be the *double arrow space*, that is, K is the set $((0, 1] \times \{0\}) \cup ([0, 1) \times \{1\})$ endowed with the order topology arising from the lexicographic order. It is well known that K is a first-countable, non-metrizable, compact Hausdorff space which is separable (hence also c.c.c.), see [6, 3.10.C]. We note that for each $0 < b < 1$ the order-interval $I_b := [(0, 1), (b, 0)]$ is clopen in K , which means that its characteristic function $f_b = \mathbf{1}_{I_b}$ is continuous. Consequently, $\{f_b\}_{0 < b < 1}$ is an uncountable 1- \prec -chain, so K fails (\mathfrak{H}) , as desired. \square

3. COMPACT HAUSDORFF SPACES ENJOYING (\mathfrak{H})

In this section we analyse certain compact spaces to show that the identity operator on a $C(K)$ -space can satisfy (\mathfrak{H}) . We note that there is no loss of generality in considering only \prec -chains consisting of non-negative functions. Indeed, if $f \prec g$ then $|f| \prec |g|$. In the remainder of this section, we shall therefore assume that every \prec -chain has this property.

For a \prec -chain F , we denote by $A(F, f)$ the set of predecessors of f , that is, $A(F, f) = \{g \in F : g \prec f\}$; furthermore we set

$$Z(F, f) = \{x \in K : f(x) \neq 0\} \setminus \bigcup \{\text{supp}(g) : g \in A(F, f)\}.$$

Lemma 3.1. *Let K be a compact Hausdorff space, let $\delta > 0$ and let $F \subset C(K)$ be a δ - \prec -chain. Then, for each $f \in F$, $Z(F, f)$ is nonempty; in particular there is $x \in Z(F, f)$ such that $f(x) \geq \delta$.*

Proof. This is clear if f has a direct predecessor g in F ; in that case

$$Z(F, f) = \{x \in K : x \notin \text{supp}(g) \text{ and } f(x) > 0\}.$$

As $\|f - g\| \geq \delta$ there must be $x \in K \setminus \text{supp}(g)$ with $f(x) \geq \delta$; hence x belongs to $Z(F, f)$.

In case where f does not have a direct predecessor, we take an increasing and cofinal (possibly transfinite) sequence $\{g_\alpha : \alpha < \theta\}$ in $A(F, f)$. For each α we pick $x_\alpha \in \text{supp}(g_{\alpha+1}) \setminus \text{supp}(g_\alpha)$ such that $g_{\alpha+1}(x_\alpha) \geq \delta$. Note that this implies that $f(x_\alpha) \geq \delta$ for all α ; therefore $f(x) \geq \delta$ for all $x \in L$, where

$$L = \bigcap_{\alpha < \theta} \overline{\{x_\beta : \beta \geq \alpha\}}.$$

On the other hand, if $g \prec f$, then $g \prec g_\alpha$ for some α and then $g(x_\beta) = 0$ for $\beta \geq \alpha$ and hence $g(x) = 0$ for $x \in L$. This yields that the intersection $L \cap \text{supp}(g)$ is empty for all $g \in A(F, f)$ and hence that $L \subseteq Z(F, f)$. \square

Corollary 3.2. *Every compact Hausdorff space which is countable satisfies (\clubsuit) .*

For an order type α we denote by α^* the reverse of α .

Proposition 3.3. *Let K be the one-point compactification of a discrete space Γ , let $\delta > 0$, and let a δ - \prec -chain $F \subset C(K)$ be given. Then there are $\alpha, \beta \leq \omega$ such that the order type of F is $\alpha + \beta^*$*

Proof. Let F be a δ - \prec -chain. Split F into $F_1 = \{f: f(\infty) = 0\}$ and $F_2 = \{f: f(\infty) > 0\}$. (Recall that we are assuming that F consists of non-negative functions only.) Note that $f \prec g$ whenever $f \in F_1$ and $g \in F_2$.

For $f \in F_1$ set $\Upsilon_f = \{x: f(x) \geq \delta\}$; this set is finite. If $f \prec g$ in F_1 then Υ_f is a proper subset of Υ_g . This implies that the order type of F_1 is at most ω .

Likewise, for $f \in F_2$ set $\Upsilon_f = \{x: f(x) = 0\}$, which is again a finite set. In this case Υ_f contains properly Υ_g whenever $f \prec g$ in F_2 . It follows that F_2 is order-isomorphic to a subset of $\{-n: n \in \omega\} \subseteq \mathbb{Z}$. \square

We have now arrived at one of the main results of this section. We shall prove that local connectedness is a sufficient condition for absence of uncountable δ - \prec -chains of functions on spaces satisfying c.c.c. A topological space is *locally connected* if each point has a neighbourhood basis consisting of connected sets. The disjoint union of finitely many copies of the unit interval is an easy example of a (linearly ordered) compact space which is locally connected, but not connected.

Theorem 3.4. *Let K be a compact Hausdorff space which is locally connected and let $\delta > 0$. The cardinality of any δ - \prec -chain in K does not exceed the cellularity of K .*

Proof. Let $F = \{f_i\}_{i \in I} \subset C(K)$ be a δ - \prec -chain for some $\delta > 0$. The sets $Z(F, f_i)$ ($i \in I$) are non-empty and pairwise disjoint. It is enough to prove that each $Z(F, f_i)$ ($i \in I$) is open, as this will immediately yield the inequality $|I| \leq c(K)$.

Fix $i \in I$ and $x \in Z(F, f_i)$; our aim is to prove that x lies in the interior of $Z(F, f_i)$. Choose an open connected neighbourhood $U \subseteq K$ of x such that $f(y) > \frac{1}{2}f(x)$ for each $y \in U$. We *claim* that $U \cap \text{supp}(f_j) = \emptyset$ for each $f_j \prec f_i$, which means that $U \subseteq Z(F, f_i)$. Suppose that this is not the case, that is, $U \cap \text{supp}(f_j) \neq \emptyset$ for some $f_j \prec f_i$. Because $\text{supp}(f_j)$ is the closure of the open set $V = \{w \in K: f_j(w) \neq 0\}$, there must be $y \in U \cap \text{supp}(f_j)$ such that $f_j(y) \neq 0$. Since U is connected, $x \in U \setminus \text{supp}(f_j)$ and $y \in U \cap V$, the set U intersects the boundary of V ; let z be an element of this intersection. Then, $z \in U$, so $f_i(z) \neq 0$. On the other hand, $z \in \text{supp}(f_j)$, yet z does not belong to V , so $0 = f_j(z) = f_i(z)$, as $f_j \prec f_i$; a contradiction. \square

Corollary 3.5. *Every locally connected compact Hausdorff space which satisfies c.c.c. also satisfies (\clubsuit) .*

Proof. This is a direct reformulation of Theorem 3.4 for spaces having countable cellularity. \square

Remark 3.6. (i) Compact spaces which are connected need not be locally connected; a standard example of such a space (which is also not path-connected) is the so-called

topologist's closed sine curve, that is, the graph of $\sin(1/x)$, $x \in (0, 1]$ with the interval $\{0\} \times [-1, 1]$ adjoined, endowed with the relative Euclidean topology. Nevertheless, it is easily seen from the very definition of the order topology that every connected linearly ordered compact space is also locally connected.

Evidently, the unit interval satisfies the assumptions of Corollary 3.5. The existence of other linearly ordered connected (hence locally connected) examples of compact spaces satisfying c.c.c. is equivalent to the negation of the Souslin hypothesis, **SH**, which is consistent with and independent of **ZFC**. Namely, under $\neg\mathbf{SH}$, one may consider the two-point compactification of a Souslin line (recall that a *Souslin line* is a non-separable, linearly ordered, connected space without end-points which satisfies c.c.c.). Consistently, there may exist non-homeomorphic Souslin lines.

(ii) There are consistent examples of non-metrisable, locally connected compact spaces which are hereditarily Lindelöf and hereditarily separable (hence also c.c.c.) (see [7] and [9]). Curiously, there may be no non-trivial convergent sequences in spaces satisfying the assumptions of Corollary 3.5 as shown by van Mill under **CH** ([16]). Assuming Jensen's diamond principle \diamond , one has an example of such a space which is moreover one-dimensional (cf. [10]).

We refer to [5, Chapter b-11] for an exposition concerning local connectedness and further examples.

Corollary 3.5 is optimal in the sense that there may exist uncountable \prec -chains, yet for any $\delta > 0$ there may be no uncountable δ - \prec -chains.

Example 3.7. Let $\Delta \subseteq [0, 1]$ be the ternary Cantor set. For each $d \in \Delta$ set $f_d: [0, 1] \rightarrow [0, 1]$ by $f_d(x) = \text{dist}(x, \Delta) \cdot \mathbf{1}_{[d, 1]}(x)$. Each function f_d ($d \in \Delta$) is continuous and the family $\{f_d\}_{d \in \Delta}$ forms an uncountable \prec -chain in $C[0, 1]$. On the other hand, it is not hard to see that it is not a δ - \prec -chain for any positive δ .

Now, we exhibit another example of a compact, totally disconnected Hausdorff space which enjoys (\clubsuit) . This is a well-known construction in point-set topology, where it is sometimes called a *ladder system space*.

We equip the ordinal number ω_1 with a topology as follows:

- we declare zero and each countable successor number to be an isolated point;
- for each non-zero limit ordinal λ we choose a set $\{\alpha_{n,\lambda}: n < \omega\} \subseteq \lambda$ of order type ω consisting of successor numbers and cofinal in λ (a *ladder*); then we define basic open neighbourhoods of λ to be of the form $U_{\lambda,m} = \{\lambda\} \cup \{\alpha_{n,\lambda}: n \geq m\}$ ($m < \omega$).

Of course, the topology on ω_1 depends on the choice of ladders but in any case the space ω_1 topologised in this manner is first countable, locally compact and Hausdorff.

Theorem 3.8. *Let K be the one-point compactification of a ladder system space on ω_1 , and let $\delta > 0$. Then every δ - \prec -chain in $C(K)$ is countable. In particular, K satisfies (\clubsuit) .*

Proof. We model the proof after that of Proposition 3.3: we take a δ - \prec -chain F and divide it into two sets: $F_1 = \{f \in F: f(\omega_1) = 0\}$ and $F_2 = \{f \in F: f(\omega_1) > 0\}$.

Let L_λ denote the ladder associated to λ , and set $L_\lambda^+ = L_\lambda \cup \{\lambda\}$. Note that a closed subset of ω_1 is compact if and only if it is covered by finitely many sets of the form L_λ^+ and possibly a finite number of isolated points.

We *claim* that F_1 is countable. For f in F_1 let $\Upsilon_f = \{x \in \omega_1: f(x) \geq \delta\}$. Each Υ_f is compact, hence countable. Lemma 3.1 implies that if $g \prec f$, then $Z(F, g) \cap \Upsilon_f \neq \emptyset$. As the sets $Z(F, f)$ and $Z(F, g)$ are disjoint, this shows that $A(F, f)$ is countable for any $f \in F_1$. Also, if $g \prec f$ in F_1 , then Υ_g is a proper subset of Υ_f .

It suffices to show that F_1 has countable cofinality. We assume, seeking of a contradiction, that $\{f_\alpha: \alpha < \omega_1\}$ is a strictly increasing sequence in F_1 . Then the sets Υ_{f_α} form a strictly increasing sequence as well so that the union $\bigcup_{\alpha < \omega_1} \Upsilon_{f_\alpha}$ is uncountable.

Given $\alpha < \omega_1$ let $\beta_\alpha > \alpha$ be such that

- $\bigcup_{\gamma \leq \alpha} \Upsilon_{f_\gamma} \subseteq \beta_\alpha$, and
- there is a $\gamma < \beta_\alpha$ such that $x > \alpha$ for some $x \in \Upsilon_{f_\gamma}$.

Next let C be a closed and unbounded set such that for each $\gamma \in C$ we have $\beta_\alpha < \gamma$ whenever $\alpha < \gamma$.

Let $\gamma \in C$ be such that $C \cap \gamma$ has order type ω^2 . We show that Υ_{f_γ} is not compact. To this end, we note that, by the above two items there is a point in Υ_{f_γ} between η and ξ , whenever η and ξ are consecutive elements of $C \cap \gamma$. This shows that the order type of $\Upsilon_{f_\gamma} \cap \gamma$ is at least ω^2 .

Let H be a finite subset of Λ . We show that the set $\Upsilon_{f_\gamma} \setminus \bigcup_{\lambda \in H} L_\lambda^+$ is infinite. This follows from the following three observations

- if $\lambda \in H \cap \gamma$, then L_λ^+ is bounded below by γ ;
- $\Upsilon_{f_\gamma} \setminus L_\gamma^+$ is cofinal in γ (because of the order types of both sets); and
- if $\lambda \in H$ is larger than γ then $L_\lambda^+ \cap \gamma$ is finite.

Thus we see that Υ_{f_γ} is not compact.

For $f \in F_2$ we consider the compact set $\Upsilon_f = \{x \in X: f(x) = 0\}$. Arguing similarly as before we conclude that F_2 is countable, yet the order of F_2 is reversed in the following sense: if $f \prec g$ in F_2 , then $\Upsilon_f \supseteq \Upsilon_g$ and there is $x \in \Upsilon_f \setminus \Upsilon_g$ such that $g(x) \geq \delta$. \square

A compact Hausdorff space is *Eberlein* if it is homeomorphic to a weakly compact subset of a Banach space. Countable compact Hausdorff spaces, the one-point compactification of a discrete space (Proposition 3.3) as well as the unit interval (*cf.* Remark 3.6 (i)) are classical examples of Eberlein compact spaces. A space K is Eberlein if and only if the Banach space $C(K)$ is weakly compactly generated (*cf.* [2, Theorem 2]). Every weakly compactly generated Banach space is necessarily Lindelöf in its weak topology. Wage observed that no ladder system space K is Eberlein ([17]), while Pol proved that the Banach space $C(K)$ is Lindelöf in its weak topology ([14]). In the light of these observations, we raise the following question.

Question 3.9. Does the class of compact Hausdorff spaces satisfying (\clubsuit) contain all Eberlein compact spaces? More generally, does every compact Hausdorff space K for which the Banach space $C(K)$ is Lindelöf in its weak topology satisfy (\clubsuit) ?

4. THE LEFT IDEAL STRUCTURE OF OPERATORS SATISFYING (\clubsuit)

Let $[A]^2$ stand for the set of all 2-element subsets of a given set A . The Dushnik and Miller partition lemma ([4]) asserts that, for every infinite regular aleph \aleph_α , and any colouring $c: [\aleph_\alpha]^2 \rightarrow \{0, 1\}$, at least one of the following conditions holds true:

- (i) there is a set $A \subset \aleph_\alpha$ with $|A| = \aleph_\alpha$ and $[A]^2 \subset c^{-1}\{0\}$;
- (ii) for every set $A \subset \aleph_\alpha$ with $|A| = \aleph_\alpha$ there are $a \in A$ and a set $B \subset A$ such that $|B| = \aleph_\alpha$ and $\{a, b\} \in c^{-1}\{1\}$ for each $b \in B$.

Proposition 4.1. *Let K be a compact Hausdorff space and let X be a Banach space. If operators $T, U: C(K) \rightarrow X$ satisfy (\clubsuit) , then so does $T + U$.*

Proof. Assume, in search of a contradiction, that there are $\delta > 0$ and a \prec -chain $\{f_i\}_{i \in I}$ in $C(K)$, where I has cardinality \aleph_1 , such that $\sup_{i \in I} \|f_i\| < \infty$ and

$$(*) \quad \|(T + U)(f_i) - (T + U)(f_j)\| \geq \delta \quad (i, j \in I, i \neq j).$$

Set

$$\mathcal{I} = \{\{i, j\} \in [I]^2 : \|U(f_i) - U(f_j)\| \leq \delta/4\}$$

and consider the partition $[I]^2 = \mathcal{I} \cup ([I]^2 \setminus \mathcal{I})$.

If assertion (i) from the partition lemma holds (where $\alpha = 1$ and $c^{-1}\{0\} = [I]^2 \setminus \mathcal{I}$), then we get a contradiction with the fact that U satisfies (\clubsuit) .

If assertion (ii) is valid (where $c^{-1}\{1\} = \mathcal{I}$), then there is an $i_0 \in I$ and a set $B \subset I$ with $|B| = \aleph_1$, such that $\{i_0, j\} \in \mathcal{I}$ for each $j \in B$. Then for every $j, k \in B$ we have

$$\|U(f_j) - U(f_k)\| \leq \|U(f_{i_0}) - U(f_j)\| + \|U(f_{i_0}) - U(f_k)\| \leq \delta/2,$$

hence our assumption $(*)$ implies that $\|T(f_j) - T(f_k)\| \geq \delta/2$ for all $j, k \in B, j \neq k$, which contradicts the condition (\clubsuit) for T . \square

Theorem 4.2. *For every compact Hausdorff space K the set of all operators $T \in \mathcal{B}(C(K))$ satisfying (\clubsuit) forms a closed left ideal of the Banach algebra $\mathcal{B}(C(K))$.*

Proof. From the very definition of (\clubsuit) it is evident that if $(T_n)_{n=1}^\infty \subset \mathcal{B}(C(K))$ norm converges to some $T \in \mathcal{B}(C(K))$ and each T_n satisfies (\clubsuit) , then T does as well.

Now, suppose that $T \in \mathcal{B}(C(K))$ satisfies (\clubsuit) and let $S \in \mathcal{B}(C(K))$. If for some $\delta > 0$ there were an uncountable chain $\{f_i\}_{i \in I}$ with $\|ST(f_i) - ST(f_j)\| > \delta$ for all $i, j \in I, i \neq j$, then for all such i, j we would also have $\|T(f_i) - T(f_j)\| > \delta/\|S\|$; a contradiction. Finally, an appeal to Proposition 4.1 completes the proof. \square

We have already noticed that for K extremally disconnected the family of operators $T: C(K) \rightarrow C(K)$ which satisfy (\clubsuit) coincides with the two-sided ideal of weakly compact operators. Examples in Section 3 demonstrate that in some particular cases every operator on a $C(K)$ -space may satisfy (\clubsuit) , which means that the left ideal of operators satisfying (\clubsuit) is again a two-sided ideal, the improper ideal $\mathcal{B}(C(K))$. This raises the following open question.

Question 4.3. Is the set of operators on a $C(K)$ -space which satisfy (\clubsuit) always a right, and hence a two-sided ideal of $\mathcal{B}(C(K))$?

Acknowledgements. The authors would like to express their gratitude to N.J. Laustsen and J.M. Lindsay for careful reading of preliminary versions of this paper.

REFERENCES

1. F. Albiac and N. J. Kalton, *Topics in Banach Space Theory*, Graduate Texts in Mathematics 233, Springer 2006.
2. D. Amir and J. Lindenstrauss, “The structure of weakly compact sets in Banach spaces”, *Ann. of Math.* 88 (1968), 35–46.
3. J. Diestel and J. J. Uhl, *Vector Measures*. With a foreword by B.J. Pettis, Mathematical Surveys, no. 15, American Mathematical Society, Providence, R.I. 1977.
4. B. Dushnik, E. W. Miller, “Partially ordered sets”, *Amer J. Math.* 63 (1941), 600–610.
5. *Encyclopedia of General Topology*, ed. K. P. Hart, J. Nagata, and J. E. Vaughan, Elsevier (2004).
6. R. Engelking, *General Topology*, 2nd edition, Sigma Ser. Pure Math., Vol. 6, Heldermann, Berlin (1989).
7. V. V. Filippov, “Perfectly normal bicompecta” (Russian), *Dokl. Akad. Nauk SSSR* 189 (1969) 736–739; English translation in *Soviet Math. Dokl.* 10 (1969) 1503–1507.
8. D. B. Goodner, “Projections in normed linear spaces”, *Trans. Amer. Math. Soc.* 69 (1950), 89–108.
9. K. Kunen, “Locally connected hereditarily Lindelöf compacta”. *Topology Appl.* 158 (2011), no. 18, 2473–2478.
10. K. Kunen and J. E. Hart, “One-dimensional locally connected S-spaces”, *Topology Appl.* 156 (2009), no. 3, 601–609.
11. R. Haydon, “A non-reflexive Grothendieck space that does not contain l^∞ ”, *Israel J. Math.* 40 (1981), 65–73.
12. T. Kania and T. Kochanek, “The ideal of weakly compactly generated operators acting on a Banach space”, to appear in *J. Operator Theory*, <http://arxiv.org/abs/1206.5424v3>.
13. L. Nachbin, “A theorem of the Hahn–Banach type for linear transformations”, *Trans. Amer. Math. Soc.* 68 (1950), 28–46.
14. R. Pol, “A function space $C(X)$ which is weakly Lindelöf but not weakly compactly generated”, *Studia Math.* 64 (1979), 279–285.
15. H. P. Rosenthal, “On relatively disjoint families of measures, with some applications to Banach space theory”, *Studia Math.* 37 (1970), 13–36.
16. J. van Mill, “A locally connected continuum without convergent sequences”, *Topology Appl.* 126 (2002), no. 1-2, 273–280.
17. M. L. Wage, *Applications of the set theory to analysis and topology*, Thesis, University of Wisconsin–Madison, 1976.

FACULTY OF ELECTRICAL ENGINEERING, MATHEMATICS AND COMPUTER SCIENCE, TU DELFT,
POSTBUS 5031, 2600 GA DELFT, THE NETHERLANDS
E-mail address: k.p.hart@tudelft.nl

DEPARTMENT OF MATHEMATICS AND STATISTICS, FYLDE COLLEGE, LANCASTER UNIVERSITY, LAN-
CASTER LA1 4YF, UNITED KINGDOM
E-mail address: t.kania@lancaster.ac.uk

INSTITUTE OF MATHEMATICS, UNIVERSITY OF SILESIA, BANKOWA 14, 40-007 KATOWICE, POLAND
E-mail address: tkochanek@math.us.edu.pl