

## Grassland productivity is limited by multiple nutrients

Philip A. Fay<sup>1\*</sup>, Suzanne M. Prober<sup>2</sup>, W. Stanley Harpole<sup>3-6</sup>, Johannes M.H. Knops<sup>7</sup>, Jonathan D. Bakker<sup>8</sup>, Elizabeth T. Borer<sup>9</sup>, Eric M. Lind<sup>9</sup>, Andrew S. MacDougall<sup>10</sup>, Eric W. Seabloom<sup>9</sup>, Peter D. Wragg<sup>9</sup>, Peter B. Adler<sup>11</sup>, Dana M. Blumenthal<sup>12</sup>, Yvonne M. Buckley<sup>13</sup>, Chengjin Chu<sup>14</sup>, Elsa E. Cleland<sup>15</sup>, Scott L Collins<sup>16</sup>, Kendi F. Davies<sup>17</sup>, Guozhen Du<sup>14</sup>, Xiaohui Feng<sup>18</sup>, Jennifer Firn<sup>19</sup>, Daniel S Gruner<sup>20</sup>, Nicole Hagenah<sup>21</sup>, Yann Hautier<sup>22</sup>, Robert W. Heckman<sup>23</sup>, Virginia L. Jin<sup>24</sup>, Kevin P. Kirkman<sup>21</sup>, Julia Klein<sup>25</sup>, Laura M. Ladwig<sup>16</sup>, Qi Li<sup>26</sup>, Rebecca L. McCulley<sup>27</sup>, Brett A. Melbourne<sup>17</sup>, Charles E. Mitchell<sup>23</sup>, Joslin L Moore<sup>28</sup>, John W. Morgan<sup>29</sup>, Anita C. Risch<sup>30</sup>, Martin Schütz<sup>30</sup>, Carly J Stevens<sup>31</sup>, David A. Wedin<sup>32</sup>, Louie H. Yang<sup>33</sup>

1. USDA-ARS Grassland Soil and Water Research Lab, Temple, TX, 76502 USA
2. CSIRO Land and Water Flagship, Private Bag 5, Wembley, WA 6913, Australia
3. Department of Ecology, Evolution, and Organismal Biology, Iowa State University, Ames, IA, 50011 USA
4. German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, Deutscher Platz 5e, D-04103 Leipzig, Germany
5. Department of Physiological Diversity, Helmholtz Center for Environmental Research – UFZ, Permoserstr. 15, 04318 Leipzig, Germany
6. Institute of Biology, Martin Luther University Halle-Wittenberg, Am Kirchtor 1, 06108 Halle (Saale), Germany
7. School of Biological Sciences, University of Nebraska, Lincoln, NE, 68588 USA
8. School of Environmental and Forest Sciences, University of Washington, Seattle, WA, 98195 USA
9. Department of Ecology, Evolution, and Behavior, University of MN, St. Paul, MN, 55108 USA
10. Department of Integrative Biology, University of Guelph, Guelph, Ontario, Canada N1G 2W1
11. Department of Wildland Resources and the Ecology Center, Utah State University, Logan, UT, 84322 USA
12. USDA-ARS Rangeland Resources Research Unit, Fort Collins, CO 80526 USA
13. School of Natural Sciences, Zoology, Trinity Centre for Biodiversity Research, Trinity College Dublin, Dublin 2, Ireland
14. Research Station of Alpine Meadow and Wetland Ecosystems, Lanzhou University, Lanzhou, 730000, China.
15. Ecology, Behavior & Evolution Section, University of California, San Diego, La Jolla, CA, 92093 USA
16. Department of Biology, MSC03-2020, University of New Mexico, Albuquerque, NM, 88003 USA
17. Department of Ecology and Evolutionary Biology, University of Colorado, Boulder, CO, 80309 USA
18. Department of Plant Biology, University of Illinois, Urbana, IL, 61801 USA
19. School of Earth, Environment and Biological Sciences, Queensland University of Technology, Brisbane, Queensland 4001, Australia
20. Department of Entomology, University of Maryland, College Park, MD, 20742 USA
21. School of Life Sciences, University of KwaZulu-Natal, Pietermaritzburg, 3209, South Africa.
22. Ecology and Biodiversity Group, Department of Biology, Utrecht University, Padualaan 8, 3584 CH Utrecht, Netherlands

23. Department of Biology, University of North Carolina at Chapel Hill, Chapel Hill, NC, 27599 USA
24. USDA-ARS Agroecosystem Management Research Unit, Lincoln, NE, 68538 USA.
25. Department of Forest, Rangeland, and Watershed Stewardship, Colorado State University, Fort Collins, CO, 80523 USA
26. Key Laboratory of Adaptation and Evolution of Plateau Biota, Northwest Institute of Plateau Biology, Chinese Academy of Sciences, Xining, 810008, China
27. Department of Plant and Soil Sciences, University of Kentucky, Lexington, KY, 40546 USA
28. School of Biological Sciences, Monash University, Victoria 3800, Australia.
29. Department of Botany, La Trobe University, Bundoora 3083, Victoria, Australia
30. Swiss Federal Institute for Forest, Snow and Landscape Research, Community Ecology, 8903 Birmensdorf, Switzerland
31. Lancaster Environment Centre, Lancaster University, Lancaster, LA1 4YQ, UK
32. School of Natural Resources, University of Nebraska, Lincoln, NE, 68583 USA
33. Department of Entomology and Nematology, University of California, Davis, CA, 95616 USA.

\*To whom correspondence should be addressed ([philip.fay@ars.usda.gov](mailto:philip.fay@ars.usda.gov))

**Terrestrial ecosystem productivity is widely accepted to be nutrient limited<sup>1</sup>. Although nitrogen is deemed a key determinant of aboveground net primary production<sup>2,3</sup>, the prevalence of co-limitation by nitrogen and phosphorus is increasingly recognized<sup>4-8</sup>. However, the extent to which terrestrial productivity is co-limited by nutrients other than nitrogen and phosphorus, such as potassium and micronutrients, has remained unclear. Here, we report results from a standardized factorial nutrient addition experiment, in which we added nitrogen (N), phosphorus (P), and potassium combined with a selection of micronutrients ( $K_{+\mu}$ ), alone or in concert, to 42 grasslands spanning five continents, and monitored aboveground net primary production. Nutrient availability limited productivity at 31 of the 42 grassland sites studied. Pairwise combinations of N, P, and  $K_{+\mu}$  co-limited aboveground net primary production at 24 of the 42 sites, and nitrogen limitation peaked in cool, high latitude sites. Our findings indicate widespread variation in the combination of nutrients that limit aboveground grassland productivity, and highlight the importance of potassium and micronutrients, and an apparent absence of limitation by the nutrients we considered at some sites. This variation in nutrient limitation must be considered when assessing the ecosystem-scale consequences of nutrient enrichment.**

1 Terrestrial ecosystem productivity is widely accepted to be nutrient limited<sup>1</sup>, and many  
2 studies have focused on limitation by a single nutrient, nitrogen (N)<sup>2,3</sup>. In grasslands, and in  
3 other systems, the role of additional nutrients is increasingly recognized. A recent meta-analysis  
4 of 1400 N and phosphorus (P) fertilization studies<sup>4</sup> showed limitation of terrestrial productivity  
5 by both N and P. Furthermore, these nutrients often were synergistically co-limiting, where  
6 together they limited productivity more than the sum of their individual limitations. This meta-  
7 analysis is the most comprehensive assessment of ecosystem nutrient limitation to date.  
8 However, the global extent and magnitude of multiple limitation by nutrients other than N and P  
9 remains poorly understood in natural systems<sup>4,5,9</sup>, including grasslands<sup>3,10</sup>, a critically endangered  
10 biome that accounts for approximately one-third of Earth's terrestrial net primary production<sup>11</sup>.  
11 More importantly, multiple nutrient limitation has not been experimentally tested in grasslands  
12 on a global scale using a standardized experimental approach. The potential for synergistic co-  
13 limitation of grasslands by multiple nutrients, or conversely, failing to account for the absence of  
14 single or multiple nutrient limitation means we may misestimate the magnitude and extent of  
15 nutrient limitation of terrestrial net primary productivity.

16 Humans now produce more reactive N than is produced from all natural terrestrial  
17 sources, primarily as fertilizers, industrial products, and through fossil fuel combustion<sup>12,13</sup>.  
18 Anthropogenic increases in atmospheric N will result in further terrestrial N deposition, altering  
19 ecosystem function<sup>12,14-17</sup> and potentially increasing limitation by other nutrients, such as P,  
20 potassium (K), or trace elements<sup>16</sup>. K occurs in high concentrations in plant tissues<sup>18</sup> and its  
21 uptake is correlated with that of other nutrients<sup>19,20</sup>. Our understanding of limitation by nutrients  
22 other than N in grasslands lags that of agro-ecosystems, where the importance of P, K, and  
23 micronutrients is better understood<sup>20</sup>.

24 Here we report the frequency, magnitude, and global extent of nutrient limitation of  
25 grassland ANPP by N, P, and K combined with micronutrients ( $K_{+\mu}$ ) at 42 grassland sites in the  
26 Nutrient Network (NutNet; Supplementary Material and <sup>21</sup>). All sites conducted the same  
27 standardized N-P- $K_{+\mu}$  addition experiment, the definitive test for nutrient limitation. This  
28 approach overcomes major constraints of previous meta-analyses<sup>4</sup>, differing nutrient addition  
29 rates and methodologies and the rarity of factorial nutrient treatments in the underlying studies<sup>5</sup>.

30 Sites were located on five continents (Supplementary Figure 1) and spanned a 23-fold  
31 range in ANPP (Supplementary Figure 2A), over 25° in absolute latitude (actual latitudes 54° N  
32 to 37° S), nearly 3,500 m in elevation, and wide ranges in mean annual precipitation (260 – 1900  
33 mm), mean annual temperature (0.3 – 22° C; Supplementary Figure 1), soil texture, and pre-  
34 treatment soil N, P, and K pools (Supplementary Table 1). The sites included both native and  
35 previously cultivated grassland, and some sites were managed with burning, grazing, or other  
36 practices. Thus, we were able to evaluate single and multiple-nutrient limitation of ANPP at  
37 global, continental, and site spatial extents, and as influenced by management, climate, and soils.

38 N, P, and  $K_{+\mu}$  were factorially applied annually to replicated 5 m<sup>2</sup> plots at the beginning  
39 of each site's growing season at rates commonly used in grassland fertilization experiments<sup>22,23</sup>.  
40 The  $K_{+\mu}$  treatment included a micronutrient mix in the first treatment year only, to avoid  
41 micronutrient toxicity. Nutrient limitation of ANPP was quantified by harvesting current year  
42 standing crop biomass for 3 to 5 years. and computing the log response ratio (LRR), the natural  
43 logarithm of the ratio of treatment plot to control plot ANPP, a metric commonly used in meta-  
44 analyses<sup>24</sup>

45 Multiple nutrient limitation of ANPP often occurs in the form of co-limitation<sup>5</sup>. Co-  
46 limitation is synergistic when the response to multiple nutrients is greater than the sum of the  
47 responses to each nutrient added individually, additive if the multiple-nutrient response equals  
48 the sum of the individual nutrient responses, and sub-additive if less than the sum of the single-  
49 nutrient responses<sup>5</sup>. Across all sites and years, the combined addition of N and P increased  
50 ANPP by an average of 40% over controls (LRR= 0.34), compared to increases of only 18%  
51 (LRR = 0.16) for N individually and 9% for P individually (LRR = 0.09; p = 0.03, Figure 1A,  
52 Supplementary Figure 3). This provides clear evidence for globally-averaged synergistic co-  
53 limitation of ANPP by N and P in these grasslands. Synergistic co-limitation by N and P across  
54 sites and years emerged from frequent occurrence of NP co-limitation at the site level. Some  
55 form of NP co-limitation averaging 67% (LRR = 0.51) occurred at 60% of the sites, including  
56 sites on all five continents, with synergistic NP co-limitation at 13 sites and sub-additive co-  
57 limitation at 12 sites, (Supplementary Table 3). Globally-averaged synergistic co-limitation of  
58 grassland ANPP by N and P contradicts the long-held perception that N is the predominant  
59 nutrient limiting grassland productivity<sup>3,10</sup> and highlights a critical role for P.

60 There was no globally-averaged single or multiple-nutrient limitation involving  $K_{+\mu}$  (p >  
61 0.32, Supplementary Table 2). However, this did not preclude frequent site-level single and  
62 multiple-nutrient  $K_{+\mu}$  limitation of grassland ANPP. ANPP was limited at 3 sites by  $K_{+\mu}$  alone  
63 and co-limited by  $NK_{+\mu}$  at 18 sites, both averaging 48% (LRR = 0.39; Figure 1A, Supplementary  
64 Table 3). Additionally, ANPP was co-limited by  $PK_{+\mu}$  at 9 sites by an average of 52% (LRR =  
65 0.42), approaching the magnitude of ANPP limitation at sites where N (57%) or P (54%)  
66 individually limited ANPP (Figure 1A). Site-level  $NK_{+\mu}$  and  $PK_{+\mu}$  co-limitation occurred in  
67 synergistic, additive, and sub-additive forms, and in total, single or multiple-nutrient limitation

68 involving  $K_{+\mu}$  occurred at 24 sites again occurring on all five continents (Supplementary Table  
69 3). This finding represents the broadest assessment to date of potential  $K_{+\mu}$  limitation of  
70 grassland ANPP, and suggests that single- and multiple-nutrient K limitation, potentially  
71 augmented by micronutrient addition during year 1, occurs more frequently than previously  
72 recognized<sup>20,25</sup>.

73 Although single and multiple-nutrient limitation of ANPP was widespread, there were  
74 nonetheless 15 sites with no evidence for multiple nutrient limitation, and at 12 of the 15 also no  
75 evidence for single-nutrient limitation (Supplementary Table 3). The presence and magnitude of  
76 nutrient limitation may depend on site climate, soil development, or fertility<sup>26,27</sup>. For these  
77 reasons, greater site-level limitation of ANPP by one nutrient may correlate with greater  
78 limitation by one or more additional nutrients. Indeed, site-level individual nutrient limitation of  
79 ANPP increased with limitation by other individual nutrients ( $R^2$  0.08 – 0.15; Figure 2A,B,D),  
80 and ANPP limitation by nutrient pairs increased with that of a third individual nutrient ( $R^2$  0.07 –  
81 0.21; Figure 2C,E,F). These correlations suggest that site-level attributes may predict the  
82 magnitude of single and multiple nutrient limitation of ANPP.

83 Sites differed in various potential qualitative and quantitative predictors of the presence  
84 and magnitude of nutrient limitation of ANPP (Supplementary Table 1). We found no evidence  
85 that nutrient limitation differed among the qualitative predictors continent, history of  
86 management with burning, grazing, and previous cultivation, and predominant soil texture  
87 (Supplementary Figure 4, Supplementary Table 4). Quantitative predictors were examined at the  
88 19 longest running sites, where estimates of site mean nutrient limitation should best correspond  
89 to site mean climate and soil attributes. N limitation increased with decreasing site mean annual  
90 temperature (MAT;  $R^2 = 0.40$ ,  $p = 0.002$ , Figure 3A) and peaked at temperate latitude sites

91 (~45°;  $R^2 = 0.38$ ,  $p = 0.009$ , Figure 3B) with lower MAT ( $R^2 = 0.66$ ,  $p < 0.0001$ , Figure 3B  
92 inset). Limitation of ANPP by other single or multiple nutrient combinations was not correlated  
93 with soil texture fractions, pre-treatment soil nutrient contents, soil pH, or climate variables ( $p >$   
94  $0.05$ , data not shown). These findings suggest that N limitation predominated at cooler  
95 temperate latitudes, as others have predicted<sup>26</sup>, and validate the power of these experiments to  
96 detect spatial trends in ecosystem nutrient limitation.

97         The magnitude of single or multiple-nutrient limitation of grassland ANPP may increase  
98 through time (Supplementary Figure 2B). To isolate the temporal signal of nutrient addition  
99 effects on ANPP, we calculated new ANPP response ratios using the pre-treatment year (i.e.,  
100 year 0) ANPP for each treatment plot rather than the control plot, to control for plot-to-plot  
101 spatial variation in ANPP. For 37 sites with no missing ANPP data in years 1-3, the ANPP  
102 response to nutrient addition increased through time in all treatments, ANPP increased each year  
103 more in treatments containing N ( $p < 0.005$ , Figure 4, Supplementary Table 2) compared to the  
104 P,  $K_{+\mu}$ , and  $PK_{+\mu}$  treatments. Thus, the larger global-averaged increase in ANPP with N than P or  
105  $K_{+\mu}$  (Figure 1A) can be explained partly by both smaller and later manifestation of P and  $K_{+\mu}$   
106 effects. The full extent of nutrient limitation of grassland ANPP, particularly for nutrients other  
107 than N, may require additional years to emerge.

108         Multiple nutrient limitation of ANPP in these grasslands was spatially and temporally  
109 varied, often included K and micronutrients, but sometimes none of the nutrients we considered.  
110 Thus, while N was an important limiting nutrient in many grasslands, P and  $K_{+\mu}$  were also  
111 limiting in many instances, both alone and in combination with each other and with N. These  
112 estimates for globally-averaged nutrient limitation in these grasslands are conservative because

113 nutrient limitation increased in successive treatment years. The magnitude of nutrient limitation  
114 may require revision upward if the increases continue in future years.

115 Multiple nutrient limitation in grasslands, especially synergistic co-limitation, makes the  
116 consequences of anthropogenic nutrient inputs in these and other systems more difficult to  
117 predict. For example, nutrient co-limitation may actually restrict the impacts of single nutrient  
118 inputs such as N deposition<sup>17</sup>, but ecosystems may respond more rapidly than expected to  
119 subsequent inputs of the co-limiting nutrient. Considering only one of the co-limiting nutrients  
120 would risk reaching erroneous conclusions about the consequences of future inputs for  
121 biodiversity or ecosystem goods/services provision<sup>14,15,28,29</sup>. It is imperative to consider multiple  
122 nutrient constraints on primary productivity in grasslands and other ecosystems.

123 **Full Methods** and any associated references are available in the online version of the paper at  
124 [www.nature.com/nature](http://www.nature.com/nature).

125 **Received** |date|; **Accepted** |date|.

126 **Published online** |date|.

127

128 **Methods**

129 *Study system*

130 The rationale, goals, and experimental protocols used at grasslands in the Nutrient  
131 Network (NutNet) are described in Borer et al.<sup>21</sup>, and are summarized here. This study used  
132 aboveground net primary productivity (ANPP) data from a standardized nutrient addition  
133 experiment conducted at 42 NutNet sites in eight countries (Australia, Canada, China,  
134 Switzerland, Tanzania, United Kingdom, USA, South Africa) on five continents (Australia [N =



135 4], Africa [N = 4], Europe [N = 5], Asia [N = 1], North America [N = 28]; Supplementary Figure  
136 1, Table 1).

137 The grassland types included alpine/montane grasslands, shortgrass, mixed, and tallgrass  
138 prairies, desert and semi-arid grasslands, old fields and pastures, savanna and shrub-steppe, and  
139 annual grasslands. Sites span wide ranges of mean annual precipitation (MAP; 252 - 1898 mm  
140  $y^{-1}$ ), mean annual temperature (MAT; 0.3 - 22.1 °C; Supplementary Figure 1), elevation (50 -  
141 3500 m), and absolute latitude (over 25°, actual latitudes 54° N to 37° S, Supplementary Table  
142 1). Site climate data were derived from the WorldClim database<sup>30</sup> (version 1.4.). Local  
143 investigators classified sites as to management practice (burned, grazed, other management) and  
144 native or anthropogenic (restored grassland and agricultural pasture) origin. Sites collected pre-  
145 treatment soil nutrient content and texture data.

146 At each site, N, P, and K were added in full factorial combination (8 treatments including  
147 control plots) to 5 x 5 m plots in a randomized complete blocks design with N=3 for most sites  
148 (range 2-6). This experimental scale is well-suited for the relatively short-statured herbaceous  
149 vegetation in grasslands. Micronutrients (6% Ca, 3% Mg, 12% S, 0.1% B, 1% Cu, 17% Fe,  
150 2.5% Mn, 0.05% Mo, and 1% Zn) were added with K only in year 1 to avoid possible  
151 micronutrient toxicity. Nutrient additions began at most sites in 2008 (N = 28), and additional  
152 sites began in 2009 (N = 9), 2010 (N = 3) and 2011 (N = 2).

### 153 ***Sampling***

154 Total ANPP ( $g\ m^{-2}\ yr^{-1}$ ) was estimated annually from clip samples of current year peak  
155 aboveground biomass from two 0.1  $m^2$  quadrats per plot. Samples were dried to constant mass at  
156 60°C.

157 *Statistical Analyses*

158 We tested for nutrient treatment effects on ANPP using linear mixed models in  
159 SAS/STAT v 9.2. First, we tested for variation among sites in nutrient main and interactive  
160 effects and their variation with treatment year (Equation 1):

$$\begin{aligned} 161 \quad \ln(\text{ANPP})_{ijkl} = & \mu + \text{nutrients}_i + \text{nutrients}(\text{site})_{ij} + \text{nutrients}(\text{year})_{ik} \\ 162 \quad & + \text{nutrients}(\text{site} \times \text{year})_{ijk} + e_{ijkl}. \quad (1) \end{aligned}$$

163 where *nutrients* refers to the eight factorial combinations of added N, P, and K. Nutrient  
164 treatments (i) were nested within site (j), within year (k), and within site x year (jk). Year was  
165 not treated as a repeated effect because each site experienced each year differently, and thus both  
166 the year effects and the covariance among years were not the same at each site. There were 42  
167 sites with three years of data, 33 sites with four years, and 22 sites with five years. Model (1) was  
168 tested using all available treatment years, and with only the first three treatment years to  
169 ascertain whether any nutrient x year interactions were caused by sites added later with fewer  
170 years of nutrient addition.

171 We tested whether ANPP responses to nutrient treatments varied with continent,  
172 management, dominant soil texture, and origin by replacing the site term in Eq. 1 with each of  
173 the site classification and soils variables, one at a time. Europe and Asia were combined in tests  
174 of continent because of the single Asian site.

175 *Calculation of nutrient effect sizes*

176 Effects sizes of the nutrient treatments were calculated and graphed as natural-log  
177 response ratios (LRR). LRRs were calculated in two ways; first, by comparing the ANPP of each

178 treatment plot to the ANPP of the control plot in the same block (2), to isolate treatment effects;  
179 and second, by compared treatment plot ANPP to that of its pre-treatment (i.e., year 0) ANPP  
180 (3). This isolated the temporal signal of nutrient addition responses.

$$181 \quad \mathbf{LRR = \ln[ANPP_{\text{trt}}/ANPP_{\text{control}}]} \quad (2)$$

$$182 \quad \mathbf{LRR = \ln[ANPP_{\text{trt}}(\text{year X})/ANPP_{\text{trt}}(\text{year 0})]} \quad (3)$$

183 We estimated a critical threshold LRR (4) for categorizing the significance of site LRRs. An  
184 LRR was considered statistically significant if it exceeded a value corresponding to the critical  
185 Z-score (1.65) at  $p = 0.05$ , computed as:

$$186 \quad \mathbf{\text{Threshold LRR} = 1/7 * (\text{LRR}_N + \text{LRR}_P + \text{LRR}_K + \dots \text{LRR}_{\text{NPK}})/\text{pooled SD.}} \quad (4)$$

### 187 *Correlations of site variables with nutrient effects*

188 We used linear regression analyses to test whether LRRs for each single nutrient and  
189 nutrient pair were predicted by site latitude (absolute value), elevation, mean annual precipitation  
190 (MAP), mean temperature annual (MAT), potential evapotranspiration (PET), soil fractions of  
191 sand, silt, and clay, pre-treatment soil N, P, and K contents, and soil pH. We presented  
192 regressions only where  $p < 0.05$ , and fit polynomial curves where indicated by evaluation of  
193 residuals.

194

195

## References

- 196 <sup>1</sup> Chapin, F. S., Matson, P. A. & Vitousek, P. M. *Principles of terrestrial ecosystem ecology* 2nd ed.  
197 (Springer, New York, 2011).
- 198 <sup>2</sup> Vitousek, P. & Howarth, R. Nitrogen limitation on land and in the sea: How can it occur?  
199 *Biogeochemistry* **13**, 87-115 (1991).
- 200 <sup>3</sup> LeBauer, D. S. & Treseder, K. K. Nitrogen limitation of net primary productivity in terrestrial  
201 ecosystems is globally distributed. *Ecology* **89**, 371-379 (2008).
- 202 <sup>4</sup> Elser, J. J. *et al.* Global analysis of nitrogen and phosphorus limitation of primary producers in  
203 freshwater, marine and terrestrial ecosystems. *Ecol. Lett.* **10**, 1135-1142 (2007).
- 204 <sup>5</sup> Harpole, W. S. *et al.* Nutrient co-limitation of primary producer communities. *Ecol. Lett.* **14**, 852-  
205 862 (2011).
- 206 <sup>6</sup> Bracken, M. E. S. *et al.* Signatures of nutrient limitation and co-limitation: responses of autotroph  
207 internal nutrient concentrations to nitrogen and phosphorus additions. *Oikos* **124**, 113-  
208 121 (2015).
- 209 <sup>7</sup> Ågren, G. I. , Wetterstedt, J. Å. & Billberger, M. F. K. Nutrient limitation on terrestrial plant growth  
210 – modeling the interaction between nitrogen and phosphorus. *New Phytol.* **194**, 953-960  
211 (2012).
- 212 <sup>8</sup> Carnicer, J. *et al.* Global biodiversity, stoichiometry and ecosystem function responses to human-  
213 induced C–N–P imbalances. *Journal of Plant Physiology* **172**, 82-91 (2015).
- 214 <sup>9</sup> Fisher, J. B. , Badgley, G. & Blyth, E. Global nutrient limitation in terrestrial vegetation. *Glob.*  
215 *Biogeochem. Cycles* **26**, GB3007 (2012).
- 216 <sup>10</sup> Hooper, D. U. & Johnson, L. C. Nitrogen limitation in dryland ecosystems: Responses to  
217 geographical and temporal variation in precipitation. *Biogeochemistry* **46**, 247-293 (1999).
- 218 <sup>11</sup> Hoekstra, J. M. , Boucher, T. M. , Ricketts, T. H. & Roberts, C. Confronting a biome crisis: global  
219 disparities of habitat loss and protection. *Ecol. Lett.* **8**, 23-29 (2005).
- 220 <sup>12</sup> Galloway, J. N. *et al.* The nitrogen cascade. *BioScience* **53**, 341-356 (2003).
- 221 <sup>13</sup> Rockstrom, J. *et al.* A safe operating space for humanity. *Nature* **461**, 472-475 (2009).
- 222 <sup>14</sup> Stevens, C. J. , Dise, N. B. , Mountford, J. O. & Gowing, D. J. Impact of nitrogen deposition on the  
223 species richness of grasslands. *Science* **303**, 1876-1879 (2004).
- 224 <sup>15</sup> Fenn, M. E. *et al.* Ecological effects of nitrogen deposition in the western United States. *BioScience*  
225 **53**, 404-420 (2003).

226 <sup>16</sup> Mahowald, N. *et al.* Global distribution of atmospheric phosphorus sources, concentrations and  
227 deposition rates, and anthropogenic impacts. *Glob. Biogeochem. Cycles* **22**, GB4026  
228 (2008).

229 <sup>17</sup> Phoenix, G. K. *et al.* Impacts of atmospheric nitrogen deposition: responses of multiple plant and  
230 soil parameters across contrasting ecosystems in long-term field experiments. *Glob.*  
231 *Change Biol.* **18**, 1197-1215 (2012).

232 <sup>18</sup> Øgaard, A. F. , Krogstad, T. & Løes, A. K. Potassium uptake by grass from a clay and a silt soil in  
233 relation to soil tests. *Acta Agr Scand B-S P* **51**, 97-105 (2001).

234 <sup>19</sup> Veresoglou, D. S. & Fitter, A. H. Spatial and temporal patterns of growth and nutrient uptake of  
235 five co-existing grasses. *J. Ecol.* **72**, 259-272 (1984).

236 <sup>20</sup> Kayser, M. & Isselstein, J. Potassium cycling and losses in grassland systems: a review. *Grass*  
237 *Forage Sci.* **60**, 213-224 (2005).

238 <sup>21</sup> Borer, E. T. *et al.* Finding generality in ecology: a model for globally distributed experiments.  
239 *Methods Ecol. Evol.* **5**, 65-73 (2014).

240 <sup>22</sup> Guevara, J. C. , Stasi, C. R. , Estevez, O. R. & Le Houerou, H. N. N and P fertilization on rangeland  
241 production in midwest Argentina. *J. Range Manage.* **53**, 410-414 (2000).

242 <sup>23</sup> Clark, C. M. & Tilman, D. Recovery of plant diversity following N cessation: effects of recruitment,  
243 litter, and elevated N cycling. *Ecology* **91**, 3620-3630 (2010).

244 <sup>24</sup> Hedges, L. V. , Gurevitch, J. & Curtis, P. S. The meta-analysis of response ratios in experimental  
245 ecology. *Ecology* **80**, 1150-1156 (1999).

246 <sup>25</sup> Olf, H. & Pegtel, D. Characterisation of the type and extent of nutrient limitation in grassland  
247 vegetation using a bioassay with intact sods. *Plant Soil* **163**, 217-224 (1994).

248 <sup>26</sup> Walker, T. W. & Syers, J. K. The fate of phosphorus during pedogenesis. *Geoderma* **15**, 1-19 (1976).

249 <sup>27</sup> Laliberte, E. *et al.* Experimental assessment of nutrient limitation along a 2-million-year dune  
250 chronosequence in the south-western Australia biodiversity hotspot. *J. Ecol.* **100**, 631-642  
251 (2012).

252 <sup>28</sup> Wedin, D. A. & Tilman, G. D. Influence of nitrogen loading and species composition on the carbon  
253 balance of grasslands. *Science* **274**, 1720-1723 (1996).

254 <sup>29</sup> Jones, L. *et al.* A review and application of the evidence for nitrogen impacts on ecosystem  
255 services. *Ecosystem Services* **7**, 76-88 (2014).

256 <sup>30</sup> Hijmans, R. J. *et al.* Very high resolution interpolated climate surfaces for global land areas. *Int. J.*  
257 *Climatol.* **25**, 1965-1978 (2005).  
258  
259

260 **Supplementary Information** is linked to the online version of the paper at  
261 [www.nature.com/nature](http://www.nature.com/nature).

262 **Acknowledgements** We thank the Minnesota Supercomputer Institute for hosting project data,  
263 the University of Minnesota Institute on the Environment for hosting Nutrient Network  
264 meetings, and each site investigator for funding their site-level operations. Network coordination  
265 and data management were supported by funds from the National Science Foundation Research  
266 Coordination Network (NSF-DEB-1042132) to E.T.B. and E.W.S., from the Long Term  
267 Ecological Research program (NSF-DEB-1234162) to the Cedar Creek LTER, and from the  
268 Institute on the Environment (DG-0001-13). P. Fay acknowledges USDA-NIFA (2010-65615-  
269 20632). USDA is an equal opportunity employer and provider.

270 **Author Contributions** P.A.F. wrote the manuscript, drafted the figures, and led the data  
271 analysis; E.M.L., developed the database; S.M.P. and W.S.H. contributed to data analysis;  
272 S.M.P., W.S.H., J.M.H.K., J.D.B., E.T.B., A.S.M., E.W.S., and P.D.W. contributed conceptual  
273 development and data interpretation. All co-authors contributed data and manuscript editing.


274 This work was generated using data from the Nutrient Network (<http://www.nutnet.org>)  
275 experiment.

276 **Author Information** The data reported in this paper are available through  
277 <http://www.nutnet.org>. Reprints and permissions information is available at  
278 [www.nature.com/reprints](http://www.nature.com/reprints). The authors declare no competing financial interests. Readers are  
279 welcome to comment on the online version of this article at [www.nature.com/nature](http://www.nature.com/nature).

280 Correspondence and requests for materials should be addressed to P.A.F.  
281 ([philip.fay@ars.usda.gov](mailto:philip.fay@ars.usda.gov)).

**Figure 1**


282



283 **Figure 1.** Nutrient limitation of aboveground net primary productivity (ANPP; log response  
 284 ratio,  $\ln[\text{treatment} \cdot \text{control}^{-1}]$ ,) by nitrogen (N), phosphorus (P), and potassium plus year 1  
 285 micronutrients ( $K+\mu$ ). a) Mean  $\pm$  SE over all years available at each site. Means for all sites are  
 286 closed symbols. Means of sites where individual nutrient treatments had significant positive  
 287 effects (defined in panel b) are open symbols. b) Site nutrient limitation of ANPP ranked by the  
 288 magnitude of limitation, averaged over all available years. Vertical dashed lines mark the effect  
 289 sizes approximating a 0.05 significance level for negative and positive nutrient effects on ANPP.  
 290 Note that sites rank in a different order in each nutrient treatment.

291

Figure 2



292

293 **Figure 2.** Correlations of nutrient limitation of aboveground net primary productivity (ANPP);  
294 log response ratios,  $\ln[\text{treatment} \cdot \text{control}^{-1}]$  among single and paired nutrients. a,b,d)

295 Correlations among single nutrients. c,e,f) Correlations of nutrient pairs with the third nutrient.


296 Coefficients of determination ( $R^2$ ) from linear regression ( $0.002 \leq P \leq 0.05$ ).

297

298



Figure 3



299

300

301 **Figure 3.** Predictors of nitrogen (N) limitation of aboveground net primary productivity (ANPP;


302 log response ratio,  $\ln[\text{treatment} \cdot \text{control}^{-1}]$ ). a) site mean annual temperature (MAT), b) site

303 latitude, in degrees away from the equator, and inset) MAT relationship with site latitude. N

304 limitation values are means of five continuous years of nutrient treatment from 19 sites.

305

Figure 4



306 **Figure 4.** Aboveground net primary productivity (ANPP) responses to factorial nitrogen (N),  
307 phosphorus (P) and potassium plus year 1 micronutrient ( $K_{+\mu}$ ) treatments by year in 37  
308 grasslands with three continuous years of ANPP data. Data represent the mean natural log  
309 response ratio of ANPP in the treatment year to ANPP in the pre-treatment year in the same  
310 plot, which controls for plot-to-plot variation.