

1 **Bright spots among the world's coral reefs**

2

3 Joshua E. Cinner^{1,*}, Cindy Huchery¹, M. Aaron MacNeil^{1,2,3}, Nicholas A.J. Graham^{1,4},
4 Tim R. McClanahan⁵, Joseph Maina^{5,6}, Eva Maire^{1,7}, John N. Kittinger^{8,9}, Christina C.
5 Hicks^{1,4,8}, Camilo Mora¹⁰, Edward H. Allison¹¹, Stephanie D'Agata^{5,7,12}, Andrew
6 Hoey¹, David A. Feary¹³, Larry Crowder⁸, Ivor D. Williams¹⁴, Michel Kulbicki¹⁵,
7 Laurent Vigliola¹², Laurent Wantiez¹⁶, Graham Edgar¹⁷, Rick D. Stuart-Smith¹⁷,
8 Stuart A. Sandin¹⁸, Alison L. Green¹⁹, Marah J. Hardt²⁰, Maria Beger⁶, Alan
9 Friedlander^{21,22}, Stuart J. Campbell⁵, Katherine E. Holmes⁵, Shaun K. Wilson^{23,24},
10 Eran Brokovich²⁵, Andrew J. Brooks²⁶, Juan J. Cruz-Motta²⁷, David J. Booth²⁸,
11 Pascale Chabanet²⁹, Charlie Gough³⁰, Mark Tupper³¹, Sebastian C.A. Ferse³², U.
12 Rashid Sumaila³³, David Mouillot^{1,7}

13

14 ¹Australian Research Council Centre of Excellence for Coral Reef Studies, James
15 Cook University, Townsville, QLD 4811 Australia

16 ²Australian Institute of Marine Science, PMB 3 Townsville MC, Townsville, QLD
17 4810 Australia

18 ³Department of Mathematics and Statistics, Dalhousie University, Halifax, NS B3H
19 3J5 Canada

20 ⁴Lancaster Environment Centre, Lancaster University, Lancaster, LA1 4YQ, UK

21 ⁵Wildlife Conservation Society, Global Marine Program, Bronx, NY 10460 USA

22 ⁶Australian Research Council Centre of Excellence for Environmental Decisions,
23 Centre for Biodiversity and Conservation Science, University of Queensland,
24 Brisbane St Lucia QLD 4074 Australia

25 ⁷MARBEC, UMR IRD-CNRS-UM-IFREMER 9190, Université Montpellier, 34095
26 Montpellier Cedex, France

27 ⁸Center for Ocean Solutions, Stanford University, CA 94305 USA

28 ⁹Conservation International Hawaii, Betty and Gordon Moore Center for Science and
29 Oceans, 7192 Kalanianaʻole Hwy, Suite G230, Honolulu, Hawaiʻi 96825 USA

30 ¹⁰Department of Geography, University of Hawaiʻi at Manoa, Honolulu, Hawaiʻi
31 96822 USA

32 ¹¹School of Marine and Environmental Affairs, University of Washington, Seattle,
33 WA 98102 USA

34 ¹²Institut de Recherche pour le Développement, UMR IRD-UR-CNRS ENTROPIE,
35 Laboratoire d'Excellence LABEX CORAIL, BP A5, 98848 Nouméa Cedex, New
36 Caledonia

37 ¹³Ecology & Evolution Group, School of Life Sciences, University Park, University
38 of Nottingham, Nottingham NG7 2RD, UK

39 ¹⁴Coral Reef Ecosystems Division, NOAA Pacific Islands Fisheries Science Center,
40 Honolulu, HI 96818 USA

41 ¹⁵UMR Entropie, Labex Corail, –IRD, Université de Perpignan, 66000, Perpignan,
42 France

43 ¹⁶EA4243 LIVE, University of New Caledonia, BPR4 98851 Noumea cedex, New
44 Caledonia

45 ¹⁷Institute for Marine and Antarctic Studies, University of Tasmania, Hobart,
46 Tasmania, 7001 Australia

47 ¹⁸Scripps Institution of Oceanography, University of California, San Diego, La Jolla,
48 CA 92093 USA

49 ¹⁹The Nature Conservancy, Brisbane, Australia

50 ²⁰Future of Fish, 7315 Wisconsin Ave, Suite 1000W, Bethesda, MD 20814, USA

51 ²¹Fisheries Ecology Research Lab, Department of Biology, University of Hawaii,
52 Honolulu, HI 96822, USA

53 ²²National Geographic Society, Pristine Seas Program, 1145 17th Street N.W.
54 Washington, D.C. 20036-4688, USA

55 ²³Department of Parks and Wildlife, Kensington, Perth WA 6151 Australia

56 ²⁴Oceans Institute, University of Western Australia, Crawley, WA 6009, Australia

57 ²⁵The Israeli Society of Ecology and Environmental Sciences, Kehilat New York 19
58 Tel Aviv, Israel

59 ²⁶Marine Science Institute, University of California, Santa Barbara, CA 93106-6150,
60 USA

61 ²⁷Departamento de Ciencias Marinas., Recinto Universitario de Mayaguez,
62 Universidad de Puerto Rico, 00680, Puerto Rico

63 ²⁸School of Life Sciences, University of Technology Sydney 2007 Australia

64 ²⁹UMR ENTROPIE, Laboratoire d'Excellence LABEX CORAIL, Institut de
65 Recherche pour le Développement, CS 41095, 97495 Sainte Clotilde, La Réunion
66 (FR)

67 ³⁰Blue Ventures Conservation, 39-41 North Road, London N7 9DP, United Kingdom

68 ³¹Coastal Resources Association, St. Joseph St., Brgy. Nonoc, Surigao City, Surigao
69 del Norte 8400, Philippines

70 ³²Leibniz Centre for Tropical Marine Ecology (ZMT), Fahrenheitstrasse 6, D-28359
71 Bremen, Germany

72 ³³Fisheries Economics Research Unit, University of British Columbia, 2202 Main
73 Mall, Vancouver, B.C., V6T 1Z4, Canada

74

75 *Correspondence to: Joshua.cinner@jcu.edu.au

76

77 **Ongoing declines among the world's coral reefs^{1,2} require novel approaches to**
78 **sustain these ecosystems and the millions of people who depend on them³. A**
79 **presently untapped approach that draws on theory and practice in human health**
80 **and rural development^{4,5} is systematically identifying and learning from the**
81 **'outliers'- places where ecosystems are substantially better ('bright spots') or**
82 **worse ('dark spots') than expected, given the environmental conditions and**
83 **socioeconomic drivers they are exposed to. Here, we compile data from more**
84 **than 2,500 reefs worldwide and develop a Bayesian hierarchical model to**
85 **generate expectations of how standing stocks of reef fish biomass are related to**
86 **18 socioeconomic drivers and environmental conditions. We then identified 15**
87 **bright spots and 35 dark spots among our global survey of coral reefs, defined as**
88 **sites that had biomass levels more than two standard deviations from**
89 **expectations. Importantly, bright spots were not simply comprised of remote**
90 **areas with low fishing pressure- they include localities where human populations**
91 **and use of ecosystem resources is high, potentially providing novel insights into**
92 **how communities have successfully confronted strong drivers of change.**
93 **Alternatively, dark spots were not necessarily the sites with the lowest absolute**
94 **biomass and even included some remote, uninhabited locations often considered**
95 **near-pristine⁶. We surveyed local experts about social, institutional, and**
96 **environmental conditions at these sites to reveal that bright spots were**
97 **characterised by strong sociocultural institutions such as customary taboos and**
98 **marine tenure, high levels of local engagement in management, high dependence**
99 **on marine resources, and beneficial environmental conditions such as deep-**
100 **water refuges. Alternatively, dark spots were characterised by intensive capture**
101 **and storage technology and a recent history of environmental shocks. Our**

102 **results suggest that investments in strengthening fisheries governance,**
103 **particularly aspects such as participation and property rights, could facilitate**
104 **innovative conservation actions that help communities defy expectations of**
105 **global reef degradation.**

106

107 *Main text*

108 Despite substantial international conservation efforts, many of the world's ecosystems
109 continue to decline^{1,7}. Most conservation approaches aim to identify and protect
110 places of high ecological integrity under minimal threat⁸. Yet, with escalating social
111 and environmental drivers of change, conservation actions are also needed where
112 people and nature coexist, especially where human impacts are already severe⁹. Here,
113 we highlight an approach for implementing conservation in coupled human-natural
114 systems focused on identifying and learning from outliers - places that are performing
115 substantially better than expected, given the socioeconomic and environmental
116 conditions they are exposed to. By their very nature, outliers deviate from
117 expectations, and consequently can provide novel insights on confronting complex
118 problems where conventional solutions have failed. This type of positive deviance, or
119 'bright spot' analysis has been used in fields such as business, health, and human
120 development to uncover local actions and governance systems that work in the
121 context of widespread failure^{10,11}, and holds much promise in informing conservation.

122

123 To demonstrate this approach, we compiled data from 2,514 coral reefs in 46
124 countries, states, and territories (hereafter 'nation/states') and developed a Bayesian
125 hierarchical model to generate expected conditions of how standing reef fish biomass
126 (a key indicator of resource availability and ecosystem functions¹²) was related to 18
127 key environmental variables and socioeconomic drivers (Box 1; Extended Data
128 Tables 1,2; Methods). A key and significant finding from our global analysis is that
129 the size and accessibility of the nearest market, more so than local or national
130 population pressure, management, environmental conditions, or national
131 socioeconomic context, was the strongest driver of reef fish biomass globally (Box 1).

132

133 Next, we identified 15 ‘bright spots’ and 35 ‘dark spots’ among the world’s coral reefs,
134 defined as sites with biomass levels more than two standard deviations higher or
135 lower than expectations from our global model, respectively (Fig. 1; Methods;
136 Extended Data Table 3). Rather than simply identifying places in the best or worst
137 condition, our bright spots approach reveals the places that most strongly defy
138 expectations. Using them to inform the conservation discourse will certainly
139 challenge established ideas of where and how conservation efforts should be focused.
140 For example, remote places far from human impacts are conventionally considered
141 near-pristine areas of high conservation value⁶, yet most of the bright spots we
142 identified occur in fished, populated areas (Extended Data Table 3), some with
143 biomass values below the global average. Alternatively, some remote places such as
144 parts of the NW Hawaiian Islands underperform (i.e. were identified as dark spots).

145

146 Detailed analysis of why bright spots can evade the fate of similar areas facing
147 equivalent stresses will require a new research agenda gathering detailed site-level
148 information on social and institutional conditions, technological innovations, external
149 influences, and ecological processes¹³ that are simply not available in a global-scale
150 analysis. As a preliminary hypothesis-generating exercise to begin uncovering why
151 bright and dark spots may diverge from expectations, we surveyed data providers and
152 other experts about the presence or absence of 10 key social and environmental
153 conditions at the 15 bright spots, 35 dark spots, and 14 average sites with biomass
154 values closest to model expectations (see Methods for details). Our survey revealed
155 that bright spots were more likely to have high levels of local engagement in the
156 management process, high dependence on coastal resources, and the presence of

157 sociocultural governance institutions such as customary tenure or taboos (Fig. 2,
158 Methods). For example, in one bright spot, Karkar Island, Papua New Guinea,
159 resource use is restricted through an adaptive rotational harvest system based on
160 ecological feedbacks, marine tenure that allows for the exclusion of fishers from
161 outside the local village, and initiation rights that limit individuals' entry into certain
162 fisheries¹⁴. Bright spots were also generally proximate to deep water, which may help
163 provide a refuge from disturbance for corals and fish¹⁵ (Fig. 2, Extended Data Fig. 6).
164 Conversely, dark spots were distinguished by having fishing technologies allowing
165 for more intensive exploitation, such as fish freezers and potentially destructive
166 netting, as well as a recent history of environmental shocks (*e.g.* coral bleaching or
167 cyclone; Fig. 2). The latter is particularly worrisome in the context of climate change,
168 which is likely to lead to increased coral bleaching and more intense cyclones¹⁶.
169
170 Our global analyses highlight two novel opportunities to inform coral reef governance.
171 The first is to use bright spots as agents of change to expand the conservation
172 discourse from the current focus on protecting places under minimal threat⁸, toward
173 harnessing lessons from places that have successfully confronted high pressures.
174 Our bright spots approach can be used to inform the types of investments and
175 governance structures that may help to create more sustainable pathways for impacted
176 coral reefs. Specifically, our initial investigation highlights how investments that
177 strengthen fisheries governance, particularly issues such as participation and property
178 rights, could help communities to innovate in ways that allow them to defy
179 expectations. Conversely, the more typical efforts to provide capture and storage
180 infrastructure, particularly where there are environmental shocks and local-scale
181 governance is weak, may lead to social-ecological traps¹⁷ that reinforce resource

182 degradation beyond expectations. Effectively harnessing the potential to learn from
183 both bright and dark spots will require scientists to increase research efforts in these
184 places, NGOs to catalyze lessons from other areas, donors to start investing in novel
185 solutions, and policy makers to ensure that governance structures foster flexible
186 learning and experimentation. Indeed, both bright and dark spots may have much to
187 offer in terms of how to creatively confront drivers of change, identify the paths to
188 avoid and those offering novel management solutions, and prioritizing conservation
189 actions. Critically, the bright spots we identified span the development spectrum from
190 low (Solomon Islands and Papua New Guinea) to high (territories of the USA and
191 UK; Fig. 1) income, showing that lessons about effective reef management can
192 emerge from diverse places.

193

194 A second opportunity stems from a renewed focus on managing the socioeconomic
195 drivers that shape reef conditions. Many social drivers are amenable to governance
196 interventions, and our comprehensive analysis (Box 1) shows how an increased policy
197 focus on social drivers such as markets and development could result in
198 improvements to reef fish biomass. For example, given the important influence of
199 markets in our analysis, reef managers, donor organisations, conservation groups, and
200 coastal communities could improve sustainability by developing interventions that
201 dampen the negative influence of markets on reef systems. A portfolio of market
202 interventions, including eco-labelling and sustainable harvesting certifications,
203 fisheries improvement projects, and value chain interventions have been developed
204 within large-scale industrial fisheries to increase access to markets for seafood that is
205 sourced sustainably²¹⁻²³. Although there is considerable scope for adapting these
206 interventions to artisanal coral reef fisheries in both local and regional markets,


207 effectively dampening the negative influence of markets may also require developing
208 novel interventions that address the range of ways in which markets can lead to
209 overexploitation. Existing research suggests that markets create incentives for
210 overexploitation not only by affecting price and price variability for reef products¹⁸, ,
211 but also by influencing people's behavior¹⁹, including their willingness to cooperate in
212 the collective management of natural resources²⁰.

213

214 The long-term viability of coral reefs will ultimately depend on international action to
215 reduce carbon emissions¹⁶. However, fisheries remain a pervasive source of reef
216 degradation, and effective local-level fisheries governance is crucial to sustaining
217 ecological processes that give reefs the best chance of coping with global
218 environmental change²⁵. Seeking out and learning from bright spots has uncovered
219 novel solutions in fields as diverse as human health, development, and business^{10,11},
220 and this approach may offer insights into confronting the complex governance
221 problems facing coupled human-natural systems such as coral reefs.

222


223 **Figures**



224

225 **Figure 1 | Bright and dark spots among the world's coral reefs.** (a) Each site's deviation from expected biomass (y-axis) along a gradient of
 226 nation/state mean biomass (x-axis). Sites with biomass values >2 standard deviations above or below expected values were considered bright and
 227 dark spots, respectively. The 15 bright and 35 dark spots are indicated with yellow and black dots respectively. Each grey vertical line represents

228 a nation/state in our analysis. Nation/states with bright or dark spots are labelled and numbered, corresponding to the numbers in panel b. There
229 can be multiple bright or dark spots in each nation/state, thus the 50 bright and dark spots are distributed among 17 nation/states. As a
230 conservative precaution, we did not consider a site a bright or dark spot if there were fewer than 5 sites sampled in a nation/state (Methods);
231 consequently there is one site with biomass levels lower than 2 SD below expectations that is not labelled as a dark spot. BIOT= British Indian
232 Ocean Territory (Chagos); PNG= Papua New Guinea; CNMI= Commonwealth of the Northern Mariana Islands; NWHI= Northwest Hawaiian
233 Islands; PRIA= Pacific Remote Island Areas. (b) Map highlighting bright spots and dark spots with large circles, and other sites in small circles.
234 Bright spots are mostly concentrated on islands of the Pacific and Southeast Asia, while dark spots are spread among every major tropical ocean
235 basin.



236

237

Figure 2 | Differences in social and environmental conditions between bright

238

spots, dark spots, and ‘average’ sites. *=p<0.05, **=p<0.01, *=p<0.001. P**

239


values are determined using Fisher’s Exact test. Intensive netting includes beach seine

240

nets, surround gill nets, and muro-ami.

Box 1

Drawing on a broad body of theoretical and empirical research in the social sciences^{24,26,27} and ecology^{2,6,28} on coupled human-natural systems, we quantified how reef fish biomass (panel a) was related to distal social drivers such as markets, affluence, governance, and population (panels b,c), while controlling for well-known environmental conditions such as depth, habitat, and productivity (panel d) (Extended Data Table 1, Methods). In contrast to many global studies of reef systems that are focused on demonstrating the severity of human impacts⁶, our examination seeks to uncover potential policy levers by highlighting the relative role of specific social drivers. Critically, the strongest driver of reef fish biomass (*i.e.* the largest standardized effect size) was our metric of potential interactions with urban centres, called market gravity²⁹ (Extended Data Fig. 1, 2, 3; Methods). Specifically, we found that reef fish biomass decreased as the size and accessibility of markets increased (Extended Data Fig. 2b, and Extended Data Fig. 3). Somewhat counter-intuitively, fish biomass was higher in places with high local human population growth rates, likely reflecting human migration to areas of better environmental quality³⁰ -a phenomenon that could result in increased degradation at these sites over time. We found a strong positive, but less certain relationship (*i.e.* a high standardized effect size, with >75% of the posterior distribution above zero) with the Human Development Index, meaning that reefs tended to be in better condition in wealthier nations/states (panel c). Our analysis also confirmed the role that marine reserves can play in sustaining biomass on coral reefs, but only when compliance is high (panel b), reinforcing the importance of fostering compliance for reserves to be successful.



Global patterns and drivers of reef fish biomass. (a) Reef fish biomass [in (log)kg/ha] among 918 study sites across 46 nations/states. For illustration purposes and to avoid the overlap of sites in a global map, we display sites as points that vary in size and colour proportional to amount of fish biomass, with small, red dots indicating low fish biomass and large, green dots indicating high biomass. b-d) Standardised effect size of local scale social drivers, nation/state scale social drivers, and environmental covariates, respectively. Parameter estimates are Bayesian posterior median values, 95% uncertainty intervals (UI; thin lines), and 50% UI (thick lines). Black dots indicate that the 95% UI does not overlap 0; Grey closed circles indicates that 75% of the posterior distribution lies to one side of 0; and grey open circles indicate that the 50% UI overlaps 0.

241 **Methods**

242

243 Scales of data

244 Our data were organized at three spatial scales: reef (n=2514), site (n=918), and
245 nation/state (n=46).

246 i) reef (the smallest scale, which had an average of 2.4 surveys/transects -
247 hereafter 'reef').

248 ii) site (a cluster of reefs). We clustered reefs together that were within 4km
249 of each other, and used the centroid of these clusters (hereafter 'sites') to
250 estimate site-level social and site-level environmental covariates
251 (Extended Data Table 1). To make these clusters, we first estimated the
252 linear distance between all reefs, then used a hierarchical analysis with the
253 complete-linkage clustering technique based on the maximum distance
254 between reefs. We set the cut-off at 4km to select mutually exclusive sites
255 where reefs cannot be more distant than 4km. The choice of 4km was
256 informed by a 3-year study of the spatial movement patterns of artisanal
257 coral reef fishers, corresponding to the highest density of fishing activities
258 on reefs based on GPS-derived effort density maps of artisanal coral reef
259 fishing activities³¹. This clustering analysis was carried out using the R
260 functions 'hclust' and 'cutree', resulting in an average of 2.7 reefs/site.

261 iii) Nation/state (nation, state, or territory). A larger scale in our analysis was
262 'nation/state', which are jurisdictions that generally correspond to
263 individual nations (but could also include states, territories, overseas
264 regions, or extremely remote areas within a state such as the northwest

265 Hawaiian Islands; Extended Data Table 2), within which sites and reefs
266 were nested for analysis.

267

268 Estimating Biomass

269 Reef fish biomass can reflect a broad selection of reef fish functioning and benthic
270 conditions^{12,32-34}, and is a key metric of resource availability for reef fisheries. Reef
271 fish biomass estimates were based on instantaneous visual counts from 6,088 surveys
272 collected from 2,514 reefs. All surveys used standard belt-transects, distance sampling,
273 or point-counts, and were conducted between 2004 and 2013. Where data from
274 multiple years were available from a single reef, we included only data from the year
275 closest to 2010. Within each survey area, reef associated fishes were identified to
276 species level, abundance counted, and total length (TL) estimated, with the exception
277 of one data provider who measured biomass at the family level. To make estimates of
278 biomass from these transect-level data comparable among studies, we:

279 i) Retained families that were consistently studied and were above a
280 minimum size cut-off. Thus, we retained counts of >10cm diurnally-active,
281 non-cryptic reef fish that are resident on the reef (20 families, 774 species),
282 excluding sharks and semi-pelagic species (Extended Data Table 4). We
283 also excluded three groups of fishes that are strongly associated with coral
284 habitat conditions and are rarely targets for fisheries (Anthiinae,
285 Chaetodontidae, and Cirrhitidae). We calculated total biomass of fishes on
286 each reef using standard published species-level length-weight relationship
287 parameters or those available on FishBase³⁵. When length-weight
288 relationship parameters were not available for a species, we used the
289 parameters for a closely related species or genus.

- 290 ii) Directly accounted for depth and habitat as covariates in the model (see
291 “environmental conditions” section below);
292 iii) Accounted for any potential bias among data providers (capturing
293 information on both inter-observer differences, and census methods) by
294 including each data provider as a random effect in our model.

295

296 Biomass means, medians, and standard deviations were calculated at the reef-scale.

297 All reported log values are the natural log.

298

299 Social Drivers

300 *1. Local Population Growth:* We created a 100km buffer around each site and used
301 this to calculate human population within the buffer in 2000 and 2010 based on the
302 Socioeconomic Data and Application Centre (SEDAC) gridded population of the
303 world database³⁶. Population growth was the proportional difference between the
304 population in 2000 and 2010. We chose a 100km buffer as a reasonable range at
305 which many key human impacts from population (e.g., land-use and nutrients) might
306 affect reefs³⁷.

307

308 *2. Management:* For each site, we determined if it was: i) unfished- whether it fell
309 within the borders of a no-take marine reserve. We asked data providers to further
310 classify whether the reserve had high or low levels of compliance; ii) restricted -
311 whether there were active restrictions on gears (e.g. bans on the use of nets, spearguns,
312 or traps) or fishing effort (which could have included areas inside marine parks that
313 were not necessarily no take); or iii) fished - regularly fished without effective

314 restrictions. To determine these classifications, we used the expert opinion of the data
315 providers, and triangulated this with a global database of marine reserve boundaries³⁸.

316

317 3. *Gravity*: We adapted the economic geography concept of *gravity*, also called
318 interactance³⁹, to examine potential interactions between reefs and: i) major urban
319 centres/markets (defined as provincial capital cities, major population centres,
320 landmark cities, national capitals, and ports); and ii) the nearest human settlements
321 (Extended Data Fig. 1). This application of the gravity concept infers that potential
322 interactions increase with population size, but decay exponentially with the effective
323 distance between two points. Thus, we gathered data on both population estimates and
324 a surrogate for distance: travel time.

325

326 *Population estimations*

327 We gathered population estimates for: 1) the nearest major markets (which
328 includes national capitals, provincial capitals, major population centres, ports,
329 and landmark cities) using the World Cities base map from ESRITM; and 2) the
330 nearest human settlement within a 500km radius using LandScanTM 2011
331 database. The different datasets were required because the latter is available in
332 raster format while the former is available as point data. We chose a 500km
333 radius from the nearest settlement as the maximum distance any non-market
334 fishing activities for fresh reef fish are likely to occur.

335

336 *Travel time calculation*

337 Travel time was computed using a cost-distance algorithm that computes the
338 least 'cost' (in minutes) of travelling between two locations on a regular raster

339 grid. In our case, the two locations were either: 1) the centroid of the site (i.e.
340 reef cluster) and the nearest settlement, or 2) the centroid of the site and the
341 major market. The cost (i.e. time) of travelling between the two locations was
342 determined by using a raster grid of land cover and road networks with the
343 cells containing values that represent the time required to travel across them⁴⁰
344 (Extended Data Table 5), we termed this raster grid a *friction-surface* (with the
345 time required to travel across different types of surfaces analogous to different
346 levels of friction). To develop the friction-surface, we used global datasets of
347 road networks, land cover, and shorelines:

- 348 - Road network data was extracted from the Vector Map Level 0
349 (VMap0) from the National Imagery and Mapping Agency's (NIMA)
350 Digital Chart of the World (DCW®). We converted vector data from
351 VMap0 to 1km resolution raster.
- 352 - Land cover data were extracted from the Global Land Cover 2000⁴¹.
- 353 -To define the shorelines, we used the GSHHS (Global Self-consistent,
354 Hierarchical, High-resolution Shoreline) database version 2.2.2.

355

356 These three friction components (road networks, land cover, and water bodies)
357 were combined into a single friction surface with a Behrmann map projection.
358 We calculated our cost-distance models in R⁴² using the *accCost* function of
359 the '*gdistance*' package. The function uses Dijkstra's algorithm to calculate
360 least-cost distance between two cells on the grid and the associated distance
361 taking into account obstacles and the local friction of the landscape⁴³. Travel
362 time estimates over a particular surface could be affected by the infrastructure
363 (e.g. road quality) and types of technology used (e.g. types of boats). These

364 types of data were not available at a global scale but could be important
365 modifications in more localised studies.

366

367 *Gravity computation*

368 i) To compute the gravity to the nearest market, we calculated the population
369 of the nearest major market and divided that by the squared travel time
370 between the market and the site. Although other exponents can be used⁴⁴, we
371 used the squared distance (or in our case, travel time), which is relatively
372 common in geography and economics. This decay function could be
373 influenced by local considerations, such as infrastructure quality (e.g. roads),
374 the types of transport technology (i.e. vessels being used), and fuel prices,
375 which were not available in a comparable format for this global analysis, but
376 could be important considerations in more localised adaptations of this study.

377 ii) To determine the gravity of the nearest settlement, we located the nearest
378 populated pixel within 500kms, determined the population of that pixel, and
379 divided that by the squared travel time between that cell and the reef site.

380 As is standard practice in many agricultural economics studies⁴⁵, an assumption in
381 our study is that the nearest major capital or landmark city represents a market.

382 Ideally we would have used a global database of all local and regional markets for
383 coral reef fish, but this type of database is not available at a global scale. As a
384 sensitivity analysis to help justify our assumption that capital and landmark cities
385 were a reasonable proxy for reef fish markets, we tested a series of candidate
386 models that predicted biomass based on: 1) cumulative gravity of all cities within
387 500km; 2) gravity of the nearest city; 3) travel time to the nearest city; 4)
388 population of the nearest city; 5) gravity to the nearest human population above 40

389 people/km² (assumed to be a small peri-urban area and potential local market); 6)
390 the travel time between the reef and a small peri-urban area; 7) the population size
391 of the small peri-urban population; 8) gravity to the nearest human population
392 above 75 people/km² (assumed to be a large peri-urban area and potential market);
393 9) the travel time between the reef and this large peri-urban population; 10) the
394 population size of this large peri-urban population; and 11) the total population
395 size within a 500km radius. Model selection revealed that the best two models
396 were gravity of the nearest city and gravity of all cities within 500km (with a 3
397 AIC value difference between them; Extended Data Table 6). Importantly, when
398 looking at the individual components of gravity models, the travel time
399 components all had a much lower AIC value than the population components,
400 which is broadly consistent with previous systematic review studies⁴⁶. Similarly,
401 travel time to the nearest city had a lower AIC score than any aspect of either the
402 peri-urban or urban measures. This suggests our use of capital and landmark cities
403 is likely to better capture exploitation drivers from markets rather than simple
404 population pressures. This may be because market dynamics are difficult to
405 capture by population threshold estimates; for example some small provincial
406 capitals where fish markets are located have very low population densities, while
407 some larger population centres may not have a market. Downscaled regional or
408 local analyses could attempt to use more detailed knowledge about fish markets,
409 but we used the best proxy available at a global scale.

410

411 *4. Human Development Index (HDI):* HDI is a summary measure of human
412 development encompassing: a long and healthy life, being knowledgeable, and having

413 a decent standard of living. In cases where HDI values were not available specific to
414 the State (e.g. Florida and Hawaii), we used the national (e.g. USA) HDI value.

415

416 *5. Population Size:* For each Nation/state, we determined the size of the human
417 population. Data were derived mainly from census reports, the CIA fact book, and
418 Wikipedia.

419

420 *6. Tourism:* We examined tourist arrivals relative to the nation/state population size
421 (above). Tourism arrivals were gathered primarily from the World Tourism
422 Organization's Compendium of Tourism Statistics.

423

424 *7. National Reef Fish Landings:* Catch data were obtained from the Sea Around Us
425 Project (SAUP) catch database (www.searoundus.org), except for Florida, which
426 was not reported separately in the database. We identified 200 reef fish species and
427 taxon groups in the SAUP catch database⁴⁷. Note that reef-associated pelagics such as
428 scombrids and carangids normally form part of reef fish catches. However, we chose
429 not to include these species because they are also targeted and caught in large
430 amounts by large-scale, non-reef operations.

431

432 *8. Voice and Accountability:* This metric, from the World Bank survey on governance,
433 reflects the perceptions of the extent to which a country's citizens are able to
434 participate in selecting their government, as well as freedom of expression, freedom
435 of association, and a free media. In cases where governance values were not available
436 specific to the Nation/state (e.g. Florida and Hawaii), we used national (e.g. USA)
437 values.

438

439 Environmental Drivers

440 *1. Depth:* The depth of reef surveys were grouped into the following categories: <4m,
441 4-10m, >10m to account for broad differences in reef fish community structure
442 attributable to a number of inter-linked depth-related factors. Categories were
443 necessary to standardise methods used by data providers and were determined by pre-
444 existing categories used by several data providers.

445

446 *2. Habitat:* We included the following habitat categories: i) Slope: The reef slope
447 habitat is typically on the ocean side of a reef, where the reef slopes down into deeper
448 water; ii) Crest: The reef crest habitat is the section that joins a reef slope to the reef
449 flat. The zone is typified by high wave energy (i.e. where the waves break). It is also
450 typified by a change in the angle of the reef from an inclined slope to a horizontal reef
451 flat; iii) Flat: The reef flat habitat is typically horizontal and extends back from the
452 reef crest for 10's to 100's of metres; iv) Lagoon / back reef: Lagoonal reef habitats
453 are where the continuous reef flat breaks up into more patchy reef environments
454 sheltered from wave energy. These habitats can be behind barrier / fringing reefs or
455 within atolls. Back reef habitats are similar broken habitats where the wave energy
456 does not typically reach the reefs and thus forms a less continuous 'lagoon style' reef
457 habitat. Due to minimal representation among our sample, we excluded other less
458 prevalent habitat types, such as channels and banks. To verify the sites' habitat
459 information, we used the Millennium Coral Reef Mapping Project (MCRMP)
460 hierarchical data⁴⁸, Google Earth, and site depth information.

461

462 3. *Productivity*: We examined ocean productivity for each of our sites in mg C / m² /
463 day (<http://www.science.oregonstate.edu/ocean.productivity/>). Using the monthly data
464 for years 2005 to 2010 (in hdf format), we imported and converted those data into
465 ArcGIS. We then calculated yearly average and finally an average for all these years.
466 We used a 100km buffer around each of our sites and examined the average
467 productivity within that radius. Note that ocean productivity estimates are less
468 accurate for nearshore environments, but we used the best available data.

469

470 Analyses

471 We first looked for collinearity among our covariates using bivariate correlations and
472 variance inflation factor estimates (Extended Data Fig. 4, Extended Data Table 7).
473 This led to the exclusion of several covariates (not described above): i) *Geographic*
474 *Basin* (Tropical Atlantic, western Indo-Pacific, Central Indo-Pacific, or eastern Indo-
475 Pacific); ii) *Gross Domestic Product* (purchasing power parity); iii) *Rule of Law*
476 (World Bank governance index); iv) *Control of Corruption* (World Bank governance
477 index); and v) *Sedimentation*. Additionally, we removed an index of climate stress,
478 developed by Maina et al.⁴⁹, which incorporated 11 different environmental
479 conditions, such as the mean and variability of sea surface temperature due to
480 repeated lack of convergence for this parameter in the model, likely indicative of
481 unidentified multi-collinearity. All other covariates had correlation coefficients 0.7 or
482 less and Variance Inflation Factor scores less than 5 (indicating multicollinearity was
483 not a serious concern). Care must be taken in causal attribution of covariates that were
484 significant in our model, but demonstrated colinearity with candidate covariates that
485 were removed during the aforementioned process. Importantly, the covariate that

486 exhibited the largest effect size in our model, market gravity, was not strongly
487 collinear with other candidate covariates.

488

489 To quantify the multi-scale social, environmental, and economic factors affecting reef
490 fish biomass we adopted a Bayesian hierarchical modelling approach that explicitly
491 recognized the three scales of spatial organization: reef (j), site (k), and nation/state (s).

492

493 In adopting the Bayesian approach we developed two models for inference: a null
494 model, consisting only of the hierarchical units of observation (i.e. intercepts-only)
495 and a full model that included all of our covariates (drivers) of interest. Covariates
496 were entered into the model at the relevant scale, leading to a hierarchical model
497 whereby lower-level intercepts (averages) were placed in the context of higher-level
498 covariates in which they were nested. We used the null model as a baseline against
499 which we could ensure that our full model performed better than a model with no
500 covariate information. We did not remove 'non-significant' covariates from the model
501 because each covariate was carefully considered for inclusion and could therefore
502 reasonably be considered as having an effect, even if small or uncertain; removing
503 factors from the model is equivalent to fixing parameter estimates at exactly zero - a
504 highly-subjective modelling decision after covariates have already been selected as
505 potentially important⁵⁰.

506

507 The full model assumed the observed, environmental-scale observations of fish
508 biomass (y_{ijks}) were modelled using a noncentral-T distribution, allowing for fatter
509 tails than typical log-normal models of reef fish biomass³².

510

$$\log(y_{ijks}) \sim \text{Noncentral}T(\mu_{ijks}, \tau_{reef}, 3.5)$$

$$\mu_{ijks} = \beta_{0jks} + \beta_{reef} X_{reef}$$

$$\tau_{reef} \sim U(0, 100)^{-2}$$

511

512 with X_{reef} representing the matrix of observed environmental-scale covariates and
 513 β_{reef} the array of estimated reef-scale parameters. The τ_{reef} (and all subsequent τ 's)
 514 were assumed common across observations in the final model and were minimally
 515 informative⁵⁰. Using a similar structure, the environmental-scale intercepts (β_{0jks})
 516 were structured as a function of site-scale covariates (X_{sit}):

517

$$\beta_{0jks} \sim N(\mu_{jks}, \tau_{sit})$$

$$\mu_{jks} = \gamma_{0ks} + \gamma_{sit} X_{sit}$$

$$\tau_{sit} \sim U(0, 100)^{-2}$$

518

519 with γ_{sit} representing an array of site-scale parameters. Building upon the hierarchy,
 520 the site-scale intercepts (γ_{0ks}) were structured as a function of state-scale covariates
 521 (X_{sta}):

522

$$\gamma_{0ks} \sim N(\mu_{ks}, \tau_{sta})$$

$$\mu_{ks} = \gamma_{0s} + \gamma_{sta} X_{sta}$$

$$\tau_{sta} \sim U(0, 100)^{-2}$$

523

524 Finally, at the top scale of the analysis we allowed for a global (overall) estimate of
 525 average log-biomass (μ_0):

526

$$\gamma_{0s} \sim N(\mu_0, \tau_{glo})$$

$$\mu_0 \sim N(0.0, 1000)$$

527 $\tau_{glo} \sim U(0, 100)^{-2}$.

528

529 The relationships between fish biomass and environmental, site, and state scale
530 drivers was carried out using the PyMC package⁵¹ for the Python programming
531 language, using a Metropolis-Hastings (MH) sampler run for 10^6 iterations, with a
532 900,000 iteration burn in, leaving 10,000 samples in the posterior distribution of each
533 parameter; these long burn-in times are often required with a complex model using
534 the MH algorithm. Convergence was monitored by examining posterior chains and
535 distributions for stability and by running multiple chains from different starting points
536 and checking for convergence using Gelman-Rubin statistics⁵² for parameters across
537 multiple chains; all were at or close to 1, indicating good convergence of parameters
538 across multiple chains.

539

540 *Overall model fit*

541

542 We conducted posterior predictive checks for goodness of fit (GoF) using Bayesian p-
543 values⁴⁰ (BpV), whereby fit was assessed by the discrepancy between observed or
544 simulated data and their expected values. To do this we simulated new data (y_i^{new}) by
545 sampling from the joint posterior of our model (θ) and calculated the Freeman-Tukey
546 measure of discrepancy for the observed (y_i^{obs}) or simulated data, given their expected
547 values (μ_i):

548

549
$$D(y|\theta) = \sum_i (\sqrt{y_i} - \sqrt{\mu_i})^2$$

550

551 yielding two arrays of median discrepancies $D(y^{obs}/\theta)$ and $D(y^{new}/\theta)$ that were then
552 used to calculate a BpV for our model by recording the proportion of times $D(y^{obs}/\theta)$
553 was greater than $D(y^{new}/\theta)$ (Extended Data Fig. 5). A BpV above 0.975 or under 0.025
554 provides substantial evidence for lack of model fit. Evaluated by the Deviance
555 Information Criterion (DIC), the full model greatly outperformed the null model
556 ($\Delta DIC=472$).

557

558 To examine homoscedasticity, we checked residuals against fitted values. We also
559 checked the residuals against all covariates included in the model, and several
560 covariates that were not included in the model (primarily due to collinearity),
561 including: 1) *Atoll* - A binary metric of whether the reef was on an atoll or not; 2)
562 *Control of Corruption*: Perceptions of the extent to which public power is exercised
563 for private gain, including both petty and grand forms of corruption, as well as
564 'capture' of the state by elites and private interests. Derived from the World Bank
565 survey on governance; 3) *Geographic Basin*- whether the site was in the Tropical
566 Atlantic, western Indo-Pacific, Central Indo-Pacific, or eastern Indo-Pacific; 4)
567 *Connectivity* – we examined 3 measures based on the area of coral reef within a 30km,
568 100km, and 600km radius of the site; 5) *Sedimentation*; 6) *Coral Cover* (which was
569 only available for a subset of the sites); 7) *Climate stress*⁴⁹; and 8) *Census method*.
570 The model residuals showed no patterns with these eight additional covariates,
571 suggesting they would not explain additional information in our model.

572

573 *Bright and dark spot estimates*

574 Because the performance of site scale locations are of substantial interest in

575 uncovering novel solutions for reef conservation, we defined bright and dark spots at
576 the site scale. To this end, we defined bright (or dark) spots as locations where
577 expected site-scale intercepts (γ_{0ks}) differed by more than two standard deviations
578 from their nation/state-scale expected value (μ_{ks}), given all the covariates present in
579 the full hierarchical model:

$$580 \quad SS_{spot} = |(\mu_{ks} - \gamma_{0ks})| > 2[SD(\mu_{ks} - \gamma_{0ks})].$$

581 This, in effect, probabilistically identified the most deviant sites, given the model,
582 while shrinking sites toward their group-level means, thereby allowing us to
583 overcome potential bias due to low and varying sample sizes that can lead to extreme
584 values from chance alone. As a conservative precaution, we did not consider a site a
585 bright or dark spot if the group-level (i.e. nation/state) mean had fewer than 5
586 estimates (sites).

587

588 *Analysing conditions at bright spots*

589 For our preliminary investigation of why bright and dark spots may diverge from
590 expectations, we surveyed data providers and other experts about key social,
591 institutional, and environmental conditions at the 15 bright spots, 35 dark spots, and
592 14 sites that performed most closely to model specifications. Specifically, we
593 developed an online survey using Survey MonkeyTM software, which we asked data
594 providers who sampled those sites to complete with input from local experts where
595 necessary. Data providers generally filled in the survey in consultation with
596 nationally-based field team members who had detailed local knowledge of the
597 socioeconomic and environmental conditions at each of the sites. Research on bright
598 spots in agricultural development¹³ highlights several types of social and

599 environmental conditions that may lead to bright spots, which we adapted and
600 developed proxies for as the basis of our survey into why our bright and dark spots
601 may diverge from expectations. These include:

- 602 i) *Social and institutional conditions.* We examined the presence of
603 customary management institutions such as taboos and marine tenure
604 institutions, whether there was a high level of engagement by local people
605 in management, whether there was high levels of dependence on marine
606 resources (whether a majority of local residents depend on reef fish as a
607 primary source of food or income). All social and institutional conditions
608 were recorded as presence/absence. Dependence on resources and
609 engagement were limited to sites that had adjacent human populations. All
610 other conditions were recorded regardless of whether there is an adjacent
611 community;
- 612 ii) *Technological use/innovation.* We examined the presence of motorised
613 vessels, intensive capture equipment (such as beach seine nets, surround
614 gill nets, and muro-ami nets), and storage capacity (i.e. freezers); and
- 615 iii) *External influences* (such as donor-driven projects). We examined the
616 presence of NGOs, fishery development projects, development initiatives
617 (such as alternative livelihoods), and fisheries improvement projects. All
618 external influences were recorded as present/absent then summarised into
619 a single index of whether external projects were occurring at the site;
- 620 iv) *Environmental/ecological processes* (e.g. recruitment & connectivity). We
621 examined whether sites were within 5km of mangroves and deep-water
622 refuges, and whether there had been any major environmental disturbances

623 such as coral bleaching, tsunami, and cyclones within the past 5 years. All
624 environmental conditions were recorded as present/absent.
625
626 To test for associations between these conditions and whether sites diverged more or
627 less from expectations, we used two complementary approaches. The link between the
628 presence/absence of the aforementioned conditions and whether a site was bright,
629 average, or dark was assessed using a Fisher's Exact Test. Then we tested whether the
630 mean deviation in fish biomass from expected was similar between sites with
631 presence or absence of the mechanisms in question (i.e. the presence or absence of
632 marine tenure/taboo) using an ANOVA assuming unequal variance. The two tests
633 yielded similar results, but provide slightly different ways to conceptualise the issue,
634 the former is correlative while the latter explains deviation from expectations based
635 on conditions, so we provide both (Figure 2, Extended Data Fig. 6).
636

637 **References**

- 638 1 Pandolfi, J. M. *et al.* Global trajectories of the long-term decline of coral reef
639 ecosystems. *Science* **301**, 955-958 (2003).
- 640 2 Bellwood, D. R., Hughes, T. P., Folke, C. & Nystrom, M. Confronting the
641 coral reef crisis. *Nature* **429**, 827-833 (2004).
- 642 3 Hughes, T. P., Bellwood, D. R., Folke, C., Steneck, R. S. & Wilson, J. New
643 paradigms for supporting the resilience of marine ecosystems. *Trends Ecol*
644 *Evol* **20**, 380-386 (2005).
- 645 4 Sternin, M., Sternin, J. & Marsh, D. in *The Hearth Nutrition Model:*
646 *Applications in Haiti, Vietnam, and Bangladesh.* (eds O Wollinka, E Keeley,
647 B Burkhalter, & N Bashir) 49-61 (VA: BASICS, 1997).
- 648 5 Pretty, J. N. *et al.* Resource-conserving agriculture increases yields in
649 developing countries. *Emviron Sci Tech* **40**, 1114-1119 (2006).
- 650 6 Knowlton, N. & Jackson, J. B. C. Shifting baselines, local impacts, and global
651 change on coral reefs. *Plos Biol* **6**, 215-220 (2008).
- 652 7 Naeem, S., Duffy, J. E. & Zavaleta, E. The functions of biological diversity in
653 an age of extinction. *Science* **336**, 1401-1406 (2012).
- 654 8 Devillers, R. *et al.* Reinventing residual reserves in the sea: are we favouring
655 ease of establishment over need for protection? *Aquat Conserv* (2014).
- 656 9 Pressey, R. L., Visconti, P. & Ferraro, P. J. Making parks make a difference:
657 poor alignment of policy, planning and management with protected-area
658 impact, and ways forward. *Philos T R Soc B* **370** (2015).
- 659 10 Pascale, R. T. & Sternin, J. Your company's secret change agents. *harvard*
660 *business review* **83**, 72-81 (2005).

- 661 11 Levinson, F. J., Barney, J., Bassett, L. & Schultink, W. Utilization of positive
662 deviance analysis in evaluating community-based nutrition programs: An
663 application to the Dular program in Bihar, India. *Food Nutr Bull* **28**, 259-265
664 (2007).
- 665 12 McClanahan, T. R. *et al.* Critical thresholds and tangible targets for
666 ecosystem-based management of coral reef fisheries. *P Natl Acad Sci USA*
667 **108**, 17230-17233 (2011).
- 668 13 Noble, A., Pretty, J., de Vries, F. P. & Bossio, D. in *Bright spots demonstrate*
669 *community successes in African agriculture* (ed F. W. T. Penning de Vries)
670 7 (International Water Management Institute, 2005).
- 671 14 Cinner, J., Marnane, M. J., McClanahan, T. R. & Almany, G. R. Periodic
672 closures as adaptive coral reef management in the Indo-Pacific. *Ecol Soc* **11**
673 (2006).
- 674 15 Lindfield, S. J., Harvey, E. S., Halford, A. R. & McIlwain, J. L. Mesophotic
675 depths as refuge areas for fishery-targeted species on coral reefs. *Coral Reefs*,
676 1-13 (2015).
- 677 16 Cinner, J. E. *et al.* A framework for understanding climate change impacts on
678 coral reef social–ecological systems. *Reg Environ Chang*, 1-14 (2015).
- 679 17 Cinner, J. E. Social-ecological traps in reef fisheries. *Global Environ Chang*
680 **21**, 835-839 (2011).
- 681 18 Schmitt, K. M. & Kramer, D. B. Road development and market access on
682 Nicaragua's Atlantic coast: implications for household fishing and farming
683 practices. *Environ Conserv* **36**, 289-300 (2009).
- 684 19 Falk, A. & Szech, N. Morals and markets. *Science* **340**, 707-711 (2013).

- 685 20 Ostrom, E. *Governing the commons: The evolution of institutions for*
686 *collective action*. (Cambridge university press, 1990).
- 687 21 O'Rourke, D. The science of sustainable supply chains. *Science* **344**, 1124-
688 1127 (2014).
- 689 22 Sampson, G. S. *et al.* Secure sustainable seafood from developing countries.
690 *Science* **348**, 504-506 (2015).
- 691 23 Van Barneveld, W., Sary, Z., Woodley, J., Miller, M. & Picou-Gill, M.
692 Towards the co-operative management of fishing in Discovery Bay, Jamaica:
693 the role of the Fisheries Improvement Project. *Proc Gulf & Caribbean*
694 *Fisheries Institute (1991)* **44**, 195-210 (1996).
- 695 24 Cinner, J. E. *et al.* Comanagement of coral reef social-ecological systems. *P*
696 *Natl Acad Sci USA* **109**, 5219-5222 (2012).
- 697 25 Graham, N. A. J., Jennings, S., MacNeil, M. A., Mouillot, D. & Wilson, S. K.
698 Predicting climate-driven regime shifts versus rebound potential in coral reefs.
699 *Nature* **518**, 94-+ (2015).
- 700 26 York, R., Rosa, E. A. & Dietz, T. Footprints on the earth: The environmental
701 consequences of modernity. *Am Sociol Rev* **68**, 279-300 (2003).
- 702 27 Lambin, E. F. *et al.* The causes of land-use and land-cover change: moving
703 beyond the myths. *Global Environ Chang* **11**, 261-269 (2001).
- 704 28 Hughes, T. P., Huang, H. & Young, M. A. L. The wicked problem of China's
705 disappearing coral reefs. *Conserv Biol* **27**, 261-269 (2013).
- 706 29 Dodd, S. C. The interactance hypothesis: a gravity model fitting physical
707 masses and human groups. *Am Sociol Rev* **15**, 245-256 (1950).

708 30 Wittemyer, G., Elsen, P., Bean, W. T., Burton, A. C. O. & Brashares, J. S.
709 Accelerated human population growth at protected area edges. *Science* **321**,
710 123-126 (2008).

711 31 Daw, T. *et al.* The spatial behaviour of artisanal fishers: Implications for
712 fisheries management and development (Fishers in Space). (WIOMSA, 2011).

713 32 MacNeil, M. A. *et al.* Recovery potential of the world's coral reef fishes.
714 *Nature* **520**, 341-344 (2015).

715 33 Mora, C. *et al.* Global human footprint on the linkage between biodiversity
716 and ecosystem functioning in reef fishes. *Plos Biol* **9** (2011).

717 34 Edwards, C. B. *et al.* Global assessment of the status of coral reef herbivorous
718 fishes: evidence for fishing effects. *P Roy Soc B-Biol Sci* **281** (2014).

719 35 Froese, R. & Pauly, D. *FishBase. World Wide Web electronic publication*,
720 <www.fishbase.org> (2014).

721 36 Center for International Earth Science Information Network (CIESIN),
722 Columbia University & Centro Internacional de Agricultura Tropical (CIAT).
723 *Gridded population of the world. Version 3 (GPWv3): centroids*,
724 <<http://sedac.ciesin.columbia.edu/gpw>> (2005).

725 37 MacNeil, M. A. & Connolly, S. R. 12 Multi-scale patterns and processes in
726 reef fish abundance. *Ecol Fish Coral Reefs*, 116 (2015).

727 38 Mora, C. *et al.* Coral reefs and the global network of marine protected areas.
728 *Science* **312**, 1750-1751 (2006).

729 39 Lukermann, F. & Porter, P. W. Gravity and potential models in economic
730 geography. *Ann Assoc Am Geog* **50**, 493-504 (1960).

731 40 Nelson, A. Travel time to major cities: A global map of Accessibility. (Ispra,
732 Italy, 2008).

733 41 Bartholomé, E. *et al.* *GLC 2000: Global Land Cover mapping for the year*
734 *2000: Project status November 2002.* (Institute for Environment and
735 Sustainability, 2002).

736 42 R: A language and environment for statistical computing (R Foundation for
737 Statistical Computing, Vienna, Austria, 2012).

738 43 Dijkstra, E. W. A note on two problems in connexion with graphs.
739 *Numerische mathematik* **1**, 269-271 (1959).

740 44 Black, W. R. An analysis of gravity model distance exponents. *Transportation*
741 **2**, 299-312 (1973).

742 45 Emran, M. S. & Shilpi, F. *The extent of the market and stages of agricultural*
743 *specialization.* Vol. 4534 (World Bank Publications, 2008).

744 46 Cinner, J. E., Graham, N. A., Huchery, C. & Macneil, M. A. Global effects of
745 local human population density and distance to markets on the condition of
746 coral reef fisheries. *Conserv Biol* **27**, 453-458 (2013).

747 47 Teh, L. S. L., Teh, L. C. L. & Sumaila, U. R. A global estimate of the number
748 of coral reef fishers. *Plos One* **8** (2013).

749 48 Andréfouët, S. *et al.* in *10th International Coral Reef Symposium* (eds Y.
750 Suzuki *et al.*) 1732-1745 (Japanese Coral Reef Society, 2006).

751 49 Maina, J., McClanahan, T. R., Venus, V., Ateweberhan, M. & Madin, J.
752 Global gradients of coral exposure to environmental stresses and implications
753 for local management. *Plos One* **6** (2011).

754 50 Gelman, A., Carlin, J. B., Stern, H. S. & Rubin, D. B. *Bayesian data analysis.*
755 Vol. 2 (Taylor & Francis, 2014).

756 51 Patil, A., Huard, D. & Fonnesbeck, C. J. PyMC: Bayesian stochastic
757 modelling in Python. *J Stat Software* **35**, 1 (2010).

758 52 Gelman, A. & Rubin, D. B. Inference from iterative simulation using multiple
759 sequences. *Stat Sci* **7**, 457-472 (1992).

760 53 Mora, C. *et al.* Management effectiveness of the world's marine fisheries. *Plos*
761 *Biol* **7** (2009).

762 54 Ravenstein, E. G. The laws of migration. *J Statist Soc London* **48**, 167-235
763 (1885).

764 55 Anderson, J. E. A theoretical foundation for the gravity equation. *Am Econ*
765 *Rev*, 106-116 (1979).

766 56 Anderson, J. E. The gravity model. (National Bureau of Economic Research,
767 2010).

768

769

770 **End Notes**

771 Supplementary Information is linked to the online version of the paper at

772 www.nature.com/nature.

773

774 **Acknowledgments**

775 The ARC Centre of Excellence for Coral Reef Studies, Stanford University, and

776 University of Montpellier funded working group meetings. Thanks to M. Barnes for

777 constructive comments.

778

779 **Author Contributions**

780 J.E.C. conceived of the study with support from M.A.M, N.A.J.G, T.R.M, J.K, C.H,

781 D.M, C.M, E.A, and C.C.H; C.H. managed the database; M.A.M. and J.E.C.

782 developed and implemented the analyses; J.E.C. led the manuscript with M.A.M, and

783 N.A.J.G. All other authors contributed data and made substantive contributions to the

784 text.

785

786 **Author Information**

787 Reprints and permissions information is available at www.nature.com/reprints. The

788 authors declare no competing financial interests. Correspondence and request for

789 materials should be addressed to J.E.C. (Joshua.cinner@jcu.edu.au). This is the

790 Social-Ecological Research Frontiers (SERF) working group contribution #11.

791

792 **Extended Data Tables**

793

794 **Extended Data Table 1 | Summary of social and environmental covariates.**

795 Further details can be found in the Supplemental Online Methods. The smallest scale

796 is the individual reef. Sites consist of clusters of reefs within 4km of each other.

797 Nation/states generally correspond to country, but can also include or territories or

798 states, particularly when geographically isolated (e.g. Hawaii).

799

Covariate	Description	Scale	Key data sources
Local population growth	Difference in local human population (i.e. 100km buffer around our sites) between 2000-2010	Site	Socioeconomic Data and Application Centre (SEDAC) gridded population of the world database ³⁶
‘Gravity’ of major markets within 500km	The population of the major market divided by the squared travel time between the reef sites and the market. This value was summed for all major markets within 500km of the site.	Site	Human population size, land cover, road networks, coastlines
‘Gravity’ of the closest human settlement	The population of the nearest human settlement divided by the squared travel time between the reef site and the settlement.	Site	Human population size, land cover, road networks, coastlines
Protection	Whether the reef is	Reef	Expert opinion, global map of

status	openly fished, restricted (e.g. effective gear bans or effort restrictions), or unfished		marine protected areas.
Human Development index	A summary measure of human development encompassing: a long and healthy life, being knowledgeable and have a decent standard of living. We used linear and quadratic functions for HDI.	Nation/state	United Nations Development Programme
Population Size	Total population size of the jurisdiction	Nation/state	World Bank, census estimates, Wikipedia
Tourism	Proportion of tourist visitors to residents	Nation/state	World Tourism Organization's Compendium of Tourism Statistics, census estimates
Voice and accountability	Perceptions of the extent to which a country's citizens are able to participate in selecting their government.	Nation/state	World Bank
Fish landings	Landings of reef fish (tons) per Km ²	Nation/state	Teh et al. ⁴⁷

	of reef		
National fisheries poaching	Results from survey of national fisheries managers about levels of compliance with national fisheries regulations	Nation/ state	Mora et al. ⁵³
Climate stress	A composite metric comprised of 11 different environmental variables that are related to coral mortality from bleaching	Site	Maina et al. ⁴⁹
Productivity	The average (2005-2010) ocean productivity in mg C / m ² / day	Site	http://www.science.oregonstate.edu/ocean.productivity/
Habitat	Whether the reef is a slop, crest, flat, or back reef/lagoon	Reef	Primary data
Depth	Depth of the ecological survey (<4m, 4.1-10m, >10m)	Reef	Primary data

801 **Extended Data Table 2 | List of ‘Nation/states’ covered in study and their**
802 **respective average biomass (plus or minus standard error)** In most cases,
803 nation/state refers to an individual country, but can also include states (e.g. Hawaii or
804 Florida), territories (e.g. British Indian Ocean Territory), or other jurisdictions. We
805 treated the NW Hawaiian Islands and Farquhar as separate ‘nation/states’ from
806 Hawaii and Seychelles, respectively, because they are extremely isolated and have
807 little or no human population. In practical terms, this meant different values for a few
808 nation/state scale indicators that ended up having relatively small effect sizes, anyway
809 (Fig. 1b): Population, tourism visitations, and in the case of NW Hawaiian Island, fish
810 landings.
811

Nation/states	Average biomass	(± SE)
American Samoa	235.93	(± 17.75)
Australia	735.01	(± 136.85)
Belize	981.16	(± 65.32)
Brazil	663.35	(± 115.17)
British Indian Ocean Territory (Chagos)	2975.58	(± 603.99)
Cayman Islands	464.09	(± 25.41)
Colombia	846.07	(± 162.49)
Commonwealth of the Northern Mariana Islands	505.54	(± 99.3)
Comoros Islands	305.62	(± 38.73)
Cuba	2107.37	(± 466.34)
Egypt	552.73	(± 70.18)
Farquhar	2665.48	(± 492.62)
Federated States of Micronesia	377.90	NA (n=1)
Fiji	1464.54	(± 144.39)
Florida	1661.35	(± 198.42)
French Polynesia	1077.20	(± 101.4)
Guam	118.98	(± 16.81)
Hawaii	380.45	(± 25.11)
Indonesia	275.76	(± 19.89)
Israel	445.16	(± 105.13)
Jamaica	275.77	(± 50.75)
Kenya	335.25	(± 65.81)
Kiribati	1219.93	(± 93.2)
Madagascar	409.48	(± 46.1)
Maldives	688.64	(± 97.07)
Marshall Islands	707.72	(± 174.38)
Mauritius	166.93	(± 73.7)
Mayotte	631.43	(± 68.25)
Mexico	1930.81	(± 737.09)

Mozambique	461.01	(± 60.14)
Netherlands Antilles	428.01	(± 53.99)
New Caledonia	1460.27	(± 143.18)
NW Hawaiian Islands	729.71	(± 46.33)
Oman	282.79	(± 70.22)
Palau	3212.26	(± 332.02)
Panama	373.78	(± 85.41)
Papua New Guinea	566.70	(± 31.76)
Philippines	202.62	NA (n=1)
Pacific Remote Island Areas (PRIA), USA	641.47	(± 79.25)
Reunion	172.32	(± 30.67)
Seychelles	446.99	(± 46.6)
Solomon Islands	1280.30	(± 216.74)
Tanzania	346.29	(± 41.51)
Tonga	1149.97	(± 151.27)
United Arab Emirates	81.35	(± 28.66)
Venezuela	1472.39	(± 496.95)

812

813 **Extended Data Table 3| List of Bright and Dark Spot locations, population status,**
 814 **and protection status.**

815

Bright or Dark	Nation/State	Location	Populated	Protection
Bright	British Indian Ocean Territory	Chagos	Unpopulated	Unfished (high compliance)
	Commonwealth of the Northern Mariana Islands	Agrihan	Unpopulated	Fished
		Guguan	Unpopulated	Fished
	Indonesia	Raja Ampat 1	Populated	Restricted
		Raja Ampat 2	Populated	Restricted
		Kalimantan	Populated	Restricted
	Kiribati	Tabueran 1	Populated	Fished
		Tabueran 2	Populated	Fished
	Papua New Guinea	Karkar	Populated	Restricted
	PRIA	Baker	Unpopulated	Restricted
		Jarvis Island	Unpopulated	Restricted
	Solomon Islands	Choiseul	Populated	Fished
		Isabel	Populated	Fished
Makira		Populated	Fished	
New Georgia		Populated	Fished	
Australia	Lord Howe	Populated	Unfished (high compliance)	
Dark	Hawaii	Hawaii	Populated	Fished
		Kauai 1	Populated	Fished
		Kauai 2	Populated	Fished
		Lanai	Populated	Fished
		Maui 1	Populated	Fished
		Maui 2	Populated	Fished
		Molokai	Populated	Fished
		Oahu 1	Populated	Fished
		Oahu 2	Populated	Fished
		Oahu 3	Populated	Fished
		Oahu 4	Populated	Fished
		Oahu 5	Populated	Fished
		Oahu 6	Populated	Fished
		Indonesia	Karimunjawa 1	Populated
Karimunjawa 2	Populated		Unfished (low compliance)	
Karimunjawa 3	Populated		Unfished (low compliance)	
Pulau Aceh	Populated		Fished	
Jamaica	Montego Bay 1	Populated	Unfished (low compliance)	
	Montego Bay 2	Populated	Fished	

	Rio Bueno	Populated	Fished
Kenya	Diani	Populated	Fished
Madagascar	Toliara	Populated	Fished
Mauritius	Anse Raie	Populated	Fished
	Grand Sable	Populated	Fished
NW Hawaii	Lisianski	Unpopulated	Unfished (high compliance)
	Pearl & Hermes 1	Unpopulated	Unfished (high compliance)
	Pearl & Hermes 2	Unpopulated	Unfished (high compliance)
Reunion	Reunion	Populated	Fished
Seychelles	Bel Ombre	Populated	Restricted
Tanzania	Bongoyo	Populated	Unfished (high compliance)
	Chapwani	Populated	Fished
	Mtwara	Populated	Fished
	Stone Town, Zanzibar	Populated	Fished
Venezuela	Chuspa	Populated	Fished

816

817 **Extended Data Table 4| List of fish families included in the study, their common**
 818 **name, and whether they are commonly targeted in artisanal coral reef fisheries.**

819 Note: Targeting of reef fishes can vary by location due to gear, cultural preferences,
 820 and a range of other considerations.

821

Fish family	Common family name	Fishery target
Acanthuridae	Surgeonfishes	Target
Balistidae	Triggerfishes	Non-target
Diodontidae	Porcupinefishes	Non-target
Ephippidae	Batfishes	Target
Haemulidae	Sweetlips	Target
Kyphosidae	Drummers	Target
Labridae	Wrasses and Parrotfish	Target >20cm
Lethrinidae	Emperors	Target
Lutjanidae	Snappers	Target
Monacanthidae	Filefishes	Non-target
Mullidae	Goatfishes	Target
Nemipteridae	Coral Breams	Target
Pinguipedidae	Sandperches	Non-target
Pomacanthidae	Angelfishes	Target >20cm
Serranidae	Groupers	Target
Siganidae	Rabbitfishes	Target
Sparidae	Porgies	Target
Synodontidae	Lizardfishes	Non-target
Tetraodontidae	Pufferfishes	Non-target
Zanclidae	Moorish Idol	Non-target

822

823 **Extended Data Table 5 | Travel time estimates by land cover type.** Adapted from
 824 Nelson⁴⁰
 825

Global Land Cover Global Class	<i>Speed associated (km/h)</i>
Tree Cover, broadleaved, deciduous & evergreen, closed; regularly flooded Tree Cover, Shrub, or Herbaceous Cover (fresh, saline, & brackish water)	1
Tree Cover, broadleaved, deciduous, open (<i>open= 15-40% tree cover</i>)	1.25
Tree Cover, needle-leaved, deciduous & evergreen, mixed leaf type; Shrub Cover, closed-open, deciduous & evergreen; Herbaceous Cover, closed-open; Cultivated and managed areas; Mosaic: Cropland / Tree Cover / Other natural vegetation, Cropland / Shrub or Grass Cover	1.6
Mosaic: Tree cover / Other natural vegetation; Tree Cover, burnt	1.25
Sparse Herbaceous or sparse Shrub Cover	2.5
Water	20
Roads	60
Track	30
Artificial surfaces and associated areas	30
Missing values	1.4

826

827 **Extended Data Table 6 | Variance Inflation Factor Scores (VIF) for continuous**
 828 **data before and after removing variables due to colinearity. X = covariate**
 829 removed.

830

Covariate	starting VIF	ending VIF
Market gravity (log)	1.9	1.5
nearest settlement gravity	1.4	1.3
Population growth	1.4	1.3
Climate stress	2.7	2.0
Ocean productivity	6.5	2.2
Sedimentation	6.0	X
Tourism	2.5	X
Control Corruption	10.5	X
GDP	8.2	X
HDI	5.5	3.3
Population size	1.9	1.8
Reef fish landings	3.1	2.2
Rule of Law	33.8	x
Voice and Accountability	3.2	3.2

831


832 **Extended Data Table 7| Model selection of potential gravity indicators and**
 833 **components.**

834

Model	Covariates	AIC	Delta AIC
M2	Gravity of nearest city	2666.4	0
M1	Gravity of all cities in 500km	2669.5	3.1
M3	Travel time to nearest city	2700.0	33.6
M5	Gravity of nearest small peri-urban area (40 people/km ²)	2703.9	37.5
M11	Total Population in 500km radius	2712.0	45.6
M9	Travel time to the nearest large peri-urban area (75 people/km ²)	2712.1	45.7
M6	Travel time to nearest small peri-urban area (40 people/km ²)	2713.8	47.4
M8	Gravity to the nearest large peri-urban area (75 people/km ²)	2722.9	56.5
M7	Population of nearest small peri-urban area (40 people/km ²)	2792.7	126.3
M4	Population of the nearest city	2812.8	146.5
M10	Population of the nearest large peri-urban area (75 people/km ²)	2822.2	155.8
M0	Intercept only	2827.7	161.27

835

836 **Extended Data Figure Legends**




837

838


839 **Extended Data Figure 1** | a) A heuristic of the gravity concept where interactions
 840 between people and reefs are a function of population size (p) and the time it takes to
 841 travel to the reef (tt). Beginning in the 1800s, the concept of ‘gravity’ has been
 842 applied to measure economic interactions, migration patterns, and trade flows^{29,54-56}.
 843 Drawing on an analogy from Newton’s Law of Gravitation, the gravity concept
 844 predicts that interactions between two points are positively related to their mass (i.e.,
 845 population) and inversely related to the distance between them. Here, we adapt the
 846 gravity concept to examine interactions between people and reefs. We posit that
 847 human interactions with a reef will be a function of the population of a place (p)
 848 divided by the squared time it takes to travel (tt) to the reefs (i.e. travel time). Thus,
 849 gravity values could be similar for places that are large but far from the reefs (e.g. p_x
 850 = 30,000 people, tt_x = 10hours) as to those with small populations that are close to the
 851 reef (e.g. p_y = 300 people, tt_y =1 hour). We used travel time instead of linear distance

852 to account for the differences incurred by travelling over different surfaces (e.g. water,
853 roads, tracks—see Methods). We developed gravity measures for the nearest human
854 settlement and for the nearest major market (defined as provincial capitals, ports, and
855 other large, populated places- see Methods). b) Gravity isoclines along gradients of
856 population size and travel time.
857

858 **Extended Data Figure 2 | Marginal relationships between reef fish biomass and**
 859 **site-level social drivers.** a) local population growth, b) market gravity, c) nearest
 860 settlement gravity, d) tourism, e) nation/state population size, f) Human development
 861 Index, g) high compliance marine reserve (0 is fished baseline), h) restricted fishing
 862 (0 is fished baseline), i) low compliance marine reserve (0 is fished baseline), j) voice
 863 and accountability, k) reef fish landings, l) ocean productivity; m) depth (-1= 0-4m,
 864 0= 4-10m, 1=>10m), n) reef flat (0 is reef slope baseline), o) reef crest flat (0 is reef
 865 slope baseline), p) lagoon/back reef flat (0 is reef slope baseline). All X variables are
 866 standardized. ** 95% of the posterior density is either a positive or negative direction
 867 (Box 1); * 75% of the posterior density is either a positive or negative direction.




868
 869




870
871
872
873
874
875
876
877
878

Extended Data Figure 3 | Market gravity and fish biomass. Relationship between market gravity and a) reef fish biomass; b) targeted reef fish biomass (using fish families targeted by artisanal fisheries specified in Extended Data Table 2); c) non-target reef fish biomass. The strong relationship between gravity and reef fish biomass is very similar for the biomass of fishes generally targeted by artisanal fisheries, but very different for non-target fishes. This suggests that the relationship between market gravity and fish biomass is primarily driven by fishing, rather than other potential human impacts of urban areas (sedimentation, nutrients, pollution, etc.).

879 **Extended Data Figure 4| Correlation plot of candidate continuous covariates before accounting for colinearity (Extended Data Table 7).**
880 Colinearity between continuous and categorical covariates (including biogeographic region, habitat, protection status, and depth) were analysed
881 using boxplots.
882






884

885 **Extended Data Figure 5 | Model fit statistics.** Bayesian p Values (BpV) for the full
886 model indicating goodness of fit, based on posterior discrepancy. Points are Freeman-
887 Tukey differences between observed and expected values, and simulated and expected
888 values. Plot shows no evidence for lack of fit between the model and the data.

889



890

891 **Extended Data Figure 6| Box plot of deviation from expected as a function of the**

892 **presence or absence of key social and environmental conditions expected to**

893 **produce bright spots.**

894

895