

1 **Cartilage Biomarkers in the Osteoarthropathy of Alkaptonuria Reveal Low**
2 **Turnover and Accelerated Ageing**

3
4 Adam M Taylor^{1,2}, Ming-Feng Hsueh³, Lakshminarayan R Ranganath⁴, James A
5 Gallagher⁴, Jane P Dillon⁴, Janet L Huebner², Jon B Catterall² & Virginia B Kraus^{2,3,5}.

6
7 ¹Lancaster Medical School, Faculty of Health & Medicine, Lancaster University,
8 Lancaster, UK; ²Duke Molecular Physiology Institute, Department of Medicine,
9 ³Department of Pathology, and ⁵Department of Orthopaedic Surgery, Duke University
10 School of Medicine, Durham, North Carolina, USA; ⁴Musculoskeletal Biology,
11 University of Liverpool, Liverpool, UK.

12
13 **Short title:** Biomarkers of cartilage ageing in alkaptonuria

14 **Corresponding Author**

15 Correspondence to Dr Adam Taylor: a.m.taylor@lancs.ac.uk

16 Dr Adam Taylor

17 Clinical Anatomy Learning Centre

18 Lancaster Medical School

19 Furness College

20 Lancaster University

21 Bailrigg, LA1 4YB

22 England, UK **Keywords:** Alkaptonuria, Osteoarthritis, Biomarkers, Ageing,

23 Glycosaminoglycan, Cartilage oligomeric matrix protein, Racemization, Ochronosis.

1 **Abstract**

2 **Objective:** Alkaptonuria (AKU) is a rare autosomal recessive disease resulting from a
3 single enzyme deficiency in tyrosine metabolism. As a result, homogentisic acid (HGA)
4 cannot be metabolized, causing systemic increases. Over time, HGA polymerizes and
5 deposits in collagenous tissues leading to ochronosis. Typically, this occurs in joint
6 cartilages leading to an early onset, rapidly progressing osteoarthropathy. We examined
7 tissue turnover in cartilage affected by ochronosis and its role in disease initiation and
8 progression.

9 **Methods:** Hip and knee cartilages were obtained with informed patient consent at
10 surgery for arthropathy due to alkaptonuria (AKU n=6; 2 knees/4 hips) and osteoarthritis
11 (OA n=12; 5 knees/7 hips); healthy non-arthritic (Non-OA n=6; 1 knee/5 hips) cartilages
12 were obtained as waste from trauma surgery. We measured cartilage concentrations
13 (normalized to dry weight) of racemized aspartate, glycosaminoglycan (GAG), cartilage
14 oligomeric matrix protein (COMP) and deamidated COMP (D-COMP). Unpaired
15 alkaptonuria, OA and Non-OA samples were compared by nonparametric Mann-Whitney
16 U test.

17 **Results:** Despite more extractable total protein from AKU cartilage than from OA or
18 Non-OA, there was significantly less extractable GAG, COMP and D-COMP in AKU
19 samples compared to OA and Non-OA comparators. Racemized Asx (Aspartate and
20 Asparagine) was significantly enriched in AKU cartilage compared with OA.

21 **Conclusions:** These novel data represent the first examination of cartilage matrix
22 components in a sample of patients with AKU, representing almost 10% of the known
23 UK alkaptonuric population. Compared with OA and Non-OA, AKU cartilage

|

1 demonstrates a very low turnover state and has low amounts of extractable matrix
2 proteins.

3

4

A M TAYLOR AUTHOR FINAL SUBMITTED COPY

1 **Introduction**

2 Alkaptonuria (AKU), a rare autosomal recessive condition, is classified as a rare form of
3 osteoarthritis (OA) [1]. AKU results from a deficiency of a single enzyme, homogentisate
4 1,2-dioxygenase (HGD), responsible for opening the benzene ring of homogentisic acid
5 (HGA), an intermediate in tyrosine metabolism. The absence or lack of HGD function
6 results in HGA accumulation in tissues. HGA is excreted in the urine, but renal excretion
7 is insufficient to completely clear the burden of HGA in AKU [2]. Elevated HGA
8 concentrations in urine, plasma and body tissues result in a classic triad of clinical
9 presentations. These include the earliest presentation, from birth, of urine darkening upon
10 standing or exposure to alkaline conditions [3, 4]. Over many years of HGA exposure,
11 the second feature occurs—termed ochronosis or darkening of collagenous tissues—often
12 seen in the ear pinna, eye sclerae and other less uniformly and readily examined areas [5,
13 6]. The process of ochronosis is not completely understood, however it's suggested that
14 HGA undergoes polymerisation to a benzoquinone intermediate before assuming its final
15 polymeric form in collagenous tissues [7]. The final presenting feature of the disease,
16 namely early age onset of severe and rapidly progressing OA, results from ochronotic
17 pigment deposition in articular cartilages whose presence alters cartilage biomechanical
18 and biochemical properties [8]. The presence of this pigment appears to induce large-
19 scale aggressive resorption of the subchondral bone plate and articular calcified cartilage
20 due to stress-shielding in AKU joints [8].

21
22 A novel ultrastructural bone turnover mechanism has been identified in both AKU and
23 OA joint tissues, providing further evidence to support the similarities between the

1 diseases [1,2&8]. In both cases, novel structures result from un-coupled bone turnover,
2 deviating from the widely accepted Frostian bone turnover model. These structures also
3 appear attached to the existing trabecular stock, without prior preparation of the
4 trabecular surface by osteoclasts [8, 9]. The presence of ochronotic pigment in hyaline
5 cartilage makes it impervious to enzymatic degradation by Tergazyme (a bacterial
6 alkaline pronase); this may prevent the type of catabolism normally observed in OA
7 cartilage [8]. Shown by ultrastructural analyses, pigment associates closely with collagen
8 fibres in joint tissues, specifically in ligamentous capsule [10] and articular cartilage [AM
9 Taylor, PhD thesis, 2011]. Along with ultrastructural evidence of a disorder of collagen
10 fibres, solid state NMR demonstrates that there is disorder of extracellular proteins in
11 ochronotic cartilage when compared to control [11]. However, little is known about the
12 effects of ochronotic pigment on other matrix components, the biological ageing process
13 and joint tissue turnover of these patients [8].

14
15 AKU is iconic in medicine as the first disease, demonstrated by Archibald Garrod in
16 1902, to conform to Mendelian inheritance [3]. The symptoms [1, 2], gene responsible
17 [12, 13] and enzyme [14] are all known and well documented; even with this knowledge,
18 a viable treatment is yet to be available to patients. Trials on a potential therapy,
19 Nitisinone have been undertaken, but the outcome was deemed not beneficial [15, 16].
20 Nitisinone can inhibit ochronosis in a mouse model [17, 18]. However, Nitisinone is not
21 without its side-effects, including elevated plasma tyrosine [19]. Moreover, Nitisinone is
22 not likely to benefit individuals with established pigmentation and tissue impregnation
23 with a large burden of cross-linked HGA. A clearer understanding of the molecular

1 changes occurring in cartilage could provide more options for treating this disease and
2 shed new light on our understanding of the disease pathogenesis and progression.

3
4 Ageing leads to non-enzymatic post-translational modification of proteins including
5 racemization and isomerization [20, 21, 22]. Racemization refers to the time-dependent
6 conversion of amino acids in proteins from the L form to D form (only L amino acids are
7 incorporated into mammalian proteins). Cartilage, a tissue with low turnover due to its
8 aneural and avascular nature, undergoes quantifiable racemization of some amino acids
9 during the lifespan [23, 24]. Recent advances in the field of biomarkers of ageing and
10 turnover have demonstrated that cartilage oligomeric matrix protein (COMP) is a useful
11 determinant of differential turnover between hip and knee, with knee cartilage showing a
12 greater turnover response than hip cartilage [20]. In AKU, where cartilage and its
13 constituent matrix components become bound by HGA polymeric pigment, we
14 considered it likely that these biomarkers of ageing and turnover would provide novel
15 insights into the pathophysiology of this iconic disease. Given that pigmented cartilage
16 appears to be impervious to enzymatic degradation, we hypothesized that cartilage
17 turnover of alkaptonuria specimens will be reduced compared to osteoarthritic and non-
18 osteoarthritic comparators.

19
20 The aim of this study was to investigate the impact of ochronotic pigment on cartilage
21 matrix turnover in joint samples from patients with AKU. Through investigation of
22 aspartate racemization, glycosaminoglycan (GAG) and COMP, we characterized the

1 turnover of tissue relative to amount of pigmentation and compared results to non-
2 pigmented tissue from Non-OA and OA samples.

4 **Materials & Methods**

5 ***Cartilage sample collection.*** Alkaptonuric knees (n=2, mean age 58.5 years, range 58-59
6 years) and hips (n=4, mean age 61.0 years, range 54-70 years) were obtained at the time
7 of surgery from individuals undergoing total joint arthroplasty to alleviate symptoms
8 associated with ochronotic osteoarthropathy. Alkaptonuric cartilage samples were
9 acquired from macroscopically ochronotic areas of tissue. All alkaptonuric joint
10 specimens showed extensive pigmentation of all zones of cartilage. One specimen
11 showed complete pigmentation of all the cartilage and absence of subchondral bone in
12 many areas. Osteophytes were present in one of the samples where cartilage absence had
13 exposed underlying osseous tissues. Pigmentation of connective tissues including
14 ligaments was seen in all samples. All AKU samples were obtained with informed patient
15 consent according to the Declaration of Helsinki and under ethical approval from the
16 Liverpool Research Ethics Committee. Waste articular cartilage specimens were obtained
17 from randomly selected total knee (n=5, mean age 68.6 years, range 54-88 years) and
18 total hip (n=7, mean age 81.9 years, range 72-90 years) arthroplasties performed at Duke
19 University Medical Center to alleviate symptoms of OA. From each arthritic joint,
20 cartilage was harvested from around the lesion (lesion cartilage). Non-arthritic control
21 cartilage samples from knee (n=1, mean age 36 years) and hip (n=5, mean age 63.4 years,
22 range 23-82 years) joints were obtained at the time of reconstructive surgery for trauma
23 from patients without evidence of OA as determined by macroscopic inspection of the

1 specimens by the surgeon. All samples from Duke University were collected under Duke
2 Institutional Review Board approval as waste surgical specimens and according to the
3 Declaration of Helsinki. Due to the early onset of the arthropathy in AKU we were
4 unable to directly age match samples. Instead, we utilized a matching strategy to ensure
5 that OA and non-OA samples encompassed the AKU sample ages. All samples in each
6 group were from different donors.

7
8 *Sample preparation.* Soluble and insoluble cartilage proteins were prepared as previously
9 described with minor modifications [20]. Cartilage was pulverized under liquid nitrogen
10 and extracted for 24 h at 4°C with gentle mixing in 4M Guanidine-HCl (Gu-HCl) in
11 sodium acetate buffer pH 4.0 with protease inhibitor cocktail (Sigma-Aldrich, St Louis,
12 MO) at a ratio of 2 ml of Gu-HCL extraction buffer per 1g of wet weight cartilage.
13 Following centrifugation and collection of the supernatant, the remaining cartilage was
14 extracted a second time with Gu-HCl extraction buffer for a further 24 h. Both first and
15 second Gu-HCl extractions were combined and stored at -80°C until needed. A 200 µl
16 aliquot of extract was buffer exchanged on an AKTA purifier 10 (GE Healthcare,
17 Pittsburgh, PA) using a HiTrap Desalting column (GE, Pittsburgh, PA) into PBS and
18 stored at -80°C constituting the soluble protein fraction. The residual Gu-HCl insoluble
19 protein fraction was washed a further 5 times in nanopure water over a 24 hr period at
20 4°C with rotation mixing to remove any residual Gu-HCl and extracted proteins. Excess
21 water was removed and the samples stored at -80°C until required for hydrolysis and
22 HPLC analysis.

23

1 *Determination of protein content.* Total protein was determined using the commercially
2 available BCATM Protein Assay Kit performed according to the manufacturer's
3 instructions (Pierce, Rockford, IL). The presence of HGA was assessed by addition of
4 10M NaOH to sample extract. No color change indicative of the presence of HGA was
5 seen in any samples, assuring the lack of interference by HGA with the BCA assay for
6 protein determination.

7
8 *Determination of GAG content.* GAG content of cartilage extracts was quantified by dye
9 binding assay with dimethylmethylene blue (Sigma, St. Louis, MO) as previously
10 described [25]. Our analyses demonstrated no confounding of the DMMB reaction by
11 HGA (unpublished results).

12
13 *Quantification of racemized amino acids.* Racemized Asx (Aspartate and Asparagine)
14 concentrations in cartilage extracts were quantified using a previously described high
15 performance liquid chromatography (HPLC) approach developed and validated in our
16 laboratory [23]. Briefly, the insoluble and soluble cartilage proteins were acid hydrolyzed
17 in 6M HCl at 105°C for 6h. The resulting free D- and L-amino acids were derivatized to
18 fluorescent compounds by addition of o-phthalaldehyde and N-tert-butyloxycarbonyl-
19 L-cysteine. The subsequent derivatives were separated by reversed-phase HPLC using a
20 C18 column and mobile phases of 0.2M acetic acid adjusted to a pH of 6.0 and an
21 acetonitrile gradient. The resulting peaks were quantified fluorometrically (ex340nm,
22 em440nm) by comparison to commercially available pure D- and L-amino acids (Fluka
23 Biochemicals).

1

2 *COMP ELISA*. Total COMP (T-COMP) concentrations of the Gu-HCl soluble protein
3 fraction of cartilage were quantified by ELISA using the mAb 16F12 for capture and
4 biotinylated mAb 17C10 for detection as previously described [26] with the following
5 minor modifications. Plates were blocked with 3% BSA/PBS/0.02% sodium azide for 2
6 hours. Streptavidin-alkaline phosphatase conjugate (ExtrAvidin, Sigma, St. Louis, MO)
7 was diluted 30,000x in 0.01% BSA/PBS/0.02% sodium azide and the phosphatase
8 substrate (4-nitrophenyl phosphate disodium salt hexahydrate, Sigma, St. Louis, MO)
9 was used as the detection agent. Plates were read at wavelength 405 nm after 20 mins of
10 incubation.

11

12 *D-COMP sandwich ELISA*. D-COMP concentrations of the Gu-HCl soluble protein
13 fraction of cartilage were determined using the D-COMP sandwich ELISA as previously
14 described [20]. Briefly, 96 well plates were coated with the deamidated COMP specific
15 6-1A12 D-COMP mAb in 0.02M sodium carbonate coating buffer, pH 9.6 overnight at
16 4°C [20]. Plates were blocked for at least 2 h with 5% w/v BSA in PBS, pH 7.4 at 37°C,
17 prior to use. Samples or standard were diluted in 0.1% w/v BSA in PBS as required prior
18 to incubation of 50 µl of sample or standard overnight at 4°C. Unbound sample was
19 discarded and the plate washed with PBS-tween wash buffer before incubation with the
20 biotinylated 17-C10 COMP detection mAb (a gift from Dr. V. Vilim). Unbound 17-C10
21 was discarded and the plate washed. Due to the lower levels of D-COMP, the D-COMP
22 ELISA sensitivity was increased using avidin poly-horseradish peroxidase (poly-HRP,
23 Thermo Scientific, Rockford, IL), added for 30 min at room temperature with gentle

1 mixing. Excess avidin-HRP was discarded, the plate washed and signal detected using
2 tetramethyl benzidine (TMB, Sigma-Aldrich, St Louis, MO) reagent after stopping with
3 2M HCl and detection at 450nm.

4

5 *Statistical analysis* was performed using GraphPad Prism Version 5 software (La Jolla,
6 CA). All results were analysed as unpaired data using the nonparametric Mann-Whitney
7 U test.

8

9 *Histological analysis of alkaptonuric samples* was undertaken by fixing joint tissues at
10 surgery in 10% PBFS for H&E staining or by mounting with tissue-tek and freezing in
11 liquid N₂ for von Kossa; 5µm and 10µm sections were cut respectively. Non-decalcified
12 cryostat sections of AKU tissue were stained in aqueous 5% silver nitrate for 30mins
13 under a bench lamp, then washed with dH₂O and a 5% solution of sodium thiosuphate
14 was applied for 1 min to terminate the reaction. Sections were dehydrated through graded
15 alcohols and mounted in DPX (Sigma Aldrich, UK)

16

17 **Results**

18

19 Our results show significantly more total protein was extractable from AKU cartilage
20 than either Non-OA and OA samples, $p=0.0022$ and 0.035 respectively (Figure 1). There
21 was no significant difference between the mg of soluble protein/ml between Non-OA and
22 OA ($p=0.4245$).

23

1 However, the mean amount of extractable GAG (as a proportion of total protein) was
2 significantly less from AKU cartilage samples compared to either Non-OA or OA
3 samples, $p=0.0022$ and $p=0.0012$ respectively (Figure 2A). Similar significant disease-
4 related differences were also reflected in the smaller hip only set; AKU vs Non-OA and
5 OA, $p=0.0159$ and $p=0.0121$ respectively (Figure 2B).

6
7 A greater amount of aged COMP protein, expressed as a proportion of deamidated to
8 total COMP (D/T ratio), was extractable from AKU cartilage compared to OA samples
9 ($p=0.0441$) (Figure 3A). There was no significant difference between Non-OA and AKU
10 cartilage ($p=0.1797$). Analysis of the hip only set showed no significant differences
11 between AKU and OA ($p=0.5556$) or Non-OA cartilages ($p=0.2303$) (Figure 3B).

12
13 Significantly less total COMP (as a proportion of total protein) was extractable from
14 AKU cartilage than either Non-OA or OA samples, $p=0.0022$ and $p=0.0009$ respectively
15 (Figure 4A). Similar significant disease-related differences were reflected in the smaller
16 hip only set; AKU vs Non-OA and OA, $p=0.0159$ and $p=0.0061$ respectively (Figure 4B).

17
18 Significantly less D-COMP (as a proportion of total protein) was extractable from AKU
19 cartilage than either Non-OA or OA samples, $p=0.043$ and $p=0.0009$ respectively (Figure
20 4C). Similar significant disease-related differences were also reflected in the smaller hip
21 only set; AKU vs Non-OA and OA, $p=0.0317$ and $p=0.0061$ respectively (Figure 4D).

22
23 A general measure of aged protein, (Asx D/D+L), representing a ratio of racemized Asx
24 (D form) to non-racemized Asx (L form) was used to further assess the biological age of

1 two cartilage fractions: the Gu-HCl extractable protein from cartilage (Figure 5 A-B),
2 primarily representing proteoglycan; and the Gu-HCl insoluble protein from cartilage
3 (Figure 5 C-D), representing collagen and collagen associated proteins. Compared to OA,
4 the soluble protein fractions from AKU cartilage were significantly enriched in aged
5 (racemized) protein represented by the mean Asx D/D+L ratio, $p=0.0076$. Compared to
6 Non-OA, the mean Asx ratio from AKU cartilage was higher but this was not statistically
7 significant ($p=0.1797$) (Figure 5A). Similar disease-related differences were not reflected
8 in the smaller OA vs AKU hip only set ($p=0.1091$) or between Non-OA and AKU
9 ($p=0.0931$) (Figure 5B). The mean Asx ratio in the insoluble cartilage fraction was
10 significantly higher for AKU vs OA for all samples ($p=0.0277$), and also trended higher
11 for AKU vs Non-OA for all samples ($p=0.0931$) (Figure 5C). The hip only set also
12 showed similar trends with higher Asx ratios in AKU vs Non-OA and OA ($p=0.0635$ and
13 0.0727 , respectively) (Figure 5D).

14 Histological examination of alkaptonuric cartilages revealed full depth pigmentation of
15 numerous areas across all samples, with many empty chondrocyte lacunae; regions with
16 retained chondrocytes displayed intracellular pigmentation (Figure 6A). Some areas of
17 calcified cartilage matrix displayed a granular like pigment, often in proximity to empty
18 chondrocyte lacunae. The bone domain showed mature bone with characteristic lamellar
19 structure and multiple osteocytes without pigmentation. In a number of areas there were
20 regions where pigmented cartilage was observed without articular calcified cartilage or
21 subchondral bone beneath it. In areas where articular cartilage was absent from the bone
22 surface, pieces of impacted cartilage were observed within the marrow space, surrounded
23 by a variety of mono- and multinucleated cells, many of which displayed intracellular

1 pigmentation (Figure 6B). Histological examination by von Kossa staining and alizarin
2 red staining (data not shown) revealed that there was no deposition of calcium salts such
3 as hydroxyapatite associated with pigment. This contrasted with the mineralised bone
4 seen beneath the cartilage (Figure 6C).

6 **Discussion**

7 Although few in absolute numbers, the AKU samples in this study represent 0.6% of the
8 currently known global population with AKU and 9.375% of the currently known UK
9 AKU population, highlighting one of the difficulties in studying rare diseases [2,4]. Here
10 we present the first data analyzing biomarkers of ageing and turnover comparing
11 alkaptonuric to OA and non-OA cartilage samples. AKU samples demonstrated an older
12 profile of biomarkers of ageing across a number of analyzed parameters, both by ratio of
13 aged (deamidated) to total COMP and by ratio of aged (racemized) aspartate to total
14 aspartate in protein. These results were consistent with lower anabolism of AKU
15 cartilage, i.e. low turnover resulting in enrichment of age-related post-translation
16 modification of cartilage protein including the higher ratios of deamidated to total COMP
17 and higher ratios of racemized aspartate to total aspartate. This was of interest
18 considering that the mean age and range of age was higher for OA than AKU, which
19 would be expected to yield more age-related changes in OA.

20
21 Previously it has been suggested that turnover in articular cartilage is greatest at the
22 superficial layer and the turnover rate decreases as depth from the articular surface
23 increases [27]. This may initially be true in AKU, but once pigmentation commences, the

1 highest rate of turnover, in the form of catabolism, appears in the deepest cartilage, with
2 resorption of hyaline articular calcified cartilage, following resorption of the subchondral
3 bone [8]. With the biochemical tools of deamidated COMP and racemized aspartate, it
4 would be possible in future to evaluate turnover by depth of cartilage to determine if
5 protein turnover reflects apparent turnover based on calcified cartilage and subchondral
6 bone pathology.

7
8 Interestingly, despite the presence of pigment and potential for matrix cross-linking via
9 pigment adducts, the amount of protein extractable from AKU cartilage was higher than
10 for OA and non-OA. Higher levels of molecular degeneration could lead to more readily
11 extractable protein. In the case of AKU, a large proportion of both non-collagen and
12 collagen bound proteins might be affected resulting in a higher amount of extractable
13 protein relative to OA and non-OA. This explanation is plausible given the trend for OA
14 cartilage to yield more extractable protein than non-OA and for AKU to yield still higher
15 amounts of extractable protein.

16
17 Until recently, little was known about the molecular events occurring in AKU cartilage
18 [8]. Most information available is representative of end stage of the disease when the
19 molecular mechanisms are almost/already complete [8]. Our observation of lower levels
20 of extractable GAG in AKU are consistent with an OA trajectory [28, 29], however, the
21 extractable amounts we observed in our AKU patient samples were significantly less than
22 for OA. Similarly, total COMP was significantly less extractable from AKU cartilage.

1 The significantly reduced amounts of extractable GAG and COMP in AKU cartilage
2 could be imagined to arise through two scenarios:

3 1) They are lost very early or steadily, due to accelerated molecular degeneration,
4 ageing, trauma, biochemical alteration of the extracellular milieu, or some
5 combination of these factors, and not replaced due to chondrocyte pathology induced
6 by the pigment. One of the symptoms of AKU is an early onset, rapidly progressing
7 osteoarthropathy. If proteoglycan in AKU is lost very early and much more rapidly
8 than general OA, this may be a factor in the accelerated disease progression observed
9 in AKU.

10 2) They are bound to the cartilage extracellular matrix by the presence of ochronotic
11 polymer and cannot be extracted. If proteoglycans are bound within the ochronotic
12 matrix, this is likely a result of the polymerization process occurring in water spaces
13 of the cartilage. HGA, which will polymerize to form ochronotic pigment, is a small
14 very highly water soluble molecule and likely binds matrix proteins that are anchored
15 in and around collagen fibres [10]. However, this latter possibility seems unlikely
16 given the ability to extract more total protein from AKU cartilage than from OA or
17 Non-OA.

18
19 Regardless of scenario, both possibilities lead to a cartilage matrix in AKU that is far
20 from normal. It has already been shown that pigmentation commences pericellularly [8].

21 Given the intricate relationship that exists between chondrocytes and their pericellular
22 matrices [30, 31], this likely exacerbates processes that drive alteration in tissue
23 composition. Alterations in the pericellular matrix have been linked to OA [32].

1 Secondly, pigmented cartilage in AKU is significantly stiffer than that seen in OA and
2 normal cartilage [8]. It is likely that the observed changes in cartilage matrix composition
3 described here precede microscopic changes in the articular cartilage and subchondral
4 bone plate and may contribute to their progression by allowing HGA to polymerize in
5 spaces once occupied by GAG, or utilize GAG as its nucleation site. There is a possibility
6 that the greater loss of GAG in the AKU allows for greater extracellular protein
7 penetration and this, at least in part, might account for higher total protein content in
8 AKU samples. This, potentially combined with an increased crosslinking in the matrix
9 binding larger amounts of protein in. One could also posit increased protein production
10 due to a cellular attempt to mount a repair response, producing more proteins
11 incorporated into the matrix. The latter explanation seems unlikely given the apparent
12 low turnover of AKU cartilage specimens based on COMP D/T and racemized Asp
13 results.

14 The histological images presented demonstrate typical changes observed previously in
15 alkaptonuric joints [8]. Cartilage is full of ochronotic pigment; existing in association
16 with collagen [10] or with other matrix proteins. The bone changes indicate altered
17 mechanical loading, with resorption of subchondral bone and calcified cartilage [9,10].
18 We did not see levels of calcium deposition consistent with mineralised tissue, compared
19 to bone within our sections. Lack of calcium associated with pigmentation is consistent
20 with our previous work showing calcium is not present in association with ochronotic
21 pigment by BSE-SEM or IKI enhanced BSE-SEM [8, 32]. These morphological changes
22 are likely contributed to by the matrix changes described here.

23

1 Whilst proteoglycan loss in these patients has physiological significance, there are
2 clinical implications. Radiographic presentation of AKU patients is almost always
3 representative of some degenerative changes that mimic OA; joint space narrowing is
4 often described, particularly in the intervertebral disc spaces [8]. Cartilage thinning is not
5 well documented, the majority of specimens obtained at surgery are mostly normal, apart
6 from their ochronosis [8]. Those that do possess what would be described as lesions,
7 appear ochronotic and are typified by mechanical damage from contact between brittle
8 cartilage on opposing surfaces, rather than enzymatic degradation and lesions typically
9 observed with OA. Because the presence of pigment initially protects cartilage from overt
10 breakdown, early on radiographs appear normal whilst cartilage is
11 pigmented/pigmented. Only when mechanical trauma dislodges cartilage may
12 radiological changes be observed; therefore, other diagnostic measures for assessing the
13 state of joints of individuals with AKU, particularly during early joint pathological
14 stages, may be more appropriate. Along with HGA [8], the damaged cartilage fragments
15 are often seen embedded in the synovium and synovial fluid aspirate [34, 35].

16
17 Our data demonstrate the alterations that occur in the cartilage of alkaptonuric patients.
18 The loss of numerous matrix proteins raises potential issues for treating AKU. Nitisinone
19 has been suggested as a potential therapy [15, 16, 18, 36] however, it should be borne in
20 mind that this would most benefit individuals without established pigmentation. These
21 data represent an advance in understanding AKU cartilage turnover that could stimulate
22 new therapeutic options. For instance, our results suggest individuals may also benefit
23 from growth factor and other anabolic treatments as there appears to be a lack of an

1 anabolic response to the pigmentation process, particularly in those who have some
2 established pigmentation.

3 **Key messages**

4 1) AKU cartilage demonstrates an older profile of ageing biomarkers compared to OA
5 and non-OA comparators.

6 2) AKU cartilage shows a low turnover profile.

7 3) Anabolic therapy may be beneficial to patients with AKU.

8 **Acknowledgements**

9 We are extremely grateful to all patients who donated their tissues as part of this study.

10 The authors would like to thank Dr Hazel Sutherland for her technical assistance with
11 histology.

12 **Disclosure statement**

13 The authors declare no conflicts of interest.

14 **Funding**

15 This work was supported by an OARSI scholarship award [to AT] and the AKU Society
16 & Rosetrees Trust. This work was also supported by National Institute of Health
17 /National Institute of Arthritis Musculoskeletal and Skin Diseases [AR050245] [to VBK]
18 and the National Institute of Health /National Institute on Aging [AG028716] [to VBK].

19 **Role of the funding source**

20 Funders had no input into study design, collection, analysis and interpretation of data; in
21 the writing of the manuscript; and in the decision to submit the manuscript for
22 publication.

23 **Author Contributions**

1 AT, JAG, LRR and VBK conceived the study.
2 AT, JBC, JPD, MFH & JLH undertook experimental procedures.
3 All authors contributed to interpreting data, drafting, reading and approving the final
4 version of the manuscript.

5 **References**

- 6 1. Kraus, VB. 2014. *Rare Osteoarthritis: Ochronosis and Kashin-Beck Disease*. In:
7 *Rheumatology, 6th edition*, Ed. MC Hochberg, AJ Silman, JS Smolen, ME
8 Weinblatt, MH Weisman (Mosby Elsevier, Philadelphia) Chapter 185:1536-1540.
- 9 2. Mistry JB, Bukhari M, Taylor AM. *Alkaptonuria*. *Rare Dis*. 2013 Dec
10 18;1:e27475. doi: 10.4161/rdis.27475.
- 11 3. Garrod, Archibald E. *The Incidence of Alkaptonuria: A Study in Chemical*
12 *Individuality*. *Lancet*, 1902. vol. ii, pp. 1616-1620.
- 13 4. Ranganath L, Taylor AM, Shenkin A, , et al. *Identification of alkaptonuria in the*
14 *general population: a United Kingdom experience describing the challenges,*
15 *possible solutions and persistent barriers*. *J Inherit Metab Dis*. 2011
16 Jun;34(3):723-30.
- 17 5. Taylor AM, Wilson PJ, Ingrams DR, Helliwell TR, Gallagher JA, Ranganath LR.
18 *Calculi and intracellular ochronosis in the submandibular tissues from a patient*
19 *with alkaptonuria*. *J Clin Pathol*. 2010 Feb;63(2):186-8.
- 20 6. Taylor AM, Batchelor TJ, Adams VL, Helliwell TR, Gallagher JA, Ranganath LR.
21 *Ochronosis and calcification in the mediastinal mass of a patient with*
22 *alkaptonuria*. *J Clin Pathol*. 2011 Oct;64(10):935-6.

- 1 7. La Du BN, Zanonni VG, Laster L, Seegmiller JE. The nature of the defect in
2 tyrosine metabolism in alcaptonuria. *J Biol Chem.* 1958 Jan;230(1):251-60.
- 3 8. Taylor AM, Boyde A, Wilson PJ, et al. The role of calcified cartilage and
4 subchondral bone in the initiation and progression of ochronotic arthropathy in
5 alkaptonuria. *Arthritis Rheum.* 2011 Dec;63(12):3887-96.
- 6 9. Taylor AM, Boyde A, Davidson JS, Jarvis JC, Ranganath LR, Gallagher JA.
7 Identification of trabecular excrescences, novel microanatomical structures,
8 present in bone in osteoarthropathies. *Eur Cell Mater.* 2012 Apr 21;23:300-9.
- 9 10. Taylor AM, Wlodarski B, Prior IA, et al. Ultrastructural examination of tissue in
10 a patient with alkaptonuric arthropathy reveals a distinct pattern of binding of
11 ochronotic pigment. *Rheumatology (Oxford).* 2010 Jul;49(7):1412-4.
- 12 11. Chow WY, Taylor AM, Reid DG, Gallagher JA, Duer MJ. Collagen atomic scale
13 molecular disorder in ochronotic cartilage from an alkaptonuria patient,
14 observed by solid state NMR. *J Inherit Metab Dis.* 2011 Dec;34(6):1137-40.
- 15 12. Pollak MR, Chou YH, Cerda JJ, et al. Homozygosity mapping of the gene for
16 alkaptonuria to chromosome 3q2. *Nat Genet.* 1993 Oct;5(2):201-4.
- 17 13. Janocha S, Wolz W, Srsen S, et al. The human gene for alkaptonuria (AKU) maps
18 to chromosome 3q. *Genomics.* 1994 Jan 1;19(1):5-8.
- 19 14. Fernández-Cañón JM, Granadino B, Beltrán-Valero de Bernabé D, et al. The
20 molecular basis of alkaptonuria *Nat Genet.* 1996 Sep;14(1):19-24.
- 21 15. Suwannarat P, O'Brien K, Perry MB, et al. Use of nitisinone in patients with
22 alkaptonuria. *Metabolism.* 2005 Jun;54(6):719-28.

- 1 16. Introne WJ, Perry MB, Troendle J, et al. A 3-year randomized therapeutic trial of
2 nitisinone in alkaptonuria. *Mol Genet Metab.* 2011 Aug;103(4):307-14.
- 3 17. Taylor AM, Preston AJ, Paulk NK, et al. Ochronosis in a murine model of
4 alkaptonuria is synonymous to that in the human condition. *Osteoarthritis*
5 *Cartilage.* 2012 Aug;20(8):880-6.
- 6 18. Preston AJ, Keenan CM, Sutherland H, et al. Ochronotic osteoarthropathy in a
7 mouse model of alkaptonuria, and its inhibition by nitisinone. *Ann Rheum Dis.*
8 2014 Jan;73(1):284-9.
- 9 19. Hughes AT, Milan AM, Christensen P, et al. Urine homogentisic acid and
10 tyrosine: simultaneous analysis by liquid chromatography tandem mass
11 spectrometry. *J Chromatogr B Analyt Technol Biomed Life Sci.* 2014 Jul
12 15;963:106-12.
- 13 20. Catterall JB, Hsueh MF, Stabler TV, et al. Protein modification by deamidation
14 indicates variations in joint extracellular matrix turnover. *J Biol Chem.* 2012 Feb
15 10;287(7):4640-51
- 16 21. Catterall JB, Barr D, Bolognesi M, Zura RD, Kraus VB. Post-translational aging
17 of proteins in osteoarthritic cartilage and synovial fluid as measured by
18 isomerized aspartate. *Arthritis Res Ther.* 2009;11(2):R55.
- 19 22. McCudden C and VB Kraus. Biochemistry of amino acid racemization and
20 clinical application to musculoskeletal disease. *Clinical Biochemistry* 2006;
21 39(12):1112-30.

- 1 23. Stabler TV, Byers SS, Zura RD, Kraus VB. Amino acid racemization reveals
2 differential protein turnover in osteoarthritic articular and meniscal cartilages.
3 *Arthritis Res Ther.* 2009;11(2):R34.
- 4 24. Catterall, JB, Zura RD, Bolognesi MP, Kraus VB. Amino acid racemization
5 reveals a high state of repair in knee compared with hip osteoarthritic cartilage.
6 *Osteoarthritis Cartilage.* 2015 Sep 28. doi: 10.1016/j.joca.2015.09.003.
- 7 25. Chandrasekhar, S., M.A. Esterman, and H.A. Hoffman, Microdetermination of
8 proteoglycans and glycosaminoglycans in the presence of guanidine
9 hydrochloride. *Anal Biochem,* 1987. 161(1): p. 103-8.
- 10 26. Vilím V, Olejárová M, Macháček S, Gatterová J, Kraus VB, Pavelka K. Serum
11 levels of cartilage oligomeric matrix protein (COMP) correlate with radiographic
12 progression of knee osteoarthritis. *Osteoarthritis Cartilage.* 2002 Sep; 10(9):707-
13 13.
- 14 27. Maroudas A. Glycosaminoglycan turn-over in articular cartilage. *Philos Trans R*
15 *Soc Lond B Biol Sci.* 1975 Jul 17; 271(912):293-313.
- 16 28. Malesmud CJ. Changes in proteoglycans in osteoarthritis: biochemistry,
17 ultrastructure and biosynthetic processing. *J Rheumatol Suppl.* 1991 Feb;27:60-
18 2.
- 19 29. Ling W, Regatte RR, Navon G, Jerschow A. Assessment of glycosaminoglycan
20 concentration in vivo by chemical exchange-dependent saturation transfer
21 (gagCEST). *Proc Natl Acad Sci U S A.* 2008 Feb 19;105(7):2266-70.
- 22 30. Vincent TL, McLean CJ, Full LE, Peston D, Saklatvala J. FGF-2 is bound to
23 perlecan in the pericellular matrix of articular cartilage, where it acts as a

- 1 *chondrocyte mechanotransducer. Osteoarthritis Cartilage. 2007 Jul;15(7):752-*
2 *63.*
- 3 31. Alexopoulos LG, Haider MA, Vail TP, Guilak F. *Alterations in the mechanical*
4 *properties of the human chondrocyte pericellular matrix with osteoarthritis. J*
5 *Biomech Eng. 2003 Jun;125(3):323-33.*
- 6 32. Guilak F, Alexopoulos LG, Upton ML, et al. *The pericellular matrix as a*
7 *transducer of biomechanical and biochemical signals in articular cartilage. Ann*
8 *NY Acad Sci. 2006 Apr; 1068:498-512*
- 9 33. Gallagher JA, Ranganath LR, Boyde A. *Lessons from rare diseases of cartilage*
10 *and bone. Curr Opin Pharmacol. 2015 Jun;22:107-14.*
- 11 34. Maathuis PG, Driessen AP. *Painted black. Ann Rheum Dis. 2002 Feb;61(2):100-*
12 *1.*
- 13 35. Damian LO, Felea I, Boloşiu C, Botar-Jid C, Fodor D, Rednic S. *A case of*
14 *alkaptonuria - ultrasonographic findings. Med Ultrason. 2013 Dec;15(4):321-5.*
- 15 36. Mistry JB, Jackson DJ, Bukhari M, Taylor AM. *A role for interleukins in*
16 *ochronosis in a chondrocyte in vitro model of alkaptonuria. Clin Rheumatol. 2015*
17 *Oct 16.*

19 **Figure Legends**

20 Figure 1: Higher total protein in AKU and OA cartilage compared to Non-OA samples.
21 *Graph showing extractable protein (mg soluble protein/ml) from AKU (n=6), OA (n=12)*
22 *and Non-OA (n=6) for independent cartilage specimens. Error bars represent the mean*
23 *of the n in each group \pm 95% CI.*

1 Figure 2: Lower GAG in AKU cartilage compared to OA and Non-OA samples. *Graph*
2 *showing extractable glycosaminoglycan (GAG) (ug GAG/mg protein) from AKU (n=6),*
3 *OA (n=12) and Non-OA (n=6) for independent cartilage specimens. A; all samples; both*
4 *hips and knees. B; hip only samples. Error bars represent the mean of the n in each*
5 *group $\pm 95\%$ CI.*

6 Figure 3: Higher D/T COMP ratio in AKU cartilage compared to OA. *Graph showing*
7 *extractable D-COMP as a proportion of total COMP (D/T) from AKU (n=6), OA (n=12)*
8 *and Non-OA (n=6) for independent cartilage specimens. A; all samples; both hips and*
9 *knees. B; hip only samples. Error bars represent the mean of the n in each group $\pm 95\%$*
10 *CI.*

11 Figure 4: Lower total COMP and lower deamidated COMP in AKU cartilage compared
12 to OA and Non-OA. *Graph showing extractable total COMP (T-COMP) (T-COMP/mg*
13 *protein) from AKU (n=6), OA (n=12) and Non-OA (n=6) cartilage specimens. A; all*
14 *samples; both hips and knees. B; hip only samples. Graph showing extractable*
15 *deamidated-COMP (D-COMP) (D-COMP/mg protein) from AKU (n=6), OA (n=12) and*
16 *Non-OA (n=6) cartilage specimens. C; all samples; both hips and knees. D; hip only*
17 *samples. Error bars represent the mean of the n in each group $\pm 95\%$ CI.*

18 Figure 5: Higher racemized Asx content in PBS and insoluble extract fractions of AKU
19 cartilage compared to OA and Non-OA samples. *Graphs showing the amount of*
20 *racemized Asx (D form) as a proportion of non-racemized Asp (L form) represented by*
21 *the D/D+L ratio in AKU, OA and Non-OA cartilage specimens. A; Asx Ratio (D/D+L) in*
22 *PBS extract of all samples; both hips and knees from AKU (n=6), OA (n=12) and Non-*
23 *OA (n=6) for independent cartilage specimens B; Asx Ratio (D/D+L) in PBS extract*

1 from hip only samples from AKU (n=4), OA (n=7) and Non-OA (n=5) for independent
2 cartilage specimens **C**; Asx Ratio (D/D+L) in Insoluble extract all samples; both hips
3 and knees from AKU (n=6), OA (n=12) and Non-OA (n=6) for independent cartilage
4 specimens **D**; Asx Ratio (D/D+L) in Insoluble extract from hip only samples from AKU
5 (n=4), OA (n=7) and Non-OA (n=5) for independent cartilage specimens Error bars
6 represent the mean of the n in each group \pm 95% CI.

7 Figure 6: H&E stained section of alkaptonuric femoral head showing pigmented articular
8 cartilage and pigmented articular cartilage impacted in the marrow. **A**; Histological
9 section showing full depth articular cartilage pigmentation with numerous empty
10 chondrocyte lacunae (arrows). There is pericellular pigmentation around chondrocyte
11 lacunae in the calcified cartilage. There are also areas where the calcified cartilage is
12 not continuous with either the subchondral bone below it or hyaline articular cartilage
13 above it. Bar = 50 μ m. **B**; Shards of pigmented cartilage can be seen impacted within the
14 marrow space of the bone marrow cavity (C). The cartilage is surrounded by
15 inflammatory cells, some of which show intracellular pigmentation (arrow). Fibrous
16 tissue response of the marrow cavity can be seen by numerous strands of matrix
17 peripherally around the cartilage (asterisk). Bar = 10 μ m. **C**; 10 micron non-decalcified
18 cryostat section of AKU tissue stained with von Kossa. Densely mineralised bone, black,
19 can be seen beneath isogenous zones of chondrocytes located in the hyaline articular
20 cartilage which shows ochronotic colouration but absence of similar black staining
21 indicative of calcium deposition.