

# Accepted Manuscript

Characterization of a novel member of genus *Iflavivirus* in *Helicoverpa armigera*

He Yuan, Pengjun Xu, Xianming Yang, Robert I. Graham, Kenneth Wilson,  
Kongming Wu

PII: S0022-2011(17)30024-1

DOI: <http://dx.doi.org/10.1016/j.jip.2017.01.011>

Reference: YJIPA 6913

To appear in: *Journal of Invertebrate Pathology*

Received Date: 1 September 2016

Revised Date: 16 January 2017

Accepted Date: 19 January 2017



Please cite this article as: Yuan, H., Xu, P., Xianming Yang, Graham, R.I., Wilson, K., Wu, K., Characterization of a novel member of genus *Iflavivirus* in *Helicoverpa armigera*, *Journal of Invertebrate Pathology* (2017), doi: <http://dx.doi.org/10.1016/j.jip.2017.01.011>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Characterization of a novel member of genus *Iflavirus* in *Helicoverpa armigera***

He Yuan<sup>1#</sup>, Pengjun Xu<sup>1,2#</sup>, Xianming Yang<sup>1</sup>, Robert I. Graham<sup>3</sup>, Kenneth Wilson<sup>4</sup> and Kongming Wu<sup>1\*</sup>

<sup>1</sup>State Key Laboratory for Biology of Plant Diseases and Insect Pests, Institute of Plant Protection, Chinese Academy of Agricultural Sciences, Beijing 100193, P.R. China

<sup>2</sup>Tobacco Research Institute, Chinese Academy of Agricultural Sciences, Qingdao, P.R. China

<sup>3</sup>Crop and Environment Sciences, Harper Adams University, Edgmond, Shropshire TF10 8NB, UK

<sup>4</sup>Lancaster Environment Centre, Lancaster University, Lancaster LA1 4YQ, UK

#These authors contributed equally to this work.

\*Corresponding author

Mailing address: State Key Laboratory for Biology of Plant Diseases and Insect Pests, Institute of Plant Protection, Chinese Academy of Agricultural Sciences, Beijing 100193, P. R. China

E-mail: [kmwu@ippcaas.cn](mailto:kmwu@ippcaas.cn)

## Abstract

The cotton bollworm, *Helicoverpa armigera*, is one of the most important agricultural pests of many economic crops worldwide. Herein, we found a novel single-strand RNA virus by RNA-Seq and Polymerase Chain Reaction (PCR) method in *H. armigera* named *Helicoverpa armigera iflavivirus* (HaIV), which possessed a genome with 10,017 nucleotides in length and contained a single large open reading frame (ORF) encoding a putative polyprotein of 3,021 amino acids with a predicted molecular mass of 344.16 kDa and a theoretical isoelectric point (pI) of 6.45. The deduced amino acid sequence showed highest similarity (61.0%) with the protein of *Lymantria dispar Iflavirus 1*. Phylogenetic analysis with putative RdRp amino acid sequences indicated that the virus clustered with members of the genus *Iflavirus*. The virus was mainly distributed in the fat body of its host and was found to be capable of both horizontal and vertical transmission. The efficiency of perorally horizontal transmission was dose dependent (100% infection rate with a viral dose of  $10^8$  copies / $\mu$ l) while vertical transmission efficiency was found to be relatively low (< 28.57%). These results suggest that we have found a novel member of genus *Iflavirus* in *H. armigera*.

**Keywords:** *Helicoverpa armigera*, iflavivirus, RNA virus, horizontal transmission, vertical transmission, tissue distribution

## 1. Introduction

Insects are susceptible to a variety of pathogens, which can result in chronic or lethal infections (Burden et al., 2003). Historically, viruses have been isolated and subsequently studied after the observation of overt disease symptoms in the host. However, there are also covert infections found within hosts displaying no obvious signs of disease, which could remain undetected (Burden et al., 2003; Kemp et al., 2011; Murillo et al., 2011). These covert persistent viral infections, such as baculoviruses, are almost ubiquitous in many lepidopteran insect species and their discovery is driving further research into the dynamics and behavior of covert infections and their role in the ecology of host populations, especially those of economic and agricultural importance (Graham et al., 2015).

The cotton bollworm, *Helicoverpa armigera*, is one of the most important agricultural pests of cotton and other important economic crops worldwide. The adult moth is highly migratory, and populations have been reported in Australia, Asia, Africa, Europe (Feng et al., 2007; Wu and Guo, 2005) and most recently from South America (Tay et al., 2013). Since the introduction of Bt-cotton into China in the 1990s, the *H. armigera* population has declined dramatically. However, several studies have reported that Bt-resistance has evolved in the field (Gunning et al., 2005; Zhang et al., 2012). Thus, other forms of biological pest control, including the use of host-specific viral pesticides, derived from baculovirus (Allaway and Payne, 1984; Chen et al., 2001; Fuxa, 2004; Sun et al., 2002; Sun et al., 2004), small RNA viruses (Christian et al., 2005) and densoviruses (El-Far et al., 2012), has attracted more attention from

researchers. Moreover, high-through-put analytical methods such as metagenomics and RNA sequencing provide sensitive and effective methods for the discovery of novel viruses and asymptomatic disease agents that may be useful as biological control products (Diatchenko et al., 1996; Ge et al., 2012; Marguerat and Bähler, 2009; Mokili et al., 2012; Radford et al., 2012; Roossinck et al., 2015), or conversely, may negatively or positively impact upon the biopesticide products being used. For example, the recent discovery of a novel densovirus (HaDV2) from healthy migratory cotton bollworms revealed that HaDV2 infection significantly increased host resistance to the host-specific baculovirus HaNPV and to the generalist biopesticide Bt toxin (Xu et al., 2014). The development of molecular tools and next generation sequencing technology paves the way for a greater understanding as to how we may manipulate the host-pathogen system, with the aim of reducing pest outbreaks and economic crop damage.

In this study, a novel virus infecting *H. armigera* named Helicoverpa armigera iflavivirus (HaIV), was discovered by RNA-Seq. Alignment and phylogenetic analysis revealed that the virus showed a high sequence identity with members of the *Iflavirus*, which is the only genus within the family *Iflaviridae*. Members of this genus possess a single positive-strand RNA genome and share several common features, including:

- (1) non-enveloped icosahedral particles measuring 30 nm in diameter;
- (2) genome translation into a polyprotein;
- (3) the viral coat proteins containing three jelly-roll domains;
- (4) a three-domain containing a superfamily III helicase, a (cysteine) proteinase with a chymotrypsin-like fold and an RNA-dependent RNA polymerase

(RdRp) (Le Gall et al., 2008). The genome of *Iflavirus* is monocistronic with one single large open reading frame (ORF) encoding a single large polyprotein. To date, only nine species of iflaviruses have been recognized by The International Committee on Taxonomy of Viruses (ICTV), including *deformed wing virus* (Lanzi et al., 2006), *Ectropis obliqua virus* (Wang et al., 2004), *Infectious flacherie virus* (Isawa et al., 1998), *Lygus lineolaris virus 1* (Perera et al., 2012), *Nilaparvata lugens honeydew virus 1* (Murakami et al., 2013), *Perina nuda virus* (Wu et al., 2002), *Sacbrood virus* (Ghosh et al., 1999), *Slow bee paralysis virus* (de Miranda et al., 2010), and *Varroa destructor virus 1* (Ongus et al., 2004), although other iflaviruses have been reported (Silva et al., 2015; Suzuki et al., 2015). Herein, we report the nucleotide sequence, genome organization, phylogeny, transmission and tissue distribution of HaIV.

## 2. Materials and Methods

### 2.1 Insect culture

A laboratory colony of *H. armigera* was originally captured in 2005 from Langfang (Hebei province, China). *H. armigera* larvae were reared on an artificial diet (Liang et al., 2008) and adult moths were cultured with a 10% sugar and 2% vitamin mix (Liang et al., 1999) at  $25 \pm 1^{\circ}\text{C}$  with a 14:10, light: dark photoperiod.

### 2.2 Transcriptome analysis and annotation

For transcriptome analysis, Illumina RNA-sequencing was conducted by Novogene (Beijing, China). Four 5<sup>th</sup> instar larvae (one day post-ecdysis) were individually collected and total RNA isolated using the TRIzol kit (Invitrogen,

Carlsbad, CA, USA), according to the manufacturer's instructions. RNA samples were dissolved in RNase-free water and used to construct the cDNA library of *H. armigera* with suitable fragments (about 200bp). Paired-end transcriptome sequencing was subsequently performed using an Illumina HiSeq™ 2000. Adaptor sequences and low-quality reads were trimmed and clean reads were used for *de novo* assembly using Trinity (Grabherr et al., 2011). The assembled contigs were annotated using BLASTx and BLASTn against the NCBI non-redundant nucleic acid database (NT) and the NCBI non-redundant protein database (NR), using a cut-off E-value of  $10^{-5}$ .

### 2.3 Cloning the iflavirus genome of *H. armigera*

Total RNA was extracted from individual adult moths reared in laboratory using the TRIzol reagent kit as described above. Single-stranded cDNA was synthesized using the FastQuant RT Kit (Tiangen, Beijing, China), according to the manufacturer's instructions. Based on the assembled sequence from RNA-seq, nine pairs of primers were subsequently designed (Table S1). The genome of the iflavirus isolated from *H. armigera* was amplified and sequenced using cDNA as template, using the following PCR program: 4 min at 94 °C; 30 s at 94 °C, 30 s at 55 °C, and 2 min at 72 °C for 40 cycles. The PCR product was purified, inserted into the pEASY-T cloning vector (TransGen, Beijing, China), and sequenced.

### 2.4 Sequence and phylogenetic analysis

The open reading frame (ORF) of the viral genome was predicted using ORF finder at NCBI (<http://www.ncbi.nlm.nih.gov/gorf/gorf.html>). The amino acid

sequences encoding non-structural proteins were compared with members of the order *Picornavirales* using Clustal W (Thompson et al., 1994). The complete nucleotide sequence of the virus described in this study was submitted to GenBank under accession number KX228231. The deduced RdRp amino acid sequence of the new virus, together with that of members of the family *Iflaviridae*, was used in phylogenetic analysis. *Acyrthosiphon pisum virus* and *Spodoptera exigua virus AKJ-2014* were used as outgroup. The phylogenetic trees were constructed using the maximum likelihood method with a bootstrap of 1,000 replicates in MEGA6 (Tamura et al., 2013). Gaps were regarded as a complete deletion unless specifically noted.

## 2.5 Virus detection and quantification

A partial sequence exhibiting high similarities with known iflaviruses was identified from the RNA-seq data. For the detection of the novel virus, a pair of specific primers, VPF1/VPR1 (Table S1) were designed to amplify a PCR product of 593bp, according to the assembled sequence of the virus in *H. armigera*. The PCR program used was as follows: 30s at 94°C, 30s at 57°C, and 30s at 72°C for 40 cycles. To assess the detection threshold of the virus assay, a 10-fold dilution series of cDNA ( $3.37 \times 10^9$  to  $3.37 \times 10^1$  copies/ $\mu$ l) containing the virus was made and tested using the VPF1/VPR1 primers. 10 $\mu$ l of each PCR product was analysed by agarose gel electrophoresis.

Copy numbers of the novel virus were quantified using a standard curve by an absolute quantification qPCR method (Wong and Medrano, 2005). For quantification, the primers and probes of the virus (VPF/VPR, Table S1) were designed. A fragment

was amplified using the primers and cloned into the pEASY-T Cloning Vector (TransGen, Beijing, China) and sequenced. The PCR program was as follows: 30s at 94°C, 30s at 54°C, and 30s at 72°C for 40 cycles. The plasmid was used for the construction of the standard curve. Virus quantification was conducted with TaqMan in 20μl reaction agent, which comprised 1μl of template DNA, 2×SuperReal PreMix (Probe, Tiangen, Beijing, China), 50×ROX Reference Dye, 0.2mM of each primer and 0.4mM probe. The thermal cycling conditions on a 7500 Real-time PCR System (Applied Biosystems) were as follows: 40 cycles of 95°C for 3s, 60°C for 30s. The standard curve equation of  $y=-0.9990x+41.6662$  ( $y$ =the logarithm of plasmid copy number to base 2,  $x=Ct$  value,  $R^2=0.9998$ ) was used to calculate the copy number of the virus (Fig. S1).

## 2.6 Transmission of the virus

Filtered liquid containing an unpurified form of the iflavirus was prepared (Xu et al., 2014). RNA was isolated and the concentration of the virus was quantified. A NONINF strain was established from a single breeding pair that was not infected with the iflavirus, according to the method described by Xu et al. (2014). An infected line (INF strain) of *H. armigera* was established by orally infecting NONINF strain larvae with the filtered liquid. Fifth-instar larvae were randomly selected to detect the infection rate of the virus.

For quantification of vertical transmission rates, four pair-types ( $\text{♀}+/\text{♂}+$ ,  $\text{♀}+/\text{♂}-$ ,  $\text{♀}/\text{♂}+$ , and  $\text{♀}/\text{♂}-$ ) were mated. Positive and negative individuals were from INF and NONINF culture strains respectively. The infection rate of the virus in offspring was

detected from 3<sup>rd</sup> instar offspring larvae.

For quantification of horizontal transmission rates, non-infected NONINF strain neonates were provided artificial diet with different concentrations of the virus: 10<sup>8</sup>, 10<sup>7</sup>, 10<sup>6</sup>, 10<sup>5</sup>, 10<sup>4</sup>, 0 (virus-free control) copies /μl for 2 days, then transferred to a 24-well plate and reared in glass tubes until eclosion. Newly eclosed adult moths were sampled to determine the infection rate of the virus under different titer regimes. This diet contamination assay (mimicking a similar mechanism in wild populations) was conducted according to the method described by Xu et al. (2014). Infection rate was detected by PCR and larval frass from the INF strain was used to quantify the virus copy number by qPCR.

## 2.7 Quantification of the virus in eggs and different tissues

To test whether vertical transmission was transovum or transovarial, we quantified the copy number of the virus in three groups of both non-treated (control) and sodium hypochlorite treated eggs (n=50 eggs per group) of *H. armigera*. The eggs were from INF strain breeding pairs, that is both females and males were infected with the virus and treated according to the method described by Xu et al. (2014). Third instar larvae originating from these eggs were also sampled to quantify the copy number of the virus.

To examine the virus infection in different body tissues of the host, infected fifth-instar larvae were dissected into foregut, midgut, hindgut, hemolymph, malpighian tubules and fat body, using a fresh scalpel for every cut to avoid virus cross contamination. In addition, infected females and males were dissected into brain,

muscle, wing, malpighian tubule, fat body, ovary/testis and gut as described above. Total RNA was extracted from the different body parts (both larval and adult stages) and cDNA was used as a template to quantify the copy numbers of the virus by qPCR. The cDNA sample of each body part was replicated three times. The copy number of the virus in each tissue was calculated, and these were summed to determine total copy numbers of the virus in each individual. The percentage of virus in each tissue in the same individual was statistically analyzed (larvae: n = 9; adult males: n= 6; adult females: n= 9) (Xu et al., 2014).

To examine virus replication in different developmental stages of the host, larvae 24h to 240h post-infection (1<sup>st</sup>-5<sup>th</sup> instar) and newly eclosed adults were sampled. Absolute quantification qPCR and the comparative  $2^{-\Delta\Delta Ct}$  method (Livak and Schmittgen, 2001) were used, respectively. *β-actin* (GenBank EU527017) was used as the reference gene to normalize the virus expression. The reaction was conducted in 20μl reaction mixtures, containing 10μl of SuperReal PreMix Plus (TIANGEN, Beijing, China), 0.6μl of primers (10μM), 1μl of sample cDNA, 0.5μl of Rox Reference Dye and 8.3μl of RNase-free ddH<sub>2</sub>O. The cycling parameters were: 95°C for 15min, followed by 40 cycles of 95°C for 10s and 62°C for 32s. To ensure reproducibility, each sample was carried out in three biological replicates and in three technical replicates.

## 2.8 Electron microscopy

Adult moths from the INF culture strain were collected. Then the virus were isolated and purified using the method of Sucrose Density Gradient Centrifugation

according to the method described by La Fauce et al (2007). Purified particles were negatively stained with 2% sodium phosphotungstate at pH 6.8 and observed with a transmission electron microscope.

### 3. Results

#### 3.1 Transcriptome analysis and annotation

Using RNA-Seq we sequenced the transcriptome of *H. armigera*, obtaining about 5 gigabases (in-depth) for each sample, and a total of 106,785 assembled contigs. Using BLASTx and BLASTn, 45609 genes (42.71% of transcripts) and 34383 genes (32.20% of transcripts), respectively, were obtained through BLAST hits, using the E-value cutoff. Because of the relatively short length of transcripts (mean size of 758.65 bp) and lack of genome reference, most of the assembled sequences could not be matched to any known genes. The E-value distribution of the best hit in the nr database showed that 57% of the mapped sequences had strong homology (smaller than 1.0E-50), whereas 43% of the homologous sequences ranged from 1.0E-5 to 1.0E-50 (Fig. S2A). Homologous genes came from several species, with 66% of the unigenes having the highest homology to genes from *Danaus plexippus*, followed by *Bombyx mori* (5%), *Tribolium castaneum* (3%), and *Helicoverpa armigera* (2%) (Fig. S2B). The RNA-seq original datasets generated in this study are available in the NCBI GEO database (accession number: GSE86914) (<https://www.ncbi.nlm.nih.gov/geo/>).

#### 3.2 The genome sequence of HaIV

The assembled contig of 1495 nucleotides (nt) in length from 103,935 reads

encoded actin of *H. armigera*, and one contig of 10008 nt in length assembled using 41719 reads showed high identity with *Lymantria dispar* iflavivirus 1. According to the reference sequence, we designed specific primers to amplify the viral genome sequence of HaIV containing the whole ORF, which was 10,017 nt in length, containing a single large ORF (between nt 733 and 9798) encoding a polyprotein of 3021 amino acids. It has a predicted molecular mass of 344.16 kDa and a theoretical isoelectric point (pI) of 6.45. The coding sequence was flanked by a 732bp 5' untranslated region (UTR) and a 219bp 3' UTR. The nucleotide base composition of the genome was 30.6%A, 35.6%U, 13.6%C and 20.2%G. The total A+U and G+C content were 66.2% and 33.8% respectively. Alignment analysis with putative amino acid sequences showed high identities with the members of genus *Iflavirus*, in which the highest identity was 61.0% with *L. dispar* iflavivirus 1 (Table S2). It contained all three conserved domains, including the three conserved domains in helicase sequences (Fig. 1A), the GXCG and GXHXXG conserved motifs in the protease sequences (Fig. 1B) and the eight conserved domains in the RdRp amino acid sequences (Fig. 1C). The conservation of HaIV polyprotein and the various segments were compared with those of *Heliconius erato* virus (HeIV) (Smith et al., 2014) and the results are summarized in Table 1.

### 3.3 Phylogenetic analysis

Neighbor-joining trees with Poisson model were constructed for the putative RdRp amino acid sequences (Fig. 2) of the family *Iflaviridae* using *Acyrthosiphon pisum* virus and *Spodoptera exigua* virus AKJ-2014 as the outgroup. The result

indicated that the HaIV clustered with members of the genus *Iflavivirus*.

### 3.4 Sensitivity of detection

An amplification product could be visualized by ethidium bromide staining when as little as  $3.37 \times 10^3$  copies/ $\mu\text{l}$  of HaIV in cDNA were used as template (Fig. S3).

### 3.5 Transmission of HaIV

The virus was capable of being vertically transmitted from both infected males and females, but the transmission efficiency was relatively low (< 28.6%) and that from infected females was higher than that from infected males (Table 2). To test whether vertical transmission of the virus was due to transovum or transovarial, quantification of INF eggs was conducted and indicated that the virus titers were 1000 times greater in non-treated eggs than in eggs treated with sodium hypochlorite solution (paired t-test:  $t = 15.649$ , d.f. = 5,  $P < 0.001$ ) (Fig. 3). 8 out of 17 individuals (47.06%) hatching from non-treated eggs contained more than  $1.0 \times 10^7$  copies/mg of HaIV, whereas larvae hatching from eggs treated with sodium hypochlorite solution contained no more than  $3.71 \times 10^2$  copies/mg of HaIV ( $n=10$ ), suggesting that transovum transmission was occurring via the surface of eggs (Fig. 4).

To examine horizontal transmission, NONINF strain neonates were exposed to different concentrations of virus. The results showed that the infection rate of NONINF strain larvae was dose-dependent, with 100% infection rates at a dose of  $10^8$  copies / $\mu\text{l}$  (Table 3). To examine the transmission efficiency through frass of larvae, we placed NONINF strain neonates in diet cells which had previously housed infected insects ( $n=12$ ) and quantified the copy number of the virus in frass of larvae. The

results showed that the frass contained no more than  $10^6$  copies /mg (Fig. 5). However, 75% (9 of 12 samples) of NONINF strain individuals could be infected via horizontal transmission by frass (Fig. 6).

### 3.6 Host tissue distribution

The copy number of the virus in different body tissues of *H. armigera* was quantified by qPCR. In both larvae and adults, the virus titers in the fat body were significantly higher than in other tissues: larvae:  $F = 11.32$ , df = 5,12,  $P < 0.001$  (Fig. 7A); adult females:  $F = 11.57$ , df = 6,56,  $P < 0.001$  (Fig. 7B); adult males:  $F = 2.89$ , df = 6,21,  $P=0.033$  (Fig. 7C). The virus was also detected in female ovaries and male testes, but at lower titers than in fat body.

### 3.7 Total levels of HaIV

Absolute quantification qPCR results showed that the copy number of HaIV increased over time, reaching the highest infection load ( $1.36 \times 10^7$  copies/mg) at 144h post infection and remaining stable after this time (Fig. 8A). In undertaking the relative quantification assay, all of the expression levels of HaIV were compared with those at 24h. The expression levels of HaIV increased over time and reached 10,000 fold at 240h, with an addition slight increase in expression during the adult stage (Fig. 8B).

### 3.8 Virus morphology

The virus particles purified from *H. armigera* were observed by using an electron microscope. They had an isometric appearance and an approximate diameter of 30nm (Fig. 9).

#### 4. Discussion

Recently, next-generation sequencing technology has provided a rapid approach to high-throughput sequence determination and allowed a wide diversity of novel viruses to be discovered (Ansorge, 2009; Mokili et al., 2012; Roossinck et al., 2015). Herein, a novel virus named *H. armigera iflavivirus* (HaIV) was detected in larvae of *H. armigera* by RNA-seq method. The genome organization of HaIV has the common features of other members within the family *Iflaviridae*. Briefly, the virus is monocistronic, with a single-stranded RNA genome, and contains a single large ORF encoding both structural and non-structural proteins. It also possesses the functional motifs of RNA helicase, protease and RdRp conserved in members of the order *Picornavirales* (van Oers, 2010). As members of the genus *Iflavirus*, the genomic sequence of HaIV is A/U rich (> 60% A/U). Based on the genomic nucleotide sequences and the amino acid sequences, phylogenetic analysis indicated HaIV clustered with the members of *Iflavirus*, suggesting that HaIV was a novel member of genus *Iflavirus* (Carrillo-Tripp et al., 2014; Silva et al., 2015; van Oers, 2010).

Viruses distribution in host tissues were diverse according to different virus species, such as HaDV2, a beneficial densovirus which could accelerate the growth rate of its insect host, was mainly distributed in fat body of *H. armigera* (Xu et al., 2014) and HaNPV, a lethal baculovirus, viral loads of which were higher in the head, legs and wings than in the abdomen and thorax (Graham et al., 2015). Previously, some members of *Iflavirus* don't appear to be lethal to their hosts, such as Kakugo virus (Fujiyuki et al., 2009) and Brevicoryne brassicae virus (Ryabov, 2007). And,

some members of *Iflavirus* are known to be harmful to their hosts by lead to developmental problems and host mortality (van Oers, 2010). For example, sacbrood virus (SBV) is mainly found in larval fat body cells of the honeybee, which may impact metabolic function of these cells, resulting in failure to pupate and ultimately causing death (Bitondi et al., 2006; Park et al., 2016). Interestingly, HaIV was also mainly distributed in the fat body both in larvae and adults while with much lower titers in both the ovary and testis, suggesting a similar function with other members of genus *Iflavirus*.

As beneficial virus, HaDV2 could be high efficiently transmitted by both the horizontal and vertical mode (Xu et al., 2014). However, the harmful virus (eg. HaNPV) could infect host through high horizontal transmission rate (86.2% mortality at a dose of  $10^7$  OBs/ml) whereas quite low vertical transmission rate ( $12.6\% \pm 2.0\%$ ) (Zhou et al., 2005). Members of *Iflavirus* can transmit horizontally and vertically. For example, DWV could vertically transmit from queens to both worker and drone offspring (Yue and Genersch, 2005), and horizontally transmit through *Varroa* mites (Bowen-Walker et al., 1999; Wilfert et al., 2016). Transovarial and horizontal transmission of SBV have also been found to occur (Shen et al., 2005). Herein, we also found that HaIV could be both horizontally and vertically transmitted, which the horizontally transmitted efficiency was dose-dependent and the vertically transmitted efficiency was quite low (< 28.57%). To examine the possibility of horizontal transmission on natural conditions, we performed diet contamination assay and the results suggested that although the dose of HaIV in frass was no more than  $10^6$ /mg,

the infection rate was 75%. The copy number of HaIV associated with eggs was significantly decreased by washing with sodium hypochlorite solution, suggesting that transovum transmission was occurring via the surface of eggs.

In conclusion, we report a novel virus isolated from the host lepidopteran *H. armigera* named *H. armigera* iflavivirus. Molecular characterization and phylogenetic analysis indicated that HaIV was a novel member of the genus *Iflavirus*. HaIV was found to be mainly distributed in the fat body of its host, and could be both horizontally and vertically transmitted but with low efficiency, suggesting a harmful factor to its host.

### Acknowledgments

This work was supported by Science Fund for Creative Research Groups of the National Science Foundation of China (No. 31321004) and the National Natural Science Foundation of China (Grant No. 31401752). KW was supported by a Biotechnology and Biological Sciences Research Council UK-China Partnering Award (ref: BB/L026821/1).

## Reference

- Allaway, G. P., Payne, C. C., 1984. Host range and virulence of five baculoviruses from lepidopterous hosts. *Ann. Appl. Biol.* 105, 29-37.
- Ansorge, W. J., 2009. Next-generation DNA sequencing techniques. *N. Biotechnol.* 25, 195-203.
- Bitondi, M. M., Nascimento, A. M., Cunha, A. D., Guidugli, K. R., Nunes, F. M., Simões, Z. L., 2006. Characterization and expression of the Hex 110 gene encoding a glutamine-rich hexamerin in the honey bee, *Apis mellifera*. *Arch. insect biochem.* 63, 57-72.
- Bowen-Walker, P., Martin, S., Gunn, A., 1999. The transmission of deformed wing virus between honeybees (*Apis mellifera* L.) by the ectoparasitic mite *Varroa jacobsoni* Oud. *J. Invertebr. Pathol.* 73, 101-106.
- Burden, J. P., Nixon, C. P., Hodgkinson, A. E., Possee, R. D., Sait, S. M., King, L. A., Hails, R. S., 2003. Covert infections as a mechanism for long-term persistence of baculoviruses. *Ecol. Lett.* 6, 524-531.
- Carrillo-Tripp, J., Krueger, E. N., Harrison, R. L., Toth, A. L., Miller, W. A., Bonning, B. C., 2014. *Lymantria dispar iflavivirus I* (LdIV1), a new model to study iflaviral persistence in lepidopterans. *J. Gen. Virol.* 95, 2285-2296.
- Chen, X., IJkel, W. F., Tarchini, R., Sun, X., Sandbrink, H., Wang, H., Peters, S., Zuidema, D., Lankhorst, R. K., Vlak, J. M., 2001. The sequence of the *Helicoverpa armigera* single nucleocapsid nucleopolyhedrovirus genome. *J. Gen. Virol.* 82, 241-257.
- Christian, P. D., Murray, D., Powell, R., Hopkinson, J., Gibb, N. N., Hanzlik, T. N., 2005. Effective control of a field population of *Helicoverpa armigera* by using the small RNA virus *Helicoverpa armigera stunt virus* (Tetraviridae: *Omegatetravirus*). *J. econ. ent.* 98,

1839-1847.

- de Miranda, J. R., Dainat, B., Locke, B., Cordoni, G., Berthoud, H., Gauthier, L., Neumann, P., Budge, G. E., Ball, B. V., Stoltz, D. B., 2010. Genetic characterization of slow bee paralysis virus of the honeybee (*Apis mellifera* L.). *J. Gen. Virol.* 91, 2524-2530.
- Diatchenko, L., Lau, Y., Campbell, A. P., Chenchik, A., Moqadam, F., Huang, B., Lukyanov, S., Lukyanov, K., Gurskaya, N., Sverdlov, E. D., 1996. Suppression subtractive hybridization: a method for generating differentially regulated or tissue-specific cDNA probes and libraries. *Proc. Natl. Acad. Sci. U S A* 93, 6025-6030.
- El-Far, M., Szelei, J., Yu, Q., Fediere, G., Bergoin, M., Tijssen, P., 2012. Organization of the ambisense genome of the *Helicoverpa armigera* densovirus. *J. Virol.* 86, 7024.
- Feng, H., Wu, K., Ni, Y. X., Cheng, D., Guo, Y., 2007. Return migration of *Helicoverpa armigera* (Lepidoptera: Noctuidae) during autumn in northern China. *B. Entomol. Res.* 95, 361-370.
- Fujiyuki, T., Matsuzaka, E., Nakaoka, T., Takeuchi, H., Wakamoto, A., Ohka, S., Sekimizu, K., Nomoto, A., Kubo, T., 2009. Distribution of Kakugo virus and its effects on the gene expression profile in the brain of the worker honeybee *Apis mellifera* L. *J. Virol.* 83, 11560-11568.
- Fuxa, J. R., 2004. Ecology of insect nucleopolyhedroviruses. *Agr. Ecosyst. Environ.* 103, 27-43.
- Ge, X., Li, Y., Yang, X., Zhang, H., Zhou, P., Zhang, Y., Shi, Z., 2012. Metagenomic analysis of viruses from bat fecal samples reveals many novel viruses in insectivorous bats in China. *J. Virol.* 86, 4620-4630.
- Ghosh, R., Ball, B., Willcocks, M., Carter, M., 1999. The nucleotide sequence of sacbrood virus of the honey bee: an insect picorna-like virus. *J. Gen. Virol.* 80, 1541-1549.

- Grabherr, M. G., Haas, B. J., Yassour, M., Levin, J. Z., Thompson, D. A., Amit, I., Adiconis, X., Fan, L., Raychowdhury, R., Zeng, Q., 2011. Full-length transcriptome assembly from RNA-Seq data without a reference genome. *Nature biotechnology*. 29, 644-652.
- Graham, R. I., Tummala, Y., Rhodes, G., Cory, J. S., Shirras, A., Grzywacz, D., Wilson, K., 2015. Development of a real-time qPCR assay for quantification of covert baculovirus infections in a major African crop pest. *Insects* 6, 746-759.
- Gunning, R. V., Dang, H. T., Kemp, F. C., Nicholson, I. C., Moores, G. D., 2005. New resistance mechanism in *Helicoverpa armigera* threatens transgenic crops expressing *Bacillus thuringiensis* Cry1Ac toxin. *Appl. Environ. Microb.* 71, 2558-2563.
- Isawa, H., Asano, S., Sahara, K., Iizuka, T., Bando, H., 1998. Analysis of genetic information of an insect picorna-like virus, infectious flacherie virus of silkworm: evidence for evolutionary relationships among insect, mammalian and plant picorna (-like) viruses. *Arch. Virol.* 143, 127-143.
- Kemp, E. M., Woodward, D. T., Cory, J. S., 2011. Detection of single and mixed covert baculovirus infections in eastern spruce budworm, *Choristoneura fumiferana* populations. *J. Invertebr. Pathol.* 107, 202-205.
- Koonin, E. V., Dolja, V. V., Morris, T. J., 1993. Evolution and taxonomy of positive-strand RNA viruses: implications of comparative analysis of amino acid sequences. *Crit. Rev. Biochem. Mol. Biology* 28, 375-430.
- Lanzi, G., de Miranda, J. R., Boniotti, M. B., Cameron, C. E., Lavazza, A., Capucci, L., Camazine, S. M., Rossi, C., 2006. Molecular and biological characterization of deformed wing virus of honeybees (*Apis mellifera* L.). *J. Virol.* 80, 4998-5009.

- Le Gall, O., Christian, P., Fauquet, C. M., King, A. M., Knowles, N. J., Nakashima, N., Stanway, G., Gorbatenya, A. E., 2008. Picornavirales, a proposed order of positive-sense single-stranded RNA viruses with a pseudo-T = 3 virion architecture. *Arch. Virol.* 153, 715-727.
- Liang, G., Tan, W., Guo, Y., 1999. An improvement in the technique of artificial rearing cotton bollworm. *Plant Protec.* 25, 15-17.
- Liang, G., Wu, K., Yu, H., Li, K., Feng, X., Guo, Y., 2008. Changes of inheritance mode and fitness in *Helicoverpa armigera* (Hubner) (Lepidoptera: Noctuidae) along with its resistance evolution to Cry1Ac toxin. *J. Invertebr. Pathol.* 97, 142-149.
- Livak, K. J., T. D. Schmittgen. 2001. Analysis of relative gene expression data using real-time quantitative PCR and the 2 (-Delta Delta C(T)) Method. *Methods.* 25: 402-408. Marguerat, S., Bähler, J., 2009. RNA-seq: from technology to biology. *Cell. Mol. Life Sci.* 67, 569-579.
- Mokili, J. L., Rohwer, F., Dutilh, B. E., 2012. Metagenomics and future perspectives in virus discovery. *Curr. Opin. Virol.* 2, 63-77.
- Murakami, R., Suetsugu, Y., Kobayashi, T., Nakashima, N., 2013. The genome sequence and transmission of an iflavirus from the brown planthopper, *Nilaparvata lugens*. *Virus Res.* 176, 179-187.
- Murillo, R., Hussey, M. S., Possee, R. D., 2011. Evidence for covert baculovirus infections in a *Spodoptera exigua* laboratory culture. *J. Gen. Virol.* 92, 1061-1070.
- Ongus, J. R., Peters, D., Bonmatin, J. M., Bengsch, E., Vlak, J. M., van Oers, M. M., 2004. Complete sequence of a picorna-like virus of the genus Iflavirus replicating in the mite *Varroa destructor*. *J. Gen. Virol.* 85, 3747-3755.
- Park, C., Kang, H., Jeong, J., Kang, I., Choi, K., Yoo, M.S., Kim, Y.H., Kang, S.W., Lim, H.Y., Yoon,

- B.S., 2016. In-situ Hybridization for the Detection of Sacbrood Virus in Infected Larvae of the Honey Bee (*Apis cerana*). *J. Comp. Pathol.* 154, 258-262.
- Perera, O. P., Snodgrass, G. L., Allen, K. C., Jackson, R. E., Becnel, J. J., O'Leary, P. F., Luttrell, R. G., 2012. The complete genome sequence of a single-stranded RNA virus from the tarnished plant bug, *Lygus lineolaris* (Palisot de Beauvois). *J. Invertebr. Pathol.* 109, 11-19.
- Radford, A. D., Chapman, D., Dixon, L., Chantrey, J., Darby, A. C., Hall, N., 2012. Application of next-generation sequencing technologies in virology. *J. Gen. Virol.* 93, 1853-1868.
- Roossinck, M. J., Martin, D. P., Roumagnac, P., 2015. Plant virus metagenomics: Advances in virus discovery. *Phytopathology* 105, 716-727.
- Ryabov, E. V., 2007. A novel virus isolated from the aphid *Brevicoryne brassicae* with similarity to Hymenoptera picorna-like viruses. *J. Gen. Virol.* 88, 2590-2595.
- Shen, M., Cui, L., Ostiguy, N., Cox-Foster, D., 2005. Intricate transmission routes and interactions between picorna-like viruses (Kashmir bee virus and sacbrood virus) with the honeybee host and the parasitic *varroa* mite. *J. Gen. Virol.* 86, 2281-2289.
- Silva, L. A., Ardisson-Araujo, D. M., Tinoco, R. S., Fernandes, O. A., Melo, F. L., Ribeiro, B. M., 2015. Complete genome sequence and structural characterization of a novel iflavirus isolated from *Opsiphanes invirae* (Lepidoptera: Nymphalidae). *J. Invertebr. Pathol.* 130, 136-140.
- Smith, G., Macias-Muñoz, A., Briscoe, A. D., 2014. Genome sequence of a novel iflavirus from mRNA sequencing of the butterfly *Heliconius erato*. *Genome Announc.* 2: e00398-14.
- Sun, X., Chen, X., Zhang, Z., Wang, H., Bianchi, F. J., Peng, H., Vlak, J. M., Hu, Z., 2002. Bollworm responses to release of genetically modified *Helicoverpa armigera* nucleopolyhedroviruses in cotton. *J. Invertebr. Pathol.* 81, 63-69.

- Sun, X., Wang, H., Sun, X., Chen, X., Peng, C., Pan, D., Jehle, J. A., van der Werf, W., Vlak, J. M., Hu, Z., 2004. Biological activity and field efficacy of a genetically modified *Helicoverpa armigera* single-nucleocapsid nucleopolyhedrovirus expressing an insect-selective toxin from a chimeric promoter. *Biol. Control* 29, 124-137.
- Suzuki, T., Takeshima, Y., Mikamoto, T., Saeki, J. D., Kato, T., Park, E. Y., Kawagishi, H., Dohra, H., 2015. Genome sequence of a novel iflavirus from mRNA sequencing of the pupa of *Bombyx mori* inoculated with *Cordyceps militaris*. *Genome Announc.* 3, e01039-15.
- Tamura, K., Stecher, G., Peterson, D., Filipski, A., Kumar, S., 2013. MEGA6: molecular evolutionary genetics analysis version 6.0. *Mol. Biol. evol.* 30, 2725-2729.
- Tay, W. T., Soria, M. F., Walsh, T., Thamazoni, D., Silvie, P., Behere, G. T., Anderson, C., Downes, S., 2013. A brave new world for an old world pest: *Helicoverpa armigera* (Lepidoptera: Noctuidae) in Brazil. *PLoS One*. 8, e80134.
- Thompson, J. D., Higgins, D. G., Gibson, T. J., 1994. CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic acids res.* 22, 4673-4680.
- van Oers, M. M., 2010. Genomics and biology of Iflaviruses. *Insect Virology*. Caister Academic Press, Norfolk. 231-250.
- Wang, X., Zhang, J., Lu, J., Yi, F., Liu, C., Hu, Y., 2004. Sequence analysis and genomic organization of a new insect picorna-like virus, *Ectropis obliqua* picorna-like virus, isolated from *Ectropis obliqua*. *J. Gen. Virol.* 85, 1145-1151.
- Wong, M. L., Medrano, J. F., 2005. Real-time PCR for mRNA quantitation. *Biotechniques* 39, 75-85.
- Wu, C., Yang, H., Lo, C., Wang, C., 2002. A *Perina nuda* cell line (NTU-Pn-HF) from pupal ovary

- that is persistently infected with a picorna-like virus (PnPV). *Appl. Entomol. zool.* 37, 171-179.
- Wu, K., Xu, G., Guo, Y., 1997. Observations on migratory activity of cotton bollworm woths across the Bohai Gulf in China. *Acta. Phytophy. Sin.* 25, 337-340.
- Xu, P., Liu, Y., Graham, R. I., Wilson, K., Wu, K., 2014. Densovirus is a mutualistic symbiont of a global crop pest (*Helicoverpa armigera*) and protects against a baculovirus and Bt biopesticide. *PLoS Pathog.* 10, e1004490.
- Yue, C., Genersch, E., 2005. RT-PCR analysis of deformed wing virus in honeybees (*Apis mellifera*) and mites (*Varroa destructor*). *J. Gen. Virol.* 86, 3419-3424.
- Zhang, H., Tian, W., Zhao, J., Jin, L., Yang, J., Liu, C., Yang, Y., Wu, S., Wu, K., Cui, J., Tabashnik, B. E., Wu, Y., 2012. Diverse genetic basis of field-evolved resistance to Bt cotton in cotton bollworm from China. *Proc. Natl. Acad. Sci. U S A* 109, 10275-10280.
- Zhou, M., Sun, X., Sun, X., Vlak, J. M., Hu, Z., van der Werf, W., 2005. Horizontal and vertical transmission of wild-type and recombinant *Helicoverpa armigera* single-nucleocapsid nucleopolyhedrovirus. *J. Invertebr. Pathol.* 89, 165-175.

**Supporting information****Table S1 Primers used in this study.****Table S2 Percentage amino acid differences between accepted and proposed iflavirus species.****Table S3 Full names and GenBank accession numbers of the viruses used in this paper.****Figure S1 The standard curve for quantifying the number of virus particles using Taqman method.****Figure S2 (A) The E-value distribution of the top matches in the nr database. (B) The species distribution of best hits in the nr database.****Figure S3 Demonstration of the detection threshold of HaIV by agarose gel electrophoresis using the VPF1/VPR1 primers.**

**Figure legends**

**Figure 1. Protein domain alignments.** Shown are alignments of the helicase (A), protease (B) and RdRp sequences (C) domains of the novel virus (HaIV) with those of the other *Iflaviridae* (ApIV, DWV, EoV, HeIV, IFV, LyLV-1, PnV, SBV and VDV-1), some *Dicistroviridae* (ABPV, BQCV, CrPV and DCV) and one *Picornaviridae* (EMCV). Conserved regions corresponding to those recognized by Koonin et al. (1993) are indicated by bars above the protein alignment. Black shading indicates 100% sequence identity and other residues that are also conserved among these sequences are shaded in gray. The full names and the accession numbers of the virus are given in Table S3.

**Figure 2. Maximum likelihood phylogenetic analysis of the family *Iflaviridae*, with *Acyrthosiphon pisum* virus and *Spodoptera exigua* virus AKJ-2014 used as the outgroup.** The phylogenograms are based on the putative RdRp amino acid sequence (Fig. 2). The Genbank accession numbers are listed behind each virus species. Numbers at nodes represent bootstrap values as percentages estimated by 1000 replicates in an analysis using Clustal\_X software. Branches with less than 50% bootstrap support were collapsed.

**Figure 3. Viral load of HaIV in cotton bollworm eggs.** Absolute quantification of HaIV copy number per ug of host RNA in eggs washed or non-washed in 1% sodium hypochlorite (n=6).

**Figure 4. Viral load of HaIV in larvae hatching from non-treated eggs (C1-C17) and eggs treated with sodium hypochlorite solution (T1-T10).**

**Figure 5. Absolute quantification of HaIV copy number per mg of frass of larvae.**

LF: larval frass.

**Figure 6. The detection of HaIV in sample used in diet contamination assay. (a)**

PCR detection of HaIV in larvae which were reared on contaminated diet for 6 days;

(b) PCR detection of HaIV in frass of larvae; (c)  $\beta$ -actin was used as internal reference gene to test the integrity of each cDNA templates.

**Figure 7. Tissue distribution of the HaIV in A) larvae, B) adult females and C)**

**adult male cotton bollworms.** Within each figure, significant differences ascribed using Tukey tests are shown using different letters. Percentage (%) = the ratio of HaIV in different tissues (per mg), as described by Xu et al. (2014) (larvae: n = 9; adult males: n = 6; adult females: n = 9). Means  $\pm$  SE.

**Figure 8. HaIV levels in different developmental stages of host were tested using two methods.** (A) absolute quantification qPCR method. (B) the comparative  $2^{-\Delta\Delta Ct}$  method.

**Figure 9. Electron microscopy image showing HaIV particles purified from an extract from INF strain (100000X).**

**Table 1.** Sequence identity and similarity in an alignment between nucleotide and amino acid of HaIV and HeIV.

HaIV sequence	Sequence identity (similarity) (%)
<b>Nucleotide sequence</b>	
3'NTR	20.91
5'UTR	35.77
<b>Amino acid sequence</b>	
Entire polyprotein	58.78
RdRp	80.22
Protease	67.50
Helicase	92.17
Rhv_like	58.92
Rhv_like	77.95
CRPV_capsid	52.78

**Table 2.** Vertical transmission efficiency of HaIV.

Individuals	Number testing + ve	Number testing - ve	Transmission efficiency (%)
Female+/Male+	6	17	26.09
Female+/Male-	4	10	28.57
Female-/Male+	1	23	4.17
Female-/Male-	0	15	0.00

Infected individuals = “+”, uninfected individuals = “-”

**Table 3.** Detection of HaIV infecting larvae dosed at a range of concentrations.



Concentrations (copy number/uL)	Number testing + ve	Number testing - ve	Infection rate (%)
10 <sup>8</sup>	11	0	100.00
10 <sup>7</sup>	10	3	76.92
10 <sup>6</sup>	9	3	75.00
10 <sup>5</sup>	8	4	66.67
10 <sup>4</sup>	4	8	33.33
0	0	12	0





Infected individuals = “+ve”, uninfected individuals = “-ve”.


ACCEPTED MANUSCRIPT


ACCEPTED MANUSCRIPT

<b>A</b>		<b>B</b>
ABPV	: BTQIIVWVPECESGRGKGSQWTPEAIDINNSLLDN-----VDEMRN-FSKNIYMRNVEQEWINYQGQNINCKDXIFGQMRDSSSNPNEF :	84
ApIV	: RKLFFVYNYNCAPEIIGKSHITTNCSAICKDQGIT-----TETSLMCVLNATSKREWIDCDRGPCLVMILAFNIIKKGP-MFEDQV :	78
BQCV	: RNPFEVTVYNGETGVGKSTIYTYPICAIILKTIITFREGNTVMLESLKQHYKEMIYVYRAAEQEFWFGYTCQLVTVFIDLNFQQVDSSANPSLEI :	91
CrPV	: KMRITCWVTCESGIGKTCIYPCICIDILREMGIVK-----FLAYKHQAYARQVTEEWIYGNGOKINVYDIAFCIKDDDKTPNFEI :	82
DCV	: RMRICOMVCESGVGKTEIVYPICIDVLREMGMIK-----KDFHHHQVYGRQVETEWIYGKGOKINVYDIAFCIKDDDKTAANFEI :	82
DWV	: BRECFTHQMLGASGIGKSYITDSICSEILRASRTP-----VTGKICVVNLPLSDWICQDFCPVLCVDMWSVETST-TLDKQI :	78
EMCV	: BCEFVVWVIRGLAGQGKSLSSQVIAQAVSFKTIGFR-----QSVYSLPLPDSDFDGYENQFAAIMDLGONPDGS----DF :	71
EoV	: KPVIFCVWVYHSGCGKGSHWCDNVLTBEGSALGIN-----TANPIYTRSPDVFWNGTYGKLIISWCFAKTTTGE-TYRKQV :	77
HaIV	: RKMFFVYNYNCAPEIIGKSHITTDCTB1CKSQDIK-----TETSLMCVLNATSKREWIDCDRGPCLVMILAFNIRKGT-MLEDQI :	78
HeIV	: RRMFFVYNYNCAPEIIGKSHITTDCTB1CKSQDIK-----TETDLMCVLNATSKREWIDCDRGPCLVMILAFNIRKGT-MLEDQI :	78
IFV	: RFEFVVWVFCGPRGVGKSTIILQQQLIILILEQHKHGP-----KFEHPNPVYTRNPTSOQWNGYGPIVLYDILGAWSAAEIDPIVAG :	82
LyLV-1	: BYELFVVCVTCEDNSNIGKSFETEYVVAIILKSGIVNN-----CES-PVYYRQPGVKWFWSAYSNGPVIWYDILWLNINNVE-NVVMQV :	78
PnV	: KPVIFCVWVYHSGCGKGSHWCDNVLTBEGSALGIN-----TANPIYTRSPDVFWNGTYGKLIISWCFAKTTTGE-TYRKQV :	77
SBV	: BYELFVVCVTCEDNSNIGKSFETEYVVAIILKSGIVNN-----PHSGATYFRMPGSRFWSGYRDGPVIVYDILWANLTERPQ-ALMQOI :	79
VDV-1	: BRECFTHQMLGASGIGKSYITDSICSEILRASRTP-----VTGKICVVNLPLSDWICQDFCPVLCVDMWSVETST-TLDKQI :	78
<b>C</b>		
ABPV	: MEIIRTANIAFYPLHMZHLERKRTKFTTSKVIIMTNSN : 122	
ApIV	: AAIFIIVSPVVLVPKIAVEDK-GRPYNPDELFLNSN : 114	
BQCV	: FEIIRSSNIFPYPLHMZSISERKANTVFQSKVILICSSNN : 129	
CrPV	: FEVIRTCNTFFQQLHMLHIALDKK-NTFSAAEFLYTNIN : 119	
DCV	: FEVIRSCNTFFQQLHMLHIALDKK-NTFSAAEFLYTNIN : 118	
DWV	: NMIFQVHSPIVLSPLPKDLELGK-KMRYNPEFIFYNNINK : 115	
EMCV	: TTFCCMVSTNTFLPNNMSLERA-GTPTTSQVLAATINL : 108	
EoV	: SEISSLIEPTFPNPFPALEDK-RKIALAWAVVYSSNK : 114	
HaIV	: AAIFNVVSPVVLVPKIAVEDK-GRPYNPDELFLNSN : 115	
HeIV	: AAIFNVVSPVVLVPKIAVEDK-GKLYNPNPDELFLNSN : 115	
IFV	: -E-MAIKSCAITLEPLEKPRIEK-ESLMTSIVVGIABNH : 118	
LyLV-1	: SEIYQIKSTSLFIPEMDHLLEK-KIRGNPLIVLLINH : 115	
PnV	: SEISSLIEPTFPNPFPALEDK-RKIALAWAVVYSSNK : 114	
SBV	: SEIYQLKSTSTFIPEMDHLLEK-KIRGNPLIVLLINH : 116	
VDV-1	: NMIFQVHSPIVLSPLPKDLELGK-KMRYNPEFIFYNNINK : 115	
<b>B</b>		
GxCG		GxHxxG
ABPV	: ILRGQLELTIMP---TNGICGAAVINETQVIR-KIAGIEHVACIARG : 43	
ApIV	: RLSECYSMYS---TVGICGSAI--LCSTL-ERIIIGIHEACT--- : 37	
BQCV	: RLREAWEISLET--ISGDCCAPFVTNSKICPGKIGIGIHTAGG--- : 41	
CrPV	: -----CNEGYAPTRIGCIGCIGIGLYNYL-KERIIGIHMIACN--- : 36	
DCV	: ILQGVYTMVYH---GDEVGCGIIL---LSRNL-QREIIIGIHVACTEGL : 40	
EMCV	: TFNHCIMKRN--TRKGWCGSAI--LADLGSSKKTILGHSAG--- : 38	
EoV	: TGTSMIGQTWLGAQSVCQGSI---MSNNL---IICFFFVAQNLT : 40	
HaIV	: RLDECYSIYQYS---GVELOGSVI---VARNL-ERIIIGIHVAAELKSG : 40	
HeIV	: RLDCYCSIYRT---CKGICGSVIL--LSSTL-ERIIIGIHEACT--- : 37	
IFV	: SIPSQYDIIITPTQAKCNGLIAF-LDKQG---NIIICFFFVAQNLT : 41	
LyLV-1	: EMCGLLWVNNFS---KKCACGGIV-MKKNCN-TRIYAHIAARG--- : 38	
PnV	: KGTNMIGAQWLGSQSVCQGSI---MSNNL---IICFFFVAQNLT : 40	
SBV	: YANDVICHDS---QQCACGSLC--FLSRS-QREIVGMHBAARG--- : 38	
VDV-1	: ILQGVYTMVYH---GDEVGCGIIL---LSRNL-QREIIIGIHVACTEGL : 40	
<b>C</b>		
I		II
ABPV	: VRTFVNIVVTLKIERRPIEVQDL-KRVESSNGVYDFSTERMVYLGIHLHMLNN---TNEVSICIVYSCQDNKTVRLKTMGPKVLAGHSSTDGSLNVCIM---E : 103	
ApIV	: -----FVCKLKTICIDIKCISIPGKTRRISSPQVYTAKEQIENDL-SYQEAF-LSAHCIGICIDSLEPQIANYCTTYGDNINAGDMNNGGPSIMLKCVERAFDI : 103	
BQCV	: KLDHYFPLTICIDIKCISIPGKTRRISSPQVYTAKEQIENDL-SYQEAF-LSAHCIGICIDSLEPQIANYCTTYGDNINAGDMNNGGPSIMLKCVERAFDI : 105	
CrPV	: IIKDVVIVTLKIERRPIEVQDL-KRVESSNGVYDFSTERMVYLGIHLHMLNN---TNEVSICIVYSCQDNKTVRLKTMGPKVLAGHSSTDGSLNVCIM---E : 108	
DCV	: KISNVIFVTLKIERRPIEVQDL-KRVESSNGVYDFSTERMVYLGIHLHMLNN---TNEVSICIVYSCQDNKTVRLKTMGPKVLAGHSSTDGSLNVCIM---E : 109	
DWV	: IKRPTIFCICLTKLPVCRICPGRTRIISIPEQTFPQRQYLDMSYSAAR-IAEHICGIDMSLEWATNATRISKXGTHILTGOMYNNGGPLGDSVVAASAEFI : 109	
EMCV	: ---EVVYQTFKLEKPLVQQA-KTRAVDWEIYQKPLGKASKFPTQPGIPEKQFVAGLSPK-DVHEATAGFVAMQGFER-WYDVSNSDSTHSVAMFRLAAEE : 104	
EoV	: ILPPTTFFFEKKEIPLLKGKD---TRINGAPHTHTDPRVLMPEIAITT---INNKIAVGIVDVSQIHWGIAVHLHG---GADDVJDEYYGGGGFGEHSQWLTVCPI : 98	
HaIV	: IRIPTFDCKLKTICIDLEKCKIPGKTRRISSPQVYTAKEQIENDL-SYQNCR---INSEHICGIDMSLEPQIANYCTTYGDNINAGDMNNGGPSIMLKCVERAFDI : 109	
HeIV	: IRCPFTICLKTICIDLEKCKIPGKTRVSVFQFTAAPEQIENDL-SYQNCR---IAEHICGIDMSLEPQIANYCTTYGDNINAGDMNNGGPSIMLKCVERAFDI : 109	
IFV	: IRPCVWVAKHKEPLRPSLELRRFFGCVTVEPLFVWLNNSRLLPMLMAFQSFPE-IEAHHAIGIAPNSGIWVRAIRTLLEGKPMQIQLDPRMNSCAIAFKEVAKAFHI : 109	
LyLV-1	: VVFATIPLTUKERLRLIAWDSLGGTRVCMSCSPLVNSRQYIPEAFCIARWLFHAGVLDANGEEWTRIARWVSRKFGLIDYNNGGFAENAVVAGAAEI : 109	
PnV	: TLPLPTTFFFEKKEIPLLKGKD---TRINGAPHTHTDPRVLMPEIAITT---INNKIAVGIVDVSQIHWGIAVHLHG---GADDVJDEYYGGGGFGEHSQWLTVCPI : 98	
SBV	: VQPITFDTUPLKIERRPIEVQDL-KRVESSNGVYDFSTERMVYLGIHLHMLNN---TNEVSICIVYSCQDNKTVRLKTMGPKVLAGHSSTDGSLNVCIM---E : 109	
VDV-1	: ---PHTFDTUPLKIERRPIEVQDL-KRVESSNGVYDFSTERMVYLGIHLHMLNN---TNEVSICIVYSCQDNKTVRLKTMGPKVLAGHSSTDGSLNVCIM---E : 107	
II		
III		IV
ABPV	: KFADIANEFYD---DGSENALIRVHLLMDVNSTHICG---SVIMMTHSCSNSFAFPNCIFINSGKRMVLCLSKRYSSALNGTKCYVMKDFSKHVSIVSGDNVIN : 208	
ApIV	: IMMWYERYDND---EERQLIRVLLS---EILHAQHICLN-VVIVGPGCISGCGSPITPFLNSLVSISYRCGKSIITN-----QNSTMHENVRILYGDVCIN : 196	
BQCV	: VLGELLSKFIN---STEDEMLQQRATIISVNSLHNINQGILQWCKSLSGCHYAIILNSFVNMVCLMEANQKYSFTTAS---SEFRECGIAGGDHVS : 205	
CrPV	: SEYEWFKIIFNDVNTEDGKRNLMICIGLWTHVHSVW-SYGT-NVIMWTHSCSNSGPNTVILVNCNISVIRWILLIARKLAPEMQS---MKKRENNSVMSYGDNCIN : 213	
DCV	: IFVWVLQKFIDIESEGKRILCICLGLWSHVHSVHYPE-NVMMWTHSCSNSGPNTVILVNCNISVIRWILLIARKLAPEMQS---MKWENEVVALLYGDNVIN : 213	
DWV	: IIDWVLHYTHEE---DNKDEMKRVMWTTMAQELAPSHIYR-LVVRVCPGCSISGPSPDTINTSCLVIRLQGTLGIDT---LPLSEFSQNWWVCGDILIN : 205	
EMCV	: FFTP-----ENGDFPLTREYELAISIAPEE-KR-LITGGL-EACGAIMSMNTIMNIIAGLYLTYKIN-----FEFDUWKVJLSQGELLA : 188	
EoV	: AVAWCKHRLN---PERNLVVRVLLS---EILYARHICLN-VVIVGPGCISGCGSPITPFLNSLVSISYRCAKVITG---TLTGFWSHYTLWVY---D : 185	
HaIV	: IMAWYERYDND---PERNLVVRVLLS---EILYARHICLN-VVIVGPGCISGCGSPITPFLNSLVSISYRCAKVITG---QFEDIMHSNKLIVGDCVCIN : 202	
HeIV	: IMAWYERYDND---EERQLIRVLLS---EILYARHICLN-VVIVGPGCISGCGSPITPFLNSLVSISYRCAKVITG---ESFEVMHNVNKFITYGDVCIN : 202	
IFV	: IVDYRKWHCS---TWEIHLANLTFLDIALAEIIVLG---DVEKVNNGVLAQHPMVSVNSVNLALLNYMIAKIKTRR---ASEFFKLTYLIVGDCVWLS : 203	
LyLV-1	: MIRWVLNMVEG---VDETELRL-VIAA---ETTQSLHIAIL-LVYRQCGSISGAPDPTQRMVIVMIAIMIAVYLYLWPH---DRTLLWETFFEELDNECIGDEVLS : 206	
PnV	: AIAWCKYHCKV---DKEYEDVVSCLL---EIQNAYHVAGC-LVYVQLVCGSISGAPDPTQRMVIVMIAIMIAVYLYLWPH---TLTGFWSHYNWY---D : 185	
SBV	: MWRVTRMERVEG---VNEIEAY-TLH---ECLNSVHVS---TLYVQKCGSISGAPDPTQRMVIVMIAIMIAVYLYLWPH---ERGQTWESFKQNVEFCDGELIN : 206	
VDV-1	: IIDWVLNYTEE---DDKDEMKRVMWTTMAQELAPSHIYR-LVVRVCPGCSISGPSPITPFLNSLVSISYRCAKVITG---LPLSEFSRHWVVCXGELIN : 203	
V		
VI		VII
ABPV	: KFADIANEFYD---DGSENALIRVHLLMDVNSTHICG---SVIMMTHSCSNSFAFPNCIFINSGKRMVLCLSKRYSSALNGTKCYVMKDFSKHVSIVSGDNVIN : 208	
ApIV	: IMMWYERYDND---EERQLIRVLLS---EILHAQHICLN-VVIVGPGCISGCGSPITPFLNSLVSISYRCGKSIITN-----QNSTMHENVRILYGDVCIN : 196	
BQCV	: VLGELLSKFIN---STEDEMLQQRATIISVNSLHNINQGILQWCKSLSGCHYAIILNSFVNMVCLMEANQKYSFTTAS---SEFRECGIAGGDHVS : 205	
CrPV	: SEYEWFKIIFNDVNTEDGKRNLMICIGLWTHVHSVW-SYGT-NVIMWTHSCSNSGPNTVILVNCNISVIRWILLIARKLAPEMQS---MKKRENNSVMSYGDNCIN : 213	
DCV	: IFVWVLQKFIDIESEGKRILCICLGLWSHVHSVHYPE-NVMMWTHSCSNSGPNTVILVNCNISVIRWILLIARKLAPEMQS---MKWENEVVALLYGDNVIN : 213	
DWV	: IIDWVLHYTHEE---DNKDEMKRVMWTTMAQELAPSHIYR-LVVRVCPGCSISGPSPDTINTSCLVIRLQGTLGIDT---LPLSEFSQNWWVCGDILIN : 205	
EMCV	: FFTP-----ENGDFPLTREYELAISIAPEE-KR-LITGGL-EACGAIMSMNTIMNIIAGLYLTYKIN-----FEFDUWKVJLSQGELLA : 188	
EoV	: AVAWCKHRLN---PERNLVVRVLLS---EILYARHICLN-VVIVGPGCISGCGSPITPFLNSLVSISYRCAKVITG---TLTGFWSHYTLWVY---D : 185	
HaIV	: IMAWYERYDND---PERNLVVRVLLS---EILYARHICLN-VVIVGPGCISGCGSPITPFLNSLVSISYRCAKVITG---QFEDIMHSNKLIVGDCVCIN : 202	
HeIV	: IMAWYERYDND---EERQLIRVLLS---EILYARHICLN-VVIVGPGCISGCGSPITPFLNSLVSISYRCAKVITG---ESFEVMHNVNKFITYGDVCIN : 202	
IFV	: IVDYRKWHCS---TWEIHLANLTFLDIALAEIIVLG---DVEKVNNGVLAQHPMVSVNSVNLALLNYMIAKIKTRR---ASEFFKLTYLIVGDCVWLS : 203	
LyLV-1	: MIRWVLNMVEG---VDETELRL-VIAA---ETTQSLHIAIL-LVYRQCGSISGAPDPTQRMVIVMIAIMIAVYLYLWPH---DRTLLWETFFEELDNECIGDEVLS : 206	
PnV	: AIAWCKYHCKV---DKEYEDVVSCLL---EIQNAYHVAGC-LVYVQLVCGSISGAPDPTQRMVIVMIAIMIAVYLYLWPH---TLTGFWSHYNWY---D : 185	
SBV	: MWRVTRMERVEG---VNEIEAY-TLH---ECLNSVHVS---TLYVQKCGSISGAPDPTQRMVIVMIAIMIAVYLYLWPH---ERGQTWESFKQNVEFCDGELIN : 206	
VDV-1	: IIDWVLNYTEE---DDKDEMKRVMWTTMAQELAPSHIYR-LVVRVCPGCSISGPSPITPFLNSLVSISYRCAKVITG---LPLSEFSRHWVVCXGELIN : 203	
VII		
VIII		IX
ABPV	: FSDEVSLWVSEMEDEAERKLGYTFLDLGKNGEVFKWKRTHEDVQYLGKPPRPRYDSK---RKVWEABCMDTLEMPNVC : 285	
ApIV	: VSDMYKDIVNTETLSEFFKEVINYEDI---DSDVIIKYRNNWVFLGKSPILHPLN-SKEFIFLAPELQSLSRKCVN : 272	
BQCV	: VPEKYLSVLQQPITPFLMSKFGMFYIETKDE-BEIDFLSLRPELDSVYLNNSVYDYES-RQRYIAFSLDVLEMPMT : 282	
CrPV	: ISDRVVEWVQGQIISQEMKQMEIKHEYDE---GAGDMWKFPSIISIEHPLKQFREWSHQ---LQRTVAPQKDQVYEMLNIN : 288	
DCV	: IIARVVEWVQGQIISQEMKQMEIKHEYDE---AAGDIDIVKSRIKEFIFLXKREFREFSE-LQRTVAPQKDQVYEMLNIN : 288	
DWV	: VSINNMIDKAVTIGKFFSQXKMFEDQ---DXSGNTVKWRTQTATFLXHGEFLKHP---TRPVEIANLDRVSLEGTNT : 280	
EMCV	: TNQG---LFDKFRASLAKTGYKIDFA---NATSTFELNSTEDVFLXKREKKEGP---LYRFVWNEAREEAMLSY : 257	
EoV	: DTRRRETAATGDEBFQECMASGVIVVNR---DRSG-----VTSFLXKREPFLIDHRDVRVMLAPRPRFEDILIN : 252	
HaIV	: VSDEYIYSETNTSEFFKKYNIYEDI---DSDNIVPYRQDUDNVIPLGKCPVRHPH-SGVVIFIAFDEQSIRKCVN : 278	
HeIV	: VSDNYKDIVNTETLSEFFKKYNIYEDI---DSDNIIKYRQDUDNVIPLGKCPKLPS-SRAVEIAFDEQSIRKCVN : 278	
IFV	: LPKQLTTEE-CRKCAEEFKYDIKWVDSKEAIIIGEKPYD8F3KFBFPLS-GESE : 255	
LyLV-1	: VSEFVAEWVHGVSINHFEARRGVIVSSA---DRLSIVPTGTCIMNTSFLXKWWVWRHEK-WIGEWHGAIDTSLEECI : 282	
PnV	: DTRRRETAATGDEQQCMASGIVVNR---DRSG-----VTSFLXKREPFLIDHRDVRVMLAPRPRFEDILIN : 251	
SBV	: VTDKYKDVWALTISQELAQYGVIALEA-NAGDEVEAYTTLN1STFLXHGBBPHEPVE-YPHLWQSAWNSLNTTQ : 282	
VDV-1	: VSDEMIDRERAVTGDFFSRYKMEFDQ---DGSNTVWRWLTQATFLXHGBLKHP---TRPVEIANLDRVSLEGTNT : 278	









ACCEPTED  
MANUSCRIPT

## Highlights

We characterized a novel iflavirus in *Helicoverpa armigera* (HaIV).

HaIV was found to be mainly distributed in the fat body of its host.

HaIV could be both horizontally and vertically transmitted but with low efficiency.

## Graphical abstract

