


Accepted Manuscript

Title: A novel dual GLP-1/GIP receptor agonist alleviates cognitive decline by re-sensitizing insulin signaling in the Alzheimer icv. STZ rat model

Authors: Lijuan Shi, Zhihua Zhang, Lin Li, Christian Hölscher



PII: S0166-4328(17)30224-3
DOI: <http://dx.doi.org/doi:10.1016/j.bbr.2017.03.032>
Reference: BBR 10777

To appear in: *Behavioural Brain Research*

Received date: 3-2-2017
Revised date: 9-3-2017
Accepted date: 21-3-2017

Please cite this article as: Shi Lijuan, Zhang Zhihua, Li Lin, Hölscher Christian. A novel dual GLP-1/GIP receptor agonist alleviates cognitive decline by re-sensitizing insulin signaling in the Alzheimer icv. STZ rat model. *Behavioural Brain Research* <http://dx.doi.org/10.1016/j.bbr.2017.03.032>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

A novel dual GLP-1 / GIP receptor agonist alleviates cognitive decline by re-sensitizing insulin signaling in the Alzheimer icv. STZ rat model

Lijuan Shi¹, Zihua Zhang¹, Lin Li¹, Christian Hölscher^{2,3}

¹Key Laboratory of Cellular Physiology, Shanxi Medical University, Taiyuan, Shanxi, PR China.

²Biomedical and Life Science, Faculty of Health and Medicine, Lancaster University, Lancaster LA1 4YQ, UK

³Neurology department, The Second Affiliated Hospital of Shanxi Medical University, Taiyuan, Shanxi, PR China

running title: *Dual agonist effects in the STZ rat model*

Corresponding author:

Prof. Lin Li

Key Laboratory of Cellular Physiology,
Shanxi Medical University,

Taiyuan, PR China

Email: linlilin999@163.com

Highlights

Novel dual GLP-1/GIP receptor agonists are superior to single GLP-1 agonists
In the icv. STZ model of neurodegeneration, a novel dual agonist was highly effective at low doses
Cognition was protected, inflammation reduced
Insulin signalling was re-sensitized, apoptotic signalling reduced

Abstract

Alzheimer's disease (AD) is a progressive neurodegenerative disorder, accompanied by memory loss and cognitive impairments, and there is no effective treatment for it at present. Since type 2 diabetes (T2DM) has been identified as a risk factor for AD, the incretins glucagon-like peptide 1 (GLP-1) and glucose dependent insulinotropic polypeptide (GIP), promising antidiabetic agents for the treatment of type 2 diabetes, have been tested in models of neurodegenerative disease including AD and achieved good results. Here we show for the first time the potential neuroprotective effect of a novel dual GLP-1/GIP receptor agonist (DA-JC4) in the icv. streptozotocin (STZ)-induced AD rat model. Treatment with DA-JC4 (10 nmol/kg ip.) once-daily for 14 days after STZ intracerebroventricular (ICV) administration significantly prevented spatial learning deficits in a Y- maze test and Morris water maze tests, and decreased phosphorylated tau levels in the rat cerebral cortex and hippocampus. DA-JC4 also alleviated the chronic inflammation response in the brain (GFAP-positive astrocytes, IBA1-positive microglia). Apoptosis was reduced as shown in the reduced ratio of pro-apoptotic BAX to anti- apoptotic Bcl-2 levels. Importantly, insulin signaling was re-sensitized as evidenced by a reduction of phospho-IRS1^{Ser1101} levels and phospho-Akt^{Ser473} up-regulation. In conclusion, the novel dual agonist DA-JC4 shows promise as a novel treatment for sporadic AD, and reactivating insulin signaling pathways may be a key mechanism that prevents disease progression in AD.

Keywords: Alzheimer's disease; Incretin; Insulin signaling; Apoptosis; Inflammation; tau phosphorylation; Type 2 diabetes mellitus

1. Introduction

Alzheimer's disease (AD) is the most common neurodegenerative disorder and it is estimated that the population of dementia patient worldwide may reach 131.5 million in 2050 due to longer life-expectancy in the industrialized nations. The pathology of AD is characterized by the presence of neurofibrillary tangles

composed of hyperphosphorylated tau protein (p-tau) and amyloid- peptide accumulation, as well as neuronal loss in different brain regions which are associated with progressive memory loss and cognitive decline [1-3]. Drug therapies for AD are currently restricted and mainly depend on three cholinesterase inhibitors and the receptor inhibitor memantine, but these drugs cannot fundamentally halt or delay the progression of the disease at present [4-7].

Type 2 diabetes mellitus (T2DM) is a risk factor for developing AD [8-10]. Patients with T2DM present with learning and memory deficits [11] and there is an increase an approximately twofold risk of AD [8, 12]. Also, AD shares many pathophysiological features with T2DM, including insulin resistance, inflammatory stress and amyloid- peptide accumulation [13]. Based on these shared features, insulin resistance is one of the key underlying mechanisms [12, 14, 15].

Incretin hormones, glucagon-like peptide 1 (GLP-1) and glucose dependent insulintropic polypeptide (GIP) which can treat T2DM have been proposed as a novel therapeutic schedule for AD. Extensive preclinical studies show good effects in animal models of AD by reducing memory loss, synapse loss and amyloid plaque load [16-19], decrease the hyperphosphorylation of protein [20-23], exert anti-inflammatory function [24-26] and reduce neuronal loss [22, 27, 28]. A pilot study testing the GLP-1 analogue liraglutide in AD patients showed good protective effects in FDG-PET brain scans [29]. Other clinical trials in patients with AD or Parkinson's disease (PD) are currently ongoing [30]. Dual GLP-1 and GIP receptor agonists have been developed to treat T2DM and have shown first positive results in patients with diabetes [31]. We have tested one of these dual agonists named DA-JC1 in the MPTP mouse model of PD with good results [32, 33]. Here, we are testing the newer dual agonist DA-JC4 that has been optimized to cross the blood-brain barrier in the icv. streptozotocin (STZ) rat model of AD. STZ desensitizes insulin signaling in the brain [34, 35] and produces a range of pathological changes that are also found in the brains of AD patients [13, 36, 37], such as inducing cognitive impairment [21, 38-40], chronic inflammation in the brain [41] and enhanced tau protein phosphorylation [22, 42].

We tested the novel dual agonist DA-JC4 in this model and evaluated memory formation, tau phosphorylation, chronic inflammation, insulin re-sensitization, growth factor and apoptosis cell signaling.

2. Materials and methods

2.1 Drugs

Streptozotocin (STZ) was purchased from Sigma-Aldrich (St Louis, MO, USA). The dual agonist DA-JC4 was obtained from ChinaPeptides (Shanghai, China). The purity of the peptide was 95% which was

confirmed by reversed-phase high performance liquid chromatography (HPLC) and the peptide was identified using matrix-assisted laser desorption/ionisation time of flight (MALDI-TOF) mass spectrometry.

2.2 Animals and experimental groups

Male Sprague-Dawley rats (210-230g) were procured from the Animal Center of Shanxi Medical University. The animals were maintained in plastic cages (5 rats per cage) with the temperature of 25 ± 2 C and the humidity of $50 \pm 10\%$ and a 12 h light-dark cycle. Animals were allowed free access to food and water. All experimental procedures were conducted in accordance with guidelines for Care and Use of Laboratory Animals and approval of the Research and Ethics Committee of the University. At the first stage of experimental design, rats were randomly divided into four groups: 1) Control group: artificial cerebral spinal fluid (aCSF) intra-cerebral ventricular (ICV) injection plus saline intraperitoneal (IP) injection; 2) DA-JC4: aCSF ICV injection plus dual agonists IP injection; 3) STZ group: STZ ICV injection plus saline IP injection; 4) STZ + DA-JC4 group: STZ ICV injection plus dual agonist IP injection.

2.3. Drug administration

Rats in the STZ group received an ICV injection of STZ (3 mg/kg body weight) dissolved in 10 μ l aCSF (NaCl 140mM; KCl 3.0mM; CaCl₂ 2.5 mM; MgCl₂ 1.2 mM; NaH₂PO₄ 1.2 mM, PH 7.4), and the rats in the sham-operation received the same volume of aCSF. After stereotaxic surgery, rats were treated with a daily IP injection of DA-JC4 (10 nmol/kg body weight) or saline for 2 weeks.

2.4. Injection of STZ

We followed the protocol published previously [43]. For stereotaxic surgery, anesthesia was induced with 10% Chloral hydrate (0.3 ml/100g, ip). Animals were placed on stereotaxic frame with the following coordinates (0.8 mm posterior to bregma; 1.5mm lateral to the sagittal suture; 3.6 mm ventral) for the left side of lateral ventricle injection. STZ injection to the lateral ventricle was carried out with a Hamilton syringe in a 10 microliter volume over 15 s. The needle remained in position for additional 10 minutes to prevent reflux. The rats in the sham-operation group were given the ICV injection of aCSF in the same manner. After surgery, the rats were put in cage individually and allowed to access to food and water freely.

2.5 Behavioral procedure

At the nineteenth day after intracerebroventricular injection of STZ, behavioral estimations were tested in time sequence. The behavioral tests were conducted between 09:00 a.m. and 01:00 p.m. Experiments were carried out in a sound-proof and air-regulated experimental room. Rats were habituated at least 30 min before each test.

2.5.1 Y-maze test

Spatial working memory was investigated by recording spontaneous alternation behavior in a Y-maze apparatus [44]. The maze consisted of three arms ($50 \times 10 \times 20 \text{ cm}^3$ and 120° apart). Each rat was placed at the terminus of one arm and allowed to move freely through the maze for 8 min. When the rat's tail was entirely within the arm, entry was considered to be complete. The apparatus was cleaned with a 10% ethanol solution and then dried with a paper towel after each trial. The total number of arm entries (N) and the sequence of entries were recorded visually. Alternation behavior was defined as successive entries into all three arms on consecutive occasions. The alternation rate (%) = [the total number of alternations/ (N-2) *100] .

2.5.2 Morris water maze test

MWM task was performed to evaluate spatial learning and memory [45]. A black circular water tank with the 150 cm diameter and 60 cm height was used. The pool surrounded by a blue curtain was filled with tap water to a depth of 30 cm, in which the temperature of water was controlled to $28 \pm 1 \text{ }^\circ\text{C}$. A transparent platform (14 cm in diameter) was located in the midpoint of the target quadrant and submerged 1.5 cm beneath the surface of the water. Movements of rats were monitored and recorded via a video camera located above the pool. Rats were placed randomly in the water facing the wall in one of the four quadrants and allowed to swim freely to find the hidden platform. If a rat failed to find the platform within 120 s at the first day, it would be guided onto the platform for 15 s. A behavior software system (Ethovision 11.5, Noldus Information Technology, Wageningen, Netherlands) was used to record the escape latency of five consecutive days, the length of time to reach the platform. To control the influence of different animal's athletic ability, the swim speed was also recorded and analyzed. On day 6, the hidden platform was removed and probe trial was performed. The percentage of swimming time spent in the target quadrant was record in this phase (n = 8–10 each group).

2.6 Immunohistochemistry

After the behavior tests, rats (n=4-5) were transcardially perfused with saline and 4% paraformaldehyde. The brains were removed and fixed in 4% paraformaldehyde for 24 h. After fixation, the tissues were dehydrated and embedded in paraffin. 5 μ m sections were cut with a microtome and stored at 20 °C until immunohistochemistry was performed.

Sections were treated with xylene and graded ethanol solutions in turn. Endogenous peroxidase activity was blocked with 3% H₂O₂ for 10 min. Then the sections were pretreated using heat mediated antigen retrieval with citrate buffer. After blocked with 5% BSA, the sections were incubated with p-Tau (Ser396) (rabbit anti- p-Tau (Ser396); 1:1000, Abcam, Cambridge, MA, USA), GFAP (rabbit anti-GFAP; 1:100; Boster Biotechnology Co., Ltd. Wuhan, China) and IBA1 (rabbit anti-IBA1; 1:100; Boster Biotechnology Co., Ltd. Wuhan, China) at 37 °C for 2 h. Then they were incubated with biotinylated goat anti-rabbit IgG at 37 °C for 30 min and followed by the avidin-biotin peroxidase complex reagent (Boster Biotechnology Co., Ltd. Wuhan, China) at 37 °C for 30 min. The peroxidase was visualized with 3, 3-diaminobenzidine (DAB) (Zhongshan Golden Bridge Biotechnology Co., Ltd. Beijing, China). The photomicrographs of all stained sections were captured with a digital camera (Motic BA210) under a Zeiss light microscope and quantitatively analyzed using Image J software (Version 6.0, developed by National Institutes of Health). Stereological rules were applied [46] and analysis was blind to treatment.

2.7 Western blot

The hippocampus and cortex tissue (n=4-5) were dissected and stored at -80 °C for immunoblot analysis. These tissues were cut into pieces in cold RIPA lysis buffer (Beyotime Institute of Biotechnology, Shanghai, China). After 2 h, tissue lysates added with phenylmethanesulfonyl fluoride (PMSF) and phosphotransferase inhibitor were fully homogenized. Then protein supernatant was taken after centrifugation (14 000 \times g for 20minutes at 4°C). Protein concentration was measured using BCA protein assay (Boster Biotechnology Co., Ltd. Wuhan, China). Samples mixed with loading buffer to the same concentration were boiled for 5 min. Samples with equivalent amounts of protein were run on 8%, 10% or 12% SDS-polyacrylamide gel and transferred onto polyvinylidene difluoride (PVDF) membrane. After blocking the membrane with 5% bovine serum albumin (Boster Biotechnology Co., Ltd. Wuhan, China) in TBST (Tris-buffered saline contains 0.05% Tween-20) for 1 h, the membranes were probed overnight at 4°C with primary antibodies that specifically detect IRS-1 (1:500; Abcam, Cambridge, UK), p-IRS-1 (Ser1101) (1:750; Cell Signaling Technology, Danvers, MA), Akt (1:1000; Cell Signaling Technology, Danvers, MA), p-Akt (Ser473, 1:2000; Cell

Signaling Technology, Danvers, MA), Bcl-2 (1:500; Bioworld Technology Co., Ltd. Nanjing, China), Bax (1:500; Bioworld Technology Co., Ltd. Nanjing, China), and β -Actin (1:5000; Abcam, Cambridge, UK), followed by labeled with secondary antibodies (goat-anti-rabbit IgG-horseradish peroxidase, HRP), 1:5000; (Abcam, Cambridge, UK) at 4 °C for 2h. The relative immunoreactive bands were captured with a chemiluminescent imaging system (Sagecreation, Beijing, China), visualized by using ECL-enhanced chemilluminescence (Boster Biotechnology Co., Ltd. Wuhan, China), and digitalized by the image system of Quantity One 4.31 (Bio-Rad, Hercules, CA, USA).

2.8. Statistical analysis

All experiments data were presented as means \pm standard error (SEM). Statistical analysis was conducted using GraphPad Prism 5 (Graph-Pad software Inc., San Diego, CA, USA). Statistical significance was considered at $P < 0.05$, the result of acquisition task in MWM was evaluated by two-way ANOVA with repeated measures, and other data were analyzed by one-way ANOVA and Student-Newman-Keuls post-hoc tests.

3. Results

3.1 DA-JC4 improved STZ-induced learning and memory impairment

In the Y-maze test, there was an overall difference between groups (one-way ANOVA, $F=3.931$, $P < 0.05$). As shown in Fig.1A, rats that received STZ alone (0.4708 ± 0.04312) exhibited significant impairments in spontaneous alternation compared with the control rats (0.6811 ± 0.03200) ($P < 0.05$). Moreover, treatment with DA-JC4 (0.6283 ± 0.06798) partially reversed the impairment induced by STZ treatment ($P < 0.05$) as shown in Newman-Keuls post-hoc tests, $n=8-10$ per group (Fig.1A).

In the Morris water maze test, the escape latency was gradually shortened during the training period. However, there is virtually no difference until on the fourth and fifth day (Figure 1.B). Based on the two-way ANOVA analysis, the effect of DA-JC4 treatment on escape latency was significant ($F=15.20$, $P<0.0001$ for groups). There was no interaction between groups and time ($F=0.5319$, $P=0.8918$). The escape latency time in the STZ group (48.61 ± 2.056) was longer than in the control rats (36.67 ± 1.548) and the major difference began to emerge on day 2. ($F=6.459$, $P<0.01$). An outstanding difference was found between the control group (22.81 ± 1.498 , 20.02 ± 1.387) and the STZ group on day 4 ($F=9.438$, $P<0.001$, 22.81 ± 1.498 vs 39.65 ± 2.793) and on day 5 ($F=3.618$, $P<0.05$, 20.02 ± 1.387 vs 26.43 ± 2.086). There was

a marked decrease in the escape latency in the STZ group and STZ + DA-JC4 group (30.65 ± 3.372 , 21.91 ± 1.243). Figure 1.C shows the exploration time of the probe trial that rats spent in the target quadrant. A one-way ANOVA analysis showed an overall difference between groups ($F=7.818$, $P<0.001$). Contrary to the escape latency, rats in the STZ group (0.2800 ± 0.01630) resulted in a decrease in spatial memory compared to the control rats (0.4217 ± 0.01696) ($P<0.001$). Rats that received both STZ and DA-JC4 (0.3920 ± 0.02713) performed better than STZ saline treated rats ($P<0.01$). In addition, there was no significant difference in swim speed among all four groups ($F=1.151$, $P=0.3422$).

3.2 DA-JC4 reduced the levels of phosphorylated tau protein

Overall differences between groups in the cortex and hippocampus were found via one-way ANOVA analysis ($F=8.497$, $p<0.01$; $F=10.38$, $P<0.001$). Fig 2. I, II show that levels of p-tau in the STZ group (0.2774 ± 0.02585 , 0.2794 ± 0.02579) were higher compared to the control group (0.1610 ± 0.01080 , 0.167 ± 0.01308) ($P<0.01$, $P<0.01$). However, the STZ + DA-JC4 group (0.1941 ± 0.01825 , 0.1902 ± 0.008825) attenuated the levels of phospho-tau compared with the control group ($P<0.01$, $P<0.01$).

3.3 DA-JC4 normalized the STZ -induced increase of the chronic inflammation response

As to the activation of astrocytes, significant differences between groups in the cortex and hippocampus were found ($F=30.83$, $P<0.0001$; $F=9.867$, $P<0.01$). There was an increase of GFAP-positive astrocytes in the STZ group (8.340 ± 0.4890 , 15.32 ± 2.348) than the control group (2.808 ± 0.3406 , 6.761 ± 0.2818) ($P<0.001$; $P<0.01$), and injection with DA-JC4 alone (2.881 ± 0.4943 , 7.174 ± 0.7706) has no effect. Rats that received both STZ and DA-JC4 (5.010 ± 0.5244 , 8.820 ± 0.7110) had reduced levels of activated astrocytes. See Fig. 3.

The immunoreactivity of IBA1-positive microglia was observed in the cortex and hippocampus as shown in Fig. 4. I and . There was no difference between control group (0.8879 ± 0.04772 , 1.244 ± 0.1184) and DA-JC4 (0.9417 ± 0.04789 , 1.217 ± 0.1018) group. The activation of microglia in the STZ group (4.895 ± 0.2610 , 4.103 ± 0.4160) was significantly higher than in the control group ($P<0.001$, $P<0.001$), and DA-JC4 administration (1.839 ± 0.2617 , 2.586 ± 0.2038) decreased the number of IBA1-positive microglia compared to the STZ only group ($P<0.001$, $P<0.01$). See Fig. 4.

3.4 DA-JC4 reversed the STZ-induced increase in apoptotic signalling

The results show that the BAX/Bcl-2 ratio did differ significantly among the four different treatment groups in the rat cortex and hippocampus (one-way ANOVA $F=26.78$, $P<0.0001$; $F=120.1$, $P<0.0001$). Compared with the controls (0.9915 ± 0.01694 , 0.8145 ± 0.03596), STZ administration (2.387 ± 0.2384 , 2.276 ± 0.1191) markedly increased this ratio ($P<0.001$, $P<0.001$). In comparison, the increases in BAX/Bcl-2 ratios induced by STZ were reversed when treated with DA-JC4 (1.192 ± 0.06932 , 0.8966 ± 0.02022). Also, injection with DA-JC4 alone (1.034 ± 0.1056 , 0.7970 ± 0.03804) did not change the level of apoptotic signaling in the brain. See Fig. 5

3.5 DA-JC4 promoted re-sensitization of insulin signaling in the brain

Total protein of Akt and IRS1 did not differ significantly in any of the groups. A one-way ANOVA analysis showed differences between groups ($p<0.001$). The levels of phosphorylated Akt and IRS1 was changed by the treatment. As seen in Fig. 6, the western blot assay revealed that the levels of phospho-IRS1^{Ser1101} in the STZ treated group (0.04414 ± 0.001197 , 0.03244 ± 0.001044) were significantly increased in both the cortex and the hippocampus, compared with the controls (0.03712 ± 0.001724 , 0.02815 ± 0.0008825) ($P<0.01$, $P<0.05$). However, rats that had received both DA-JC4 and STZ (0.04034 ± 0.0006476 , 0.02869 ± 0.0009052) exhibited lower levels of phospho-IRS1^{Ser1101} in the brain than the STZ only rats ($P<0.05$, $P<0.05$).

In contrast with the change of phospho-IRS1^{Ser1101}, STZ treatment markedly suppressed the expression of phospho-Akt^{Ser473} (0.1872 ± 0.02649 , 0.1540 ± 0.02694) in both the cortex and the hippocampus as compared to the control group (0.3356 ± 0.02061 , 0.2929 ± 0.01382) ($P<0.01$, $P<0.01$). However, DA-JC4 (0.2940 ± 0.005428 , 0.2114 ± 0.01248) moderately attenuated the STZ-induced inactivation of phospho-Akt^{Ser473} ($P<0.05$, $P<0.05$).

4. Discussion

In the present study, we investigated the effect of the novel GLP-1/GIP dual receptor agonist DA-JC4 for the first time in the rat model of STZ- induced cognitive impairment. ICV. administration of STZ at low dosage (3 mg/kg body weight) induces learning and memory deficits [21, 38-40, 43]. Consistent with previous findings, we found that learning and memory impairment were induced by the STZ treatment. We also found that DA-JC4 effectively attenuated the spatial working memory deficit induced by STZ. Consistent with previous findings, the phosphorylation of tau protein was enhanced by icv. administration of STZ [42].

DA-JC4 treatment prevented or reversed the phosphorylation of tau. The chronic inflammation response in the brain is a critical element of AD progression [47, 48]. Chronic inflammation generates free radicals and increases the release of pro-inflammatory cytokines that are detrimental to neurons [49, 50]. Consistent with previous findings, the chronic inflammation response was clearly triggered by the treatment with STZ [41]. DA-JC4 effectively reduced the chronic inflammation response induced by STZ, and the ratio of BAX/Bcl-2 was much improved, demonstrating that cellular apoptotic signaling had been reduced by the novel drug. In addition, insulin signaling as shown in IRS-1 phosphorylation levels and Akt phosphorylation furthermore was re-sensitized by DA-JC4. Levels of pIRS-1 were reduced and of pAkt were enhanced to normalise second messenger cell signalling. The novel dual agonist showed good effects in improving these key biomarkers of neuropathological processes.

Insulin de-sensitization has been observed in the brains of people with AD [51-53] and we previously demonstrated that the GLP-1 analogue liraglutide was able to reverse this [54]. Importantly, brain insulin de-sensitization correlates well with cognitive decline [15, 55] and with tau protein phosphorylation levels [56], and the application of insulin via nasal spray showed good improvements in AD patients [57, 58].

The chronic inflammation response in the brain plays a key role in the progression of neurodegenerative disorders [59], and a reduction of this process most likely also plays a part in the overall improvement of the pathophysiology. GLP-1 has anti-inflammatory properties [60], and we have shown previously that single GLP-1 or GIP receptor agonists can reduce the chronic inflammation response in the brain [16, 17, 26]. Novel dual GLP-1/GIP receptor agonists have the advantage of activating two signaling pathways, and have been shown to be superior to single GLP-1 receptor agonists [31, 61]. We previously reported that another dual agonists named DA-JC1 showed good neuroprotective effects in the MPTP mouse model of Parkinson's disease [33, 62]. That dual agonist was not as effective as a single GLP-1 receptor agonist [63] and had to be injected at a higher dose (50nmol/kg ip). In this study, we used the much lower dose of DA-JC4 (10 nmol/kg ip) for the evaluation of the neuroprotective effects of DA-JC4 in the Alzheimer Rat Model induced by STZ, which compares well to the effective dose of liraglutide of 25nmol/kg ip. [26]. This first study shows promise that such dual agonists may be superior to single incretin analogues, but further dose response tests need to be conducted in order to find the most potent drug dose for this dual agonist. This is of importance as liraglutide has shown first neuroprotective effects in a pilot clinical study of AD patients. Brain activity and energy turnover was assessed by ¹⁸F-DG-PET brain imaging, and after 6 months of treatment, brain scans revealed that there was significant deterioration in cortical activity in the placebo group. In contrast, the liraglutide treated group showed no deterioration at all [29]. A larger phase II trial is

currently ongoing (clinical trial ID NCT01843075). Furthermore, another GLP-1 receptor agonist, exendin-4, has shown good protective effects in a pilot study in patients with Parkinson's disease [64]. It is therefore of interest to develop more efficient drugs that can activate not just GLP-1 but also the sister incretin signaling pathway GIP, which has shown neuroprotective effects on its own in preclinical studies of AD (Duffy and Holscher, 2013; Faivre and Holscher, 2013) and PD [65, 66]. Novel dual agonists therefore may be superior in the clinic for treating chronic neurodegenerative disorders such as AD or PD.

5. Acknowledgements

This study was supported by the Shanxi Science and Technology Department (2010081062) and a grant to C.H. under the '100 foreign talents' of the Shanxi province government. C.H. is a named inventor on a patent application by Lancaster University, covering the use of novel dual GLP-1/GIP receptor agonists as treatments for neurodegenerative disorders. The other authors do not declare a conflict of interest.

References

- [1] Mu Y, Gage FH. Adult hippocampal neurogenesis and its role in Alzheimer's disease. *Molecular neurodegeneration*. 2011;6:85.
- [2] Tanzi RE, Bertram L. Twenty years of the Alzheimer's disease amyloid hypothesis: a genetic perspective. *Cell*. 2005;120:545-55.
- [3] Ono M, Watanabe H, Kitada A, Matsumura K, Ihara M, Saji H. Highly Selective Tau-SPECT Imaging Probes for Detection of Neurofibrillary Tangles in Alzheimer's Disease. *Scientific reports*. 2016;6:34197.
- [4] Roberson ED, Mucke L. 100 years and counting: prospects for defeating Alzheimer's disease. *Science*. 2006;314:781-4.
- [5] Blennow K, de Leon MJ, Zetterberg H. Alzheimer's disease. *Lancet*. 2006;368:387-403.
- [6] Atri A. Effective pharmacological management of Alzheimer's disease. *The American journal of managed care*. 2011;17 Suppl 13:S346-55.
- [7] Godyn J, Jonczyk J, Panek D, Malawska B. Therapeutic strategies for Alzheimer's disease in clinical trials. *Pharmacological reports : PR*. 2016;68:127-38.
- [8] Haan MN. Therapy Insight: type 2 diabetes mellitus and the risk of late-onset Alzheimer's disease. *Nat Clin Pract Neurol*. 2006;2:159-66.
- [9] Kopf D, Frolich L. Risk of incident Alzheimer's disease in diabetic patients: a systematic review of prospective trials. *Journal of Alzheimer's disease : JAD*. 2009;16:677-85.
- [10] Sims-Robinson C, Kim B, Rosko A, Feldman EL. How does diabetes accelerate Alzheimer disease pathology? *Nature reviews Neurology*. 2010;6:551-9.
- [11] Awad N, Gagnon M, Messier C. The relationship between impaired glucose tolerance, type 2 diabetes, and cognitive function. *Journal of clinical and experimental neuropsychology*. 2004;26:1044-80.
- [12] Ohara T, Doi Y, Ninomiya T, Hirakawa Y, Hata J, Iwaki T, et al. Glucose tolerance status and risk of dementia in the community: The Hisayama Study. *Neurology*. 2011;77:1126-34.
- [13] Li L, Holscher C. Common pathological processes in Alzheimer disease and type 2 diabetes: a review. *Brain Res Rev*. 2007;56:384-402.
- [14] van der Heide LP, Ramakers GM, Smidt MP. Insulin signaling in the central nervous system: learning to survive. *Progress in*

- neurobiology. 2006;79:205-21.
- [15] Freiherr J, Hallschmid M, Frey WH, 2nd, Brunner YF, Chapman CD, Holscher C, et al. Intranasal insulin as a treatment for Alzheimer's disease: a review of basic research and clinical evidence. *CNS drugs*. 2013;27:505-14.
- [16] Duffy AM, Holscher C. The incretin analogue D-Ala2GIP reduces plaque load, astrogliosis and oxidative stress in an APP/PS1 mouse model of Alzheimer's disease. *Neuroscience*. 2013;228:294-300.
- [17] Faivre E, Holscher C. D-Ala2GIP facilitated synaptic plasticity and reduces plaque load in aged wild type mice and in an Alzheimer's disease mouse model. *Journal of Alzheimer's disease : JAD*. 2013;35:267-83.
- [18] McClean PL, Holscher C. Liraglutide can reverse memory impairment, synaptic loss and reduce plaque load in aged APP/PS1 mice, a model of Alzheimer's disease. *Neuropharmacology*. 2014;76:57-67.
- [19] Li Y, Duffy K, Ottinger M, Ray B, Bailey J, Holloway H, et al. GLP-1 Receptor Stimulation Reduces Amyloid-beta Peptide Accumulation and Cytotoxicity in Cellular and Animal Models of Alzheimer's Disease. *Journal of Alzheimer's disease : JAD*. 2010;19:1205-19.
- [20] Xiong H, Zheng C, Wang J, Song J, Zhao G, Shen H, et al. The neuroprotection of liraglutide on Alzheimer-like learning and memory impairment by modulating the hyperphosphorylation of tau and neurofilament proteins and insulin signaling pathways in mice. *Journal of Alzheimer's disease : JAD*. 2013;37:623-35.
- [21] Chen Y, Tian Z, Liang Z, Sun S, Dai CL, Lee MH, et al. Brain gene expression of a sporadic (icv-STZ Mouse) and a familial mouse model (3xTg-AD mouse) of Alzheimer's disease. *PloS one*. 2012;7:e51432.
- [22] Gao C, Liu Y, Jiang Y, Ding J, Li L. Geniposide ameliorates learning memory deficits, reduces tau phosphorylation and decreases apoptosis via GSK3beta pathway in streptozotocin-induced alzheimer rat model. *Brain pathology*. 2014;24:261-9.
- [23] Hansen HH, Barkholt P, Fabricius K, Jelsing J, Terwel D, Pyke C, et al. The GLP-1 receptor agonist liraglutide reduces pathology-specific tau phosphorylation and improves motor function in a transgenic hTauP301L mouse model of tauopathy. *Brain research*. 2015;1634:158-70.
- [24] Solmaz V, Cinar BP, Yigitturk G, Cavusoglu T, Taskiran D, Erbas O. Exenatide reduces TNF-alpha expression and improves hippocampal neuron numbers and memory in streptozotocin treated rats. *European journal of pharmacology*. 2015;765:482-7.
- [25] McClean PL, Parthasarathy V, Faivre E, Holscher C. The diabetes drug liraglutide prevents degenerative processes in a mouse model of Alzheimer's disease. *The Journal of neuroscience*. 2011;31:6587-94.
- [26] Parthasarathy V, Holscher C. The type 2 diabetes drug liraglutide reduces chronic inflammation induced by irradiation in the mouse brain. *European journal of pharmacology*. 2013;700:42-50.
- [27] Holscher C. The role of GLP-1 in neuronal activity and neurodegeneration. *Vitamins and hormones*. 2010;84:331-54.
- [28] Yoshino Y, Ishisaka M, Tsujii S, Shimazawa M, Hara H. Glucagon-like peptide-1 protects the murine hippocampus against stressors via Akt and ERK1/2 signaling. *Biochemical and biophysical research communications*. 2015;458:274-9.
- [29] Gejl M, Gjedde A, Egefjord L, Møller A, Hansen SB, Vang K, et al. In Alzheimer's Disease, Six-Month Treatment with GLP-1 Analogue Prevents Decline of Brain Glucose Metabolism: Randomized, Placebo-Controlled, Double-Blind Clinical Trial. *Frontiers in aging neuroscience*. 2016;8:1-10.
- [30] Holscher C. GLP-1 and GIP analogues as novel treatments for Alzheimer's and Parkinson's disease. *Cardiovasc Endocrinol*. 2016;5:93-8.
- [31] Finan B, Ma T, Ottaway N, Muller TD, Habegger KM, Heppner KM, et al. Unimolecular dual incretins maximize metabolic benefits in rodents, monkeys, and humans. *Science translational medicine*. 2013;5:209ra151.
- [32] Cao L, Li D, Feng P, Li L, Xue G, Li G, et al. A novel dual GLP-1 and GIP incretin receptor agonist is neuroprotective in a mouse model of Parkinson's disease by reducing chronic inflammation in the brain. *Neuroreport*. 2016;37:384-91.
- [33] Ji C, Xue GF, Lijun C, Feng P, Li D, Li L, et al. A novel dual GLP-1 and GIP receptor agonist is neuroprotective in the MPTP mouse model of Parkinson's disease by increasing expression of BDNF. *Brain research*. 2016;1634:1-11.
- [34] Plaschke K, Kopitz J, Siegelin M, Schliebs R, Salkovic-Petrisic M, Riederer P, et al. Insulin-resistant brain state after intracerebroventricular streptozotocin injection exacerbates Alzheimer-like changes in Tg2576 AbetaPP-overexpressing mice. *Journal of Alzheimer's disease : JAD*. 2010;19:691-704.

Dual agonist effects in the STZ rat model

- [35] Plaschke K, Muller D, Hoyer S. Insulin-resistant brain state (IRBS) changes membrane composition of fatty acids in temporal and entorhinal brain cortices of rats: relevance to sporadic Alzheimer's disease? *Journal of neural transmission*. 2010;117:1419-22.
- [36] Dhull DK, Jindal A, Dhull RK, Aggarwal S, Bhateja D, Padi SS. Neuroprotective effect of cyclooxygenase inhibitors in ICV-STZ induced sporadic Alzheimer's disease in rats. *Journal of molecular neuroscience : MN*. 2012;46:223-35.
- [37] Ponce-Lopez T, Liy-Salmeron G, Hong E, Meneses A. Lithium, phenserine, memantine and pioglitazone reverse memory deficit and restore phospho-GSK3beta decreased in hippocampus in intracerebroventricular streptozotocin induced memory deficit model. *Brain research*. 2011;1426:73-85.
- [38] Lin F, Jia J, Qin W. Enhancement of beta-amyloid oligomer accumulation after intracerebroventricular injection of streptozotocin, which involves central insulin signaling in a transgenic mouse model. *Neuroreport*. 2014;25:1289-95.
- [39] Singh JC, Kakalij RM, Kshirsagar RP, Kumar BH, Komakula SS, Diwan PV. Cognitive effects of vanillic acid against streptozotocin-induced neurodegeneration in mice. *Pharmaceutical biology*. 2015;53:630-6.
- [40] Salkovic-Petrisic M, Osmanovic-Barilar J, Knezovic A, Hoyer S, Mosetter K, Reutter W. Long-term oral galactose treatment prevents cognitive deficits in male Wistar rats treated intracerebroventricularly with streptozotocin. *Neuropharmacology*. 2014;77:68-80.
- [41] Rai S, Kamat PK, Nath C, Shukla R. A study on neuroinflammation and NMDA receptor function in STZ (ICV) induced memory impaired rats. *Journal of neuroimmunology*. 2013;254:1-9.
- [42] Grunblatt E, Salkovic-Petrisic M, Osmanovic J, Riederer P, Hoyer S. Brain insulin system dysfunction in streptozotocin intracerebroventricularly treated rats generates hyperphosphorylated tau protein. *Journal of neurochemistry*. 2007;101:757-70.
- [43] Agrawal R, Tyagi E, Shukla R, Nath C. Insulin receptor signaling in rat hippocampus: a study in STZ (ICV) induced memory deficit model. *European neuropsychopharmacology : the journal of the European College of Neuropsychopharmacology*. 2011;21:261-73.
- [44] Baluchnejadmojarad T, Roghani M. Chronic epigallocatechin-3-gallate ameliorates learning and memory deficits in diabetic rats via modulation of nitric oxide and oxidative stress. *Behavioural brain research*. 2011;224:305-10.
- [45] Morris R. Developments of a water-maze procedure for studying spatial learning in the rat. *Journal of neuroscience methods*. 1984;11:47-60.
- [46] Gengler S, McClean P, McCurtin R, Gault V, Holscher C. Val(8)GLP-1 rescues synaptic plasticity and reduces dense core plaques in APP/PS1 mice. *Neurobiology of aging*. 2012;33:265-76.
- [47] Qin L, Chong T, Rodriguez R, Pugazhenth S. Glucagon-Like Peptide-1-Mediated Modulation of Inflammatory Pathways in the Diabetic Brain: Relevance to Alzheimer's Disease. *Current Alzheimer research*. 2016;13:1346-55.
- [48] Ji C, Xue GF, Li G, Li D, Holscher C. Neuroprotective effects of glucose-dependent insulinotropic polypeptide in Alzheimer's disease. *Reviews in the neurosciences*. 2016;27:61-70.
- [49] Tansey MG, Goldberg MS. Neuroinflammation in Parkinson's disease: its role in neuronal death and implications for therapeutic intervention. *Neurobiology of disease*. 2010;37:510-8.
- [50] Martino Adami PV, Galeano P, Wallinger ML, Quijano C, Rabossi A, Pagano ES, et al. Worsening of memory deficit induced by energy-dense diet in a rat model of early-Alzheimer's disease is associated to neurotoxic Abeta species and independent of neuroinflammation. *Biochimica et biophysica acta*. 2016;1863:731-43.
- [51] Moloney AM, Griffin RJ, Timmons S, O'Connor R, Ravid R, O'Neill C. Defects in IGF-1 receptor, insulin receptor and IRS-1/2 in Alzheimer's disease indicate possible resistance to IGF-1 and insulin signalling. *Neurobiology of aging*. 2010;31:224-43.
- [52] Talbot K, Wang HY, Kazi H, Han LY, Bakshi KP, Stucky A, et al. Demonstrated brain insulin resistance in Alzheimer's disease patients is associated with IGF-1 resistance, IRS-1 dysregulation, and cognitive decline. *The Journal of clinical investigation*. 2012;122:1316-38.
- [53] Holscher C. Diabetes as a risk factor for Alzheimer's disease: insulin signalling impairment in the brain as an alternative model of Alzheimer's disease. *Biochemical Society transactions*. 2011;39:891-7.
- [54] Long-Smith CM, Manning S, McClean PL, Coakley MF, O'Halloran DJ, Holscher C, et al. The diabetes drug liraglutide

Dual agonist effects in the STZ rat model

- ameliorates aberrant insulin receptor localisation and signalling in parallel with decreasing both amyloid-beta plaque and glial pathology in a mouse model of Alzheimer's disease. *Neuromolecular medicine*. 2013;15:102-14.
- [55] Townsend M, Mehta T, Selkoe DJ. Soluble Abeta inhibits specific signal transduction cascades common to the insulin receptor pathway. *The Journal of biological chemistry*. 2007;282:33305-12.
- [56] Escribano L, Simon AM, Gimeno E, Cuadrado-Tejedor M, Lopez de Maturana R, Garcia-Osta A, et al. Rosiglitazone rescues memory impairment in Alzheimer's transgenic mice: mechanisms involving a reduced amyloid and tau pathology. *Neuropsychopharmacology : official publication of the American College of Neuropsychopharmacology*. 2010;35:1593-604.
- [57] Novak V, Milberg W, Hao Y, Munshi M, Novak P, Galica A, et al. Enhancement of vasoreactivity and cognition by intranasal insulin in type 2 diabetes. *Diabetes care*. 2014;37:751-9.
- [58] Holscher C. First clinical data of the neuroprotective effects of nasal insulin application in patients with Alzheimer's disease. *Alzheimer's & dementia : the journal of the Alzheimer's Association*. 2014;10:S33-S7.
- [59] Holmes C, Cunningham C, Zotova E, Woolford J, Dean C, Kerr S, et al. Systemic inflammation and disease progression in Alzheimer disease. *Neurology*. 2009;73:768-74.
- [60] Dozier KC, Cureton EL, Kwan RO, Curran B, Sadjadi J, Victorino GP. Glucagon-like peptide-1 protects mesenteric endothelium from injury during inflammation. *Peptides*. 2009;30:1735-41.
- [61] Jalewa J, Sharma MK, Holscher C. Novel incretin analogues improve autophagy and protect from mitochondrial stress induced by rotenone in SH-SY5Y cells. *Journal of neurochemistry*. 2016;139:55-67.
- [62] Cao L, Li D, Feng P, Li L, Xue GF, Li G, et al. A novel dual GLP-1 and GIP incretin receptor agonist is neuroprotective in a mouse model of Parkinson's disease by reducing chronic inflammation in the brain. *Neuroreport*. 2016;27:384-91.
- [63] Liu W, Jalewa J, Sharma M, Li G, Li L, Holscher C. Neuroprotective effects of lixisenatide and liraglutide in the MPTP mouse model of Parkinson's disease. *Neuroscience*. 2015;303:42-50.
- [64] Aviles-Olmos I, Dickson J, Kefalopoulou Z, Djamshidian A, Kahan J, Fmedsci PE, et al. Motor and Cognitive Advantages Persist 12 Months After Exenatide Exposure in Parkinson's Disease. *Journal of Parkinson's disease*. 2014;4:337-44.
- [65] Li Y, Liu W, Li L, Holscher C. Neuroprotective effects of a GIP analogue in the MPTP Parkinson's disease mouse model. *Neuropharmacology*. 2016;101:255-63.
- [66] Li Y, Liu W, Li L, Holscher C. D-Ala2-GIP-glu-PAL is neuroprotective in a chronic Parkinson's disease mouse model and increases BDNF expression while reducing neuroinflammation and lipid peroxidation. *European journal of pharmacology*. 2016;DOI:10.1016/j.ejphar.2016.11.050.

Figure captions

Figure 1. DA-JC4 improved STZ-induced learning and memory impairments. **A.** Spontaneous alternation behavior was significantly decreased in the STZ group compared with the control group (compared with controls, $*=P<0.05$), and DA-JC4 treatment prevented the STZ-induced learning deficit (compared with STZ group, $\#=P<0.05$). **B.** Escape latency was significantly increased in the STZ group compared to the control group on day 2, 4, 5 (compared with control, $*=P<0.05$; $**=P<0.01$; $***=P<0.001$), and DA-JC4 treatment partially attenuated the impairment induced by STZ. **C.** STZ treatment significantly decreased the percent of time each rat spent in the target quadrant ($P<0.001$), and DA-JC4 treatment available prevented the STZ-induced memory deficit ($P<0.01$). **D.** Swim speed did not differ between groups. **E.** Sample tracks in the probe test.

Figure 2. DA-JC4 reduced the levels of phospho-tau protein in the rat cerebral cortex () and hippocampus () (**A–D**, scale bar = 100 μm ; **E–H**, scale bar = 50 μm). The expression of phospho-tau was increased following treatment with STZ ($**=P<0.01$; compared with control), and reduced by DA-JC4 ($\#=P<0.01$, compared with STZ group).

Figure 3. DA-JC4 attenuated the STZ - induced the activation of astrocytes in the rat cerebral cortex () and hippocampus () (**A–D**, scale bar = 100 μm). Quantification of the area of GFAP positive stain in the rat cerebral cortex () and hippocampus () demonstrated that STZ injection induced astrogliosis ($***=P<0.001$; $**=P<0.01$ compared to controls). DA-JC4 partially reversed this ($####=P<0.001$; $###=P<0.01$ compared with STZ group).

Figure 4. DA-JC4 attenuated the STZ- induced activation of microglia in the rat cortex () and hippocampus () (**A–D**, scale bar = 100 μm). Quantification of the area of IBA-1 positive stain in the cerebral cortex () and hippocampus () revealed that STZ injection induced microgliosis ($***=P<0.001$ compared to controls). DA-JC4 partially reversed this ($####=P<0.001$; $##=P<0.01$ compared with the STZ group).

Figure 5. DA-JC4 reduced the STZ-induced apoptotic signaling in the cortex and hippocampus. **A.** Western blot assay of BAX and Bcl-2 in response to STZ and DA-JC4. **B.** Western blot quantification of protein levels of BAX and Bcl-2. A one-way ANOVA found overall differences between groups. Post-hoc analyses showed differences compared to controls ($***=P<0.001$ compared to controls; $###=P<0.001$ compared to STZ group). Data are represented as mean \pm SEM, and show data of 4 blotting repetitions.

Figure 6. DA-JC4 re-sensitized insulin signaling in the brain. The effects of DA-JC4 on protein expression levels of p-Akt and p-IRS-1 in the cortex and the hippocampus as measured by western blot assay (**A**, **C**). Quantification of p-Akt and p-IRS-1 in the cortex and the hippocampus (**B**, **D**). A one-way ANOVA found overall differences for B ($F=8.020$, $P<0.01$; $F=13.69$, $P<0.01$) and D ($F=7.724$, $P<0.01$; $F=6.841$, $P<0.01$). Post-hoc analyses showed differences compared to controls ($*=P<0.05$; $**=P<0.01$). Data are represented as mean \pm SEM, and show data of 4 blotting repetitions.




Fig 1




Fig 2




Fig 3




Fig 4




Fig 5

Fig 6