

**Heritability of hippocampal subfield volumes using a twin
and non-twin siblings design**

Journal:	<i>Human Brain Mapping</i>
Manuscript ID	HBM-16-0938.R2
Wiley - Manuscript type:	Research Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Patel, Sejal; Centre for Addiction And Mental Health, ; University of Toronto, Institute of Medical Science Park, Min Tae; Douglas Mental Health Research Institute, Cerebral Imaging Centre Devenyi, Gabriel; Douglas Mental Health University Institute, Cerebral Imaging Centre Patel, Raihaan; Douglas Mental Health University Institute , Cerebral Imaging Centre Maselli, Mario; Sunnybrook Health Sciences Centre, Neurology Knight, Jo; Lancaster University Faculty of Health and Medicine; Centre for Addiction And Mental Health; University of Toronto, Institute of Medical Science Chakravarty, Mallar; Douglas Mental Health University Institute, Cerebral Imaging Centre; McGill University, Psychiatry
Keywords:	Extended twin design, Hippocampal subfield segmentation, Univariate and Bivariate model, Genetic correlation, Genetic variation, Quantitative phenotypes

SCHOLARONE™
Manuscripts

Heritability of hippocampal subfield volumes using a twin and non-twin siblings design

Sejal Patel^{1,2*}, Min Tae M. Park³, Gabriel A. Devenyi³, Raihaan Patel³, Mario Masellis⁴, Jo Knight^{1,2,5,a}, M. Mallar Chakravarty^{3,6,a*}

¹ Campbell Family Mental Health Research Institute, Centre for Addiction and Mental Health, Toronto, ON, Canada

² Institute of Medical Science, University of Toronto, Toronto, ON, Canada

³ Cerebral Imaging Centre, Douglas Mental Health University Institute, McGill University, Verdun, QC, Canada

⁴ Department of Neurology, Sunnybrook Health Sciences Centre, Toronto, ON, Canada

⁵ Lancaster Medical School, Faculty of Health and Medicine, Lancaster University, Lancaster, UK

⁶ Department of Psychiatry, McGill University, Montreal, QC, Canada

* Corresponding authors at:

Sejal Patel, Campbell Family Mental Health Research Institute, Centre for Addiction and Mental Health, 250 College Street, Toronto, ON, M5T 1R8, Canada

Email address: Sejal.Patel@camh.ca (S. Patel)

M. Mallar Chakravarty, Cerebral Imaging Centre, Douglas Mental Health University Institute, McGill University, 6875 Boulevard LaSalle, Verdun, QC, H4H 1R3 Verdun, Canada

Email address: mallar@cobralab.ca (M.M. Chakravarty).

^a M. Mallar Chakravarty and Jo Knight contributed equally as senior authors to this manuscript.

Short Title: Heritability of hippocampal subfield

Keywords: Genetic variation, Genetic correlation, Quantitative phenotypes, Extended twin design, Hippocampal subfield segmentation, Univariate and bivariate model

ABSTRACT

The hippocampus is composed of distinct subfields linked to diverse functions and disorders.

The subfields can be mapped using high-resolution magnetic resonance images, and their volumes can potentially be used as quantitative phenotypes for genetic investigation of

hippocampal function. We estimated the heritability of hippocampus subfield volumes of 465 subjects from the Human Connectome Project (twins and non-twin siblings) using two methods.

The first used a univariate model to estimate heritability with and without adjustment for total brain volume (TBV) and ipsilateral hippocampal volume to determine if heritability was uniquely attributable to subfield volume rather than confounds that attributed to global volumes.

We observed the right: subiculum, cornu ammonis 2/3, and cornu ammonis 4/dentate gyrus subfields had the highest significant heritability estimates after adjusting for ipsilateral hippocampal volume. In the second analysis, we used a bivariate model to investigate the shared heritability and genetic correlation of the subfield volumes with TBV and ipsilateral hippocampal volume. Genetic correlation demonstrates shared genetic architecture between phenotypes and shared heritability is what proportion of the genetic architecture of one trait is shared by the other. Highest genetic correlations were between subfield volumes and ipsilateral hippocampal volume than with TBV. The pattern was opposite for shared heritability suggesting that subfields share greater proportion of the genetic architecture with TBV than with ipsilateral hippocampal volume. The relationship between the genetic architecture of TBV, hippocampal volume, and of individual subfields should be accounted for when using hippocampal subfield volumes as quantitative phenotypes for imaging genetics studies.

Box 1: Key Heritability Concepts

- Heritability estimates the amount of genetic variation that is seen in a phenotypic trait within a population. Total variation includes genetics [A], common environment [C] and unique environment [E]. Heritability is the proportion of the variation that can be attributed to genetics.
- Unique heritability accounts for the genetic variation out of total variation (genetic and common and unique environment) of one trait only. This can be calculated in a univariate model where only one phenotype is being analyzed. In a univariate model, factors that may confound heritability estimates such as total brain volume can be controlled for within the model.
- In a bivariate model, the genetic variation is accounted for between two phenotypes instead of focusing on only one trait. In these models we can calculate the genetic correlation between the two phenotypes and the shared heritability. Genetic correlation is the genetic relationship between the two traits. It indicates how similar the genetic variation (genetic architecture) is between both traits. Shared heritability measures the amount of variation from the genetic correlation that is common from trait one and present in trait two. Both genetic correlation and shared heritability estimates should be considered when examining heritability estimates in bivariate model.

INTRODUCTION

The hippocampus plays a key role in cognitive functioning, influencing processes such as learning, episodic and working memory (Braskie et al., 2013). Structural variation within the hippocampal structure and function has been implicated in neurodegenerative and neuropsychiatric disorders such as Alzheimer's disease (Braak and Braak, 1991; Mouiha and Duchesne, 2011), depression (Bremner et al., 2000; Campbell et al., 2004; Treadway et al., 2015) and schizophrenia (Altshuler et al., 1998; Bogerts et al., 1993; Haukvik et al., 2015; Tamminga et al., 2010). Although the hippocampus is often referred to in neuroimaging studies as a single unitary structure, it is composed of several different subfields. While there are varying definitions for the delineation and the nomenclature of these subfields, there is a general consensus in the neuroimaging research discipline that subfields can be resolved using magnetic resonance imaging (MRI) techniques to include some combination of cornu ammonis (CA) 1-4, dentate gyrus (DG), subiculum and the molecular layers composed of the stratum: radiatum, lacunosum and moleculare (SRLM) (Duvernoy, 2005). Subfields within the hippocampus differ from each other in terms of connection to other regions of the brain, their cytoarchitectonic structure, and their role in memory formation and cognitive function (Amunts et al., 2005; Duvernoy, 2005; Fatterpekar et al., 2002; La Joie et al., 2010; Mai et al., 2008; Mueller et al., 2007; Mueller and Weiner, 2009; Voineskos et al., 2015; Yang et al., 2013).

As improved techniques for automated mapping of the hippocampal subfields emerge (Iglesias et al., 2015; Pipitone et al., 2014; Van Leemput et al., 2008; Yushkevich et al., 2010) in concert with large-scale consortia dedicated to genome-wide association analysis (Thompson et al., 2014), it is critical to determine whether the individual subfields can be used as quantitative traits in such studies. Prior to using volumetric estimates of the hippocampal subfields as quantitative

1
2
3 phenotypes, we must first investigate whether these volumes are indeed heritable. Heritability
4
5 estimates are defined by the proportion of observed variation in a trait that can be attributed to
6
7 inherited genetic factors (Jacquard, 1983) and can be estimated using twin studies, family studies
8
9 and genome wide association data from large unrelated populations based on three factors:
10
11 genetics (A), common (C) and unique environment (E). Typically, structural equation modeling
12
13 (SEM) is used in twin studies to estimate variation of the phenotype of interest (Neale and
14
15 Cardon, 1992).
16
17
18
19

20 While several other studies on the heritability of brain structures exist (Baare et al., 2001;
21
22 Pennington et al., 2000; Sullivan et al., 2001), there is only one study that examines the
23
24 heritability of the hippocampal subfields (Whelan et al., 2015). Whelan et al., (2015) used
25
26 standard T1-weighted MRI data from the Queen Twins Imaging (QTIM) and also investigated
27
28 the reliability of automatically segmented human hippocampus with a newly developed
29
30 FreeSurfer subfield segmentation tool. Whelan et al., (2015) demonstrated moderate 4T QTIM
31
32 test-retest reliability scores of ICC (intraclass correlation coefficient) in the range of 0.50-0.86,
33
34 where the CA1 region had the highest ICC of 0.86 and hippocampal-amygdaloid transition
35
36 (HATA) area had the lowest ICC of 0.5. In addition Whelan et al., (2015) found high heritability
37
38 estimates ranging from 0.67 (HATA region) to 0.85 (molecular layer of DG) of hippocampal
39
40 subregion volumes. Our study differs from the study of Whelan et al., based on the previously
41
42 validated segmentation technique (Pipitone et al., 2014) and the use of high-resolution and
43
44 contrast of MRI scans, from the Human Connectome Project (HCP) (Van Essen et al., 2013).
45
46 Further, when exploring heritability it is important to consider the relationship of the
47
48 hippocampal subfields, with overall volume of the brain and or the entire hippocampus; both of
49
50 which vary between individuals (Blatter et al., 1995). Recent studies that have examined the
51
52
53
54
55
56
57
58
59
60

S. Patel and M.M. Chakravarty

1
2
3 heritability of neuroanatomical volume measurements have not adjusted for total brain volume
4 (TBV) or volumes of larger structures encompassing the region of interest. Baare et al,
5
6 demonstrated that the whole brain is highly heritable, estimated to be 90% (Baare et al., 2001),
7
8 therefore not accounting for TBV on regions of interest can influence the heritability estimates of
9
10 the target regions resulting in inaccurate heritability scores.
11
12

13
14 In our study, we explored how heritability of individual hippocampal subfields is influenced by
15
16 both TBV and ipsilateral (left or right) hippocampus volume to the subfield being examined
17
18 using two different methods. Our first approach was using a univariate model examining the
19
20 heritability of a single phenotype at a time; in our case we examined the heritability of
21
22 hippocampal subfields. First we adjusted hippocampal subfield volume for TBV, then we
23
24 adjusted for only ipsilateral hippocampal volume and finally the last step was adjusting for both
25
26 TBV and ipsilateral hippocampal volumes. In our second approach we used a bivariate model
27
28 where we examined the heritability of two structures together instead of one structure alone. We
29
30 went on to quantify the relationship of the shared genetic variation between hippocampal
31
32 subfield volumes with TBV and ipsilateral hippocampal volume by examining the genetic
33
34 correlation and shared heritability in a bivariate model. Both models give us heritability
35
36 estimates which can reveal the genetic architecture of a phenotype by itself but also in relation to
37
38 other phenotypes (see Box 1 for definitions of heritability terms). In the univariate model, the
39
40 heritability estimates are based on the genetic variation of one phenotype. Heritability estimates
41
42 in our univariate model are based on genetic variation of subfield volumes alone. We removed
43
44 the genetic influence of TBV and hippocampal volume which provided us with unique
45
46 heritability estimates for subfield volumes. On the other hand, the bivariate model allows us to
47
48 account for the genetic influence of TBV and hippocampal volume by estimating both the
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 genetic correlation and shared heritability between the two phenotypes. Genetic correlation
4
5 measures the genetic relationship in terms of how similar the genetic architecture is between the
6
7 two phenotypes. Shared heritability measures the shared genetic variation between the two
8
9 phenotypes as a proportion of the heritability of one phenotype. Using both the univariate and
10
11 bivariate model to investigate the genetic variation within the hippocampal subfields allows us to
12
13 **identify** potential quantitative phenotypes to be used in imaging genetic studies.
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review

METHODS

Human Connectome Project Dataset

Data from the Human Connectome project (HCP) was used for this study. The aim of the HCP is to investigate properties of human brain connectivity and function. In order to better understand the interaction of brain circuits and human behaviour, structural and functional properties of neuroanatomical structures as well as genetic factors can be studied. The HCP investigators are recruiting 1200 healthy twin and non-twin sibling adults. The HCP consortium aims to have a healthy population that represents the ethnic and racial composition of US and diversity in terms of behavioural, ethnic and socioeconomic status. Sibling relationships were removed if the individuals within the relationship had neuropsychiatric, neurodevelopmental or neurological disorders. Individuals having other illnesses such as diabetes and high blood pressure were excluded. Premature twins (born before 34 week gestation) and non-twins (born before 37 weeks gestation) were excluded. Individuals who smoked or who were overweight were included in the study. Also individuals who have not experienced severe symptoms but have a history of heavy drinking or recreational drug use were included. Reason for including individuals who smoked, who are overweight or use of recreational substances can be used for future psychiatric studies (Van Essen et al., 2013). For more information on the inclusion and exclusion criteria, see supplemental Table S1 of Van Essen DC et al. (2013).

The sample used for this study contained 542 individuals of which imaging data was available on 511 individuals (data release June 2014). Imaging data were collected using a Siemens 3 Tesla (T) Skyra scanner which has been modified with a Siemens SC72 gradient coil to increase the maximum gradient strength from 40 mT/m to 100 mT/m (Van Essen et al., 2013; Van Essen et al., 2012). For our study we used the 3T, high-resolution T1 weighted MRI data (0.7mm

1
2
3 isotropic voxel dimensions). Acquisition parameters are: inversion time = 1000ms, echo time =
4
5 2.14ms, repetition time = 2400ms, acquisition time = 7min 40sec, flip angle = 8 degrees and
6
7
8 field of view = 224mm x 224mm (Van Essen et al., 2012).
9

10 After quality control of the segmented imaging data and removal of families with one individual
11 and no siblings (described below), the final sample consisted of 100 monozygotic (MZ) twins, 94
12
13 dizygotic (DZ) twins and 271 non-twin siblings (277 women and 188 men with age range of 22-
14
15 36 years old and the average age was 29.24 (\pm 3.49 SD) years old). Average handedness for our
16
17 sample was 65.39 (\pm 46.42 SD). The scale for handedness ranges from -100 (left-hand dominant)
18
19 to 100 (right-hand dominant). The measure of fluid intelligence was represented using Raven's
20
21 Progressive Matrices test where the final score represents the number of correct responses out of
22
23 24 questions. Demographic information is summarized in Table I. Handedness and fluid
24
25 intelligence measures are not used in subsequent heritability calculations, but simply to
26
27 demonstrate that MZ, DZ, and non-twin sibling subgroups are well matched to one another and
28
29 represent a healthy population.
30
31
32
33
34
35
36

37 **Image Processing**

38
39 Hippocampal subfield segmentation was estimated using Multiple Automatically Generated
40
41 Templates (MAGeT Brain) (Chakravarty et al., 2013; Pipitone et al., 2014). The segmentation
42
43 procedure consists of three steps. First, five independent high-resolution MRI atlases (0.3mm
44
45 isotropic voxels) of the hippocampus and hippocampal subfields containing detailed
46
47 segmentations of the left and right CA1, CA2/CA3, CA4/DG, subiculum, SRLM are used as
48
49 inputs (Winterburn et al., 2013). Then, a "template library" is generated where a subset of
50
51 individuals from the HCP dataset is first selected (n=21; a number demonstrated as being optimal
52
53 in previous work (Pipitone et al., 2014)). The 21 templates were selected to represent the HCP
54
55
56
57
58
59
60

S. Patel and M.M. Chakravarty

1
2
3 dataset (12 females, 9 males, aged range: 22 to 36 years). Each template undergoes a model-
4 based segmentation procedure with each atlas yielding five candidate labels for each template.
5
6 The next step is similar to a regular multi-atlas segmentation strategy, where each subject is
7
8 nonlinearly matched to template, thereby growing the number of candidate segmentations to 105
9
10 (5 atlases \times 21 templates) for each subject. The 105 labels per subject are then fused using a
11
12 majority vote (Collins and Pruessner, 2010), a method which was previously demonstrated to be
13
14 accurate relative to weighted voting. Pipitone et al (2014) showed that using weighted voting
15
16 methods did not significantly improve MAgE Brain segmentation when compared to using
17
18 majority vote labeling (Bhagwat et al., 2016; Pipitone et al., 2014). Recent work by Bhagwat et
19
20 al (2016) has also demonstrated accuracies to be homologous to Joint Label Fusion (Wang et al.,
21
22 2013) when used within the MAgE Brain framework (Bhagwat et al., 2016). In Winterburn et
23
24 al (2013) protocol for human hippocampal subfields segmentation, intra reliability was measured
25
26 via test-retest and results ranged from 0.64 in the CA2/CA3 region to 0.83 mean Kappa in the
27
28 CA4/DG region. The second lowest test-retest was the SRLM region with mean Kappa of 0.71.
29
30 The subiculum and the CA1 region had a Kappa mean of 0.75 and 0.78 respectively.
31
32
33
34
35
36
37
38
39
40 Quality control (QC) of hippocampal segmentation was completed for 511 subjects with imaging
41
42 data and 502 passed QC. Images were rated on a 3 point scale of 0, 0.5 and 1, where 1 was a pass
43
44 with very negligible errors, 0.5 was satisfactory with few errors but still considered pass and
45
46 images scored 0 is a fail and the subject was removed for the final analysis. Subjects scored as 1
47
48 had less than 5 slices with minor errors whereas subjects that contained minor errors found in 5
49
50 to 10 slices were assigned 0.5. Subjects that contain errors in more than 10 slices unilaterality
51
52 were scored 0. Minor errors included small deviation from the correct segmentation within
53
54 subjects which are scored 1 and 0.5. Major deviation from the correct segmentation where large
55
56
57
58
59
60

1
2
3 portions of the hippocampus are missing or segmented in the wrong location on the slice was
4
5 scored as 0. In the supplementary section Figure S1 A-C show examples of subjects scored 0, 0.5
6
7 and 1.
8
9

10
11 MAGEt Brain was implemented using the publicly available pipeline

12
13 (<https://github.com/CobraLab/MAGEtBrain>) and atlases

14
15 (<https://github.com/CobraLab/atlas/tree/master/hippocampus-subfields>). In addition, TBV was

16
17 extracted using the automated BEaST pipeline (Eskildsen et al., 2012) along with minc-bpipe-

18
19 library and within BEaST output all subjects passed QC. Nonlinear registration operations were

20
21 performed using Automatic Normalization Tools (ANTs) (Avants et al., 2008). All images were

22
23 converted to the MINC format and MAGEt Brain was implemented using tools included in

24
25 MINC (<http://www.bic.mni.mcgill.ca/ServicesSoftware/ServicesSoftwareMincToolKit>),

26
27 including a version of ANTs adapted to work with MINC-tools.
28
29
30
31
32
33
34
35

36 37 **Method 1: Heritability of subfields after adjusting for ipsilateral hippocampal volume and** 38 39 **total brain volume using a univariate model** 40

41
42 Before adjusting for TBV and ipsilateral hippocampal volume on hippocampal subfield volumes,

43
44 we examined the correlation between hippocampal subfields volumes with ipsilateral

45
46 hippocampal volume and TBV; we also investigated correlation between left and right

47
48 hippocampal subfield volumes. Pearson's correlation along with a p-value was calculated using

49
50 R 3.2.1 statistical computing software (R Core Development Team, 2013). Hippocampal subfield

51
52 volumes were adjusted for: 1) sex and age alone, hereafter called 'unadjusted', 2) TBV, 3)

53
54 ipsilateral hippocampal volume, 4) ipsilateral hippocampal volume and TBV together. A
55
56
57
58
59
60

S. Patel and M.M. Chakravarty

1
2
3 residual approach in R was used for each adjustment on the hippocampal subfields volumes and
4
5 then heritability was calculated for each type of adjustment in a univariate ACE model
6
7
8 (calculating the: genetics [A], common environment [C] and unique environment [E] variation).
9
10
11 The resulting analysis examines the influence of overall brain and hippocampus volume on
12
13 subfield heritability measures.

14
15
16 Broad sense heritability (H^2) of hippocampal subfield volumes was estimated using structural
17
18 equation modeling implemented with the OpenMx 2.3.1 (Neale et al., 2015) package within R.
19
20 Following the imaging QC described above, the number of subjects was reduced to 465 healthy
21
22 subjects from 502, based on removing subjects in which the family only had one individual. The
23
24 sample included: 100 MZ twins, 94 DZ twins and 271 siblings (from 177 families, 96 families
25
26 had siblings added to the twin pair and 81 families consisted of non twin siblings only).
27
28

29
30 Heritability is the ratio of variance of a phenotypic measurement in which the numerator is the
31
32 variation attributable to genetics [A] and the denominator is the total observed variation (genetics
33
34 [A], common environment [C] and unique environment [E]). **In classical twin studies the**
35
36 **model assumes that twins (both MZ and DZ twins) are raised together therefore the**
37
38 **common environment is based on twin status and is the same for each individual within the**
39
40 **twin pair. When siblings are introduced the environment is also assumed to be identical but**
41
42 **this may be a less valid assumption. However, heritability estimates were similar whether**
43
44 **or not we included siblings lending weight to the robustness of the test. Therefore common**
45
46 **environment [C] was assumed to be identical within a family.** MZ twins share 100% of their
47
48 genes whereas DZ twins share 50% of segregating genes, similar to DZ twins, non-twin siblings
49
50 share on average 50% of their genes. To account for the genetic [A] component on the twins and
51
52 non-twin siblings in the model, MZ twins had a coefficient of 1 and DZ twins and non-twin
53
54
55
56
57
58
59
60

1
2
3 siblings a coefficient of 0.5. In the dataset, some families had data for only non-twin siblings; in
4
5 these cases a coefficient of 0.5 was used for both family members. Heritability was calculated for
6
7 volume of both ipsilateral hippocampal subfields (CA1, CA2/CA3, CA4/DG, subiculum and
8
9 SRLM) along with TBV and ipsilateral hippocampal volumes.
10
11

12 A full univariate ACE model was applied to each hippocampal subfield and compared against a
13
14 CE model to examine the significance of the genetic variation when “A” is removed from the full
15
16 model using a likelihood ratio test. A significant p-value ($p < 0.05$) for the difference in fit of the
17
18 two models indicates that the “A” component plays an important role in explaining the variance
19
20 of the trait thereby demonstrating the significance of the heritability. This was applied to all 4
21
22 models 1) unadjusted subfield volumes, 2) adjusted for TBV, 3) adjusted for ipsilateral
23
24 hippocampal volume, 4) adjusted for TBV and ipsilateral hippocampal volume together.
25
26 Confidence intervals of 95% on the heritability estimates were calculated for each hippocampal
27
28 subfield in all 4 models within the univariate model.
29
30
31
32
33

34 For both univariate model and bivariate models (shown below) we simply observed the
35
36 heritability estimate of the subfields to see which regions are heritable within the hippocampus
37
38 instead of statistically comparing the heritability estimates between regions. Since the subfield
39
40 volumes are highly correlated with TBV and ipsilateral hippocampal volume the tests would not
41
42 be independent and applying Bonferroni correction would be too stringent. Therefore the p
43
44 values reported when comparing the full ACE model against the CE model are uncorrected
45
46
47 values.
48
49
50
51
52
53
54
55
56
57
58
59
60

Method 2: Heritability using a bivariate model

A bivariate Cholesky decomposition ACE model was used to examine the shared genetic variation (H^2) and the genetic correlation (r_g) between the hippocampal subfield volumes with TBV and ipsilateral hippocampal volume. Genetic correlation measures the shared genetics effects (genetic architecture) between the two volumes. Shared heritability estimates the proportion of genetic variation of one trait that is present in the other. Therefore two structures can have a high genetic correlation such as 0.8 but a low shared heritability of 0.55. This means that there is a large overlap in genes that may influence the variation on both structures but the proportion of shared genetic variation between the structures is low. For the calculation of shared heritability, the order of the phenotypic variables under study is important within the model. We selected the larger global structure as the reference trait while controlling for sex and age. Volumes of the hippocampus subfields and TBV were normalized using z-scores. To examine the significance of heritability scores the bivariate ACE model was compared to a bivariate CE model and confidence intervals (95%) on the heritability scores were calculated.

RESULTS

Segmentation and volumes of hippocampal subfields

From the 511 images segmented using MAGeT Brain, 502 passed quality control (example of a segmented hippocampus in Figure 1). Table II shows the average volumes and standard

deviation of each hippocampal subfield including left and right hippocampal volume and TBV.

From the 5 subfield regions and as expected, the CA1 region showed the largest average volume (left CA1 = 771.51mm^3 , ± 92.15 SD, right CA1 = 836.47mm^3 , ± 98.14 SD) and the CA2/CA3

subfield was the smallest (left CA2/CA3 = 147.91mm^3 , ± 25.11 SD, right CA2/CA3 =

179.22mm^3 , ± 25.18 SD). Table II also includes average volumes and standard deviation for MZ,

DZ and non-twin siblings. Higher total brain volume found in non-twin siblings compared to

both twin groups is accounted by having a higher ratio of females whereas the ratio between

males and females are the same in the non-twin sibling group. It has been shown that females

have smaller TBV compared to males (Kretschmann et al., 1979; Swaab and Hofman, 1984),

which explains the differences in average volumes between both the twins groups and non-twin

siblings.

Hippocampal subfield correlation with TBV and ipsilateral hippocampal volume

Both the left and right hippocampal subfield volumes were highly positively correlated with left and right hippocampal volume respectively (Table III). The highest correlation coefficient

between hippocampus subfields and ipsilateral hippocampal volume was the right and left CA1

and SRLM region ($r = 0.96$, $p < 0.001$) and the lowest correlation coefficients were left CA2/CA3

($r = 0.56$, $p < 0.001$) and right subiculum ($r = 0.60$, $p < 0.001$). In contrast the correlation

coefficient was lower between hippocampal subfields and TBV but still positive. The highest

S. Patel and M.M. Chakravarty

1
2
3 coefficient between hippocampus subfield and TBV was left and right CA1 ($r = 0.69$, $r = 0.66$
4
5 respectively) and lowest correlation coefficient was the left and right CA2/CA3 region ($r = 0.32$,
6
7 $r = 0.47$ respectively). In addition, the left and right hippocampal volume was highly correlated
8
9 with TBV ($r = 0.72$, $r = 0.70$ respectively).
10
11

12 13 14 15 **Univariate Model: Heritability Estimates of hippocampal subfields volumes**

16
17 Unadjusted hippocampal subfield volumes demonstrated moderate to high heritability estimates.
18
19 The highest heritability estimate within hippocampal subfield volumes was the right subiculum
20
21 at 85% and the lowest estimate was the left CA2/CA3 at 38% (Table IV, Figure 2). Observed
22
23 heritability estimates of left and right hippocampal volume were 80% and 81%, respectively, and
24
25 the heritability of TBV was 92%. When adjusted for TBV and ipsilateral hippocampal volume
26
27 the heritability of subfield volumes ranged from 38% (left CA1) to 79% (right subiculum) (Table
28
29 IV, Figure 2) and the heritability of left and right hippocampal volume was lower with estimates
30
31 of 65% and 63% respectively when adjusting for TBV. When both TBV and ipsilateral
32
33 hippocampal volume were adjusted for, the observed heritability estimate of each subfield
34
35 volume was similar to only adjusting for ipsilateral hippocampal volume (Table IV, Figure 2).
36
37 Overall, a trend of lower heritability scores of the hippocampal subfields was seen when
38
39 adjusting for ipsilateral hippocampal volume compared to adjusting for only TBV. For example,
40
41 unadjusted left CA2/CA3 volumes (Table IV, Figure 2) had a heritability estimate of 38% but
42
43 after adjusting for TBV and left hippocampal volume separately, the estimate was higher (41%
44
45 and 53%, respectively). When the left CA2/CA3 volume was adjusted for both left hippocampal
46
47 volume and TBV together the heritability estimate was similar to when adjusting for left
48
49 hippocampal volume only (52%). Similarly, heritability of the left subiculum unadjusted volume
50
51
52
53
54
55
56
57
58
59
60

1
2
3 was 50%, but when adjusted for TBV the estimate was higher (63%). However adjusting for the
4
5 left hippocampal volume along with TBV, the heritability estimate of left subiculum volume was
6
7 also higher than unadjusted left subiculum volume but lower than accounting for only TBV.
8
9

10 When the ACE model is compared to the CE model, the heritability was significant for all
11
12 regions whether or not the subfield volume was adjusted for TBV ($p < 0.05$). When subfield
13
14 volumes were adjusted for ipsilateral hippocampal volume all but one subfield had a significant
15
16 heritability estimate (the left CA1 region $p=0.09$) (Table IV).
17
18
19

20 21 22 **Bivariate model: Shared heritability estimates of hippocampal subfields volumes with TBV** 23 24 **and ipsilateral hippocampal volume** 25 26

27 In the bivariate model, both the genetic correlation and heritability was measured between
28
29 hippocampal subfield volumes with TBV and ipsilateral hippocampal volume. The shared and
30
31 unique heritability estimates were significant ($p < 0.05$) when the full ACE bivariate model was
32
33 compared to the CE model across all subfields with both the TBV and ipsilateral hippocampal
34
35 volume.
36
37

38 The genetic correlation (r_g) between TBV and hippocampal subfield volumes (Table V) was
39
40 positive and slightly higher than the Pearson correlation performed in the univariate model
41
42 (Table III). The shared heritability between TBV and subfield was high ranging from 86% in left
43
44 subiculum to 99% heritability in right subiculum and left CA2/CA3 region (Table VI, Figure 3).
45
46
47

48 Overall the genetic correlation between ipsilateral hippocampal volume and hippocampal
49
50 subfield volumes showed a higher positive genetic correlation trend compared to TBV and
51
52 hippocampal subfield. The genetic correlation of left hippocampal volume and left hippocampal
53
54 subfield volumes ranged from 0.75 in left CA2/CA3 to 0.98 in left CA1 and left SRLM. The
55
56
57
58
59
60

S. Patel and M.M. Chakravarty

1
2
3 range in the right was from 0.66 in the right subiculum to 0.97 in the right CA1 and SRLM
4
5 region (Table V). The shared heritability between left hippocampal and left hippocampal
6
7 subfield ranged from 77% in the CA2/CA3 region to 93% in the left subiculum region (Table
8
9 VII, Figure 4). For the right, the range was 83% in the CA1 region to 94% in the right subiculum
10
11 region (Table VII, Figure 4).
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review

DISCUSSION

In this manuscript, we used 465 healthy twin and non-twin siblings from the HCP dataset to estimate the heritability of hippocampal subfield volumes. Overall in the univariate ACE model, unadjusted hippocampal subfield volumes showed moderate to high heritability estimates ranging from 38% (left CA2/CA3) to 85% (right subiculum). Furthermore, within each subfield the right subfields showed higher heritability estimates compared to left subfields when adjusting for both TBV and ipsilateral hippocampal volume. However the confidence intervals between left and right corresponding subfields overlap suggesting that the amount of genetic variation is similar. A trend of lower heritability estimates was seen when adjusting for both TBV and ipsilateral hippocampal volume. The TBV and ipsilateral hippocampal volume was also highly heritable. Interestingly, hippocampal subfield heritability was often lower when adjusting for ipsilateral hippocampal volume compared to adjusting for only TBV alone. This trend suggests that once you have accounted for the hippocampal volume, the overall brain volume minimally influences the outputs of hippocampal subfield heritability measures. These results suggest when investigating heritability estimate of small brain structures such as hippocampal subfields it is important to account for the influence of larger structures surrounding the region of interest. In addition, these results also provide indirect evidence of the shared heritability between each of the subfields and TBV and ipsilateral hippocampal volume.

To further investigate how TBV and ipsilateral hippocampal volume influence heritability estimates of hippocampal subfield volumes we used a bivariate model to examine the shared genetics variance within the two structures. Shared heritability between ipsilateral hippocampal volume and hippocampal subfields was quite high (77% - 94%) (Table VII), albeit lower than shared heritability between TBV and hippocampal subfields volume (86% - 99%) (Table VI).

S. Patel and M.M. Chakravarty

1
2
3 However, overall the genetic correlation between ipsilateral hippocampal volume and
4
5 hippocampal subfields volume was higher (0.66 - 0.98) than TBV and subfield volumes (0.49 -
6
7 0.77) (Table V), which is similar to the general correlation without accounting for the genetic
8
9 effects (Table III). The highest phenotypic and genetic correlation between TBV and
10
11 hippocampal subfield region was the CA1 region (0.69 and 0.77 respectively). Similarly,
12
13 Harding et al., (1998) also reported high correlation between CA1 and cerebral volume of 0.68
14
15 when exploring the variation in the number of hippocampal neurons with age and brain volume
16
17 (Harding et al., 1998). These correlations suggest a possible similarity in genetic etiologies
18
19 governing the size between CA1 region and total brain which can be seen in the high genetic
20
21 correlation observed in our study. Overall the genetic correlation trend suggests that there is less
22
23 overlap in genetic effects (genetic variation) that influence TBV and hippocampal subfield
24
25 volume than those that influence ipsilateral hippocampal volume and subfields based on genetic
26
27 correlation estimates. However in terms of the proportion of the genetic correlation estimates, the
28
29 shared heritability between subfield volume and TBV demonstrates a greater proportion of
30
31 shared genetic variation from the genetics correlation than ipsilateral hippocampal volume. This
32
33 could be due to the fact that the genetic architecture related to total brain volume is simpler than
34
35 that of the ipsilateral hippocampal volume and warrants more study. For example there can be
36
37 different genetic signaling mechanisms for both TBV and ipsilateral hippocampal development,
38
39 where the types of genes and gene expression can differ (Cipriani et al., 2016; Roy et al., 2000;
40
41 Stein et al., 2012). In addition, the role of genes that influence brain development processes
42
43 (Miller et al., 2014) is another area of research that can be used to understand the genetic
44
45 architecture. To tease apart these differences, more studies are required comparing the
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 development of the brain as a whole and the subfields of the hippocampus at pre-natal stages to
4
5 adult hood.
6

7
8 The recent study by Whelan et al., (2015) also analyzed heritability of hippocampal subfields
9
10 using the Queen Twins Imaging (QTIM) dataset (T1-weighted MRI from 132 MZ twins pairs
11
12 and 232 DZ twin pairs (Whelan et al., 2015)). The differences between our study and the Whelan
13
14 study relate to the quality of the dataset used and the study design. A significant advantage in the
15
16 use of the HCP dataset is that it not only includes twins but also their non-twin siblings. Families
17
18 within our data consisted of twin pairs, non-twin siblings and twin pairs with their non-twin
19
20 sibling. Posthuma et al., (2000) have shown that within a family, adding one or two non-twin
21
22 siblings along with their twin siblings is an advantage in heritability analysis by reducing the
23
24 sample size needed and increase the statistical power to detect heritability (Posthuma and
25
26 Boomsma, 2000). Also the HCP dataset has high-resolution MRI T1-weighted scans with voxel
27
28 dimensions of 0.7x 0.7x 0.7mm, (whereas QTIM data has a voxel dimension of 0.94 x 0.98 x
29
30 0.98mm). Implicitly, the specific subfields can actually be resolved using HCP data whereas this
31
32 has yet to be proven to be the case in more standard T1-weighted acquisition (Pipitone et al.,
33
34 2014). Furthermore, in our study the heritability estimates are presented bilaterally and we
35
36 explore the effects of ipsilateral hippocampal volume and TBV on subfield volume heritability,
37
38 whereas only the overall heritability of each subfield volumes was described by Whelan et al.,
39
40 (2015). This is a critical distinction as we clearly show a significant impact of ipsilateral
41
42 hippocampal volume and a moderate impact of TBV on subfield heritability. We note here that
43
44 our findings report similar results for the heritability of the hippocampus, where Whelan et al.,
45
46 (2015) reported an estimate of 88% heritability in total hippocampal volume and our study
47
48 estimates slightly lower heritability of 80% and 81% for left and right hippocampal volume
49
50
51
52
53
54
55
56
57
58
59
60

S. Patel and M.M. Chakravarty

1
2
3 respectively. Similarly across subfields the heritability estimates were lower in our study
4
5 compared to Whelan et al., (2015). Heritability estimates for unadjusted left (77%) and right
6
7 (80%) CA4/DG were similar to estimates reported by Whelan et al., for the CA4 region (79%)
8
9 and the granule cells of the DG region (82%). Our paper also extends the analysis through the
10
11 use of the bivariate model. This approach has previously only been used to examine total brain
12
13 volume with total hippocampal volumes (DeStefano et al., 2009). Lastly, the segmentation
14
15 process differs between both studies. We used MAGeT Brain, the Whelan study used FreeSurfer
16
17 (which was previously validated against our own Winterburn protocol (Iglesias et al., 2015)),
18
19 which results in differences between the hippocampal subfield definitions and the methodology
20
21 used to define these subfields. In our study we define 5 subfields which included the CA1,
22
23 CA2/CA3, CA4/DG, SRLM and subiculum (Winterburn et al., 2013). Whereas, subfield
24
25 delineation in the Whelan et al., study did not include the CA2 region in their heritability
26
27 estimates. However, the Whelan paper examined other regions which we did not, such as the
28
29 fimbria (white matter tracts) and hippocampal-amygdaloid transition area. The framework
30
31 provided in our study allows us to examine the role of TBV and overall hippocampal volume to
32
33 the heritability of individual subfields, which will allow for further investigation with respect to
34
35 newer subfield definitions that are the product of the ongoing subfield segmentation
36
37 harmonization effort (Yushkevich et al., 2015a).
38
39
40
41
42
43
44
45

46
47 The use of twin/non-twin sibling HCP design and dataset presents limitations due to variation in
48
49 intrauterine environment and pre/postnatal complications. Unfortunately, to the best of our
50
51 knowledge, this information is not provided for HCP subjects. As part of the exclusion process
52
53 individuals that were premature (see method section) was excluded but other prenatal or
54
55 postnatal information was not collected which may affect hippocampal volume and in turn affect
56
57
58
59
60

1
2
3 heritability estimates. For example the CA1 region is hypoxia-sensitive and prenatal hypoxia
4 may be related to greater neuronal loss in the CA1 region compared to other regions (Kuchna,
5
6 1994).
7
8

9
10 The HCP scans used in this study are of high-resolution and –contrast with 0.7mm isotropic
11 voxels. **Figure 1 and figure S2 in the supplementary section demonstrate that the internal**
12 **structure of the hippocampus can be distinguished in the sagittal and coronal view.** While
13 work from our group has assessed the feasibility of subfield segmentation using standard T1-
14 weighted images (Amaral et al., 2016; Pipitone et al., 2014), as has work from the FreeSurfer
15 group (Iglesias et al., 2015; Iglesias et al., 2016) (although see refs for criticism with respect to
16 FreeSurfer subfield segmentations (de Flores et al., 2015; Wisse et al., 2014)), it is important to
17 note that there are other options that are often used for subfield level segmentations. **Other**
18 **prevalent techniques use T2-weighted images that are high-resolution in the coronal plane**
19 **and low-resolution outside of this plane (Goubran et al., 2016; Yushkevich et al., 2015b).**
20
21

22 However it is likely that many of the well-documented limitations inherent to other subfield
23 segmentation techniques (Yushkevich et al., 2015a), are also relevant here. These limitations
24 include ambiguity in the identification of the CA2/3 definition and the CA1/subiculum border.
25 A limitation found in MRI-based neuroanatomical studies is that it cannot account for all
26 cytoarchitectonic features within the subfields of the hippocampus. In our segmentation process
27 a combination of intensity, contrast, and geometric rules based on known cytoarchitectonic
28 definitions were used to define the hippocampal subfields (Winterburn et al., 2013); however
29 there are specific boundaries where there is less certainty across subfield protocols. For example,
30 defining the boundary between the CA1 region and subiculum is difficult and is one of the most
31 variable delineations in hippocampal subfield definitions (Yushkevich et al., 2015a). This is not
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

S. Patel and M.M. Chakravarty

1
2
3 only true at the level and resolution offered by MRI data, but even when attempting to identify
4 this boundary at the microscopic-level using histological data (Adler et al., 2014). This
5 heterogeneity in the ability to identify the subiculum will logically contribute to possible
6 measurement errors and higher uncertainty in structure-specific heritability estimates. The
7 ambiguity in this boundary at the level of subiculum/CA1 delineation may have also contributed
8 to the lack of significance in our left CA1 heritability estimates after adjusting for TBV and left
9 hippocampal volume together in the ACE univariate model. However when we did not adjust for
10 TBV or left hippocampal volume our heritable estimates for the CA1 region are significant
11 (comparing univariate ACE against CE model) for both right and left CA1 volume, 73% and
12 72% respectively. Similarly, Whelan et al. (2015) reported a slightly higher and significant
13 heritability estimate of 84% for CA1 volume. The CA2/CA3 and SRLM subfields are also
14 regions that are difficult to define based on the thin and complex structure on T1 weighted
15 images. Winterburn et al (2013) showed low test-retest results of the manual segmentation
16 protocol for the CA2/CA3 and the SRLM region. Also Pipitone et al., (2014) showed that the
17 CA2/CA3 and SRLM region are less reliably reproduced segmentation on standard $1 \times 1 \times 1 \text{ mm}^3$
18 T1-weighted images out of the five subfields. In addition, Wisse et al (2016) demonstrated lower
19 accuracy through automated segmentation of hippocampal subfields on 7T images on small
20 subfield such as the CA2/CA3 regions which are undersegmented. In our study the heritability
21 was higher in the left CA2/CA3 region when adjusting for total hippocampal volume compared
22 to no adjustment which is the opposite trend compared to other subfields regions in the ACE
23 univariate model. These limitations in defining the CA2/CA3 and SRLM region can potentially
24 influence the accuracy of the heritability scores. Therefore higher resolution images, the use of
25 T2 weighted images, and methodological improvements to segmentation techniques may further
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 help with defining these regions. As such, direct comparison of results between studies need to
4
5 be made with caution and a detailed understanding of these different protocols to define the
6
7 subfields.
8
9

10
11 There are different types of segmentation tools available and used to segment the hippocampus
12
13 such as FreeSurfer (Fischl et al., 2002) and Automatic Segmentation of Hippocampal Subfields
14
15 (ASHS (Yushkevich et al., 2015b)). Comparisons have been done between FreeSurfer and
16
17 MAGeT Brain by Pipitone et al., (2014) demonstrating the robustness of MAGeT brain on whole
18
19 hippocampal segmentations. A recent comparison of automated segmentation approaches,
20
21 including MAGeT Brain, ASHS, FreeSurfer and a Bayesian inference model (Van Leemput et
22
23 al., 2009) has not been done. Therefore in future research a comparison study on automatic
24
25 segmentations protocols will be of great value. Furthermore, it will be interesting to compare the
26
27 automated segmentation methods on the HCP dataset, along with calculating heritability
28
29 estimates of hippocampal subfield volumes across each segmentation protocol to evaluate the
30
31 reliability in the heritability scores.
32
33
34
35
36
37

38 In conclusion, we have demonstrated the heritability of hippocampal subfield volumes using the
39
40 HCP data in a twin and non-twin sibling design using a univariate and bivariate model. The
41
42 univariate model allowed us to examine the heritability of each subfield itself adjusting for TBV
43
44 and ipsilateral hippocampal volume, whereas in the bivariate model we were able to examine the
45
46 shared heritability and genetic correlation between two traits. The univariate model demonstrated
47
48 the heritability of the subfield volumes was lower but significant compared to the heritability of
49
50 the left or right hippocampal volume. From the bivariate model, shared heritability between
51
52 many subfields and the hippocampal volume was high and significant. Identifying subfields that
53
54 have significant and high heritability estimates such as the right subiculum, right CA2/CA3 and
55
56
57
58
59
60

S. Patel and M.M. Chakravarty

1
2
3 right CA4/DG demonstrates their utility as quantitative phenotypes in neurological and
4
5
6 psychiatric illnesses.
7

8
9 The use of both univariate and bivariate models in the future allows for the examination of
10
11 different aspects of the genetic architecture on the target traits. The univariate model
12
13 demonstrates the heritability of hippocampal subfield volumes in isolation after removing the
14
15 influence of TBV and ipsilateral hippocampal volume. However, the bivariate model allows us
16
17 to capture the influence one trait has on another by examining the shared heritability between the
18
19 two traits. Our univariate model has shown that volumes of smaller target structures are
20
21 influenced by larger structures that contain the smaller targeted structure and this was quantified
22
23 by shared heritability estimates in the bivariate model. Therefore it is important to look at the
24
25 heritability of a structure in isolation but also in relation to other neuroanatomical structures to
26
27 get a full understanding of the genetic architecture and genetic interaction found within and
28
29 between brain structures. As segmentation and image acquisition techniques improve, sample
30
31 sizes available will also further improve heritability and genome-wide association analysis. Our
32
33 work provides a basis for similar ongoing studies, such as those pursued through the ENIGMA
34
35 consortium (Stein et al., 2012; Thompson et al., 2014). Therefore the data presented in this
36
37 manuscript further motivates the association between the genetic basis of the structure and
38
39 function of hippocampal subfields in cases of normal brain function and dysfunction.
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

ACKNOWLEDGMENTS

Data were provided [in part] by the Human Connectome Project, WU-Minn Consortium (Principal Investigators: David Van Essen and Kamil Ugurbil; 1U54MH091657) funded by the 16 NIH Institutes and Centers that support the NIH Blueprint for Neuroscience Research; and by the McDonnell Center for Systems Neuroscience at Washington University. Computations were performed on the gpc supercomputer at the SciNet HPC Consortium (Loken et al., 2010). SciNet is funded by: the Canada Foundation for Innovation under the auspices of Compute Canada; the Government of Ontario; Ontario Research Fund - Research Excellence; and the University of Toronto. MMC is funded by the National Sciences and Engineering Research Council, Canadian Institutes for Health, Weston Brain Institute, Michael J. Fox Foundation for Parkinson's Research, Alzheimer's Society, and Brain Canada. JK holds the Joanne Murphy Professorship in Behavioral Science and is also funded by the Canadian Institutes for Health. SP is funded by the Cadsby Foundation and Canadian Institutes for Health. The authors report no conflict of interest.

References

- 1
2
3
4
5
6 Adler DH, Pluta J, Kadivar S, Craige C, Gee JC, Avants BB, Yushkevich PA (2014): Histology-derived volumetric annotation of the human hippocampal subfields in postmortem MRI. *Neuroimage*, 84:505-23.
- 7
8
9
10 Altshuler LL, Bartzokis G, Grieder T, Curran J, Mintz J (1998): Amygdala enlargement in
11 bipolar disorder and hippocampal reduction in schizophrenia: an MRI study
12 demonstrating neuroanatomic specificity. *Arch Gen Psychiatry*, 55:663-4.
- 13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
- Amaral RS, Park MT, Devenyi GA, Lynn V, Pipitone J, Winterburn J, Chavez S, Schira M, Lobaugh NJ, Voineskos AN et al. (2016): Manual segmentation of the fornix, fimbria, and alveus on high-resolution 3T MRI: Application via fully-automated mapping of the human memory circuit white and grey matter in healthy and pathological aging. *Neuroimage*.
- Amunts K, Kedo O, Kindler M, Pieperhoff P, Mohlberg H, Shah NJ, Habel U, Schneider F, Zilles K (2005): Cytoarchitectonic mapping of the human amygdala, hippocampal region and entorhinal cortex: intersubject variability and probability maps. *Anat Embryol (Berl)*, 210:343-52.
- Avants BB, Epstein CL, Grossman M, Gee JC (2008): Symmetric diffeomorphic image registration with cross-correlation: evaluating automated labeling of elderly and neurodegenerative brain. *Med Image Anal*, 12:26-41.
- Baare WF, Hulshoff Pol HE, Boomsma DI, Posthuma D, de Geus EJ, Schnack HG, van Haren NE, van Oel CJ, Kahn RS (2001): Quantitative genetic modeling of variation in human brain morphology. *Cerebral cortex*, 11:816-24.
- Bhagwat N, Pipitone J, Winterburn JL, Guo T, Duerden EG, Voineskos AN, Lepage M, Miller SP, Pruessner JC, Chakravarty MM (2016): Manual-Protocol Inspired Technique for Improving Automated MR Image Segmentation during Label Fusion. *Front Neurosci*, 10:325.
- Blatter DD, Bigler ED, Gale SD, Johnson SC, Anderson CV, Burnett BM, Parker N, Kurth S, Horn SD (1995): Quantitative volumetric analysis of brain MR: normative database spanning 5 decades of life. *AJNR Am J Neuroradiol*, 16:241-51.
- Bogerts B, Lieberman JA, Ashtari M, Bilder RM, Degreef G, Lerner G, Johns C, Masiar S (1993): Hippocampus-amygdala volumes and psychopathology in chronic schizophrenia. *Biol Psychiatry*, 33:236-46.
- Braak H, Braak E (1991): Neuropathological stageing of Alzheimer-related changes. *Acta Neuropathol*, 82:239-59.
- Braskie MN, Toga AW, Thompson PM (2013): Recent advances in imaging Alzheimer's disease. *J Alzheimers Dis*, 33 Suppl 1:S313-27.
- Bremner JD, Narayan M, Anderson ER, Staib LH, Miller HL, Charney DS (2000): Hippocampal volume reduction in major depression. *Am J Psychiatry*, 157:115-8.
- Campbell S, Marriott M, Nahmias C, MacQueen GM (2004): Lower hippocampal volume in patients suffering from depression: a meta-analysis. *Am J Psychiatry*, 161:598-607.
- Chakravarty MM, Steadman P, van Eede MC, Calcott RD, Gu V, Shaw P, Raznahan A, Collins DL, Lerch JP (2013): Performing label-fusion-based segmentation using multiple automatically generated templates. *Hum Brain Mapp*, 34:2635-54.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
- Cipriani S, Nardelli J, Verney C, Delezoide AL, Guimiot F, Gressens P, Adle-Biassette H (2016): Dynamic Expression Patterns of Progenitor and Pyramidal Neuron Layer Markers in the Developing Human Hippocampus. *Cereb Cortex*, 26:1255-71.
- Collins DL, Pruessner JC (2010): Towards accurate, automatic segmentation of the hippocampus and amygdala from MRI by augmenting ANIMAL with a template library and label fusion. *Neuroimage*, 52:1355-66.
- de Flores R, La Joie R, Landeau B, Perrotin A, Mezenge F, de La Sayette V, Eustache F, Desgranges B, Chetelat G (2015): Effects of age and Alzheimer's disease on hippocampal subfields: comparison between manual and FreeSurfer volumetry. *Hum Brain Mapp*, 36:463-74.
- DeStefano AL, Seshadri S, Beiser A, Atwood LD, Massaro JM, Au R, Wolf PA, DeCarli C (2009): Bivariate heritability of total and regional brain volumes: the Framingham Study. *Alzheimer Dis Assoc Disord*, 23:218-23.
- Duvernoy HM. (2005) *The human hippocampus: functional anatomy, vascularization and serial sections with MRI*. Springer Science & Business Media.
- Eskildsen SF, Coupe P, Fonov V, Manjon JV, Leung KK, Guizard N, Wassef SN, Ostergaard LR, Collins DL (2012): BEaST: brain extraction based on nonlocal segmentation technique. *Neuroimage*, 59:2362-73.
- Fatterpekar GM, Naidich TP, Delman BN, Aguinaldo JG, Gultekin SH, Sherwood CC, Hof PR, Drayer BP, Fayad ZA (2002): Cytoarchitecture of the human cerebral cortex: MR microscopy of excised specimens at 9.4 Tesla. *AJNR Am J Neuroradiol*, 23:1313-21.
- Fischl B, Salat DH, Busa E, Albert M, Dieterich M, Haselgrove C, van der Kouwe A, Killiany R, Kennedy D, Klaveness S et al. (2002): Whole brain segmentation: automated labeling of neuroanatomical structures in the human brain. *Neuron*, 33:341-55.
- Goubran M, Bernhardt BC, Cantor-Rivera D, Lau JC, Bliston C, Hammond RR, de Ribaupierre S, Burneo JG, Mirsattari SM, Steven DA et al. (2016): In vivo MRI signatures of hippocampal subfield pathology in intractable epilepsy. *Hum Brain Mapp*, 37:1103-19.
- Harding AJ, Halliday GM, Kril JJ (1998): Variation in hippocampal neuron number with age and brain volume. *Cereb Cortex*, 8:710-8.
- Haukvik UK, Westlye LT, Morch-Johnsen L, Jorgensen KN, Lange EH, Dale AM, Melle I, Andreassen OA, Agartz I (2015): In vivo hippocampal subfield volumes in schizophrenia and bipolar disorder. *Biol Psychiatry*, 77:581-8.
- Iglesias JE, Augustinack JC, Nguyen K, Player CM, Player A, Wright M, Roy N, Frosch MP, McKee AC, Wald LL et al. (2015): A computational atlas of the hippocampal formation using ex vivo, ultra-high resolution MRI: Application to adaptive segmentation of in vivo MRI. *Neuroimage*, 115:117-37.
- Iglesias JE, Van Leemput K, Augustinack J, Insausti R, Fischl B, Reuter M (2016): Bayesian longitudinal segmentation of hippocampal substructures in brain MRI using subject-specific atlases. *Neuroimage*, 141:542-55.
- Jacquard A (1983): Heritability: one word, three concepts. *Biometrics*, 39:465-77.
- Kretschmann HJ, Schleicher A, Wingert F, Zilles K, Loblich HJ (1979): Human brain growth in the 19th and 20th century. *J Neurol Sci*, 40:169-88.
- Kuchna I (1994): Quantitative studies of human newborns' hippocampal pyramidal cells after perinatal hypoxia. *Folia Neuropathol*, 32:9-16.

S. Patel and M.M. Chakravarty

- 1
2
3 La Joie R, Fouquet M, Mezenge F, Landeau B, Villain N, Mevel K, Pelerin A, Eustache F,
4 Desgranges B, Chetelat G (2010): Differential effect of age on hippocampal subfields
5 assessed using a new high-resolution 3T MR sequence. *Neuroimage*, 53:506-14.
- 6
7 Loken C, Gruner D, Groer L, Peltier R, Bunn N, Craig M, Henriques T, Dempsey J, Yu C-H,
8 Chen J. (SciNet: lessons learned from building a power-efficient top-20 system and data
9 centre). In; 2010. IOP Publishing. p 012026.
- 10
11 Mai J, Paxinos G, Voss T (2008) *Atlas of the Human Brain*, 3rd Edn. Elsevier/Academic Press:
12 London, UK.
- 13
14 Miller JA, Ding SL, Sunkin SM, Smith KA, Ng L, Szafer A, Ebbert A, Riley ZL, Royall JJ,
15 Aiona K et al. (2014): Transcriptional landscape of the prenatal human brain. *Nature*,
16 508:199-206.
- 17
18 Mouiha A, Duchesne S (2011): Hippocampal atrophy rates in Alzheimer's disease: automated
19 segmentation variability analysis. *Neurosci Lett*, 495:6-10.
- 20
21 Mueller SG, Stables L, Du AT, Schuff N, Truran D, Cashdollar N, Weiner MW (2007):
22 Measurement of hippocampal subfields and age-related changes with high resolution
23 MRI at 4T. *Neurobiol Aging*, 28:719-26.
- 24
25 Mueller SG, Weiner MW (2009): Selective effect of age, Apo e4, and Alzheimer's disease on
26 hippocampal subfields. *Hippocampus*, 19:558-64.
- 27
28 Neale M, Cardon L. (1992) *Methodology for genetic studies of twins and families*. Springer
29 Science & Business Media.
- 30
31 Neale MC, Hunter MD, Pritikin JN, Zahery M, Brick TR, Kirkpatrick RM, Estabrook R, Bates
32 TC, Maes HH, Boker SM (2015): *OpenMx 2.0: Extended Structural Equation and
33 Statistical Modeling*. *Psychometrika*.
- 34
35 Pennington BF, Filipek PA, Lefly D, Chhabildas N, Kennedy DN, Simon JH, Filley CM,
36 Galaburda A, DeFries JC (2000): A twin MRI study of size variations in human brain.
37 *Journal of cognitive neuroscience*, 12:223-32.
- 38
39 Pipitone J, Park MT, Winterburn J, Lett TA, Lerch JP, Pruessner JC, Lepage M, Voineskos AN,
40 Chakravarty MM (2014): Multi-atlas segmentation of the whole hippocampus and
41 subfields using multiple automatically generated templates. *Neuroimage*, 101:494-512.
- 42
43 Posthuma D, Boomsma DI (2000): A note on the statistical power in extended twin designs.
44 *Behav Genet*, 30:147-58.
- 45
46 R Core Development Team (2013) *A language and environment for statistical computing*. R
47 Foundation for Statistical Computing, Vienna, Austria.
- 48
49 Roy NS, Wang S, Jiang L, Kang J, Benraiss A, Harrison-Restelli C, Fraser RA, Couldwell WT,
50 Kawaguchi A, Okano H et al. (2000): In vitro neurogenesis by progenitor cells isolated
51 from the adult human hippocampus. *Nat Med*, 6:271-7.
- 52
53 Stein JL, Medland SE, Vasquez AA, Hibar DP, Senstad RE, Winkler AM, Toro R, Appel K,
54 Bartecek R, Bergmann O et al. (2012): Identification of common variants associated with
55 human hippocampal and intracranial volumes. *Nat Genet*, 44:552-61.
- 56
57 Sullivan EV, Pfefferbaum A, Swan GE, Carmelli D (2001): Heritability of hippocampal size in
58 elderly twin men: equivalent influence from genes and environment. *Hippocampus*,
59 11:754-62.
- 60
61 Swaab DF, Hofman MA (1984): Sexual differentiation of the human brain. A historical
62 perspective. *Prog Brain Res*, 61:361-74.
- 63
64 Tamminga CA, Stan AD, Wagner AD (2010): The hippocampal formation in schizophrenia. *Am
65 J Psychiatry*, 167:1178-93.

- 1
2
3 Thompson PM, Stein JL, Medland SE, Hibar DP, Vasquez AA, Renteria ME, Toro R, Jahanshad
4 N, Schumann G, Franke B et al. (2014): The ENIGMA Consortium: large-scale
5 collaborative analyses of neuroimaging and genetic data. *Brain Imaging Behav*, 8:153-82.
6
7 Treadway MT, Waskom ML, Dillon DG, Holmes AJ, Park MT, Chakravarty MM, Dutra SJ,
8 Polli FE, Iosifescu DV, Fava M et al. (2015): Illness progression, recent stress, and
9 morphometry of hippocampal subfields and medial prefrontal cortex in major depression.
10 *Biol Psychiatry*, 77:285-94.
11
12 Van Essen DC, Smith SM, Barch DM, Behrens TE, Yacoub E, Ugurbil K (2013): The WU-Minn
13 Human Connectome Project: an overview. *Neuroimage*, 80:62-79.
14
15 Van Essen DC, Ugurbil K, Auerbach E, Barch D, Behrens TE, Bucholz R, Chang A, Chen L,
16 Corbetta M, Curtiss SW et al. (2012): The Human Connectome Project: a data acquisition
17 perspective. *Neuroimage*, 62:2222-31.
18
19 Van Leemput K, Bakkour A, Benner T, Wiggins G, Wald LL, Augustinack J, Dickerson BC,
20 Golland P, Fischl B (2008): Model-based segmentation of hippocampal subfields in ultra-
21 high resolution in vivo MRI. *Med Image Comput Comput Assist Interv*, 11:235-43.
22
23 Van Leemput K, Bakkour A, Benner T, Wiggins G, Wald LL, Augustinack J, Dickerson BC,
24 Golland P, Fischl B (2009): Automated segmentation of hippocampal subfields from
25 ultra-high resolution in vivo MRI. *Hippocampus*, 19:549-57.
26
27 Voineskos AN, Winterburn JL, Felsky D, Pipitone J, Rajji TK, Mulsant BH, Chakravarty MM
28 (2015): Hippocampal (subfield) volume and shape in relation to cognitive performance
29 across the adult lifespan. *Hum Brain Mapp*, 36:3020-37.
30
31 Wang H, Suh JW, Das SR, Pluta JB, Craige C, Yushkevich PA (2013): Multi-Atlas
32 Segmentation with Joint Label Fusion. *IEEE Trans Pattern Anal Mach Intell*, 35:611-23.
33
34 Whelan CD, Hibar DP, van Velzen LS, Zannas AS, Carrillo-Roa T, McMahon K, Prasad G,
35 Kelly S, Faskowitz J, deZubiracay G et al. (2015): Heritability and reliability of
36 automatically segmented human hippocampal formation subregions. *Neuroimage*,
37 128:125-137.
38
39 Winterburn JL, Pruessner JC, Chavez S, Schira MM, Lobaugh NJ, Voineskos AN, Chakravarty
40 MM (2013): A novel in vivo atlas of human hippocampal subfields using high-resolution
41 3 T magnetic resonance imaging. *Neuroimage*, 74:254-65.
42
43 Wisse LE, Biessels GJ, Geerlings MI (2014): A Critical Appraisal of the Hippocampal Subfield
44 Segmentation Package in FreeSurfer. *Front Aging Neurosci*, 6:261.
45
46 Yang X, Goh A, Chen SH, Qiu A (2013): Evolution of hippocampal shapes across the human
47 lifespan. *Hum Brain Mapp*, 34:3075-85.
48
49 Yushkevich PA, Amaral RS, Augustinack JC, Bender AR, Bernstein JD, Boccardi M, Bocchetta
50 M, Burggren AC, Carr VA, Chakravarty MM et al. (2015a): Quantitative comparison of
51 21 protocols for labeling hippocampal subfields and parahippocampal subregions in in
52 vivo MRI: towards a harmonized segmentation protocol. *Neuroimage*, 111:526-41.
53
54 Yushkevich PA, Pluta JB, Wang H, Xie L, Ding SL, Gertje EC, Mancuso L, Kliot D, Das SR,
55 Wolk DA (2015b): Automated volumetry and regional thickness analysis of hippocampal
56 subfields and medial temporal cortical structures in mild cognitive impairment. *Hum*
57 *Brain Mapp*, 36:258-87.
58
59 Yushkevich PA, Wang H, Pluta J, Das SR, Craige C, Avants BB, Weiner MW, Mueller S
60 (2010): Nearly automatic segmentation of hippocampal subfields in in vivo focal T2-
weighted MRI. *Neuroimage*, 53:1208-24.

FIGURE LEGENDS

Figure 1. Hippocampal subfield segmentation, T1 sagittal scan. Left image is non-segmented and right image is segmented. Hippocampal subfields: CA1, CA2/CA3, CA4 and DG, SRLM, Subiculum.

Figure 2. Heritability scores (H^2) of left (L) and right (R) hippocampal subfields volumes, L and R hippocampal volumes and total brain volume (TBV) with 95% confidence intervals (sample size $N = 465$). Pink bars indicate unadjusted subfield volume, whereas green bars indicates TBV adjustment, blue bars indicates ipsilateral (left or right) hippocampal volume adjustment and purple bars indicates ipsilateral hippocampal and TBV adjustment on subfield volumes.

Figure 3. Bivariate heritability scores (H^2) between TBV and left (L) and right (R) hippocampal subfields volumes, with 95% confidence intervals (sample size $N = 465$). Pink bars indicate heritability estimates of TBV, whereas green bars indicates heritability estimates of sub-regions (hippocampal subfields and whole hippocampal volumes) and blue bars indicates shared heritability estimates between TBV and sub-regions.

Figure 4. Bivariate heritability scores (H^2) between ipsilateral hippocampal volume and left (L) and right (R) hippocampal subfields volumes, with 95% confidence intervals (sample size $N = 465$). Pink bars indicate heritability estimates of ipsilateral hippocampal volume, whereas green bars indicates heritability estimates of sub-regions (hippocampal subfield volumes) and blue bars indicates shared heritability estimates between ipsilateral hippocampal volume and sub-regions.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 1. Hippocampal subfield segmentation, T1 sagittal scan. Left image is non-segmented and right image is segmented. Hippocampal subfields: CA1, CA2/CA3, CA4 and DG, SRLM, Subiculum.

166x64mm (150 x 150 DPI)

Peer Review




Figure 2. Heritability scores (H^2) of left (L) and right (R) hippocampal subfields volumes, L and R hippocampal volumes and total brain volume (TBV) with 95% confidence intervals (sample size $N = 465$). Pink bars indicate unadjusted subfield volume, whereas green bars indicates TBV adjustment, blue bars indicates ipsilateral (left or right) hippocampal volume adjustment and purple bars indicates ipsilateral hippocampal and TBV adjustment on subfield volumes.

165x66mm (150 x 150 DPI)




Figure 3. Bivariate heritability scores (H^2) between TBV and left (L) and right (R) hippocampal subfields volumes, with 95% confidence intervals (sample size $N = 465$). Pink bars indicate heritability estimates of TBV, whereas green bars indicates heritability estimates of sub-regions (hippocampal subfields and whole hippocampal volumes) and blue bars indicates shared heritability estimates between TBV and sub-regions.

165x89mm (150 x 141 DPI)




Figure 4. Bivariate heritability scores (H^2) between ipsilateral hippocampal volume and left (L) and right (R) hippocampal subfields volumes, with 95% confidence intervals (sample size $N = 465$). Pink bars indicate heritability estimates of ipsilateral hippocampal volume, whereas green bars indicates heritability estimates of sub-regions (hippocampal subfield volumes) and blue bars indicates shared heritability estimates between ipsilateral hippocampal volume and sub-regions.

165x89mm (150 x 140 DPI)

TABLE I. Demographic breakdown of monozygotic twins (MZ), dizygotic twins (DZ) and non-twin siblings from the subset data of the HCP, including averages and standard deviation (\pm SD). Average fluid intelligence is a measure of number of correct responses out of 24 questions.

	N	Average Age (year \pm SD)	Age Range	Gender Female : Male	Average handedness (\pm SD)	Average fluid intelligence (\pm SD)
MZ	100	29.97 (3.11)	22-36	74 : 26	72.4 (41.76)	16.26 (4.64)
DZ	94	29.86 (2.93)	22-35	67 : 27	63.19 (46.06)	16.71 (4.81)
Non-twin siblings	271	28.75 (3.72)	22-36	136 : 135	63.56 (48.06)	16.12 (5.11)
Total	465	29.24 (3.49)	22-36	277 : 188	65.39 (46.42)	16.27 (4.95)

TABLE II. Average volume and standard deviation (\pm SD) of unadjusted left and right hippocampal subfields in monozygotic twins (MZ), dizygotic twins (DZ) and non-twin siblings.

Region	MZ average volume (mm³ \pm SD) N=100	DZ average volume (mm³ \pm SD) N=94	Non twin sibling average volume (mm³ \pm SD) N=271	Total Sample average volume mm³ (\pm SD) N=465
Left CA1	755.71 (89.43)	761.50 (74.00)	780.82 (97.77)	771.51 (92.15)
Right CA1	821.41(90.97)	826.97 (87.04)	845.33 (103.51)	836.47 (98.14)
Left CA2CA3	145.9705 (22.55)	146.69 (21.56)	149.04 (27.10)	147.91(25.12)
Right CA2CA3	174.82(22.61)	174.13 (24.04)	182.61 (25.98)	179.22 (25.18)
Left CA4DG	623.61 (70.02)	621.75 (55.12)	649.06 (73.27)	638.07 (70.34)
Right CA4GD	615.00(65.91)	616.62 (55.99)	649.88 (72.54)	635.67 (70.00)
Left SRLM	577.99 (67.90)	581.60 (61.98)	589.51 (73.93)	585.44 (70.42)
Right SRLM	543.56 (64.33)	546.66 (63.48)	558.37 (68.70)	552.81 (66.94)
Left Subiculum	331.43 (46.22)	327.84 (43.27)	335.89 (50.83)	333.31 (48.43)
Right Subiculum	329.45(43.63)	327.48 (38.08)	333.92 (45.87)	331.66 (43.92)
Left Hippocampus	2434.72 (253.57)	2439.39 (216.75)	2504.33 (277.15)	2476.23 (262.56)
Right Hippocampus	2484.25 (249.13)	2491.87(233.02)	2570.11 (277.67)	2535.82 (265.78)
Total brain	1344941.77 (138663.72)	1351379.79 (117637.94)	1393633.69 (150431.05)	1374620.66 (143368.64)

TABLE III. Pearson Correlations (r) between subfield with ipsilateral hippocampal volume and total brain volume (TBV). All subfield volumes showed significant correlation between ipsilateral hippocampal volume and TBV ($p < 0.001$).

	Left Hippocampal Volume	TBV
Left CA1	0.96	0.69
Left CA2CA3	0.56	0.32
Left CA4DG	0.91	0.64
Left SRLM	0.96	0.60
Left Subiculum	0.58	0.63
Left Hippocampus	1.00	0.72
	Right Hippocampal Volume	TBV
Right CA1	0.96	0.66
Right CA2CA3	0.74	0.47
Right CA4DG	0.89	0.63
Right SRLM	0.96	0.60
Right Subiculum	0.60	0.61
Right Hippocampus	1.00	0.71

TABLE IV. Heritability estimates of left and right hippocampal subfield volumes with 95% confidence intervals (sample size N = 465). Significant heritability p values are represented by * indicates $p < 0.05$, and ** indicates $p < 0.001$ when ACE model is compared to CE model. Heritability estimates highlighted in yellow are not significant, but show a trend towards significance.

Region	No Volume Adjustment (only adjusted for sex and age)		Total brain volume adjustment		ipsilateral hippocampal volume adjustment		ipsilateral hippocampal and total brain volume adjustment	
	H ² (%)	95% confidence interval	H ² (%)	95% confidence interval	H ² (%)	95% confidence interval	H ² (%)	95% confidence interval
Left CA1	73**	55-83	56**	33-72	39**	21-55	39	21-55
Right CA1	70**	50-83	54**	29-71	59*	42-72	59*	42-72
Left CA2CA3	38*	18-58	41**	21-59	53*	35-67	52*	34-67
Right CA2CA3	70**	51-82	68**	51-80	67*	52-78	66*	50-78
Left CA4DG	77**	58-87	65**	41-81	57**	38-73	57**	37-73
Right CA4DG	80**	65-88	72**	54-83	65**	50-76	65**	50-76
Left SRLM	64**	42-78	53**	30-70	47*	30-62	40*	22-56
Right SRLM	74**	56-84	64**	43-78	59**	42-72	55**	38-69
Left Subiculum	50*	16-77	63**	26-75	60*	20-72	57**	22-71
Right Subiculum	85**	68-90	76**	62-84	82**	71-88	78**	65-86
Left Hippocampus	80**	66-88	65**	43-80				
Right Hippocampus	81**	66-89	63**	41-78				
Total brain	92**	75-95						

TABLE V. Bivariate genetic correlation (r_g) between hippocampal subfield volumes with ipsilateral hippocampal volume and total brain volume (TBV). All subfield volumes showed significant correlation between ipsilateral hippocampal volume and TBV ($p < 0.001$)

	Left Hippocampal Volume	TBV
Left CA1	0.98	0.77
Left CA2CA3	0.75	0.49
Left CA4DG	0.94	0.69
Left SRLM	0.98	0.70
Left Subiculum	0.76	0.71
Left Hippocampus		0.77
	Right Hippocampal Volume	TBV
Right CA1	0.97	0.77
Right CA2CA3	0.79	0.55
Right CA4DG	0.93	0.69
Right SRLM	0.97	0.68
Right Subiculum	0.66	0.67
Right Hippocampus		0.78

TABLE VI. Bivariate heritability estimates between TBV and sub-regions (left and right hippocampal subfield volumes and whole hippocampal volumes) with 95% confidence intervals (sample size N = 465). All values were significant at $p < 0.001$, when ACE model is compared to CE model.

Region	TBV		Sub-region		Shared (TBV and Sub-region)	
	H ² (%)	95% confidence interval	H ² (%)	95% confidence interval	H ² (%)	95% confidence interval
Left CA1	95	87-97	77	62-86	96	84-100
Right CA1	96	86-97	76	62-86	97	87-100
Left CA2CA3	95	84-97	52	24-68	99	80-100
Right CA2CA3	95	85-97	75	56-84	95	82-100
Left CA4DG	95	87-97	79	63-88	95	83-100
Right CA4DG	96	86-97	85	73-91	97	87-100
Left SRLM	95	86-97	70	53-81	95	81-100
Right SRLM	96	86-97	79	65-87	98	87-100
Left Subiculum	93	82-97	62	29-82	86	63-100
Right Subiculum	95	85-97	88	75-92	99	86-100
Left Hippocampus	95	87-97	83	70-90	96	85-99
Right Hippocampus	96	86-97	84	73-91	98	89-100

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

Table VII: Bivariate heritability estimates between ipsilateral hippocampal volume and sub-regions (left and right hippocampal subfield volumes) with 95% confidence intervals (sample size N = 465). All values were significant at p<0.001, when ACE model is compared to CE model.

Region	Ipsilateral Hippocampal Volume		Sub-region		Shared (Ipsilateral Hippocampal Volume and Sub-region)	
	H ² (%)	95% confidence interval	H ² (%)	95% confidence interval	H ² (%)	95% confidence interval
Left CA1	86	76-92	82	69-89	86	75-92
Right CA1	85	73-91	78	62-87	83	69-90
Left CA2CA3	80	64-89	43	9-67	77	45-98
Right CA2CA3	86	74-92	76	53-85	84	67-93
Left CA4DG	87	76-92	83	70-90	87	76-93
Right CA4DG	87	77-92	87	75-92	89	79-94
Left SRLM	85	75-91	75	62-84	82	70-89
Right SRLM	87	76-92	81	69-88	85	74-91
Left Subiculum	81	67-89	63	32-82	93	67-100
Right Subiculum	84	72-91	88	75-92	94	80-100