

A CONSTRUCTIVE CHARACTERISATION OF CIRCUITS IN THE SIMPLE $(2, 1)$ -SPARSE MATROID

T.A. MCCOURT AND A. NIXON

ABSTRACT. A simple graph $G = (V, E)$ is a $(2, 1)$ -circuit if $|E| = 2|V|$ and $|E(H)| \leq 2|V(H)| - 1$ for every proper subgraph H of G . Motivated, in part, by ongoing work to understand unique realisations of graphs on surfaces, we derive a constructive characterisation of $(2, 1)$ -circuits. The characterisation uses the well known 1-extension and X -replacement operations as well as several summation moves to glue together $(2, 1)$ -circuits over small cutsets.

1. INTRODUCTION

For a finite (multi)graph $G = (V, E)$ let $i_G(X)$ denote the number of edges induced by $X \subset V$. The graph G is (k, ℓ) -tight if $|E| = k|V| - \ell$ and $i_G(X) \leq k|V| - \ell$ for all $X \subset V$. For $\ell < 2k$ the (edge sets of) (k, ℓ) -tight graphs form the bases of a matroid [10, 18]. These matroids are examples of count matroids (see [4]); we refer to them as the (k, ℓ) -sparse matroid, and when loops and multiple edges are prohibited, as the simple (k, ℓ) -sparse matroid.

A simple (respectively multi-) graph G is a (k, ℓ) -circuit if G is the graph induced by a circuit in the simple (k, ℓ) -sparse matroid (respectively the (k, ℓ) -sparse matroid). Equivalently a graph G is a (k, ℓ) -circuit if $|E| = k|V| - \ell + 1$ and $i_G(X) \leq k|V| - \ell$ for all $X \subsetneq V$.

A constructive characterisation of a class of graphs is a method of building all graphs in the class from certain base graphs by elementary operations. Constructive characterisations of classes of graphs (or other objects) from small base cases by elementary local transformations are natural and relevant to a variety of problems. In particular in the field of combinatorial optimization [9]. As particular motivation for us we seek a means to understand when a generic realisation of a graph on a surface is globally rigid (unique up to isometries), see [7, 8] for details on the geometry of this problem. A related construction for circuits in the $(2, 3)$ -sparse matroid [1] was a vital aspect of the characterisation of global rigidity in the plane [6] and we expect that the construction here, along with the construction for circuits in the simple $(2, 2)$ -sparse matroid, will be crucial in establishing analogues for global rigidity on surfaces supporting either two (e.g. the cylinder) or one (e.g. the cone or torus) tangentially acting isometries. See [11, Conjecture 5.7] and [7, Conjecture 9.1].

Constructive characterisations of (k, ℓ) -tight (multi)graphs are known, for example when $k = \ell$ [17] and more generally when $l \leq k$ [3]. For simple graphs the requirement that each intermediate graph be simple means that new constructions are required. When $k = 2$ constructions are known for several classes [12, 13, 14].

For (k, ℓ) -circuits less is known, however some general constructions, such as when $k = \ell$ can be extracted from work on tree packings [5]. In this paper we are interested in simple (k, ℓ) -circuits. For $k = 1$ the problem is easy. When $k = 2$, and $\ell = 3$, there is an elegant result of Berg and Jordán [1]. It is easy to check that the minimum degree, $\delta(G)$, in a




FIGURE 1. Base graphs on six vertices.

$(2, 3)$ -circuit G is equal to three. They proved first that every $(2, 3)$ -circuit which was 3-connected contains an *admissible* degree three vertex, that is a degree three vertex which can be removed and an edge added between its neighbours in such a way that the resulting graph is a $(2, 3)$ -circuit. To complete their constructive characterisation they observed that whenever G was not 3-connected there is a 2-separation $\{x, y\}$ and $xy \notin E$. Such a graph can be broken into two smaller $(2, 3)$ -circuits by separating over this cut and adding the edge xy to both parts. (This is the inverse of the well known 2-sum operation.)

For $(2, 2)$ -circuits one must distinguish between the multigraph and simple graph cases. For multigraphs a constructive characterisation occurs as a special case of a characterisation of highly k -tree connected graphs [5]. For simple graphs there is a constructive characterisation in [11]. It is a nontrivial extension of Berg and Jordán's result in the following senses: firstly since K_4 can occur as a subgraph of any $(2, 2)$ -circuit, it is non-trivial to preserve simplicity; secondly the connectivity level required to guarantee admissibility is higher (requiring 3-connectivity and essential 4-edge-connectivity); and thirdly when these connectivity assumptions do not hold there are several separation moves required.

In this paper we consider simple $(2, 1)$ -circuits and in particular, we prove a constructive characterisation of such circuits where every intermediate graph is also simple. At a high level the scheme is analogous to the above cases: we establish the connectivity level required to guarantee an admissible vertex and when this fails we define separation moves to pull apart $(2, 1)$ -circuits into smaller $(2, 1)$ -circuits. However, in the $(2, 1)$ -circuit case there are numerous complications to previous cases. Firstly observe that $(2, 1)$ -circuits need not have $\delta(G) = 3$, thus we are forced to consider a degree four operation. Secondly since K_4 is not a base in the simple $(2, 1)$ -sparse matroid we separate out two cases in our result guaranteeing an admissible vertex of degree three. Thirdly, due to the connectivity level required to guarantee an admissible node, there are six different separations that can occur.

1.1. Terminology. We now define the set \mathcal{G} of base graphs for our constructive characterisation. Note first that K_5 is the unique $(2, 1)$ -circuit on at most five vertices. There are five distinct graphs formed from deleting three edges from K_6 . Deleting a one factor leaves a 4-regular graph which has an admissible vertex by Theorem 1.2. Deleting a degree three star does not give a $(2, 1)$ -circuit. The other three possibilities are the complements of $G57$, $G59$ and $G60$ as listed in [16] (see also Figure 1).

There are 65 non-isomorphic graphs formed from deleting seven edges from K_7 ; of these 34 are $(2, 1)$ -circuits. Two of these are 4-regular and hence contain an admissible node by Theorem 1.2. Of the remainder note that if there is a node not in a copy of K_4 then either adding a missing edge creates a K_5 subgraph or that node is admissible. This observation allows us to easily spot that 29 of the remaining $(2, 1)$ -circuits contain an admissible node. The remaining three do not contain admissible nodes; they are the complements of $G293$, $G308$ and $G312$ as listed in [16] (see also Figure 2).




FIGURE 2. Base graphs on seven vertices.




FIGURE 3. Base graphs on eight vertices.




FIGURE 4. Base graphs on nine vertices.

A further five base graphs of orders eight and nine arise in the proof of the recursive construction; there are four on eight vertices (graphs S_1 , S_2 , S_3 and S_4 in Figure 3) and there is one on nine vertices (graph S_5 in Figure 4).

Hence we define \mathcal{G} to be the set $\{K_5, \overline{G57}, \overline{G59}, \overline{G60}, \overline{G293}, \overline{G308}, \overline{G312}, S_1, S_2, S_3, S_4, S_5\}$.

For a graph G let $\delta(G)$ denote the minimum degree of a vertex in G . A k -edge-cutset is a set of k edges whose removal disconnects a graph. If one of the components, of the resulting disconnected graph, contains only a single vertex then we say that the k -edge-cutset is *trivial*. Otherwise it is *non-trivial*. A graph is *essentially k -edge-connected* if it has no non-trivial $(k - 1)$ -edge-cutsets. Since $(2, 1)$ -circuits have $\delta(G) \in \{3, 4\}$ this will be a useful notion for us.

Let $G = (V, E)$ be a graph and let G' be formed from G by deleting an edge xy from E and adding a new vertex v and edges xv, yv, zv for $x, y, z \in V$. This operation is known as a 1-extension (elsewhere it is also referred to as the Henneberg 2 move [15, 17, 18]). The inverse is known as a 1-reduction. Also let G'' be formed from G by deleting two non-adjacent edges xy, zw from E and adding a new vertex v and edges xv, yv, zv, wv for $x, y, z, w \in V$. This operation is known as X -replacement [15, 18].

As outlined above, when a circuit is sufficiently connected we will show that either a 1-reduction or an inverse X -replacement can be applied to form a new circuit. In the cases where a circuit is not sufficiently connected we will apply inverse ‘summation’ moves to reduce to a smaller circuit.

There will remain a handful of cases where the circuit, G say, is invariant under these inverse ‘summation’ moves. In order to deal with these cases we will construct a new circuit G^* which is a, sufficiently connected, multigraph. We will then show that there exists a 1-reduction or inverse X -replacement that can be applied to G^* to yield a new circuit. Finally we use this to show that an analogous 1-reduction or X -replacement could have been applied in G to yield a new circuit.

Let $M^*(2, 1)$ denote the set of all multigraphs that are $(2, 1)$ -circuits and let $M(2, 1)$ denote the set of all simple graphs that are $(2, 1)$ -circuits. The multigraph circuits that we construct in the manner referred to above are all elements of the subset $\mathcal{M} \subseteq M^*(2, 1)$ defined as follows.

Let \mathcal{M}' be the subset of $M^*(2, 1)$ where $G \in \mathcal{M}'$ if and only if

- G is 3-connected;
- the maximum edge multiplicity is three;
- all the vertices in G incident with a multiple edge have degree greater than three;
- if G contains a loop, then the vertex incident with the loop has degree greater than three and it is not incident with any multiple edges;
- if $G = (V, E)$ contains a vertex, v say, incident to two loops then $V = \{v\}$; and
- if there are any triple edges in G , then they are incident with vertices of degree greater than or equal to five, and hence, G must contain vertices of degree three.

Then $\mathcal{M} = M(2, 1) \cup \mathcal{M}'$.

Note that $M(2, 1) \subset \mathcal{M} \subset M^*(2, 1)$. With this in mind, letting $G \in \mathcal{M}$, we call

- a vertex of degree three in G a *node*;
- a node v of G *admissible* if there is a 1-reduction removing v that results in a $(2, 1)$ -circuit in which the added edge is between previously non-adjacent vertices;
- a degree four vertex v in G *admissible* if there is an inverse X -replacement removing v that results in a $(2, 1)$ -circuit in which the added edges are between pairs of previously non-adjacent vertices.

As $M(2, 1) \subset \mathcal{M}$, in the case where $G \in M(2, 1)$ the resulting circuit is also in $M(2, 1)$. (The reader may find the motivation behind the results in Section 3 clearer if these results are thought of in the restricted case of $M(2, 1)$; we will not need the general case until the end of the proof of Theorem 1.1.) We will use \mathcal{G}^* to refer to the set of base graphs in \mathcal{M} . The elements of $\mathcal{G}^* \setminus M(2, 1)$ will be derived in Section 3.

1.2. Results. Our main result is as follows.

Theorem 1.1. *A simple graph G is a $(2, 1)$ -circuit if and only if it can be generated recursively from $G \in \mathcal{G}$ by applying 1-extensions and X -replacements sequentially within connected components and taking sums of connected components.*

In order to prove the theorem we first establish admissibility when the minimum degree is 4.

Theorem 1.2. *Every essentially 5-edge-connected $(2, 1)$ -circuit G not equal to K_5 with $\delta(G) = 4$ contains an admissible vertex.*

From there, the key technical step to proving Theorem 1.1 is to establish admissibility, in the $\delta(G) = 3$ case, when G is sufficiently connected.

Theorem 1.3. *Let $G = (V, E)$ be a 3-connected $(2, 1)$ -circuit with $\delta(G) = 3$ and $G \notin \mathcal{G}$. Suppose either*

- (i) *G is essentially 5-edge-connected and there is no proper critical set or*
- (ii) *G is essentially 4-edge-connected and there is a proper critical set.*

Then G contains an admissible node.

Proper critical sets are defined in Section 3 and the reason for splitting the above theorem into these two cases is purely technical.

1.3. Outline of paper. In Section 3 we prove Theorem 1.3 establishing that whenever a $(2, 1)$ -circuit has $\delta(G) = 4$ then either it has an admissible vertex or it has a small separating set of edges. In Section 2 we prove Theorem 1.2 establishing the analogue for the case when $\delta(G) = 3$. In Section 4 we deal with graphs that are $(2, 1)$ -circuits but fail to be sufficiently connected. In Section 5 we combine our results to prove Theorem 1.1.

2. X-REPLACEMENT

An *X-replacement* is the deletion of two non-adjacent edges ab, cd and the addition of a vertex v adjacent to a, b, c, d . The inverse operation is the deletion of a degree four vertex and the addition of two non-adjacent edges between the neighbours. For $G = (V, E)$, define $f(G) = 2|V| - |E|$, so that a graph G is a $(2, 1)$ -circuit if and only if $f(G) = 0$ and every proper subgraph H satisfies $f(H) \geq 1$. If G is a $(2, 1)$ -circuit and $v \in V$, then we say that v is *admissible* if there is an inverse X-replacement on v resulting in a $(2, 1)$ -circuit.

Lemma 2.1. *A graph G is a $(2, 1)$ -circuit with $\delta(G) = 4$ if and only if G is connected and 4-regular.*

Proof. Let $G = (V, E)$. If G is a $(2, 1)$ -circuit with $\delta(G) = 4$, then G is 4-regular and a simple counting argument implies that G is connected. Conversely, if G is connected and 4-regular then $|E| = 2|V|$. Suppose $H = (V', E')$ is a proper subgraph of G with $|E'| \geq 2|V'|$ then H has average degree at least four. Since H is proper and G is connected we contradict the 4-regularity of G . \square

With this lemma, the following theorem can be extracted from [2]. For completeness we provide an alternate proof.

Theorem 1.3. *Every essentially 5-edge-connected $(2, 1)$ -circuit $G \in M(2, 1)$ not equal to K_5 with $\delta(G) = 4$ contains an admissible vertex.*

Proof. Let $G = (V, E)$.

Claim 2.2. *Let $v \in V$. Then v is non-admissible if and only if every possible inverse X-replacement on v results in a non-simple graph.*

Proof of Claim. If every inverse X-replacement on v creates a multigraph, then clearly v is non-admissible. For the converse let G' be the result of the inverse X-replacement. By Lemma 2.1 we need to check that 4-regularity and connectedness are preserved. The first of these is clear and if G' is not connected then v is a cut-vertex, but since v has degree four this contradicts the assumption that G is essentially 5-edge-connected. \square

It follows that G' is not simple if and only if in G , v is (a) contained in a copy of K_4 or (b) there is $x \in N(v)$ (the neighbour set of v) adjacent to every other vertex in $N(v)$.

We next show that no subgraph of G is isomorphic to K_4 . Suppose otherwise, i.e., that H is a subgraph of G isomorphic to K_4 . As G is 4-regular there is a 4-edge-cutset between H and $G - H$, since $G \neq K_5$ this cutset is non-trivial.

We complete the proof by showing that either u or one of its neighbours are admissible. (In fact, either u or three of its neighbours are admissible.) Let $N(u) = w, x, y, z$ and note that u is not in a copy of K_4 . Hence if u is not admissible then, say, $xy, xz, xw \in E$. Let $N(y) = \{u, x, r, s\}$ and note that, since u is not in a subgraph isomorphic to K_4 , $r, s \in V - \{u, x, y, z, w\}$. By 4-regularity ur, xs are not in E . Thus Claim 2.2 implies y is admissible. \square

3. THE 1-EXTENSION OPERATION

In this section we prove Theorem 1.3. We start with some facts (Lemmas 3.1 to 3.6) about $(2, 1)$ -circuits that can be proved by simple counting arguments. See [1] or [11] for similar results. Crucial to the problem is that we must retain simplicity throughout the recursive construction. We achieve this, in Subsection 3.2, by establishing conditions for a $(2, 1)$ -circuit with $\delta(G) = 3$ to have nodes not contained in subgraphs isomorphic to K_4 . Finally in Subsection 3.3 we loosely follow the method established in [1] to deduce admissibility for some node.

3.1. Preliminaries. Let $G = (V, E)$ be a $(2, 1)$ -circuit. A subset $X \subset V$ is *critical* if $i(X) = 2|X| - 1$ and is *semi-critical* if $i(X) = 2|X| - 2$ and for all $X' \subset X$ we have $i(X') \leq 2|X'| - 2$. We are particularly interested in special kinds of critical sets. We define a critical set $X \subset V$ to be: *v-critical* for a node $v \in V$ if X contains exactly two neighbours of v but not v itself; *node-critical* if X is v -critical for some node v (where $N(v) = \{x, y, z\}$) in V such that $x, y \in X$ and $d(z) \geq 4$; and *proper* if $|X| < |V| - 1$. Note that node-critical sets are proper, but proper critical sets need not be node-critical. Suppose that $A, B \subseteq V$; then we denote the number of edges $uv \in E$ such that $u \in A - B$ and $v \in B - A$ by $d(A, B)$.

Lemma 3.1. *Let $G = (V, E)$ be a $(2, 1)$ -circuit. Then G is connected and 2-edge-connected.*

Lemma 3.2. *Let $G = (V, E)$ be a $(2, 1)$ -circuit. Let $X, Y \subset V$ be critical sets, let $|X \cap Y| \geq 1$ and let $|X \cup Y| \leq |V| - 1$. Then $X \cap Y$ and $X \cup Y$ are critical sets and $d(X, Y) = 0$.*

Lemma 3.3. *Let $G = (V, E)$ be a $(2, 1)$ -circuit. Let v be a node with $N(v) = \{u, w, z\}$. Then removing v and adding uw is not admissible if and only if there exists a critical set $X \subset V$ with $u, w \in X$ and $v, z \notin X$ or $uw \in E$.*

Lemma 3.4. *Let $G = (V, E)$ be a 3-connected $(2, 1)$ -circuit containing a node v with $N(v) = \{w, u, z\}$, $uz \notin E, wz, wu \in E$. Then v is admissible.*

Lemma 3.5. *Let $G = (V, E)$ be a $(2, 1)$ -circuit containing a node v with $N(v) = \{w, u, z\}$, $uz, wu \notin E, wz \in E$. Then v is admissible.*

Let $V_3 = \{v \in V : d(v) = 3\}$. Let $V_3^* \subset V_3$ be the subset of degree three vertices not contained in copies of K_4 .

Lemma 3.6. *Let G be a $(2, 1)$ -circuit. Then $G[V_3]$ is a (possibly empty) forest.*

3.2. Finding Nodes. We now establish connectivity conditions that guarantee that, in a 3-connected, essentially 5-edge-connected (2,1)-circuit G with $\delta(G) = 3$, there exist nodes not contained in copies of K_4 . Since (the vertex set of) K_4 is semi-critical, this is significantly harder than the corresponding result in [11]. We will first prove two lemmas for (2,1)-circuits that contain proper critical sets, before proving their (simpler) analogues for (2,1)-circuits that contain no proper critical sets. We make this distinction as we can give a weaker edge-connectivity assumption for (2,1)-circuits that contain proper critical sets, than for (2,1)-circuits that do not (see Theorem 1.3).

3.2.1. Proper critical sets. We first consider the case when there exists at least one proper critical set.

Lemma 3.7. *Let $G \in \mathcal{M}$ be essentially 4-edge-connected. Suppose that G contains a proper critical set X . Then $|V_3 - X| \geq 2$ (and hence $\delta(G) = 3$).*

Proof. Let E' be the set of edges between X and $V - X$. The sum of the degrees of vertices in X is $4|X| - 2 + |E'|$. The total sum of the degrees of vertices in G is $4|V|$. So the sum of the degrees in $V - X$ is $4(|V| - |X|) + 2 - |E'|$. Hence, as $\delta(G) \geq 3$ and the average degree of G is four, there are at least two nodes that are not in X . \square

Lemma 3.8. *Let $G \in \mathcal{M}$ be 3-connected and essentially 4-edge-connected. Let X be a critical set and let $\mathcal{W} = \{W_i \mid 1 \leq i \leq n : W_i \text{ is critical or semi-critical}\}$, where $X \notin \mathcal{W}$. Let $Y = V - (\cup_{i=1}^n W_i) - X$. Suppose that either*

- (1) $|Y| \geq 2$; or
- (2) $\mathcal{A} := G[X] \cup (\cup_{i=1}^n G[W_i])$ is disconnected.

Then Y contains two nodes.

Proof. Since $G \in \mathcal{M}$, with vertices of V labelled as $v_1, v_2, \dots, v_{|V|}$, we have

$$(3.1) \quad \sum_{i=1}^{|V|} (4 - d_G(v_i)) = 0.$$

Let A_1, \dots, A_m be the vertex sets of the connected components of \mathcal{A} . For any pair of semi-critical sets $W_i, W_j \in \mathcal{W}$ we have

$$(3.2) \quad \begin{aligned} 2(|W_i| + |W_j|) - 3 &= 2|W_i \cup W_j| - 1 + 2|W_i \cap W_j| - 2 \\ &\geq i(W_i \cup W_j) + i(W_i \cap W_j) \\ &= i(W_i) + i(W_j) + d(W_i, W_j) \\ &= 2(|W_i| + |W_j|) - 4 + d(W_i, W_j) \end{aligned}$$

so either $d(W_i, W_j) = 0$ or equality holds and $d(W_i, W_j) = 1$. Hence, whenever W_i and W_j have non-empty intersection, the union is semi-critical or critical. Similarly if W_i is critical, W_j is semi-critical and they have a non-empty intersection then the union is critical. (Note two critical sets with a non-empty intersection gives V , contradicting (1) and (2).)

Reordering if necessary, let A_1, \dots, A_r be critical sets and let A_{r+1}, \dots, A_m be semi-critical sets. Denote the vertices of A_j by $u_1, \dots, u_{|A_j|}$ and define

$$f_j = \sum_{i=1}^{|A_j|} (4 - d_{G[A_j]}(u_i)).$$

Hence, for $1 \leq j \leq r$, $f_j = 2$ and for $r + 1 \leq j \leq m$, $f_j = 4$. As either (1) or (2) holds and there are no non-trivial 3-edge-cutsets, there exist four edges $x_\ell y_\ell$, $1 \leq \ell \leq 4$, where $x_\ell \in A_j$

and $y_\ell \in V - A_j$, where the $x_\ell \in A_j$ (respectively the $y_\ell \in V - A_j$) are not necessarily all distinct. Define

$$g_j = \sum_{i=1}^{|A_j|} (4 - d_G(u_i)).$$

Hence, for $1 \leq j \leq r$, $g_j \leq -2$ and, for $r + 1 \leq j \leq m$, $g_j \leq 0$. Now

$$(3.3) \quad \sum_{i=1}^m g_j = -2r \leq -2$$

since $r \geq 1$. The result follows by comparing Equations (3.1) and (3.3). \square

Lemma 3.9. *Let $G = (V, E) \in \mathcal{M}$ be 3-connected and essentially 4-edge-connected. Suppose that G contains a proper critical set X . Then $|V_3^* - X| \geq 2$.*

Proof. If G contains no copies of K_4 , then $V_3^* = V_3$ and Lemma 3.7 yields the result.

So suppose that B_1, \dots, B_n are the vertex sets of all the copies of K_4 which are not subsets of X . Let $Y = V - (\cup_{i=1}^n B_i) - X$ and $\mathcal{A} = G[X] \cup (\cup_{i=1}^n G[B_i])$. If $|Y| \geq 2$ or \mathcal{A} is disconnected, then we may apply Lemma 3.8 to yield the result.

Hence, suppose that $|Y| \leq 1$ and \mathcal{A} is connected. So, without loss of generality, there exists a $t \geq 1$, such that $X \cap B_i \neq \emptyset$, for all $1 \leq i \leq t$. For each of these B_i , and a subset $B' \subset B_i$, we have $i(B') = 2|B'| - \alpha$ where $\alpha = 2$ when $|B'| \in \{1, 4\}$ and $\alpha = 3$ when $|B'| \in \{2, 3\}$. Thus, using a similar calculation to Equation (3.2) we have: if $|X \cap B_i| = 3$, then $X \cup B_i = V$, contradicting the fact that X is a proper critical set, and if $|X \cap B_i| = 2$, then $X \cup B_i = V$, contradicting 3-connectivity. So we have that $|X \cap B_i| = 1$. If $d(X, B_i) > 0$ or the multiplicity of any of the edges in $G[B_i]$ is greater than one, then $X \cup B_i = V$, contradicting 3-connectivity. So $d(X, B_i) = 0$ and $G[B_i]$ is simple, for $1 \leq i \leq t$.

We define an auxiliary graph G^\dagger as follows. The vertex set is X, B_1, \dots, B_n and there is an edge between two vertices if and only if the corresponding sets in G intersect.

Claim 3.10. *The graph G^\dagger is a tree.*

Proof. First note that a path in G^\dagger either corresponds to a critical or a semi-critical set in G .

Suppose that G^\dagger contains a cycle. Let $C = (c_0, c_1, \dots, c_\ell)$ be such a cycle. Without loss of generality c_0 corresponds to one of the B_i , so let P be the path obtained by deleting c_0 from C .

Suppose P corresponds to a critical set $P' = \cup_{i=1}^\ell c_i$ in G , that is $i(P') = 2|P'| - 1$. Now let $C' = \cup_{i=0}^\ell c_i$, note that C' contains, at most, an additional two vertices from P' and $G[C']$ contains an additional six edges from $G[P']$, a contradiction.

Suppose P corresponds to a semi-critical set $P' = \cup_{i=1}^\ell c_i$ in G , that is $i(P') = 2|P'| - 2$. Then X is not a vertex of P . Now let $C' = \cup_{i=0}^\ell c_i$, note that, as P' is not critical, $G[C']$ contains an additional two vertices and an additional six edges from $G[P']$. Hence $C' = V$, but does not contain X , a contradiction.

The claim follows from the connectivity of \mathcal{A} . \square

As B_1 exists it follows from the claim that \mathcal{A} contains a cut-vertex, v . As \mathcal{A} is connected the vertex set of \mathcal{A} either corresponds to a critical set in G or it is V . In the first case v is part of a cut-pair in G (as $|Y| \leq 1$) and in the second case v is a cut-vertex in G . In either case we have contradicted the 3-connectivity of G . \square

3.2.2. *No proper critical sets.* Now we deal with the case when there are no proper critical sets.

Lemma 3.11. *Let $G = (V, E) \in \mathcal{M}$ be essentially 5-edge-connected. Suppose V contains no proper critical sets, let W_1, \dots, W_k , with $k \geq 1$, be the vertex sets of all the copies of K_4 in G and let $Y = V - \cup_{i=1}^k W_i$. Suppose that either*

- (1) $|Y| \geq 2$; or
- (2) $\cup_{i=1}^k G[W_i]$ is disconnected.

Then G contains an admissible node.

Proof. If any of the W_i are critical sets (i.e., $G[W_i]$ contains multi-edges or loops), then, as V contains no proper critical sets, $|V| = 5$ and neither (1) nor (2) hold. So, since G is a (2,1)-circuit, Equation (3.1) holds and W_i is semi-critical, for each $1 \leq i \leq k$. Let A_1, \dots, A_r be the vertex sets of the connected components of $\cup_{i=1}^k G[W_i]$. Observe that each of A_1, \dots, A_m is semi-critical.

Denote the vertices of A_j by $u_1, \dots, u_{|A_j|}$. Then

$$4 = \sum_{i=1}^{|A_j|} (4 - d_{G[A_j]}(u_i)).$$

There are no non-trivial 4-edge-cutsets so, as at least one of (1) or (2) holds, there exists five edges $x_\ell y_\ell$, $1 \leq \ell \leq 5$, where $x_\ell \in A_j$ and $y_\ell \in V - A_j$ (where the x_ℓ are not necessarily distinct and nor are the y_ℓ). Then

$$(3.4) \quad \sum_{j=1}^r \sum_{i=1}^{|A_j|} (4 - d_G(u_i)) \leq -1.$$

Comparing Equations (3.1) and (3.4) gives that $|Y|$ contains a node v . Now v is not in a copy of K_4 and since there are no proper critical sets, we can apply Lemma 3.3 to deduce that v is admissible. \square

3.3. Admissibility.


Lemma 3.12. *Let $G = (V, E)$ be a (2,1)-circuit. Suppose $v \in V_3^*$ is a non-admissible node with $N(v) = \{x, y, z\}$. Then there exists two v -critical sets X, Y such that $X \cup Y = V - v$. Moreover, we may choose X, Y such that $z \in X \cap Y$.*

Proof. By Lemmas 3.4 and 3.5, since v is not admissible and $v \in V_3^*$, there are no edges between x, y and z . Now 3.3 implies there exists critical sets X containing x, z but not y, v and Y containing y, z but not x, v . It follows from Lemma 3.2 and the definition of a (2,1)-circuit that $X \cup Y$ is critical, hence $X \cup Y = V - v$. \square

The following has essentially the same proof as the corresponding results for (2,3) and (2,2)-circuits, [1, Lemma 3.3] and [11, Lemma 2.10]; however we obtain a slightly stronger result.

Lemma 3.13. *Let $G = (V, E)$ be a (2,1)-circuit with $\delta(G) = 3$. Let $v \in V$ be a node with $N(v) = \{x, y, z\}$, $d(z) \geq 4$ and suppose there are no edges between neighbours of v . Let X be v -critical on x, y . Suppose either*

- (1) *there is a non-admissible node $u \in V - X - v$ where $d_{G[V_3^*]}(u) = 2$ with no edges between its neighbours, precisely one neighbour w in X and w is a node; or*

FIGURE 5. Circuits in $\mathcal{G}^* \cap (\mathcal{M} \setminus M(2,1))$.

(2) there is a non-admissible node $u \in V - X - v$ where $d_{G[V_3^*]}(u) = 0$ or 1 with no edges between its neighbours.

Then, in G , there is a node-critical set X' for a node with no edges between its neighbours such that $X \subsetneq X'$.

Proof. Suppose (1) holds and let $N(u) = \{w, p, q\}$. As $d_{G[V_3^*]}(u) = 2$ we may assume that $d_G(p) = 3$ and $d_G(q) \geq 4$. Since u is non-admissible and $wp \notin E$, by Lemma 3.3, there exists a u -critical set, Y say, on w and p . Now, $w \in X \cup Y$, and $u, q \notin X \cup Y$, so by Lemma 3.2, $X' := X \cup Y$ is node-critical for u . As $p \notin X$, $X \subsetneq X'$.

So, suppose that (2) holds. Let r be a neighbour of u of minimum degree in G , by Lemma 3.12, there exist u -critical sets Y_1 and Y_2 whose intersection contains r and $Y_1 \cup Y_2 = V - u$. Hence, without loss of generality, $X \cap Y_1 \neq \emptyset$. As $|N(u) \cap X| \leq 3$, we consider four cases. If $|N(u) \cap X| = 0$, then the set $X' = X \cup Y_1$ is node-critical and, as $|N(u) \cap X| = 0$, we have that $X \subsetneq X'$.

Suppose that $|N(u) \cap X| = 1$, say $N(u) \cap X = \{s\}$. If $s \in Y_1$, then set $X' = X \cup Y_1$ is node-critical and, as $N(u) - (X \cup Y_1) \neq \emptyset$, we have that $X \subsetneq X'$. So, assume that $s \notin Y_1$, that is $s \in Y_2 - Y_1$; hence $X' = X \cup Y_2$ is node-critical and, as $N(u) - (X \cup Y_2) \neq \emptyset$, we have that $X \subsetneq X'$.

If $|N(u) \cap X| = 2$, then the set $X' = X \cup \{u\}$ is node-critical and hence $X \subsetneq X'$. Finally, if $|N(u) \cap X| = 3$, then $V = X \cup \{u\}$, but $v \notin X \cup \{u\}$, a contradiction. \square

Before we prove Theorem 1.3 we establish the following lemma to deal with some small circuits.

Lemma 3.14. *Let $G = (V, E) \in \mathcal{M} \setminus \mathcal{G}^*$ be 3-connected and essentially 5-edge-connected with $|V| \leq 8$, $\delta(G) = 3$ and no proper critical sets.*




FIGURE 6. Potential base graphs on seven vertices.




FIGURE 7. Potential base graphs on eight vertices.

Suppose that G contains a copy of K_4 . Let W_1, \dots, W_k , with $k \geq 1$, be the vertex sets of all copies of K_4 in G and let $Y = V - \cup_{i=1}^k W_i$. Further suppose that $\mathcal{A} = \cup_{i=1}^k G[W_i]$ is connected and $|Y| \leq 1$. Then G contains an admissible node.

Proof. First note that, as G contains a copy of K_4 , $|V| \geq 4$.

Suppose that $|V| = 4$. In this case $V = W_1$ and as G is a circuit, $G \in \mathcal{M}'$ (so G is a multigraph). There are five possibilities for a circuit on four vertices: R_4, R_5, R_6, R_7, R_8 (see Figure 5) all of which are contained in \mathcal{G}^* .

Suppose that $|V| = 5$. Either there are two vertex sets underlying copies of K_4 or there is one set and an additional vertex of degree three. Note that these cases are equivalent, hence there are five possibilities for a such a circuit on five vertices: $K_5, R_9, R_{10}, R_{11}, R_{12}$ (see Figure 5) all of which are contained in \mathcal{G}^* .

Suppose that $|V| = 6$. As $|Y| \leq 1$, $k \geq 2$ and any pair of distinct W_i, W_j intersect in either two or three vertices. Consider W_1 and W_2 . If $|W_1 \cap W_2| = 2$, then $V = W_1 \cup W_2$, and as G is 3-connected it must be isomorphic to $\overline{G59} \in \mathcal{G}$. So suppose that $|W_1 \cap W_2| = 3$. Then there exists a vertex $v \notin W_1 \cup W_2$. As $W_1 \cup W_2$ is a critical set v has degree three and hence G is either isomorphic to $\overline{G57}$ or $\overline{G60}$, in either case $G \in \mathcal{G}$.

Suppose that $|V| = 7$. As $|Y| \leq 1$ and \mathcal{A} is connected, $k \geq 2$ and any two W_i, W_j intersect in either one, two or three vertices. However, as G does not contain a proper critical set $|W_i \cap W_j| \neq 3$. If $|W_1 \cap W_2| = 1$, then as G is 3-connected it must be isomorphic to either $\overline{G308}$ or $\overline{G312}$. In either case $G \in \mathcal{G}$. So suppose that $|W_1 \cap W_2| = 2$. Then there exists a vertex $v \notin W_1 \cup W_2$. As $W_1 \cup W_2$ is a critical set v has degree three and as G is 3-connected it is isomorphic to one of the two graphs illustrated in Figure 6. In either case Lemma 3.5 or 3.4 implies the existence of an admissible node.

Finally, suppose that $|V| = 8$. As \mathcal{A} is connected and $|Y| \leq 1$, $k \geq 2$ and for any pair W_i, W_j , where $1 \leq i < j \leq k$, $|W_i \cap W_j| = 0$ or 1 and there exists a pair, say W_1, W_2 , where $|W_1 \cap W_2| = 1$. Then there exists a vertex $v \notin W_1 \cup W_2$ and as G is 3-connected in this case it must be simple and isomorphic to one of the three graphs illustrated in Figure 7; in each case Lemma 3.5 or 3.4 implies the existence of an admissible node. \square

We can now prove Theorem 1.3.

Theorem 1.2. *Let $G = (V, E) \in \mathcal{M}$ be 3-connected with $\delta(G) = 3$ and $G \notin \mathcal{G}^*$. Suppose either*

- (i) *G is essentially 5-edge-connected and there is no proper critical set or*
- (ii) *G is essentially 4-edge-connected and there is a proper critical set.*

Then G contains an admissible node.

Proof. Suppose that (i) holds. There exists a vertex v of degree three. If v is not contained in a copy of K_4 , then, as there are no proper critical sets, we can apply Lemma 3.3 to show that v is admissible.

So we may assume that G contains a copy of K_4 . Let W_1, \dots, W_k , with $k \geq 1$, be the vertex sets of all the copies of K_4 in G and let $Y = V - \cup_{i=1}^k W_i$. By Lemma 3.11 we may assume that $\cup_{i=1}^k G[W_j]$ is connected and $|Y| \leq 1$. If G has less than 9 vertices, then, by Lemma 3.14, it is either in \mathcal{G}^* or has an admissible node.

So suppose G has at least 9 vertices. Since there are no proper critical sets, the only options for two copies of K_4 , with vertex sets W_i and W_j say, are that $W_i \cap W_j = \emptyset$ or $|W_i \cap W_j| = 1$ and $d(W_i, W_j) = 0$.

We define an auxiliary graph G^\ddagger as follows. The vertex set is W_1, \dots, W_n and there is an edge between two vertices if and only if the corresponding sets in G intersect.

Claim 3.15. *The graph G^\ddagger is either a tree or a cycle.*

Proof. We show that if G^\ddagger contains a cycle this must be the whole graph; then, as \mathcal{A} is connected, the claim follows. As $W_i \cap W_j = \emptyset$ or $|W_i \cap W_j| = 1$ and $d(W_i, W_j) = 0$ for $1 \leq i < j \leq n$ a path in G^\ddagger corresponds to a semi-critical set in G .

Suppose that G^\ddagger contains a cycle. Let $C = (c_0, c_1, \dots, c_\ell)$ and let P be the path obtained by deleting c_0 from C . Consider the set of vertices $P' = \cup_{i=1}^\ell c_i$ in G , then $i(P') = 2|P'| - 2$. Now let $C' = \cup_{i=1}^\ell c_i$ and note that $G[C']$ contains an additional two vertices and an additional six edges from $G[P']$. Hence $C' = V$. \square

If G^\ddagger is a cycle, then $|Y| = 0$ and every edge of G is contained in one of the K_4 's so G has a cut-pair, a contradiction. Hence, G^\ddagger is a tree.

As $|V| \geq 9$ and \mathcal{A} is connected, G^\ddagger has at least three vertices. As $W_i \cap W_j = \emptyset$ or $|W_i \cap W_j| = 1$ and $d(W_i, W_j) = 0$ for $1 \leq i < j \leq n$, $\cup_{i=1}^k W_i$ is a semi-critical set in G . Hence $|Y| \neq 0$, thus $|Y| = 1$. Denote the vertex in Y as v , as $\cup_{i=1}^k W_i$ is a semi-critical set, $d(v) = 4$. As G^\ddagger is a tree, G contains a cut-pair (the vertex v and one of the vertices in the intersection of a pair W_i, W_j that intersect); a contradiction.

Now, suppose that (ii) holds. Let $\mathcal{X} = \{X \cup V \mid X \text{ is a node-critical set in } G\}$. Suppose that $\mathcal{X} = \emptyset$. By Lemma 3.9, $|V_3^*| \geq 2$ and, by Lemma 3.6, $G[V_3^*]$ is a forest. Let v be a leaf vertex of $G[V_3^*]$, then v is also a leaf in $G[V_3]$. Thus v has a neighbour of degree greater than three, and as v is not contained in a copy of K_4 and $\mathcal{X} = \emptyset$, by Lemma 3.3, v must be admissible.

So we may assume that $\mathcal{X} \neq \emptyset$. By Lemma 3.9 we have $|V_3^*| \geq 2$. By Lemmas 3.5 and 3.4, we may assume that each vertex in V_3^* has no edges between the neighbours.

Choose $X \in \mathcal{X}$ to be a maximal node-critical set chosen over all nodes in V_3^* . Suppose X is node-critical for v on x, y where $N(v) = \{x, y, z\}$ and $d_G(z) \geq 4$. By Lemma 3.9, $V - X - v$ contains a node $u \in V_3^*$ and, by Lemma 3.6, we may choose u to be a leaf in $G[V_3^* - X - v]$.

By the maximality of X , each vertex $t \in V - X - v - z$ has at most one neighbour in X . Hence $d_{G[V_3^*]}(u) \leq 2$, so $d_{G[V_3]}(u) \leq 2$. If $d_{G[V_3^*]}(u) = 2$, then, since u has precisely one neighbour w in X and since u is a leaf in $G[V_3^* - X - v]$, it follows that w is a node.




FIGURE 8. Illustration of the sum move for Case (1).

Thus Lemma 3.13 Part (1) and the maximality of $|X|$ imply that u is an admissible node. If $d_{G[V_3^*]}(u) = 0$ or 1 , then Lemma 3.13 Part (2) and the maximality of $|X|$ imply that u is admissible. \square

4. SUM MOVES

In this section we reduce the connectivity assumptions from the previous sections by defining operations that ‘pull apart’ (2,1)-circuits into two smaller (2,1)-circuits.

Consider a graph $G = (V, E)$ that is either not 3-connected or has a non-trivial 4-edge-cutset. We will provide a series of lemmas that consider the cases where G satisfies one of the following:

- (1) a cut-vertex x , $A, B \subset V$ such that $A \cap B = \{x\}$ and $A \cup B = V$ such that $i(A) = 2|A| - 1$ and $i(B) = 2|B| - 1$;
- (2) a cut-pair x, y (where neither are cut-vertices), $xy \notin E$, $A, B \subset V$ such that $A \cap B = \{x, y\}$ and $A \cup B = V$ such that either
 - (a) $i(A) = 2|A| - 2$ and $i(B) = 2|B| - 2$, or
 - (b) $i(A) = 2|A| - 1$ and $i(B) = 2|B| - 3$;
- (3) a cut-pair x, y (where neither are cut-vertices), $xy \in E$, $A, B \subset V$ such that $A \cap B = \{x, y\}$ and $A \cup B = V$ such that $i(A) = 2|A| - 1$ and $i(B) = 2|B| - 2$;
- (4) a 3-edge-cutset $\{x_i y_i : x_i \in A, y_i \in B, 1 \leq i \leq 3\}$, where $|\{x_i, y_i \mid 1 \leq i \leq 3\}| = 6$, for $A, B \subset V$, such that $A \cap B = \emptyset$ and $A \cup B = V$ such that $i(A) = 2|A| - 1$ and $i(B) = 2|B| - 2$; or
- (5) a 4-edge-cutset $\{x_i y_i : x_i \in A, y_i \in B, 1 \leq i \leq 4\}$ for $A, B \subset V$, such that $A \cap B = \emptyset$ and $A \cup B = V$ such that $i(A) = 2|A| - 2$ and $i(B) = 2|B| - 2$.

Lemma 4.1. *Let $G = (V, E)$ be a graph satisfying (1) in the list above. Let a_i, b_i, c_i, d_i , for $i = A, B$ be eight distinct vertices not in V . Then G is a (2,1)-circuit if and only if the graphs $G_i = G[i] \cup (K_5(a_i, b_i, c_i, d_i, x) - e)$, for $i = A, B$, are (2,1)-circuits. (See Figure 8.)*

Note that the choice of the missing edge in the $K_5 - e$ is arbitrary.

Proof. Suppose G_A and G_B are (2,1)-circuits. Assume for a contradiction that G is not. Then there is a proper subset $X \subset V$ with $i(X) = i(A \cap X) + i(B \cap X) \geq 2|X|$. However $i(A \cap X) \leq 2|A \cap X| - 1$, $i(B \cap X) \leq 2|B \cap X| - 1$ and $|A \cap X| + |B \cap X| \leq |X| + 1$ so

$$2|X| \geq 2(|A \cap X| + |B \cap X|) - 2 \geq i(A \cap X) + i(B \cap X) \geq 2|X|.$$

Hence, we have equality throughout and in particular $i(A \cap X) = 2|A \cap X| - 1$, $i(B \cap X) = 2|B \cap X| - 1$ and $(A \cap X) \cap (B \cap X) = \{x\}$. Now $i((A \cap X) \cup \{a_A, b_A, c_A, d_A\}) = 2|A \cap X| - 1 + 9 = 2(|A \cap X| + 4)$, so, as G_A is a (2,1)-circuit, we have that $(A \cap X) = A$. Similarly $(B \cap X) = B$. Hence $X = V$, a contradiction.

Conversely, suppose G is a (2,1)-circuit. Then for any $X \subset A$ we have $i(X) \leq 2|X| - 1$. Let $Y \subset \{a_i, b_i, c_i, d_i, x\}$, then $i(Y) \leq 2|Y| - 1$. As x is a cut-vertex, $i(X \cup Y) \leq 2|X \cup Y|$, since $|X \cup Y| \geq |X| + |Y| - 1$. Note equality holds if and only if $X = A$ and $Y = \{a_i, b_i, c_i, d_i, x\}$. \square



FIGURE 9. Illustration of the sum move for Case (2a).

Lemma 4.2. *Let $G = (V, E)$ be a graph satisfying (2) (a) in the list above. Let a_i, b_i, c_i, d_i , for $i = A, B$ be eight distinct vertices not in V . Then G is a $(2, 1)$ -circuit if and only if the graphs $G_i = G[i] \cup (K_4(a_i, b_i, c_i, d_i) \cup \{xa_i, yb_i, xc_i, yd_i\})$ for $i = A, B$, are $(2, 1)$ -circuits. (See Figure 9.)*

Note that the proof holds for other choices of the 4-edge-cuts in G_A and G_B , provided two edges are incident with x , two are incident with y , simplicity is maintained and we do not choose the edge xy .

Proof. Suppose G_A and G_B are $(2, 1)$ -circuits. Assume for a contradiction that G is not. Then there is a proper subset $X \subset V$, containing at least one of x or y , with $i(X) = i(A \cap X) + i(B \cap X) \geq 2|X|$. First suppose that $A \cap X$ contains both x and y . Then $i(A \cap X) \leq 2|A \cap X| - 2$ with equality if and only if $X \cap A = A$; similarly $i(B \cap X) \leq 2|B \cap X| - 2$; and $|A \cap X| + |B \cap X| \leq |X| + 2$ so

$$2|X| \geq 2(|A \cap X| + |B \cap X|) - 4 \geq i(A \cap X) + i(B \cap X) \geq 2|X|.$$

Hence, we have equality throughout and in particular $i(A \cap X) = 2|A \cap X| - 2$, $i(B \cap X) = 2|B \cap X| - 2$. As noted above, this implies that $(A \cap X) = A$ and $(B \cap X) = B$. Hence $X = V$, a contradiction.

Now suppose $A \cap X$ contains x but not y . Then $|A \cap X| + |B \cap X| \leq |X| + 1$. Thus

$$2|X| \geq 2(|A \cap X| + |B \cap X|) - 2 \geq i(A \cap X) + i(B \cap X) \geq 2|X|.$$

Hence, we have equality throughout and in particular $i(A \cap X) = 2|A \cap X| - 1$ and $i(B \cap X) = 2|B \cap X| - 1$. This is a contradiction as $(A \cap X) \cup (B \cap X)$ would induce a $(2, 1)$ -circuit not containing y .

Conversely, suppose G is a $(2, 1)$ -circuit. First note that for any subset $X \subset A$ containing x but not y and any subset $Y \subset \{a_A, b_A, c_A, d_A\}$ we have $i(X \cup Y) \leq 2|X \cup Y| - 1$. Now for any $X \subset A$ containing x and y we have $i(X) \leq 2|X| - 2$ with equality if and only if $X = A$. Let $Y \subset \{a_i, b_i, c_i, d_i\}$, then $i(Y) \leq 2|Y| - 2$ with equality if and only if $Y = \{a_i, b_i, c_i, d_i\}$. Thus $i(X \cup Y) \leq 2|X| - 2 + 2|Y| - 2 + 4 = 2|X \cup Y|$ where equality holds if and only if $X = A$ and $Y = \{a_i, b_i, c_i, d_i\}$. \square

Lemma 4.3. *Let G be a 4-regular $(2, 1)$ -circuit which contains a cut-pair $\{x, y\}$ such that one component of $G[V \setminus \{x, y\}]$ is isomorphic to K_4 on the vertices a, b, c, d ; and $\{xa, xb, yc, yd\} \in E$ where $z_1, z_2 \in V \setminus \{a, b, c, d, y\}$ are neighbours of x . Then x is admissible.*

Proof. The edges $z_1a, z_2b \notin G$; so performing an inverse X-replacement on x that introduces the edges z_1a and z_2b yields a connected 4-regular graph. Hence, by Lemma 2.1, the resulting graph is a $(2, 1)$ -circuit. \square

Lemma 4.4. *Let $G = (V, E)$ be a graph satisfying (2) (b) in the list above. Let a_B, b_B, c_B, d_B, e_B be five distinct vertices not in V . Then G is a $(2, 1)$ -circuit if and only if the graphs $G_A = G[A] \cup xy$ and $G_B = G[B] \cup (K_5(a_B, b_B, c_B, d_B, e_B) - f) \cup \{xa_B, xb_B, yc_B, yd_B\}$ are $(2, 1)$ -circuits. (See Figure 10.)*




FIGURE 10. Illustration of the sum move for Case (2b).




FIGURE 11. Illustration of the sum move for Case (3).

Note that the proof holds for other choices of the 4-edge-cuts in G_A and G_B , provided two edges are incident with x , two are incident with y , simplicity is maintained and we do not choose the edge xy . Moreover the choice of the missing edge in $K_5 - f$ is arbitrary.

Proof. Suppose G_A and G_B are (2,1)-circuits. Assume for a contradiction that G is not. Then there is a proper subset $X \subset V$, containing at least one of x or y , with $i(X) = i(A \cap X) + i(B \cap X) \geq 2|X|$. First suppose that $A \cap X$ contains both x and y . Then $i(A \cap X) \leq 2|A \cap X| - 1$ with equality if and only if $X \cap A = A$. Similarly $i(B \cap X) \leq 2|B \cap X| - 3$ and $|A \cap X| + |B \cap X| \leq |X| + 2$ so

$$2|X| \geq 2(|A \cap X| + |B \cap X|) - 4 \geq i(A \cap X) + i(B \cap X) \geq 2|X|.$$

Hence, we have equality throughout. Hence, $(A \cap X) = A$. Moreover, $i(B \cap X) = 2|B \cap X| - 3$. Now $i((B \cap X) \cup \{a_B, b_B, c_B, d_B, e_B\}) = 2|B \cap X| - 3 + 13 = 2(|B \cap X| + 5)$, so, as G_B is a (2,1)-circuit, we have that $(B \cap X) = B$. Hence $X = V$, a contradiction.

Now suppose $A \cap X$ contains x but not y . Then $|A \cap X| + |B \cap X| \leq |X| + 1$. Thus

$$2|X| \geq 2(|A \cap X| + |B \cap X|) - 2 \geq i(A \cap X) + i(B \cap X) \geq 2|X|.$$

Hence, we have equality throughout and in particular $i(A \cap X) = 2|A \cap X| - 1$ and $i(B \cap X) = 2|B \cap X| - 1$. This is a contradiction as $(A \cap X) \cup (B \cap X)$ would induce a (2,1)-circuit not containing y .

Conversely, suppose G is a (2,1)-circuit. Clearly G_A is a (2,1)-circuit. Note that for any subset $X \subset B$ containing x but not y we have $i(X) \leq 2|X| - 2$ and so for any subset $Y \subset \{a_B, b_B, c_B, d_B, e_B\}$ we have $i(X \cup Y) \leq 2|X \cup Y| - 1$.

Now for any $X \subset B$ containing x and y we have $i(X) \leq 2|X| - 3$ with equality if and only if $X = B$. Let $Y \subset \{a_B, b_B, c_B, d_B, e_B\}$, then $i(Y) \leq 2|Y| - 1$ with equality if and only if $Y = \{a_B, b_B, c_B, d_B, e_B\}$. Thus $i(X \cup Y) \leq 2|X| - 3 + 2|Y| - 1 + 4 = 2|X \cup Y|$ where equality holds if and only if $X = B$ and $Y = \{a_B, b_B, c_B, d_B, e_B\}$. \square

We emphasise that in the following lemma that the edge xy is included in the count for $i_G(A)$ and $i_G(B)$.

Lemma 4.5. *Let $G = (V, E)$ be a graph satisfying (3) in the list above. Let $a_A, b_A, c_A, d_A, a_B, b_B, c_B, d_B, e_B$ be nine distinct vertices not in V . Then G is a (2,1)-circuit if and only if the graphs $G_A = (G[A] - xy) \cup K_4(a_A, b_A, c_A, d_A) \cup \{xa_A, xb_A, yc_A, yd_A\}$ and $G_B = (G[B] - xy) \cup (K_5(a_B, b_B, c_B, d_B, e_B) - f) \cup \{xa_B, xb_B, yc_B, yd_B\}$ are (2,1)-circuits. (See Figure 11.)*




FIGURE 12. Illustration of the sum move for Case (4).

Note again that the proof holds for other choices of the 4-edge-cuts in G_A and G_B , provided two edges are incident with x , two are incident with y , simplicity is maintained and we do not choose the edge xy . Moreover the choice of the missing edge in $K_5 - f$ is arbitrary.

Proof. Suppose G_A and G_B are $(2, 1)$ -circuits. Assume, for a contradiction, that G is not. Then there is a proper subset $X \subset V$, containing at least one of x or y , with either:

- $i_G(X) = i_{G_A}(A \cap X) + i_{G_B}(B \cap X) - 1 \geq 2|X|$, if $x, y \in X$; or
- $i_G(X) = i_{G_A}(A \cap X) + i_{G_B}(B \cap X) \geq 2|X|$, if $x \in X$ and $y \notin X$.

First suppose that $x, y \in X$. Then $i_G(A \cap X) \leq 2|A \cap X| - 1$ with equality if and only if $X \cap A = A$. Similarly, $i_G(B \cap X) \leq 2|B \cap X| - 2$ and $|A \cap X| + |B \cap X| = |X| + 2$ so

$$2|X| = 2(|A \cap X| + |B \cap X|) - 4 \geq i_G(A \cap X) + i_G(B \cap X) - 1 \geq 2|X|.$$

Hence, we have equality throughout. So, $(A \cap X) = A$. Also $i_G(B \cap X) = 2|B \cap X| - 2$. Now $i_{G_B}((B \cap X) \cup \{a_B, b_B, c_B, d_B, e_B\}) = 2|B \cap X| - 3 + 13 = 2(|B \cap X| + 5)$, so, as G_B is a $(2, 1)$ -circuit, we have that $(B \cap X) = B$. Hence $X = V$, a contradiction.

Now, suppose that $x \in X$ but $y \notin X$. Then $|A \cap X| + |B \cap X| \leq |X| + 1$. Thus

$$2|X| \geq 2(|A \cap X| + |B \cap X|) - 2 \geq i_{G_A}(A \cap X) + i_{G_B}(B \cap X) \geq 2|X|.$$

Hence, we have equality throughout and in particular $i_G(B \cap X) = i_{G_B}(B \cap X) = 2|B \cap X| - 1$. Then $(B \cap X) \cup \{a_B, b_B, c_B, d_B, e_B\}$ would induce a $(2, 1)$ -circuit in G_B not containing y , a contradiction.

Conversely, suppose G is a $(2, 1)$ -circuit. For any subset $X \subset A$ containing x but not y and any subset $Y \subset \{a_A, b_A, c_A, d_A\}$ we have $i_{G_A}(X \cup Y) \leq 2|X \cup Y| - 1$. Now, for any $X \subset A$ containing x and y we have $i_G(X) \leq 2|X| - 1$ with equality if and only if $X = A$. Let $Y \subset \{a_A, b_A, c_A, d_A\}$, then $i_{G_A}(Y) \leq 2|Y| - 2$ with equality if and only if $Y = \{a_A, b_A, c_A, d_A\}$. Thus $i_{G_A}(X \cup Y) \leq 2|X| - 2 + 2|Y| - 2 + 4 = 2|X \cup Y|$ where equality holds if and only if $X = A$ and $Y = K_4(a_A, b_A, c_A, d_A)$.

For any subset $X \subset B$ containing x but not y we have that $i_G(X) \leq 2|X| - 2$ and for any subset $Y \subset \{a_B, b_B, c_B, d_B, e_B\}$ we have $i_{G_A}(X \cup Y) \leq 2|X \cup Y| - 1$, otherwise we contradict G being a $(2, 1)$ -circuit.

Now for any $X \subset B$ containing x and y we have $i_G(X) \leq 2|X| - 2$ with equality if and only if $X = B$. Let $Y \subset \{a_B, b_B, c_B, d_B, e_B\}$, then $i_{G_A}(Y) \leq 2|Y| - 1$ with equality if and only if $Y = \{a_B, b_B, c_B, d_B, e_B\}$. Thus $i_{G_A}(X \cup Y) \leq 2|X| - 2 - 1 + 2|Y| - 1 + 4 = 2|X \cup Y|$ where equality holds if and only if $X = B$ and $Y = \{a_B, b_B, c_B, d_B, e_B\}$. \square

Lemma 4.6. *Let $G = (V, E)$ be a graph satisfying (4) in the list above. Let $a_A, a_B, b_B, c_B, d_B, e_B$ be six distinct vertices not in V . Then G is a $(2, 1)$ -circuit if and only if the graphs $G_A = G[A] \cup a_A \cup \{x_1 a_A, x_2 a_A, x_3 a_A\}$ and $G_B = G[B] \cup K_5(a_B, b_B, c_B, d_B, e_B) - f \cup \{y_1 r_1, y_2 r_2, y_3 r_3\}$, for $r_i \in \{a_B, b_B, c_B, d_B, e_B\}$, are $(2, 1)$ -circuits. (See Figure 12.)*

Note that we may choose all the r_i 's to be equal giving a stronger final characterisation as then we only have one move for this case.




FIGURE 13. Illustration of a sum move for Case (5).

Proof. Suppose G_A and G_B are (2,1)-circuits. Let $X \subset A$ and $Y \subset B$. First suppose X contains x_1, x_2, x_3 and Y contains y_1, y_2, y_3 , then $i(Y) \leq 2|Y| - 2$. As $i(X) \leq 2|X| - 1$, $i(X \cup Y) \leq 2|X \cup Y| - 3 + 3 = 2|V|$, with equality if and only if $X \cup Y = V$. Next suppose X contains x_1 and x_2 but not x_3 and Y contains y_1 and y_2 but not y_3 . Then $i(X) \leq 2|X| - 1$ and, as $i_{G_B} = \{a_B, b_B, c_B, d_B, e_B\} = 9$, $i(Y) \leq 2|Y| - 2$. Thus $i(X \cup Y) \leq 2|X \cup Y| - 3 + 2$. Now suppose X contains x_1, x_2, x_3 and Y contains y_1 but not y_2, y_3 . Then $i(X) \leq 2|X| - 1$ and $i(Y) \leq 2|Y| - 1$. Thus $i(X \cup Y) \leq 2|X \cup Y| - 2 + 1$. The remaining cases can be proved similarly.

Conversely, suppose G is a (2,1)-circuit. Then for any $X \subset A$ we have $i(X) \leq 2|X| - 1$ and when $x_1, x_2, x_3 \in X$ we have equality if and only if $X = A$. It follows that G_A is a (2,1)-circuit.

Let $Y \subset \{a_B, b_B, c_B, d_B, e_B\}$, then $i(Y) \leq 2|Y| - 1$ with equality if and only if $Y = \{a_B, b_B, c_B, d_B, e_B\}$. Let $X \subset B$. If $|X \cap \{x_1, x_2, x_3\}| = 0$ or 1 , then $i(X) \leq 2|X| - 1$, thus $i(X \cup Y) \leq 2|X| - 1 + 2|Y| - 1 + 1 = 2|X \cup Y| - 1$. If $|X \cap \{x_1, x_2, x_3\}| = 2$, then $i(X) \leq 2|X| - 2$, otherwise $X \cup A$ would be a (2,1)-circuit that does not contain all of B , thus $i(X \cup Y) \leq 2|X| - 2 + 2|Y| - 1 + 2 = 2|X \cup Y| - 1$. If $|X \cap \{x_1, x_2, x_3\}| = 3$, then $i(X) \leq 2|X| - 2$, with equality if and only if $X = B$, thus $i(X \cup Y) \leq 2|X| - 2 + 2|Y| - 1 + 3 = 2|X \cup Y|$, with equality if and only if $X = B$ and $Y = \{a_B, b_B, c_B, d_B, e_B\}$. \square

Recall that, e.g. in Theorem 1.3, we only want to deal with non-trivial 4-edge cutsets when there are no proper critical sets. We utilise this fact in the next lemma.

Lemma 4.7. *Let $G = (V, E)$ be a graph satisfying (5) in the list above. Let $a_A, b_A, c_A, d_A, a_B, b_B, c_B, d_B$ be eight distinct vertices not in V and let \mathcal{E} denote a set of four edges of the form xy where $x \in A, y \in B$. For $I \in \{A, B\}$, if there are exactly:*

- 3 distinct x 's in I , then let $G_I = G[I] \cup K_4(a_I, b_I, c_I, d_I) \cup \mathcal{F}_1$, where \mathcal{F}_1 is a set of four edges such that each has exactly one end vertex in I , the degree of each vertex in $G_I[I]$ is equal to their degree in $G[I]$ and between them they have two, three or four distinct end vertices in $\{a_I, b_I, c_I, d_I\}$, call the graphs in these cases type-2, type-3 and type-4 respectively and denote this set of end vertices by F_I ;
- four distinct x 's in I , then let $G_I = G[I] \cup \{a_I\} \cup \{x_1 a_I, x_2 a_I, x_3 a_I, x_4 a_I\}$;

Then we have the following.

- (1) If G_I , for each $I \in \{A, B\}$, is either a type-3 or type-4 (2,1)-circuits with no proper critical sets or type-2 (2,1)-circuits with the unique proper critical set $I \cup F_I$, then G is a (2,1)-circuit.
- (2) Suppose G is a (2,1)-circuit with no proper critical sets. Then if G_I , for $I \in \{A, B\}$, is type-3 or type-4, then G_I is a (2,1)-circuit with no proper critical sets; and if G_I , for $I \in \{A, B\}$, is type-2, then it is a (2,1)-circuit with the proper critical set $I \cup F_I$.

(See Figure 13 for an example.)

Proof. (1) Let $X \subset A$ and $Y \subset B$. Then, as $F \cap I = \emptyset$, $i(X) \leq 2|X| - 2$ and $i(Y) \leq 2|Y| - 2$. Thus $i(X \cup Y) \leq 2|X \cup Y| - 4 + d(X, Y)$. We have that $i(X \cup Y) = 2|X \cup Y|$ if and only if $i(X) = 2|X| - 2$, $i(Y) = 2|Y| - 2$ and $d(X, Y) = 4$. This holds if and only if $X = A$ and $Y = B$ and hence $X \cup Y = V$. (Note that, if $d(X, Y) < 4$ then $i(X \cup Y) < 2|X \cup Y|$.)

(2) Suppose G is a $(2, 1)$ -circuit with no proper critical sets.

First we consider the case where there are exactly three distinct x 's in A . Then for any $X \subset A$ we have $i(X) \leq 2|X| - 2$. This implies that $i(X \cup \{a_A, b_A, c_A, d_A\}) \leq 2|X \cup \{a_A, b_A, c_A, d_A\}|$ with equality if and only if X contains all the x 's in \mathcal{E} and hence $X = A$. Hence, G_A is a $(2, 1)$ -circuit. Suppose that G_A contains a proper critical set. Then this critical set must either be of the form $X \cup \{a_A, b_A, c_A, d_A\}$ where $X \subsetneq A$ or of the form $X \cup F_A$ where $|F_A| = 2$ and $X \subseteq A$. If the critical set is of the form $X \cup \{a_A, b_A, c_A, d_A\}$ where $X \subsetneq A$, then $i(X \cup B) = 2|X \cup B| - 1$ and $|X \cup B| < |V| - 1$ (since $|X| < |A| - 1$), contradicting the fact that G does not contain any proper critical sets. If $|F_A| = 2$, then $A \cup F_A$ is a proper critical set. The case where there are exactly three distinct y 's in B is identical.

Next we consider the case where there are four distinct x 's in A . Then for any $X \subset A$ we have $i(X) \leq 2|X| - 2$. This implies that $i(X \cup \{a_A\}) \leq 2|X \cup \{a_A\}|$ with equality if and only if X contains all the x 's in \mathcal{E} and hence $X = A$. Suppose that G_A contains a proper critical set. Then this critical set must be of the form $X \cup \{a_A\}$ where $X \subsetneq A$ and either: X contains exactly three of the x 's, in which case $i(X) = 2|X| - 2$; or X contains all four of the x 's, in which case $i(X) = 2|X| - 3$. In either case $i(X \cup B) = 2|X \cup B| - 1$, contradicting the fact that G does not contain any proper critical sets. The case where there are four distinct y 's in B is identical. \square

5. A RECURSIVE CONSTRUCTION

We need one final elementary lemma.

Lemma 5.1. *Let G be a $(2, 1)$ -circuit and let G' be formed from G by a 1-extension or an X -replacement. Then G' is a $(2, 1)$ -circuit.*

Now we can prove our main result.

Theorem 1.1. *A simple graph G is a $(2, 1)$ -circuit if and only if it can be generated recursively from $H \in \mathcal{G}$ by applying 1-extensions and X -replacements sequentially within connected components and taking sums of connected components.*

Proof. (\Leftarrow) Suppose G is recursively generated by applying 1-extensions and X -replacements sequentially within connected components and taking sums of connected components. Then Lemmas 5.1, 4.1, 4.2, 4.4, 4.5, 4.6 and 4.7 together imply that G is a $(2, 1)$ -circuit.

(\Rightarrow) Suppose that $G \in M(2, 1)$.

- If G is essentially 5-edge-connected and $\delta(G) = 4$, then Theorem 1.2 implies that there exists an admissible vertex (an inverse X -replacement can be performed at this vertex).
- If G is 3-connected, essentially 5-edge-connected, contains no proper critical sets and $\delta(G) = 3$, then Theorem 1.3 (i) (restricted to $M(2, 1)$) implies that there exists an admissible node (a 1-reduction can be performed at this node).
- If G is 3-connected, essentially 4-edge-connected, contains a proper critical set and $\delta(G) = 3$, then Theorem 1.3 (ii) (restricted to $M(2, 1)$) implies that there exists an admissible node (a 1-reduction can be performed at this node).

The cases left to consider are:





FIGURE 14. The interaction between cases in the proof of Theorem 1.1 .

- (1) $\delta(G) = 4$ and G is not essentially 5-edge-connected; or
- (2) $\delta(G) = 3$ and either:
 - (a) G is 3-connected, contains proper critical sets and non-trivial 3-edge-cutsets;
 - (b) G is 3-connected, contains no proper critical sets and is not essentially 5-edge-connected.
 - (c) G is not 3-connected.

We illustrate how these cases interact during the following argument in Figure 14.

Case 1: $\delta(G) = 4$ and G is not essentially 5-edge-connected.

FIGURE 15. The graphs T_1 and T_2 .

First, suppose that G contains a cut-vertex or a cut-pair. Then G satisfies either (1), (2a), (2b) or (3) in the list at the start of Section 4. If G satisfies (1), then we can apply Lemma 4.1 and reduce to two new (2,1)-circuits both with minimum degree three. If G satisfies (2a), then we can apply Lemma 4.2 to reduce to two new (2,1)-circuits unless one (or both) of A and B are isomorphic to the graph T_1 (the cut pair in G correspond to the degree two vertices in T_1) shown in Figure 15. (If both of A and B are isomorphic to T_1 we can apply two inverse X-replacements to achieve the graph $S_1 \in \mathcal{G}$, see Figure 3.)

Hence, any potential counterexample arising from this case has no cut-vertices and any cut pairs, $\{x, y\}$ say, satisfy: xy is not an edge and at least one of the ‘sides’ of the cut-pair is isomorphic to T_1 . Then, by Lemma 4.3, we can perform an inverse X-replacement at x .

If G satisfies (2b), then we can apply Lemma 4.4 to reduce to two new (2,1)-circuits (one has strictly fewer vertices than G and the other has minimum degree 3). If G satisfies (3), then we can apply Lemma 4.5 to obtain two new (2,1)-circuits (these two circuits do not contain the edge xy , so we have moved to (2,1)-circuits satisfying (2a) or (2b)).

Next, suppose that G contains a non-trivial 3-edge-cutset but no cut-vertices or cut-pairs. Then we can apply Lemma 4.6 to achieve two new (2,1)-circuits both with minimum degree three.

Finally, suppose that G is 3-connected and contains a non-trivial 4-edge-cutset, but no non-trivial 3-edge-cutset.

Claim 5.2. G contains no proper critical sets.

Proof of Claim. As $\delta(G) = 4$, we have that G is 4-regular. Suppose that G did contain a proper critical set S , then

$$\sum_{v \in S} d_{G[S]}(v) = 4|S| - 2.$$

However, as G is 4-regular this means there must be either a cut-vertex or a cut-pair in G , contradicting the fact that G is 3-connected. \square

Thus we can apply Lemma 4.7 to achieve two new (2,1)-circuits one with strictly fewer vertices than G and we can apply the lemma in such a way that the other has minimum degree three.

Case 2a: $\delta(G) = 3$, G is 3-connected, contains proper critical sets and non-trivial 3-edge-cutsets.

Suppose that G contains a non-trivial 3-edge-cutset, then we can apply Lemma 4.6 to form two new (2,1)-circuits. When we apply Lemma 4.6 we choose to do so such that the resulting circuits are strictly smaller and (by setting $r_1 = r_2$) not 3-connected respectively.

Case 2b: $\delta(G) = 3$, G is 3-connected, contains no proper critical sets and is not essentially 5-edge-connected.

As G does not contain a proper critical set it does not contain a non-trivial 3-edge cutset. Hence G contains a non-trivial 4-edge cutset and we can apply Lemma 4.7 to form two new (2,1)-circuits. We can apply Lemma 4.7 so that either the resulting circuits have strictly fewer vertices than G or, by the freedom in the choice of edges when applying Lemma 4.7, we may assume that they are not 3-connected.

Case 2c: $\delta(G) = 3$ and G is not 3-connected.

Suppose that there exists a cut-vertex in G , then we can apply Lemma 4.1 and reduce to two smaller (2,1)-circuits unless one (or both) of the sides of the cut are isomorphic to $K_5 - e$. If both sides are isomorphic to $K_5 - e$, we can, if necessary, reapply Lemma 4.1 until we obtain one graph, $S_5 \in \mathcal{G}$ (see Figure 4).

Now, suppose that there exists a cut-pair in G . Then G satisfies (2a), (2b) or (3). If G contains a cut-pair that satisfies:

- (2a), then we can apply Lemma 4.2 to obtain two new smaller (2,1)-circuits, unless one side of the cut-pair is isomorphic to T_1 (if both sides were isomorphic to T_1 , we would have $\delta(G) = 4$);
- (2b), then we can apply Lemma 4.4 to obtain two new smaller (2,1)-circuits unless one of the sides of the cut-pair is isomorphic to T_2 ; or
- (3), then we can apply Lemma 4.5 to obtain two new (2,1)-circuits (the edge xy occurs in G but in neither of the resulting circuits). The new (2,1)-circuits are either smaller or we have moved to case (2a) or (2b).

Assume G is a counter-example to the theorem (we will show that no such G exists), then G has:

- (I) every cut-vertex has one side isomorphic to $K_5 - e$; and
- (II) every cut-pair $\{x, y\}$ satisfies $xy \notin E$ and has one side isomorphic to T_1 or T_2 .

Associate the following multigraph G^* with G : if (I) occurs in G , replace the $K_5 - e$ with a loop at the cut-vertex; if (II) occurs in G , replace any occurrences of T_1 with a double edge between the vertices of the cut-pair, and any occurrences of T_2 with a triple edge between the vertices of the cut-pair. Note that:

- all the vertices in G^* incident with a multiple edge have degree greater than three and that G^* is 3-connected (otherwise we could have reduced G using one of Lemmas 4.1, 4.2, 4.4 or 4.5);
- if G^* contains a loop, then the vertex incident with the loop has degree greater than three and it is not incident with any multiple edges (else both sides of the 2-cut are not isomorphic to T_1 or T_2);
- if $G^* = (V, E)$ contains a vertex, v say, incident to two loops then $V = \{v\}$.
- if there are any triple edges in G^* , then they are incident to vertices of degree greater than or equal to five, and hence, G^* must contain vertices of degree three.

Hence $G^* \in \mathcal{M}$. See Figure 16 for an illustration of the construction of G^* .

We will consider two cases, when $\delta(G^*) = 4$ (and hence G^* is 4-regular) and when $\delta(G^*) = 3$.

First, suppose that $\delta(G^*) = 4$ and $G^* \neq R_1$, hence G^* is 4-regular and, as it is 3-connected, contains no proper critical sets (see Claim 5.2). As G^* is 4-regular and 3-connected it cannot contain a loop or a triple edge. If G^* contains a double edge, then it results from a copy of T_1 in G ; so, applying the same argument as in the proof of Lemma 4.3, we can perform an inverse X-replacement in G , contradicting G being a counterexample.




FIGURE 16. Illustration of the construction of the multigraph G^* .

So, suppose that $\delta(G^*) = 3$. We will show that there is an admissible node v in G^* . Recall that G^* is 3-connected and that any vertex of degree three is not incident with a loop or a multiple edge.

Claim 5.3. *We may assume that the multigraph $G^* \in \mathcal{M} \setminus M(2,1)$ constructed from a counterexample G does not contain any non-trivial 3-edge-cutsets.*

Proof of Claim. If G^* contains a non-trivial 3-edge-cutset, then, as G^* is 3-connected, G also contains a non-trivial 3-edge-cutset. In which case we can apply Lemma 4.6 to form two (2,1)-circuits. These circuits are both nonisomorphic to G (contradicting our assumption that G is a counterexample) unless one side of the edge-cutset is isomorphic to $K_5 - e$. Moreover, we can choose the cut-edges so that they are all incident with the same vertex in the copy of $K_5 - e$. At this stage we have a cut-vertex v with one side isomorphic to $K_5 - e$ and in our construction of G^* this is replaced by a loop on v . \square

Suppose that G^* contains a proper critical set. As G^* is essentially 4-edge-connected, as long as $G^* \notin \mathcal{G}^*$, by Theorem 1.3 (ii), G^* has an admissible node, and, by construction, this node is also admissible in G .

Finally, suppose that G^* does not contain a proper critical set.

Claim 5.4. *If the (2,1)-multi-circuit G^* (containing no proper critical sets) constructed from the counter-example G contains a non-trivial 4-edge-cutset, then either one of the sides of the cut is a double edge or, from G , we can construct a new counter-example which contains a proper critical set.*

Proof of Claim. We know that G^* is essentially 4-edge-connected. If G^* contains a non-trivial 4-edge-cutset, then:

- if a non-trivial 4-edge-cutset is formed from four single edges, then these edges form a non-trivial 4-edge-cutset in G ;
- if a non-trivial 4-edge-cutset is formed from two double edges, let the endvertices of one double edge be x_A and y_A and the other be x_B and y_B , these double edges resulted from two copies of T_1 , T_A and T_B say, in G where the degree two vertices of T_A (respectively T_B) are x_A and y_A (respectively x_B and y_B). Therefore there is a non-trivial 4-edge-cutset in G consisting of the edges incident with x_A and x_B that are contained in T_A and T_B .

- if a non-trivial 4-edge-cutset is formed from two single edges and a double edge between vertices x and y say, then the two single edges combined with the two edges incident with x in the copy of T_1 that results in the double edge form a non-trivial 4-edge-cutset.

(As G^* is 3-connected it cannot contain a 4-edge-cutset comprised of a triple edge and a single edge.) So, suppose that G^* contains a non-trivial 4-edge-cutset. As G^* contains no proper critical sets, G contains no proper critical sets. Hence we could have applied Lemma 4.7 to G and reduced to two (2,1)-circuits. These circuits are non-isomorphic to G unless one or both sides of the cutset is isomorphic to K_4 . If both sides are isomorphic to K_4 we either have $G = S_1 \in \mathcal{G}$ or a graph that we can construct by summing S_1 with itself using Lemma 4.7.

If only one side is isomorphic to K_4 a further application of Lemma 4.7 guarantees that the K_4 contains a proper two vertex cut, $\{x, y\}$ say, this yields a proper critical set. \square

We may assume that, in G^* , if it contains a non-trivial 4-edge cutset then one of the sides of the cut is a double edge.

For any such 4-edge-cut in G^* replace the two vertices forming one component with a single vertex z , call the resulting graph G^{**} . This either results in two vertices with four edges between them (which is in \mathcal{G}^*) or z is a degree four vertex with either three or four neighbours (as G^{**} is 3-connected). If z has four neighbours, then we have reduced to a case with one less non-trivial 4-edge-cutset.

So assume that z has three neighbours, say u is the neighbour with the double edge, then the degree of u is greater than three (otherwise we contradict the fact G^* was essentially 4-edge connected). If u has degree four then repeat the above over the double edge between u and z , which is one side of a non-trivial 4-edge cutset. Hence we may assume that the degree of u is at least 5. In this manner we can eliminate all non-trivial 4-edge cutsets, thus we may assume that G^{**} is essentially 5-edge-connected. If $G^{**} \notin \mathcal{G}^*$, then Theorem 1.3 (i) implies that G^{**} contains an admissible node, and, by construction, this node is also admissible in G .

All that remains is to consider the (2,1)-circuits R_i , where $0 \leq i \leq 12$. First note that, as they would reduce to R_0 , neither R_9 or R_{10} will arise as the multigraph generated from a minimal counter example. The graphs that would yield R_2 , R_8 and R_{11} all contain admissible nodes. The graphs that would yield R_1 , R_6 , R_7 and R_{12} all contain admissible vertices (at which an X -replacement can be performed). The graphs that yield R_0 are either isomorphic to S_5 or can be reduced to copies of S_5 through an application of Lemma 4.1. Any graph that generates R_3 contains a cut pair to which Lemma 4.5 can be applied to yield two (2,1)-circuits, each on fewer vertices. Any graph that generates R_4 contains a cut pair to which Lemma 4.5 can be applied to yield two new circuits, G_1 and G_2 say, where, without loss of generality, G_1 has an admissible vertex and Lemma 4.5 can be applied to G_2 to yield two (2,1)-circuits that both have admissible nodes. Finally, any graph that generates R_5 contains an admissible vertex. \square

6. CONCLUDING REMARKS

1. We expect that our characterisation will be useful for the problem of characterising the global rigidity of realisations of graphs¹ on surfaces of revolution (such as the cone). In

¹A graph is realised using a map $p : V \rightarrow \mathbb{R}^3$, such a realisation is globally rigid if any other map q inducing the same edge lengths is congruent to p . A realisation on a surface is a realisation in \mathbb{R}^3 in which each vertex is restricted to move only on the surface it lies on.




FIGURE 17. A 3-connected $(2, 1)$ -circuit where no degree 4 is admissible, and every admissible node results in a $(2, 1)$ -circuit which is not 3-connected.

[14], rigidity of such frameworks was, generically, shown to be equivalent to the graph being $(2, 1)$ -tight. In [7, Conjecture 1] it was conjectured that the graphs which are generically globally rigid on the cone are those which are 2-connected and contain a spanning subgraph which is $(2, 1)$ -tight. It is not hard to prove that the graphs with $|E| = 2|V|$ that are 2-connected with a spanning $(2, 1)$ -tight subgraph are exactly the $(2, 1)$ -circuits. Thus the constructive characterisation of $(2, 1)$ -circuits in this paper is likely to be useful in developing a combinatorial characterisation of global rigidity. See also [8, 11] for analogous results for the cylinder, additional results in this direction and warnings of the additional complication for the cone.

2. In [1] it was proved that all 3-connected $(2, 3)$ -circuits can be generated from K_4 by 1-extensions. (This is non-trivial since admissible nodes need not result in smaller circuits which are 3-connected even when the original circuit is 3-connected.) For $(2, 1)$ -circuits we now briefly comment on such extensions. It is easy to see that if G is 2-connected and v is an admissible node in G then the resulting $(2, 1)$ -circuit G' is 2-connected. Moreover it is easy to check that the same holds for inverse X -replacement and for our sum moves (clearly we do not consider the sum move for a graph with a cut-vertex). Thus we instantly have the following result. Let \mathcal{G}^2 denote all the graphs in \mathcal{G} which are 2-connected.

Corollary 6.1. *Let $G = (V, E)$ be a simple graph. Then G is a 2-connected $(2, 1)$ -circuit if and only if G can be generated from some graphs in \mathcal{G}^2 by 1-extensions, X -replacements and sum moves.*

However we do not know a corresponding statement for 3-connected $(2, 1)$ -circuits. Indeed Figure 17 illustrates a $(2, 1)$ -circuit which is 3-connected and essentially 4-edge-connected for which each admissible 1-reduction results in a $(2, 1)$ -circuit which is not 3-connected.

REFERENCES

- [1] A. Berg and T. Jordan, A Proof of Connelly's Conjecture on 3-connected Circuits of the Rigidity Matroid, *Journal of Combinatorial Theory, Series B*, 88, (2003), 77–97.
- [2] F. Bories, J-L Jolivet and J-L Fouquet, Construction of 4-regular graphs, *Annals of discrete mathematics*, 17 (1983) 99–118
- [3] Z. Fekete and L. Szego, A note on $[k, l]$ -sparse graphs, in: A. Bondy, et al. (Eds.), *Graph Theory in Paris, 2007*, 169–177.
- [4] A. Frank, *Connections in Combinatorial Optimization*, Oxford University Press 2011.
- [5] A. Frank and L. Szego, Constructive characterisations for packing and covering with trees, *Discrete Appl. Math.* 131 (2003) 347–371.
- [6] B. Jackson and T. Jordan, Connected Rigidity Matroids and Unique Realisations of Graphs, *Journal of Combinatorial Theory Series B*, vol. 94, (2005), 1–29.

- [7] B. Jackson, T. McCourt and A. Nixon, Necessary Conditions for the Generic Global Rigidity of Frameworks on Surfaces, *Discrete and Comput. Geom.* 52:2, (2014), 344–360.
- [8] B. Jackson and A. Nixon, Stress matrices and global rigidity of frameworks on surfaces, *Discrete and Comput. Geom.* 54:3, (2015) 586–609.
- [9] R. Kovacs and L. Vegh, Constructive characterization theorems in combinatorial optimization, in *Combinatorial Optimization and Discrete Algorithms* (ed. S. Iwata), RIMS Kokyuroku Bessatsu B23, (2010), 147–169.
- [10] A. Lee and I. Streinu, Pebble Game Algorithms and Sparse Graphs, *Discrete Mathematics*, 308, 8, (2008), 1425–1437.
- [11] A. Nixon, A Constructive Characterisation of Circuits in the Simple $(2, 2)$ -sparsity Matroid, *European J. Comb.*, 42, (2014), 92–106.
- [12] A. Nixon and J. Owen, An inductive construction of $(2, 1)$ -tight graphs, *Contrib. Discrete Math.*, 9:2, (2014), 1–16.
- [13] A. Nixon, J. Owen and S. Power, Rigidity of frameworks supported on surfaces, *SIAM J. Discrete Math.*, 26:4, (2012), 1733–1757.
- [14] A. Nixon, J. Owen and S. Power, A characterization of generically rigid frameworks on surfaces of revolution, *SIAM J. Discrete Math.*, 28:4, (2014), 2008–2028.
- [15] A. Nixon and E. Ross, One brick at a time: a survey of inductive constructions in rigidity theory, in *Rigidity and Symmetry*, New York: Springer, (2014) 303–324.
- [16] R. Read and R. Wilson, *An Atlas of Graphs*, Oxford Science Publications, The Clarendon Press Oxford University Press, New York, (1998).
- [17] T.-S. Tay, Henneberg’s method for bar and body frameworks, *Struct. Topol.* 17 (1991) 53–58.
- [18] W. Whiteley, Matroids for Discrete Applied Geometry, in *Matroid Theory*, J. Bonin, J. Oxley and B. Servatius (eds.), *Contemporary Mathematics* 197, AMS, (1996), 171–311.

SCHOOL OF COMPUTING, ELECTRONICS AND MATHEMATICS, PLYMOUTH UNIVERSITY, PLYMOUTH, PL4 8AA, U.K.

E-mail address: `thomas.mccourt@plymouth.ac.uk`

DEPARTMENT OF MATHEMATICS AND STATISTICS, LANCASTER UNIVERSITY, LANCASTER, LA1 4YF, U.K.

E-mail address: `a.nixon@lancaster.ac.uk`